

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/38825> holds various files of this Leiden University dissertation

Author: Pulyakhina, Irina

Title: A telescope for the RNA universe : novel bioinformatic approaches to analyze RNA sequencing data

Issue Date: 2016-04-21

Bibliography

- [1] J. D. Watson et al. Molecular structure of nucleic acids; a structure for deoxyribose nucleic acid. *Nature*, 171(4356):737–738, Apr 1953.
- [2] A. J. Shatkin. Capping of eucaryotic mRNAs. *Cell*, 9(4 PT 2):645–653, Dec 1976.
- [3] P. Danecek et al. High levels of RNA-editing site conservation amongst 15 laboratory mouse strains. *Genome Biol.*, 13(4):26, 2012.
- [4] J. D. Alfonzo et al. The mechanism of U insertion/deletion RNA editing in kinetoplastid mitochondria. *Nucleic Acids Res.*, 25(19):3751–3759, Oct 1997.
- [5] G. J. Arts et al. Mechanism and evolution of RNA editing in kinetoplastida. *Biochim. Biophys. Acta*, 1307(1):39–54, Jun 1996.
- [6] H. Tilgner et al. Deep sequencing of subcellular RNA fractions shows splicing to be predominantly co-transcriptional in the human genome but inefficient for lncRNAs. *Genome Res.*, 22(9):1616–1625, Sep 2012.
- [7] S. W. Roy et al. The evolution of spliceosomal introns: patterns, puzzles and progress. *Nat. Rev. Genet.*, 7(3):211–221, Mar 2006.
- [8] A. J. Taggart et al. Large-scale mapping of branch-points in human pre-mRNA transcripts in vivo. *Nat. Struct. Mol. Biol.*, 19(7):719–721, Jul 2012.
- [9] A. Corvelo et al. Genome-wide association between branch point properties and alternative splicing. *PLoS Comput. Biol.*, 6(11):e1001016, 2010.
- [10] A. A. Patel et al. Splicing double: insights from the second spliceosome. *Nat. Rev. Mol. Cell Biol.*, 4(12):960–970, Dec 2003.
- [11] A. J. Matlin et al. Understanding alternative splicing: towards a cellular code. *Nat. Rev. Mol. Cell Biol.*, 6(5):386–398, May 2005.
- [12] Q. Pan et al. Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nat. Genet.*, 40(12):1413–1415, Dec 2008.
- [13] G. Di Segni et al. Cis- and trans-splicing of mRNAs mediated by tRNA sequences in eukaryotic cells. *Proc. Natl. Acad. Sci. U.S.A.*, 105(19):6864–6869, May 2008.
- [14] D. L. Black. Mechanisms of alternative pre-messenger RNA splicing. *Annu. Rev. Biochem.*, 72:291–336, 2003.
- [15] M. Sammeth et al. A general definition and nomenclature for alternative splicing events. *PLoS Comput. Biol.*, 4(8):e1000147, 2008.
- [16] S. Djebali et al. Landscape of transcription in human cells. *Nature*, 489(7414):101–108, Sep 2012.
- [17] B. Ewing et al. Analysis of expressed sequence tags indicates 35,000 human genes. *Nat. Genet.*, 25(2):232–234, Jun 2000.
- [18] O. Olsvik et al. Use of automated sequencing of polymerase chain reaction-generated amplicons to identify three types of cholera toxin subunit B in *Vibrio cholerae* O1 strains. *J. Clin. Microbiol.*, 31(1):22–25, Jan 1993.
- [19] J. C. Venter et al. The sequence of the human genome. *Science*, 291(5507):1304–1351, Feb 2001.
- [20] C. A. Maher et al. Transcriptome sequencing to detect gene fusions in cancer. *Nature*, 458(7234):97–101, Mar 2009.
- [21] F. Sanger et al. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. U.S.A.*, 74(12):5463–5467, Dec 1977.
- [22] F. Sanger et al. A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase. *J. Mol. Biol.*, 94(3):441–448, May 1975.
- [23] D. R. Bentley et al. Accurate whole human genome sequencing using reversible terminator chemistry. *Nature*, 456(7218):53–59, Nov 2008.
- [24] B. Ewing et al. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.*, 8(3):186–194, Mar 1998.
- [25] A. Mortazavi et al. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods*, 5(7):621–628, Jul 2008.
- [26] K. D. Hansen et al. Biases in Illumina transcriptome sequencing caused by random hexamer priming. *Nucleic Acids Res.*, 38(12):e131, Jul 2010.
- [27] Y. Chu et al. RNA sequencing: platform selection, experimental design, and data interpretation. *Nucleic Acid Ther.*, 22(4):271–274, Aug 2012.

- [28] J. R. ten Bosch et al. Keeping up with the next generation: massively parallel sequencing in clinical diagnostics. *J Mol Diagn*, 10(6):484–492, Nov 2008.
- [29] T. Tucker et al. Massively parallel sequencing: the next big thing in genetic medicine. *Am. J. Hum. Genet.*, 85(2):142–154, Aug 2009.
- [30] M. J. Fullwood et al. Next-generation DNA sequencing of paired-end tags (PET) for transcriptome and genome analyses. *Genome Res.*, 19(4):521–532, Apr 2009.
- [31] R. Morin et al. Profiling the HeLa S3 transcriptome using randomly primed cDNA and massively parallel short-read sequencing. *BioTechniques*, 45(1):81–94, Jul 2008.
- [32] S. W. Roy et al. When good transcripts go bad: artifactual RT-PCR ‘splicing’ and genome analysis. *Bioessays*, 30(6):601–605, Jun 2008.
- [33] J. Cocquet et al. Reverse transcriptase template switching and false alternative transcripts. *Genomics*, 88(1):127–131, Jul 2006.
- [34] R. M. Mader et al. Reverse transcriptase template switching during reverse transcriptase-polymerase chain reaction: artificial generation of deletions in ribonucleotide reductase mRNA. *J. Lab. Clin. Med.*, 137(6):422–428, Jun 2001.
- [35] C. W. Fuller et al. The challenges of sequencing by synthesis. *Nat. Biotechnol.*, 27(11):1013–1023, Nov 2009.
- [36] Y. Zhang et al. PASSion: a pattern growth algorithm-based pipeline for splice junction detection in paired-end RNA-Seq data. *Bioinformatics*, 28(4):479–486, Feb 2012.
- [37] M. Hamada et al. Probabilistic alignments with quality scores: an application to short-read mapping toward accurate SNP/indel detection. *Bioinformatics*, 27(22):3085–3092, Nov 2011.
- [38] G. A. Heap et al. Genome-wide analysis of allelic expression imbalance in human primary cells by high-throughput transcriptome resequencing. *Hum. Mol. Genet.*, 19(1):122–134, Jan 2010.
- [39] H. Li et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16):2078–2079, Aug 2009.
- [40] W. J. Kent et al. The human genome browser at UCSC. *Genome Res.*, 12(6):996–1006, Jun 2002.
- [41] H. Thorvaldsdottir et al. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief. Bioinformatics*, 14(2):178–192, Mar 2013.
- [42] J. T. Robinson et al. Integrative genomics viewer. *Nat. Biotechnol.*, 29(1):24–26, Jan 2011.
- [43] F. Finotello et al. Reducing bias in RNA sequencing data: a novel approach to compute counts. *BMC Bioinformatics*, 15 Suppl 1:S7, 2014.
- [44] W. Zheng et al. Bias detection and correction in RNA-Sequencing data. *BMC Bioinformatics*, 12:290, 2011.
- [45] E. E. Khrameeva et al. Biases in read coverage demonstrated by interlaboratory and interplatform comparison of 117 mRNA and genome sequencing experiments. *BMC Bioinformatics*, 13 Suppl 6:S4, 2012.
- [46] S. Schwartz et al. Detection and removal of biases in the analysis of next-generation sequencing reads. *PLoS ONE*, 6(1):e16685, 2011.
- [47] A. Roberts et al. Improving RNA-Seq expression estimates by correcting for fragment bias. *Genome Biol.*, 12(3):R22, 2011.
- [48] J. F. Degner et al. Effect of read-mapping biases on detecting allele-specific expression from RNA-sequencing data. *Bioinformatics*, 25(24):3207–3212, Dec 2009.
- [49] V. Sousa et al. Understanding the origin of species with genome-scale data: modelling gene flow. *Nat. Rev. Genet.*, 14(6):404–414, Jun 2013.
- [50] C. A. Meyer et al. Identifying and mitigating bias in next-generation sequencing methods for chromatin biology. *Nat. Rev. Genet.*, 15(11):709–721, Nov 2014.
- [51] D. Sims et al. Sequencing depth and coverage: key considerations in genomic analyses. *Nat. Rev. Genet.*, 15(2):121–132, Feb 2014.
- [52] C. Kulahoglu et al. Quantitative transcriptome analysis using RNA-seq. *Methods Mol. Biol.*, 1158:71–91, 2014.
- [53] P. Jain et al. Augmenting transcriptome assembly by combining de novo and genome-guided tools. *PeerJ*, 1:e133, 2013.
- [54] R. Liu et al. Comparisons of computational methods for differential alternative splicing detection using RNA-seq in plant systems. *BMC Bioinformatics*, 15(1):364, Dec 2014.
- [55] A. Roberts et al. Identification of novel transcripts in annotated genomes using RNA-Seq. *Bioinformatics*, 27(17):2325–2329, Sep 2011.
- [56] D. Hiller et al. Simultaneous isoform discovery and quantification from RNA-seq. *Stat Biosci*, 5(1):100–118, May 2013.
- [57] R. K. Varshney et al. Next-generation sequencing technologies and their implications for crop genetics and breeding. *Trends Biotechnol.*, 27(9):522–530, Sep 2009.
- [58] Y. Li et al. RNA-Seq Based De Novo Transcriptome Assembly and Gene Discovery of *Cistanche deserticola* Fleshy Stem. *PLoS ONE*, 10(5):e0125722, 2015.

- [59] D. Powell et al. De novo transcriptome analysis of the banana shrimp (*Fenneropenaeus merguensis*) and identification of genes associated with reproduction and development. *Mar Genomics*, Apr 2015.
- [60] K. S. Arun-Chinnappa et al. De novo assembly of a genome-wide transcriptome map of *Vicia faba* (L.) for transfer cell research. *Front Plant Sci*, 6:217, 2015.
- [61] S. Anders et al. Detecting differential usage of exons from RNA-seq data. *Genome Res.*, 22(10):2008–2017, Oct 2012.
- [62] C. Trapnell et al. Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat. Biotechnol.*, 31(1):46–53, Jan 2013.
- [63] S. Danckwardt et al. 3' end mRNA processing: molecular mechanisms and implications for health and disease. *EMBO J.*, 27(3):482–498, Feb 2008.
- [64] B. Tian et al. A large-scale analysis of mRNA polyadenylation of human and mouse genes. *Nucleic Acids Res.*, 33(1):201–212, 2005.
- [65] E. de Klerk et al. RNA sequencing: from tag-based profiling to resolving complete transcript structure. *Cell. Mol. Life Sci.*, 71(18):3537–3551, Sep 2014.
- [66] V. Le Texier et al. AltTrans: transcript pattern variants annotated for both alternative splicing and alternative polyadenylation. *BMC Bioinformatics*, 7:169, 2006.
- [67] V. D'mello et al. Alternative mRNA polyadenylation can potentially affect detection of gene expression by affymetrix genechip arrays. *Appl. Bioinformatics*, 5(4):249–253, 2006.
- [68] G. Ji et al. A classification-based prediction model of messenger RNA polyadenylation sites. *J. Theor. Biol.*, 265(3):287–296, Aug 2010.
- [69] Z. Xia et al. Dynamic analyses of alternative polyadenylation from RNA-seq reveal a 3'-UTR landscape across seven tumour types. *Nat Commun*, 5:5274, 2014.
- [70] H. Zhang et al. PolyA-DB: a database for mammalian mRNA polyadenylation. *Nucleic Acids Res.*, 33(Database issue):D116–120, Jan 2005.
- [71] J. Y. Lee et al. PolyA-DB 2: mRNA polyadenylation sites in vertebrate genes. *Nucleic Acids Res.*, 35(Database issue):D165–168, Jan 2007.
- [72] G. Ji et al. Predictive modeling of plant messenger RNA polyadenylation sites. *BMC Bioinformatics*, 8:43, 2007.
- [73] P. de la Grange et al. A new advance in alternative splicing databases: from catalogue to detailed analysis of regulation of expression and function of human alternative splicing variants. *BMC Bioinformatics*, 8:180, 2007.
- [74] Z. H. Zhang et al. A comparative study of techniques for differential expression analysis on RNA-Seq data. *PLoS ONE*, 9(8):e103207, 2014.
- [75] R. Chandramohan et al. Benchmarking RNA-Seq quantification tools. *Conf Proc IEEE Eng Med Biol Soc*, 2013:647–650, 2013.
- [76] S. Wesolowski et al. A Comparison of Methods for RNA-Seq Differential Expression Analysis and a New Empirical Bayes Approach. *Biosensors (Basel)*, 3(3):238–258, Sep 2013.
- [77] M. D. Robinson et al. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1):139–140, Jan 2010.
- [78] S. Anders et al. Differential expression analysis for sequence count data. *Genome Biol.*, 11(10):R106, 2010.
- [79] G. K. Smyth. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol*, 3:Article3, 2004.
- [80] F. Rapaport et al. Comprehensive evaluation of differential gene expression analysis methods for RNA-seq data. *Genome Biol.*, 14(9):R95, 2013.
- [81] A. Brennicke et al. RNA editing. *FEMS Microbiol. Rev.*, 23(3):297–316, Jun 1999.
- [82] A. A. Su et al. A-to-I and C-to-U editing within transfer RNAs. *Biochemistry Mosc.*, 76(8):932–937, Aug 2011.
- [83] A. M. Kiran et al. Darned in 2013: inclusion of model organisms and linking with Wikipedia. *Nucleic Acids Res.*, 41(Database issue):D258–261, Jan 2013.
- [84] E. T. Wang et al. Alternative isoform regulation in human tissue transcriptomes. *Nature*, 456(7221):470–476, Nov 2008.
- [85] H. J. Mardon et al. A role for exon sequences in alternative splicing of the human fibronectin gene. *Nucleic Acids Res.*, 15(19):7725–7733, Oct 1987.
- [86] S. A. Filichkin et al. Genome-wide mapping of alternative splicing in *Arabidopsis thaliana*. *Genome Res.*, 20(1):45–58, Jan 2010.
- [87] A. Pandya-Jones et al. Co-transcriptional splicing of constitutive and alternative exons. *RNA*, 15(10):1896–1908, Oct 2009.
- [88] D. L. Bentley. Coupling mRNA processing with transcription in time and space. *Nat. Rev. Genet.*, 15(3):163–175, Mar 2014.
- [89] Y. L. Khodor et al. Nascent-seq indicates widespread cotranscriptional pre-mRNA splicing in *Drosophila*. *Genes Dev.*, 25(23):2502–2512, Dec 2011.
- [90] C. Svensson et al. Splicing of adenovirus 2 early region 1A mRNAs is non-sequential. *J. Mol. Biol.*, 165(3):475–495, Apr 1983.

- [91] K. M. Lang et al. In vitro splicing pathways of pre-mRNAs containing multiple intervening sequences? *Mol. Cell. Biol.*, 7(10):3428–3437, Oct 1987.
- [92] D. Baralle et al. Splicing in action: assessing disease causing sequence changes. *J. Med. Genet.*, 42(10):737–748, Oct 2005.
- [93] Y. W. Huang et al. Identification of a porcine DC-SIGN-related C-type lectin, porcine CLEC4G (LSECtin), and its order of intron removal during splicing: comparative genomic analyses of the cluster of genes CD23/CLEC4G/DC-SIGN among mammalian species. *Dev. Comp. Immunol.*, 33(6):747–760, Jun 2009.
- [94] U. Schwarze et al. Redefinition of exon 7 in the COL1A1 gene of type I collagen by an intron 8 splice-donor-site mutation in a form of osteogenesis imperfecta: influence of intron splice order on outcome of splice-site mutation. *Am. J. Hum. Genet.*, 65(2):336–344, Aug 1999.
- [95] U. Schwarze et al. Splicing defects in the COL3A1 gene: marked preference for 5' (donor) splice-site mutations in patients with exon-skipping mutations and Ehlers-Danlos syndrome type IV. *Am. J. Hum. Genet.*, 61(6):1276–1286, Dec 1997.
- [96] K. Wang et al. MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic Acids Res.*, 38(18):e178, Oct 2010.
- [97] E. Ullu et al. Temporal order of RNA-processing reactions in trypanosomes: rapid trans splicing precedes polyadenylation of newly synthesized tubulin transcripts. *Mol. Cell. Biol.*, 13(1):720–725, Jan 1993.
- [98] H. Suzuki et al. Nested introns in an intron: evidence of multi-step splicing in a large intron of the human dystrophin pre-mRNA. *FEBS Lett.*, 587(6):555–561, Mar 2013.
- [99] A. Katolik et al. Regiospecific Solid-Phase Synthesis of Branched Oligoribonucleotides That Mimic Intronic Lariat RNA Intermediates. *J. Org. Chem.*, Jan 2014.
- [100] S. Bennett. Solexa Ltd. *Pharmacogenomics*, 5(4):433–438, Jun 2004.
- [101] Z. Wang et al. RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev. Genet.*, 10(1):57–63, Jan 2009.
- [102] J. C. Marioni et al. RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays. *Genome Res.*, 18(9):1509–1517, Sep 2008.
- [103] M. J. Fullwood, C. L. Wei, E. T. Liu, and Y. Ruan. Next-generation dna sequencing of paired-end tags (pet) for transcriptome and genome analyses. *Genome Res.*, 2009.
- [104] I. Gazzoli et al. Non-sequential and recursive splicing of the dystrophin transcript. *Manuscript in preparation*.
- [105] P. Flicek et al. Ensembl 2011. *Nucleic Acids Res.*, 39(Database issue):D800–806, Jan 2011.
- [106] J. M. Ruijter et al. Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Res.*, 37(6):e45, Apr 2009.
- [107] T. D. Wu et al. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics*, 26(7):873–881, Apr 2010.
- [108] C. Trapnell et al. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics*, 25(9):1105–1111, May 2009.
- [109] M. T. Dimon et al. HMMSplicer: a tool for efficient and sensitive discovery of known and novel splice junctions in RNA-Seq data. *PLoS ONE*, 5(11):e13875, 2010.
- [110] M. Guttman et al. Ab initio reconstruction of cell type-specific transcriptomes in mouse reveals the conserved multi-exonic structure of lincRNAs. *Nat. Biotechnol.*, 28(5):503–510, May 2010.
- [111] Y. Katz et al. Analysis and design of rna sequencing experiments for identifying isoform regulation. *Nat. Biotechnol.*, 7(12):1009–1015, Dec 2010.
- [112] A. R. Hatton et al. Generation of alternative Ultra-bithorax isoforms and stepwise removal of a large intron by resplicing at exon-exon junctions. *Mol. Cell*, 2(6):787–796, Dec 1998.
- [113] T. A. Cooper et al. RNA and disease. *Cell*, 136(4):777–793, Feb 2009.
- [114] M. B. Coelho et al. Regulation of alternative pre-mRNA splicing. *Methods Mol. Biol.*, 1126:55–82, 2014.
- [115] M. C. Wahl et al. The spliceosome: design principles of a dynamic RNP machine. *Cell*, 136(4):701–718, Feb 2009.
- [116] H. Shen et al. RS domain-splicing signal interactions in splicing of U12-type and U2-type introns. *Nat. Struct. Mol. Biol.*, 14(7):597–603, Jul 2007.
- [117] G. E. Parada et al. A comprehensive survey of non-canonical splice sites in the human transcriptome. *Nucleic Acids Res.*, 42(16):10564–10578, 2014.
- [118] Y. Barash et al. Deciphering the splicing code. *Nature*, 465(7294):53–59, May 2010.
- [119] H. Suzuki et al. Characterization of RNase R-digested cellular RNA source that consists of lariat and circular RNAs from pre-mRNA splicing. *Nucleic Acids Res.*, 34(8):e63, 2006.
- [120] T. Takahara et al. Delay in synthesis of the 3' splice site promotes trans-splicing of the preceding 5' splice site. *Mol. Cell*, 18(2):245–251, Apr 2005.

- [121] J. Singh et al. Rates of in situ transcription and splicing in large human genes. *Nat. Struct. Mol. Biol.*, 16(11):1128–1133, Nov 2009.
- [122] R. F. Luco et al. Epigenetics in alternative pre-mRNA splicing. *Cell*, 144(1):16–26, Jan 2011.
- [123] S. Shukla et al. Co-transcriptional regulation of alternative pre-mRNA splicing. *Biochim. Biophys. Acta*, 1819(7):673–683, Jul 2012.
- [124] M. J. Moore et al. Pre-mRNA processing reaches back to transcription and ahead to translation. *Cell*, 136(4):688–700, Feb 2009.
- [125] R. Perales et al. "Cotranscriptionality": the transcription elongation complex as a nexus for nuclear transactions. *Mol. Cell*, 36(2):178–191, Oct 2009.
- [126] D. Y. Vargas et al. Single-molecule imaging of transcriptionally coupled and uncoupled splicing. *Cell*, 147(5):1054–1065, Nov 2011.
- [127] J. M. Gudas et al. Ordered splicing of thymidine kinase pre-mRNA during the S phase of the cell cycle. *Mol. Cell. Biol.*, 10(10):5591–5595, Oct 1990.
- [128] M. J. Tsai et al. Processing of high molecular weight ovalbumin and ovomucoid precursor RNAs to messenger RNA. *Cell*, 22(1 Pt 1):219–230, Nov 1980.
- [129] M. de la Mata et al. First come, first served revisited: factors affecting the same alternative splicing event have different effects on the relative rates of intron removal. *RNA*, 16(5):904–912, May 2010.
- [130] S. Shepard et al. The peculiarities of large intron splicing in animals. *PLoS ONE*, 4(11):e7853, 2009.
- [131] J. M. Burnette et al. Subdivision of large introns in *Drosophila* by recursive splicing at nonexonic elements. *Genetics*, 170(2):661–674, Jun 2005.
- [132] J. F. Conklin et al. Stabilization and analysis of intron lariats in vivo. *Methods*, 37(4):368–375, Dec 2005.
- [133] S. Ott et al. Intrasplicing—analysis of long intron sequences. *Pac Symp Biocomput*, pages 339–350, 2003.
- [134] N.P. Kandul et al. Large introns in relation to alternative splicing and gene evolution: a case study of *Drosophila bruno-3*. *BMC Genet.*, 10:67, 2009.
- [135] T. Kameyama et al. Re-splicing of mature mRNA in cancer cells promotes activation of distant weak alternative splice sites. *Nucleic Acids Res.*, 40(16):7896–7906, Sep 2012.
- [136] C. N. Tennyson et al. The human dystrophin gene requires 16 hours to be transcribed and is cotranscriptionally spliced. *Nat. Genet.*, 9(2):184–190, Feb 1995.
- [137] C. H. Zhu et al. Cellular senescence in human myoblasts is overcome by human telomerase reverse transcriptase and cyclin-dependent kinase 4: consequences in aging muscle and therapeutic strategies for muscular dystrophies. *Aging Cell*, 6(4):515–523, Aug 2007.
- [138] D. U. Kemaladewi et al. Dual exon skipping in myostatin and dystrophin for Duchenne muscular dystrophy. *BMC Med Genomics*, 4:36, 2011.
- [139] I. Pulyakhina et al. SplicePie: a novel analytical approach for the detection of alternative, non-sequential and recursive splicing. *Nucleic Acids Res.*, Mar 2015.
- [140] T. R. Mercer et al. Targeted sequencing for gene discovery and quantification using RNA CaptureSeq. *Nat Protoc*, 9(5):989–1009, May 2014.
- [141] A. Gnirke et al. Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted sequencing. *Nat. Biotechnol.*, 27(2):182–189, Feb 2009.
- [142] F. H. Nasim et al. A Sequential splicing mechanism promotes selection of an optimal exon by repositioning a downstream 5' splice site in preprotachykinin pre-mRNA. *Genes Dev.*, 4(7):1172–1184, Jul 1990.
- [143] O. Kessler et al. Order of intron removal during splicing of endogenous adenine phosphoribosyltransferase and dihydrofolate reductase pre-mRNA. *Mol. Cell. Biol.*, 13(10):6211–6222, Oct 1993.
- [144] M. Yang et al. Splicing of mouse p53 pre-mRNA does not always follow the "first come, first served" principle and may be influenced by cisplatin treatment and serum starvation. *Mol. Biol. Rep.*, 39(9):9247–9256, Sep 2012.
- [145] I. A. Swinburne et al. Intron length increases oscillatory periods of gene expression in animal cells. *Genes Dev.*, 22(17):2342–2346, Sep 2008.
- [146] K. M. Neugebauer. On the importance of being cotranscriptional. *J. Cell. Sci.*, 115(Pt 20):3865–3871, Oct 2002.
- [147] N. J. Proudfoot. Dawdling polymerases allow introns time to splice. *Nat. Struct. Biol.*, 10(11):876–878, Nov 2003.
- [148] M. Aebi et al. Precision and orderliness in splicing. *Trends. Genet.*, 3:102–107, Mar 1987.
- [149] A. P. Monaco et al. An explanation for the phenotypic differences between patients bearing partial deletions of the DMD locus. *Genomics*, 2(1):90–95, Jan 1988.
- [150] A. Aartsma-Rus. Antisense-mediated modulation of splicing: therapeutic implications for Duchenne muscular dystrophy. *RNA Biol*, 7(4):453–461, 2010.
- [151] A. Aartsma-Rus et al. Functional analysis of 114 exon-internal AONs for targeted DMD exon skipping: indication for steric hindrance of SR protein binding sites. *Oligonucleotides*, 15(4):284–297, Dec 2005.
- [152] S. D. Wilton et al. Antisense oligonucleotide-induced exon skipping across the human dystrophin gene transcript. *Mol. Ther.*, 15(7):1288–1296, Jul 2007.

- [153] A. Aartsma-Rus et al. Antisense-induced multiexon skipping for Duchenne muscular dystrophy makes more sense. *Am. J. Hum. Genet.*, 74(1):83–92, Jan 2004.
- [154] M. Koenig et al. The molecular basis for Duchenne versus Becker muscular dystrophy: correlation of severity with type of deletion. *Am. J. Hum. Genet.*, 45(4):498–506, Oct 1989.
- [155] J. T. Den Dunnen et al. Topography of the Duchenne muscular dystrophy (DMD) gene: FIGE and cDNA analysis of 194 cases reveals 115 deletions and 13 duplications. *Am. J. Hum. Genet.*, 45(6):835–847, Dec 1989.
- [156] C. Nobile et al. Exon-intron organization of the human dystrophin gene. *Genomics*, 45(2):421–424, Oct 1997.
- [157] A. Aartsma-Rus et al. Theoretic applicability of antisense-mediated exon skipping for Duchenne muscular dystrophy mutations. *Hum. Mutat.*, 30(3):293–299, Mar 2009.
- [158] Y. Aoki et al. Bodywide skipping of exons 45–55 in dystrophic mdx52 mice by systemic antisense delivery. *Proc. Natl. Acad. Sci. U.S.A.*, 109(34):13763–13768, Aug 2012.
- [159] C. R. Sibley et al. Recursive splicing in long vertebrate genes. *Nature*, 521(7552):371–375, May 2015.
- [160] T. Lu et al. Gene regulation and DNA damage in the ageing human brain. *Nature*, 429(6994):883–891, Jun 2004.
- [161] S. Hekimi et al. Genetics and the specificity of the aging process. *Science*, 299(5611):1351–1354, Feb 2003.
- [162] S. A. McCarroll et al. Comparing genomic expression patterns across species identifies shared transcriptional profile in aging. *Nat. Genet.*, 36(2):197–204, Feb 2004.
- [163] W. M. Passtoors et al. Genomic studies in ageing research: the need to integrate genetic and gene expression approaches. *J. Intern. Med.*, 263(2):153–166, Feb 2008.
- [164] M. Jung et al. Longitudinal epigenetic and gene expression profiles analyzed by three-component analysis reveal down-regulation of genes involved in protein translation in human aging. *Nucleic Acids Res.*, May 2015.
- [165] F. Licastro et al. Innate immunity and inflammation in ageing: a key for understanding age-related diseases. *Immun Ageing*, 2:8, May 2005.
- [166] M. Gheorghe et al. Major aging-associated RNA expressions change at two distinct age-positions. *BMC Genomics*, 15:132, 2014.
- [167] Y. N. Wei et al. Transcript and protein expression decoupling reveals RNA binding proteins and miRNAs as potential modulators of human aging. *Genome Biol.*, 16:41, 2015.
- [168] J. M. Johnson et al. Genome-wide survey of human alternative pre-mRNA splicing with exon junction microarrays. *Science*, 302(5653):2141–2144, Dec 2003.
- [169] J. Takeda et al. H-DBAS: human-transcriptome database for alternative splicing: update 2010. *Nucleic Acids Res.*, 38(Database issue):86–90, Jan 2010.
- [170] R. Blekhman et al. Sex-specific and lineage-specific alternative splicing in primates. *Genome Res.*, 20(2):180–189, Feb 2010.
- [171] J. A. Calarco et al. Global analysis of alternative splicing differences between humans and chimpanzees. *Genes Dev.*, 21(22):2963–2975, Nov 2007.
- [172] A. Battle et al. Characterizing the genetic basis of transcriptome diversity through RNA-sequencing of 922 individuals. *Genome Res.*, 24(1):14–24, Jan 2014.
- [173] T. Lappalainen et al. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature*, 501(7468):506–511, Sep 2013.
- [174] J. Ule et al. Nova regulates brain-specific splicing to shape the synapse. *Nat. Genet.*, 37(8):844–852, Aug 2005.
- [175] B. J. Blencowe. Alternative splicing: new insights from global analyses. *Cell*, 126(1):37–47, Jul 2006.
- [176] B. K. Dredge et al. The splice of life: alternative splicing and neurological disease. *Nat. Rev. Neurosci.*, 2(1):43–50, Jan 2001.
- [177] L. Buee et al. Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res. Brain Res. Rev.*, 33(1):95–130, Aug 2000.
- [178] L. W. Harries et al. Human aging is characterized by focused changes in gene expression and deregulation of alternative splicing. *Aging Cell*, 10(5):868–878, Oct 2011.
- [179] P. Mazin et al. Widespread splicing changes in human brain development and aging. *Mol. Syst. Biol.*, 9:633, 2013.
- [180] S. Scholtens et al. Cohort Profile: LifeLines, a three-generation cohort study and biobank. *Int J Epidemiol*, Dec 2014.
- [181] www.lumc.nl/org/ouderengeneeskunde/research/study-populations/LeidenLongevity-study/. Accessed: 2015-06-01.
- [182] A. Dobin et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1):15–21, Jan 2013.
- [183] H. Konig et al. Splicing segregation: the minor spliceosome acts outside the nucleus and controls cell proliferation. *Cell*, 131(4):718–729, Nov 2007.

- [184] W. Y. Tarn et al. Highly diverged U4 and U6 small nuclear RNAs required for splicing rare AT-AC introns. *Science*, 273(5283):1824–1832, Sep 1996.
- [185] S. L. Hall et al. Requirement of U12 snRNA for in vivo splicing of a minor class of eukaryotic nuclear pre-mRNA introns. *Science*, 271(5256):1716–1718, Mar 1996.
- [186] W. Y. Tarn et al. A novel spliceosome containing U11, U12, and U5 snRNPs excises a minor class (AT-AC) intron in vitro. *Cell*, 84(5):801–811, Mar 1996.
- [187] T. S. Alioto. U12DB: a database of orthologous U12-type spliceosomal introns. *Nucleic Acids Res.*, 35(Database issue):D110–115, Jan 2007.
- [188] M. Q. Zhang. Statistical features of human exons and their flanking regions. *Hum. Mol. Genet.*, 7(5):919–932, May 1998.
- [189] R. K. Bradley et al. Alternative splicing of RNA triplets is often regulated and accelerates proteome evolution. *PLoS Biol.*, 10(1):e1001229, Jan 2012.
- [190] M. Hiller et al. Widespread occurrence of alternative splicing at NAGNAG acceptors contributes to proteome plasticity. *Nat. Genet.*, 36(12):1255–1257, Dec 2004.
- [191] A. Busch et al. Extensive regulation of NAGNAG alternative splicing: new tricks for the spliceosome? *Genome Biol.*, 13(2):143, 2012.
- [192] A. E. Jaffe et al. Accounting for cellular heterogeneity is critical in epigenome-wide association studies. *Genome Biol.*, 15(2):R31, 2014.
- [193] J. J. Turunen et al. The significant other: splicing by the minor spliceosome. *Wiley Interdiscip Rev RNA*, 4(1):61–76, 2013.
- [194] J. Yang et al. Association of DNA methylation in the brain with age in older persons is confounded by common neuropathologies. *Int. J. Biochem. Cell Biol.*, May 2015.
- [195] C. Kohler et al. Release of adenylate kinase 2 from the mitochondrial intermembrane space during apoptosis. *FEBS Lett.*, 447(1):10–12, Mar 1999.
- [196] G. A. Bruns et al. Adenylate kinase 2, a mitochondrial enzyme. *Biochem. Genet.*, 15(5-6):477–486, Jun 1977.
- [197] T. H. Ho et al. Muscleblind proteins regulate alternative splicing. *EMBO J.*, 23(15):3103–3112, Aug 2004.
- [198] C. de Martel et al. Global burden of cancers attributable to infections in 2008: a review and synthetic analysis. *Lancet Oncol.*, 13(6):607–615, Jun 2012.
- [199] I. P. Tomlinson et al. Loss of heterozygosity on chromosome 11 q in breast cancer. *J. Clin. Pathol.*, 48(5):424–428, May 1995.
- [200] S. E. Shackney et al. Aneuploidy in breast cancer: a fluorescence in situ hybridization study. *Cytometry*, 22(4):282–291, Dec 1995.
- [201] A. A. Owainati et al. Tumour aneuploidy, prognostic parameters and survival in primary breast cancer. *Br. J. Cancer*, 55(4):449–454, Apr 1987.
- [202] D. Hanahan et al. Hallmarks of cancer: the next generation. *Cell*, 144(5):646–674, Mar 2011.
- [203] S. Sen. Aneuploidy and cancer. *Curr Opin Oncol*, 12(1):82–88, Jan 2000.
- [204] E. Manchado et al. Killing cells by targeting mitosis. *Cell Death Differ.*, 19(3):369–377, Mar 2012.
- [205] D. J. Gordon et al. Causes and consequences of aneuploidy in cancer. *Nat. Rev. Genet.*, 13(3):189–203, Mar 2012.
- [206] N. C. Levitt et al. Caretaker tumour suppressor genes that defend genome integrity. *Trends Mol Med*, 8(4):179–186, Apr 2002.
- [207] K. A. Kwei et al. Genomic instability in breast cancer: pathogenesis and clinical implications. *Mol Oncol*, 4(3):255–266, Jun 2010.
- [208] A. Balmain et al. The genetics and genomics of cancer. *Nat. Genet.*, 33 Suppl:238–244, Mar 2003.
- [209] B. Orsetti et al. Genomic and expression profiling of chromosome 17 in breast cancer reveals complex patterns of alterations and novel candidate genes. *Cancer Res.*, 64(18):6453–6460, Sep 2004.
- [210] G. L. Ottesen et al. DNA aneuploidy in early breast cancer. *Br. J. Cancer*, 72(4):832–839, Oct 1995.
- [211] G. Marx. Possible function found for breast cancer genes. *Science*, 276(5312):531–532, Apr 1997.
- [212] E. Rossi et al. High-level detection of gene amplification and chromosome aneuploidy in extracted nuclei from paraffin-embedded tissue of human cancer using FISH: a new approach for retrospective studies. *Eur J Histochem*, 49(1):53–58, 2005.
- [213] S. M. Johnson et al. Sporadic breast cancer in young women: prevalence of loss of heterozygosity at p53, BRCA1 and BRCA2. *Int. J. Cancer*, 98(2):205–209, Mar 2002.
- [214] B. J. Miller et al. Pooled analysis of loss of heterozygosity in breast cancer: a genome scan provides comparative evidence for multiple tumor suppressors and identifies novel candidate regions. *Am. J. Hum. Genet.*, 73(4):748–767, Oct 2003.
- [215] H. Schwarzenbach et al. A critical evaluation of loss of heterozygosity detected in tumor tissues, blood serum and bone marrow plasma from patients with breast cancer. *Breast Cancer Res.*, 9(5):R66, 2007.
- [216] C. O’Keefe et al. Copy neutral loss of heterozygosity: a novel chromosomal lesion in myeloid malignancies. *Blood*, 115(14):2731–2739, Apr 2010.

- [217] M. Tuna et al. Prognostic value of acquired uniparental disomy (aUPD) in primary breast cancer. *Breast Cancer Res. Treat.*, 132(1):189–196, Feb 2012.
- [218] P. A. Futreal et al. BRCA1 mutations in primary breast and ovarian carcinomas. *Science*, 266(5182):120–122, Oct 1994.
- [219] M. Janatova et al. Novel somatic mutations in the BRCA1 gene in sporadic breast tumors. *Hum. Mutat.*, 25(3):319, Mar 2005.
- [220] U. S. Khoo et al. Somatic mutations in the BRCA1 gene in Chinese sporadic breast and ovarian cancer. *Oncogene*, 18(32):4643–4646, Aug 1999.
- [221] Z. Haitian et al. Mutation screening of the BRCA1 gene in sporadic breast cancer in southern Chinese populations. *Breast*, 17(6):563–567, Dec 2008.
- [222] F. M. Chen et al. High frequency of somatic missense mutation of BRCA2 in female breast cancer from Taiwan. *Cancer Lett.*, 220(2):177–184, Apr 2005.
- [223] M. Morley et al. Genetic analysis of genome-wide variation in human gene expression. *Nature*, 430(7001):743–747, Aug 2004.
- [224] H. S. Lo et al. Allelic variation in gene expression is common in the human genome. *Genome Res.*, 13(8):1855–1862, Aug 2003.
- [225] S. A. Smith et al. Allele losses in the region 17q12-21 in familial breast and ovarian cancer involve the wild-type chromosome. *Nat. Genet.*, 2(2):128–131, Oct 1992.
- [226] R. S. Cornelis et al. High allele loss rates at 17q12-q21 in breast and ovarian tumors from BRCA1-linked families. The Breast Cancer Linkage Consortium. *Genes Chromosomes Cancer*, 13(3):203–210, Jul 1995.
- [227] A. Osorio et al. Loss of heterozygosity analysis at the BRCA loci in tumor samples from patients with familial breast cancer. *Int. J. Cancer*, 99(2):305–309, May 2002.
- [228] M. E. Thompson et al. Decreased expression of BRCA1 accelerates growth and is often present during sporadic breast cancer progression. *Nat. Genet.*, 9(4):444–450, Apr 1995.
- [229] C. A. Wilson et al. Localization of human BRCA1 and its loss in high-grade, non-inherited breast carcinomas. *Nat. Genet.*, 21(2):236–240, Feb 1999.
- [230] S. Staff et al. Haplo-insufficiency of BRCA1 in sporadic breast cancer. *Cancer Res.*, 63(16):4978–4983, Aug 2003.
- [231] P. Van Loo et al. Allele-specific copy number analysis of tumors. *Proc. Natl. Acad. Sci. U.S.A.*, 107(39):16910–16915, Sep 2010.
- [232] P. P. Wegman et al. Biological significance of allele specific loss of the p53 gene in breast carcinomas. *Breast Cancer Res. Treat.*, 118(1):15–20, Nov 2009.
- [233] H. Yan et al. Small changes in expression affect predisposition to tumorigenesis. *Nat. Genet.*, 30(1):25–26, Jan 2002.
- [234] A. T. Maia et al. Extent of differential allelic expression of candidate breast cancer genes is similar in blood and breast. *Breast Cancer Res.*, 11(6):R88, 2009.
- [235] T. Nguyen-Dumont et al. Detecting differential allelic expression using high-resolution melting curve analysis: application to the breast cancer susceptibility gene CHEK2. *BMC Med Genomics*, 4:39, 2011.
- [236] C. Gao et al. Identifying breast cancer risk loci by global differential allele-specific expression (DASE) analysis in mammary epithelial transcriptome. *BMC Genomics*, 13:570, 2012.
- [237] A. Kallioniemi et al. Comparative genomic hybridization for molecular cytogenetic analysis of solid tumors. *Science*, 258(5083):818–821, Oct 1992.
- [238] Y. Li et al. Deconvolving tumor purity and ploidy by integrating copy number alterations and loss of heterozygosity. *Bioinformatics*, 30(15):2121–2129, Aug 2014.
- [239] B. Li et al. A general framework for analyzing tumor subclonality using SNP array and DNA sequencing data. *Genome Biol.*, 15(9):473, 2014.
- [240] T. V. Hansen et al. High-density SNP arrays improve detection of HER2 amplification and polyploidy in breast tumors. *BMC Cancer*, 15:35, 2015.
- [241] M. Perez-Enciso et al. Sequence- vs. chip-assisted genomic selection: accurate biological information is advised. *Genet. Sel. Evol.*, 47:43, 2015.
- [242] F. Kaveh et al. Allele-specific disparity in breast cancer. *BMC Med Genomics*, 4:85, 2011.
- [243] G. Ha et al. Integrative analysis of genome-wide loss of heterozygosity and monoallelic expression at nucleotide resolution reveals disrupted pathways in triple-negative breast cancer. *Genome Res.*, 22(10):1995–2007, Oct 2012.
- [244] Q. Zhao et al. Systematic detection of putative tumor suppressor genes through the combined use of exome and transcriptome sequencing. *Genome Biol.*, 11(11):R114, 2010.
- [245] O. Vierimaa et al. Pituitary adenoma predisposition caused by germline mutations in the AIP gene. *Science*, 312(5777):1228–1230, May 2006.
- [246] K. M. Lee et al. Genetic polymorphisms of selected DNA repair genes, estrogen and progesterone receptor status, and breast cancer risk. *Clin. Cancer Res.*, 11(12):4620–4626, Jun 2005.
- [247] H. S. Lo et al. Allelic variation in gene expression is common in the human genome. *Genome Res.*, 13(8):1855–1862, Aug 2003.

- [248] D. C. Koboldt et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res.*, 22(3):568–576, Mar 2012.
- [249] M. S. Lawrence et al. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature*, 505(7484):495–501, Jan 2014.
- [250] L. A. Garraway et al. Lessons from the cancer genome. *Cell*, 153(1):17–37, Mar 2013.
- [251] P. Creixell et al. Pathway and network analysis of cancer genomes. *Nat. Methods*, 12(7):615–621, Jun 2015.
- [252] E. Garcion et al. RNA mutagenesis and sporadic prion diseases. *J. Theor. Biol.*, 230(2):271–274, Sep 2004.
- [253] O. D. Iancu et al. Utilizing RNA-Seq data for de novo coexpression network inference. *Bioinformatics*, 28(12):1592–1597, Jun 2012.
- [254] F. M. Giorgi et al. Comparative study of RNA-seq and microarray-derived coexpression networks in *Arabidopsis thaliana*. *Bioinformatics*, 29(6):717–724, Mar 2013.
- [255] N. Calabriso et al. Multiple anti-inflammatory and anti-atherosclerotic properties of red wine polyphenolic extracts: differential role of hydroxycinnamic acids, flavonols and stilbenes on endothelial inflammatory gene expression. *Eur J Nutr*, Feb 2015.
- [256] J. Hogan et al. Transcriptional profiles underpin microsatellite status and associated features in colon cancer. *Gene*, Feb 2015.
- [257] R. Al-Bahrani et al. Differential SIRT1 Expression in Hepatocellular Carcinomas and Cholangiocarcinoma of the Liver. *Ann. Clin. Lab. Sci.*, 45(1):3–9, Jan 2015.
- [258] J. C. Betts et al. Gene expression changes caused by the p38 MAPK inhibitor dilmapiomod in COPD patients: analysis of blood and sputum samples from a randomized, placebo-controlled clinical trial. *Pharmacol Res Perspect*, 3(1):e00094, Feb 2015.
- [259] E. L. van Dijk et al. Library preparation methods for next-generation sequencing: tone down the bias. *Exp. Cell Res.*, 322(1):12–20, Mar 2014.
- [260] J. Durtschi et al. VarBin, a novel method for classifying true and false positive variants in NGS data. *BMC Bioinformatics*, 14 Suppl 13:S2, 2013.
- [261] S. M. Teo et al. Statistical challenges associated with detecting copy number variations with next-generation sequencing. *Bioinformatics*, 28(21):2711–2718, Nov 2012.
- [262] J. D. Roberts et al. Fidelity of two retroviral reverse transcriptases during DNA-dependent DNA synthesis in vitro. *Mol. Cell. Biol.*, 9(2):469–476, Feb 1989.
- [263] Z. Su et al. Next-generation sequencing and its applications in molecular diagnostics. *Expert Rev. Mol. Diagn.*, 11(3):333–343, Apr 2011.
- [264] I. Braslavsky et al. Sequence information can be obtained from single DNA molecules. *Proc. Natl. Acad. Sci. U.S.A.*, 100(7):3960–3964, Apr 2003.
- [265] T. Raz et al. RNA sequencing and quantitation using the Helicos Genetic Analysis System. *Methods Mol. Biol.*, 733:37–49, 2011.
- [266] F. Ozsolak et al. Single-molecule direct RNA sequencing without cDNA synthesis. *Wiley Interdiscip Rev RNA*, 2(4):565–570, 2011.
- [267] J. Quick et al. A reference bacterial genome dataset generated on the MinION(TM) portable single-molecule nanopore sequencer. *Gigascience*, 3:22, 2014.
- [268] E. A. Manrao et al. Reading DNA at single-nucleotide resolution with a mutant MspA nanopore and phi29 DNA polymerase. *Nat. Biotechnol.*, 30(4):349–353, Apr 2012.
- [269] C. Merstorf et al. Wild type, mutant protein unfolding and phase transition detected by single-nanopore recording. *ACS Chem. Biol.*, 7(4):652–658, Apr 2012.
- [270] D. S. Talaga et al. Single-molecule protein unfolding in solid state nanopores. *J. Am. Chem. Soc.*, 131(26):9287–9297, Jul 2009.
- [271] M. M. Mohammad et al. Controlling a single protein in a nanopore through electrostatic traps. *J. Am. Chem. Soc.*, 130(12):4081–4088, Mar 2008.
- [272] J. Nivala et al. Unfoldase-mediated protein translocation through an alpha-hemolysin nanopore. *Nat. Biotechnol.*, 31(3):247–250, Mar 2013.
- [273] Y. Motorin et al. RNA nucleotide methylation. *Wiley Interdiscip Rev RNA*, 2(5):611–631, 2011.
- [274] E. Picardi et al. Uncovering RNA Editing Sites in Long Non-Coding RNAs. *Front Bioeng Biotechnol*, 2:64, 2014.
- [275] L. A. Crews et al. An RNA editing fingerprint of cancer stem cell reprogramming. *J Transl Med*, 13(1):370, Dec 2015.
- [276] M. M. Ahasan et al. APOBEC3A and 3C decrease human papillomavirus 16 pseudovirion infectivity. *Biochem. Biophys. Res. Commun.*, 457(3):295–299, Feb 2015.
- [277] X. Hu et al. RNA over-editing of BLCAP contributes to hepatocarcinogenesis identified by whole-genome and transcriptome sequencing. *Cancer Lett.*, 357(2):510–519, Feb 2015.
- [278] L. Kang et al. Genome-wide identification of RNA editing in hepatocellular carcinoma. *Genomics*, 105(2):76–82, Feb 2015.
- [279] C. M. Wei et al. Methylated nucleotides block 5' terminus of HeLa cell messenger RNA. *Cell*, 4(4):379–386, Apr 1975.

- [280] F. Rottman et al. Sequences containing methylated nucleotides at the 5' termini of messenger RNAs: possible implications for processing. *Cell*, 3(3):197–199, Nov 1974.
- [281] P. Narayan et al. An in vitro system for accurate methylation of internal adenosine residues in messenger RNA. *Science*, 242(4882):1159–1162, Nov 1988.
- [282] T. Csepány et al. Sequence specificity of mRNA N6-adenosine methyltransferase. *J. Biol. Chem.*, 265(33):20117–20122, Nov 1990.
- [283] P. Narayan et al. Context effects on N6-adenosine methylation sites in prolactin mRNA. *Nucleic Acids Res.*, 22(3):419–426, Feb 1994.
- [284] Y. Fu et al. Gene expression regulation mediated through reversible m6A RNA methylation. *Nat. Rev. Genet.*, 15(5):293–306, May 2014.
- [285] K. M. Lonergan et al. Predicted editing of additional transfer RNAs in *Acanthamoeba castellanii* mitochondria. *Nucleic Acids Res.*, 21(18):4402, Sep 1993.
- [286] L. Agranat et al. The editing enzyme ADAR1 and the mRNA surveillance protein hUpf1 interact in the cell nucleus. *Proc. Natl. Acad. Sci. U.S.A.*, 105(13):5028–5033, Apr 2008.
- [287] D. Speijer. Does constructive neutral evolution play an important role in the origin of cellular complexity? Making sense of the origins and uses of biological complexity. *Bioessays*, 33(5):344–349, May 2011.
- [288] Y. Shu et al. Predicting A-to-I RNA editing by feature selection and random forest. *PLoS ONE*, 9(10):e110607, 2014.
- [289] H. Q. Zhao et al. Profiling the RNA editomes of wild-type *C. elegans* and ADAR mutants. *Genome Res.*, 25(1):66–75, Jan 2015.
- [290] J. M. Toung et al. Detection theory in identification of RNA-DNA sequence differences using RNA-sequencing. *PLoS ONE*, 9(11):e112040, 2014.
- [291] M. A. Huntley et al. Dissecting gene expression at the blood-brain barrier. *Front Neurosci*, 8:355, 2014.
- [292] J. Gong et al. Comprehensive analysis of human small RNA sequencing data provides insights into expression profiles and miRNA editing. *RNA Biol*, 11(11):1375–1385, Nov 2014.
- [293] S. Haider et al. Integrated analysis of transcriptomic and proteomic data. *Curr. Genomics*, 14(2):91–110, Apr 2013.
- [294] L. Nie et al. Integrative analysis of transcriptomic and proteomic data: challenges, solutions and applications. *Crit. Rev. Biotechnol.*, 27(2):63–75, 2007.
- [295] M. Hecker et al. Gene regulatory network inference: data integration in dynamic models-a review. *BioSystems*, 96(1):86–103, Apr 2009.
- [296] S. Rogers. Statistical methods and models for bridging Omics data levels. *Methods Mol. Biol.*, 719:133–151, 2011.
- [297] S. P. Gygi et al. Correlation between protein and mRNA abundance in yeast. *Mol. Cell. Biol.*, 19(3):1720–1730, Mar 1999.
- [298] G. Chen et al. Discordant protein and mRNA expression in lung adenocarcinomas. *Mol. Cell Proteomics*, 1(4):304–313, Apr 2002.
- [299] L. E. Pascal et al. Correlation of mRNA and protein levels: cell type-specific gene expression of cluster designation antigens in the prostate. *BMC Genomics*, 9:246, 2008.
- [300] A. Ghazalpour et al. Comparative analysis of proteome and transcriptome variation in mouse. *PLoS Genet.*, 7(6):e1001393, Jun 2011.
- [301] N. T. Ingolia et al. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science*, 324(5924):218–223, Apr 2009.
- [302] C. Gustafsson et al. Codon bias and heterologous protein expression. *Trends Biotechnol.*, 22(7):346–353, Jul 2004.
- [303] T. Steijger et al. Assessment of transcript reconstruction methods for RNA-seq. *Nat. Methods*, 10(12):1177–1184, Dec 2013.
- [304] H. Wang et al. Detection of allelic imbalance in gene expression using pyrosequencing. *Methods Mol. Biol.*, 373:157–176, 2007.
- [305] M. Pirinen et al. Assessing allele-specific expression across multiple tissues from RNA-seq read data. *Bioinformatics*, Mar 2015.
- [306] K. R. Kukurba et al. Allelic expression of deleterious protein-coding variants across human tissues. *PLoS Genet.*, 10(5):e1004304, May 2014.
- [307] H. Roger et al. [Non-familial Duchenne-Erb myopathy appearing after encephalitis in late childhood; association with mental deficiency and cutaneous dystrophy]. *Bull Mem Soc Med Hop Paris*, 67(7-8):279–281, 1951.
- [308] E. A. Cheeseman et al. The sex ratio of mutation rates of sex-linked recessive genes in man with particular reference to Duchenne type muscular dystrophy. *Ann. Hum. Genet.*, 22(3):235–243, May 1958.
- [309] C. L. Bladen et al. The TREAT-NMD DMD Global database: Analysis of More Than 7000 Duchenne Muscular Dystrophy Mutations. *Hum. Mutat.*, Jan 2015.
- [310] A. De Sandre-Giovannoli et al. Lamin a truncation in Hutchinson-Gilford progeria. *Science*, 300(5628):2055, Jun 2003.
- [311] B. P. O’Sullivan et al. Cystic fibrosis. *Lancet*, 373(9678):1891–1904, May 2009.

- [312] V. Waters et al. Cystic fibrosis microbiology: Advances in antimicrobial therapy. *J. Cyst. Fibros.*, Feb 2015.
- [313] A. Warshel et al. Theoretical studies of enzymic reactions: dielectric, electrostatic and steric stabilization of the carbonium ion in the reaction of lysozyme. *J. Mol. Biol.*, 103(2):227–249, May 1976.
- [314] M. T. Panteva et al. Multiscale methods for computational RNA enzymology. *Meth. Enzymol.*, 553:335–374, 2015.
- [315] D. Xu et al. AIDA: Ab Initio Domain Assembly for Automated Multi-domain Protein Structure Prediction and Domain-Domain Interaction Prediction. *Bioinformatics*, Feb 2015.
- [316] J. M. Chandonia et al. The impact of structural genomics: expectations and outcomes. *Science*, 311(5759):347–351, Jan 2006.
- [317] D. Baker et al. Protein structure prediction and structural genomics. *Science*, 294(5540):93–96, Oct 2001.
- [318] S. Goldsmith-Fischman et al. Structural genomics: computational methods for structure analysis. *Protein Sci.*, 12(9):1813–1821, Sep 2003.
- [319] S. A. Lesley et al. Structural genomics of the *Thermotoga maritima* proteome implemented in a high-throughput structure determination pipeline. *Proc. Natl. Acad. Sci. U.S.A.*, 99(18):11664–11669, Sep 2002.
- [320] M. O. Duff et al. Genome-wide identification of zero nucleotide recursive splicing in *Drosophila*. *Nature*, 521(7552):376–379, May 2015.
- [321] Z. K. Hassan-Smith et al. Gender-specific differences in skeletal muscle 11B-HSD1 expression across healthy aging. *J. Clin. Endocrinol. Metab.*, page jc20151516, May 2015.