Phylogenetic and taxonomic studies in Macaranga, Mallotus and other acalyphoid genera (Euphorbiaceae s.s.)

Kulju, K.K.M.

REFERENCES

ADANSON, M. 1763. Familles des plantes 2. Vincent, Paris.

AIRY SHAW, H.K. 1963. Notes on Malaysian and other Asiatic Euphorbiaceae. Kew Bull. 16: 341–372.

AIRY SHAW, H.K. 1965. Notes on Malaysian and other Asiatic Euphorbiaceae. Kew Bull. 19: 299–328.

AIRY SHAW, H.K. 1966. Malaysian and other Asiatic Euphorbiaceae. Kew Bull. 20: 379–415.

AIRY SHAW, H.K. 1968. Malesian and other Asiatic Euphorbiaceae. Kew Bull. 21: 379–400.

AIRY SHAW, H.K. 1969. Notes on Malesian and other Asiatic Euphorbiaceae. Kew Bull. 23: 1–131.

AIRY SHAW, H.K. 1971. Notes on Malesian and other Asiatic Euphorbiaceae. Kew Bull. 25: 473–553.

AIRY SHAW, H.K. 1972. The Euphorbiaceae of Siam. Kew Bull. 26: 191–363.

AIRY SHAW, H.K. 1974. Noteworthy Euphorbiaceae from tropical Asia. Hook. Ic. Pl. 38: 3716.

AIRY SHAW, H.K. 1975. The Euphorbiaceae of Borneo. Kew Bull. Add. Ser. 4.

AIRY SHAW, H.K. 1980a. The Euphorbiaceae of New Guinea. Kew Bull. Add. Ser. 8.

AIRY SHAW, H.K. 1980b. New Euphorbiaceae from New Guinea. Kew Bull. 34: 591–606.

AIRY SHAW, H.K. 1980c. A partial synopsis of the Euphorbiaceae–Platylobeae of Australia (excluding Phyllanthus, Euphorbia and Calycopeplus). Kew Bull. 35: 577–700.

AIRY SHAW, H.K. 1981. The Euphorbiaceae of Sumatra. Kew Bull. 36: 239–374.

AIRY SHAW, H.K. 1982. The Euphorbiaceae of Central Malesia (Celebes, Moluccas, Lesser Sunda Is.). Kew Bull. 37: 1–40.

AIRY SHAW, H.K. 1983. An alphabetical enumeration of the Euphorbiaceae of the Philippine islands. Royal Botanic Gardens, Kew.

AKHMETIEV, M.A. & S.V. VIKULIN. 1995. Peltate leaves *Macaranga* Thouin (Euphorbiaceae) in the Cenozoic of Eastern Sikhote-Alin' [in Russian]. Paleontol. J. 1995(2): 151–156.

ALEJANDRO, G.D., S.G. RAZAFIMANDIMBISON & S. LIEDE-SCHUMANN. 2005. Polyphyly of *Mussaenda* inferred from ITS and trnT-F data and its implications for generic limits in Mussaendeae (Rubiaceae). Amer. J. Bot. 92: 544–557.

ALFARO, M.E., S. ZOLLER & F. LUTZONI. 2003. Bayes or bootstrap? A simulation study comparing the performance of Bayesian Markov chain Monte Carlo sampling and bootstrapping in assessing phylogenetic confidence. Mol. Biol. Evol. 20: 255–266.

ANGIOSPERM PHYLOGENY GROUP (APG). 1998. An ordinal classification for the families of flowering plants. Ann. Missouri Bot. Gard. 85: 531–553.

ANGIOSPERM PHYLOGENY GROUP (APG II). 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. Bot. J. Linn. Soc. 141: 399–436.

AUSTIN, J.J., E.N. ARNOLD & C.G. JONES. 2004. Reconstructing and island radiation using ancient and recent DNA: the extinct and living day geckos (*Phelsuma*) of the Mascarene islands. Mol. Phylog. Evol. 31: 109–122.

BACKER, C.A. & R.C. BAKHUIZEN VAN DEN BRINK F. 1964. Flora of Java 1. P. Noordhoff, Groningen.

BAILLON, H. 1858. Étude générale du groupe des Euphorbiacées. V. Masson, Paris.

BAILLON, H. 1860. Species Euphorbiacearum. A. Euphorbiacées Africaines. Adansonia 1: 58–87.

BÄNFER, G., B. FIALA & K. WEISING. 2004. AFLP analysis of phylogenetic relationships among myrmecophytic species of *Macaranga* (Euphorbiaceae) and their allies. Plant Syst. Evol. 249:

213–231.
BÄNFER, G., U. MOOG, B. FIALA, M. MOHAMED, K. WEISING & F.R. BLATTNER. 2006. A chloroplast genealogy of myrmecophytic *Macaranga* species (Euphorbiaceae) in Southeast Asia reveals hybridisation, vicariance and long-distance dispersal. Mol. Ecol. 15: 4409–4424.
BEDDOME, R.H. 1861. Contributions to the botany of southern India. Madras J. Lit. Sci. 2, 22: 70–71.
BEDDOME, R.H. 1872. Flora sylvatica 2. Adelphi Press, Madras.
BENTHAM, G. & J.D. HOOKER. 1880. Genera plantarum 3, 1. L. Reeve & Co, London.
BERRY, P.E., A.L. HIPP, K.J. WURDACK, B. VAN EE & R. RIINA. 2005. Molecular phylogenetics of the giant genus *Croton* and tribe Crotoneae (Euphorbiaceae sensu stricto) using ITS and *trnL-trnF* DNA sequence data. Amer. J. Bot. 92: 1520–1534.
BLATTNER, F.R., K. WEISING, G. BÄNFER, U. MASCHWITZ & B. FIALA. 2001. Molecular analysis of phylogenetic relationships among myrmecophytic *Macaranga* species (Euphorbiaceae). Mol. Phylog. Evol. 19: 331–344.
BLUME, C. L. 1826. Bijdragen tot de flora van Nederlandsch Indië, 12de stuk. Lands Drukkerij, Batavia.
BOLLENDORFF, S.M., P.C. VAN WELZEN & J.W.F. SLIK. 2000. A taxonomic revision of *Mallotus* section *Polyadenii* (Euphorbiaceae). Blumea 45: 319–340.
BRANDIS, D. 1906. Indian trees. A. Constable & Co., London.
BURKILL, I.H. 1966. A dictionary of the economic products of the Malay Peninsula, repr., 1. Crown Agents for Overseas Governments and Administrations, London.
BURMAN, N.L. 1768. Flora Indica. C. Haak & J. Schreuder, Leiden & Amsterdam.
CASARETTO, J. 1843. Novarum stirpium brasiliensium decades. Joannis Ferrandi, Genuae.
CHAKRABARTY, T. & M. GANGOPADHYAY. 1988a. A new species of *Cleidion* Bl. (Euphorbiaceae) from Burma. J. Econ. Taxon. Bot. 12: 473–474.
CHAKRABARTY, T. & M. GANGOPADHYAY. 1988b. Notes on some Asiatic Euphorbiaceae 2. J. Econ. Taxon. Bot. 12: 491–493.
CHASE, M.W., S. ZMARZTY, M.D. LLEDO, K.J. WURDACK, S.M. SWENSEN & M.F. FAY. 2002. When in doubt, put it in *Flacourtiaceae*: a molecular phylogenetic analysis based on plastid *rbcL* DNA sequences. Kew Bull. 57: 141–181.
CHASE, M.W., D.E. SOLTIS, R.G. OLMSTEAD, D. MORGAN, D.H. LES, B.D. MISHLER, M.R. DUVALL, R.A. PRICE, H.G. HILLS, Y.L. QIU, K.A. KRON, J.H. RETTIG, E. CONTI, J.D. PALMER, J.R. MANHART, K.J. SYTSMA, H.J. MICHAELS, W.J. KRESS, K.G. KAROL, W.D. CLARK, M. HEDREN, B.S. GAUT, R.K. JANSEN, K.J. KIM, C.F. WIMPEE, J.F. SMITH, G.R. FURNIER, S.H. STRAUSS, Q.Y. XIANG, G.M. PLUNKETT, P.S. SOLTIS, S.M. SWENSEN, S.E. WILLIAMS, P.A. GADEK, C.J. QUINN, L.E. EGUIARTE, E. GOLENBERG, G.H. LEARN, S.W. GRAHAM, S.C.H. BARRETT, S. DAYANANDAN & V.A. ALBERT. 1993. Phylogenetics of seed plants - an analysis of nucleotide-sequences from the plastid gene *rbcL*. Ann. Missouri Bot. Gard. 80: 528–580.
COOKE, T. 1906. The flora of the presidency of Bombay 2, 3. Taylor & Franchis, London.
CORNER, E.H.J. 1976. The seeds of the dicotyledons. Cambridge University Press, Cambridge.
CRAIB, W.G. 1911. Contributions to the flora of Siam. Bull. Misc. Inf. 1911: 385–474.
CRONQUIST, A. 1988. The evolution and classification of flowering plants. New York Botanical Garden, New York.
CROWE, T. 1994. Morphometrics, phylogenetic models and cladistics: Means to an end or much to do about nothing? Cladistics 10: 77–84.
DALZELL, N.A. 1851. Contributions to the botany of western India. Hooker's J. Bot. Kew Gard. Misc. 3: 229.
DALZELL, N.A. & A. GIBSON. 1861. Bombay flora. Education Society's Press, Byculla.
DARLU, P. & G. LECOINTRE. 2002. When does the incongruence length difference test fail? Mol. Biol. Evol. 19: 432–437.
DAVIES, S.J. 2001. Systematics of *Macaranga* sects. *Pachystemon* and *Pruinosae* (Euphorbiaceae). Harv. Pap. Bot. 6: 371–448.
DAVIES, S.J., S.K.Y. LUM, R. CHAN & L.K. WANG. 2001. Evolution of myrmecophytism in western Malesian *Macaranga* (Euphorbiaceae). Evolution 55: 1542–1559.

DAVIS, C.C. & M.W. CHASE. 2004. Elatinaceae are sister to Malpighiaceae; Peridiscaceae belong to Saxifragales. Amer. J. Bot. 91: 262–273.

DAVIS, C.C., O.W. CAMPBELL, K.J. WURDACK, C.A. JARAMILLO & M.J. DONOGHUE. 2005. Explosive radiation of Malphigiales supports a Mid-Cretaceous origin of modern tropical rain forests. Am. Nat. 165: E36–E65.

DAVIS, C.C., M. LATVIS, D.L. NICKRENT, K.J. WURDACK & D.A. BAUM. 2007. Floral gigantism in Rafflesiaceae. Science 315: 1812–1812.

DAVIS, J.I., D.W. STEVENSON, G. PETERSEN, O. SEBERG, L.M. CAMPBELL, J.V. FREUDENSTEIN, D.H. GOLDMAN, C.R. HARDY, F.A. MICHELANGELI, M.P. SIMMONS, C.D. SPECHT, F. VERGARA-SILVA & M. GANDOLFO. 2004. A phylogeny of the monocots, as inferred from *rbcL* and *atpA* sequence variation, and a comparison of methods for calculating jackknife and bootstrap values. Systematic Botany 29: 467–510.

DAYANANDAN, S., P.S. ASHTON, S.M. WILLIAMS & R.B. PRIMACK. 1999. Phylogeny of the tropical tree family Dipterocarpaceae based on nucleotide sequences of the chloroplast *rbcL* gene. Amer. J. Bot. 86: 1182–1190.

DE GUZMAN, E.D., R.M. UMALI & E.D. SOTALBO. 1986. Guide to Philippine flora and fauna 3. JMC Press, Quezon City.

DE LAET, J., S. FARRIS & P. GOLOBOFF. 2004. Treatment of multiple trees in resampling analyses. Cladistics 20: 590–590.

DE LOUREIRO, J. 1790. Flora Cochinchinensis 1. J de Loureiro, Ulyssipone.

DINERSTEIN, E. & C.M. WEMMER. 1988. Fruits Rhinoceros eat: dispersal of *Trewia nudiflora* (Euphorbiaceae) in lowland Nepal. Ecology 69: 1768–1774.

DOLPHIN, K., R. BELSHAW, C.D.L. ORME & D.L.J. QUICKE. 2000. Noise and incongruence: Interpreting results of the incongruence length difference test. Mol. Phylog. Evol. 17: 401–406.

DOUADY, C.J., F. DELSUC, Y. BOUCHER, W.F. DOOLITTLE & E.J.P. DOUZERY. 2003. Comparison of Bayesian and maximum likelihood bootstrap measures of phylogenetic reliability. Mol. Biol. Evol. 20: 248–254.

EICHHORN, K. 2006. Plant diversity after rain-forest fires in Borneo. Blumea Suppl. 18.

EMSHWILLER, E. & J.J. DOYLE. 1999. Chloroplast-expressed glutamine synthetase (*ncpGS*): Potential utility for phylogenetic studies with an example from *Oxalis* (Oxalidaceae). Mol. Phylog. Evol. 12: 310–319.

EMSHWILLER, E. & J.J. DOYLE. 2002. Origins of domestication and polyploidy in Oca (*Oxalis tuberosa*: Oxalidaceae). 2. Chloroplast-expressed glutamine synthetase data. Amer. J. Bot. 89: 1042–1056.

ENDLICHER, S.L. 1843. Generum plantarum, Suppl. 3. Fr. Beck, Wien.

ESSER, H.-J. 2003. Fruit characters in Malesian Euphorbiaceae. Telopea 10: 169–177.

FARRIS, J.S. 1990. Phenetics in camourlage. Cladistics 6: 91–100.

FARRIS, J.S., M. KALLERSJÖ, A.G. KLUGE & C. BULT. 1994. Testing significance of incongruence. Cladistics 10: 315–319.

FARRIS, J.S., V.A. ALBERT, M. KALLERSJÖ, D. LIPSCOMB & A.G. KLUGE. 1996. Parsimony jackknifing outperforms neighbor-joining. Cladistics 12: 99–124.

FAY, M.F., S.M. SWENSEN & M.W. CHASE. 1997. Taxonomic affinities of *Medusagyne oppositifolia* (Medusagynaceae). Kew Bull. 52: 111–120.

FELSENSTEIN, J. 1985. Confidence-limits on phylogenies - an approach using the bootstrap. Evolution 39: 783–791.

FIALA, B. 1996. Ants benefit pioneer trees: The genus *Macaranga* as an example of ant-plant associations in dipterocarp forest ecosystems. In: E. Schulte & D. Schöne (eds.), Dipterocarp forest ecosystems: towards sustainable management. World Scientific, Singapore.

FIGGE, R.M., M. SCHUBERT, H. BRINKMANN & R. CERFF. 1999. Glyceraldehyde-3-phosphate dehydrogenase gene diversity in Eubacteria and Eukaryotes: Evidence for intra- and inter-kingdom gene transfer. Mol. Biol. Evol. 16: 429–440.

FLYNN, J.J. & M.A. NEDBAL. 1998. Phylogeny of the Carnivora (Mammalia): Congruence vs incompatibility among multiple data sets. Mol. Phylog. Evol. 9: 414–426.

FORSTER, G. 1786. Florulae insularum australium prodromus. J.C. Dieterich, Göttingen.

FORSTER, P.I. 1999. A taxonomic revision of *Mallotus* Lour. (Euphorbiaceae) in Australia. Austrobaileya 5: 457–497.

FREUDENSTEIN, J.V., C. VAN DEN BERG, D.H. GOLDMAN, P.J. KORES, M. MOLVRAY & M.W. CHASE. 2004. An expanded plastid DNA phylogeny of Orchidaceae and analysis of jackknife branch support strategy. Amer. J. Bot. 91: 149–157.

GAGNEPAIN, F. & L. BEILLE. 1925/1926. Euphorbiaceae. In: M.H. Lecomte (ed.), Flore générale de l´Indo-Chine 5: 229–673. Masson & Cie, Paris.

GAMBLE, J.S. 1925. Flora of the presidency of Madras 2, 7. West Newman & Co. & Adlard & Son, London.

GARCIA-CRUZ, J. & V. SOSA. 2006. Coding quantitative character data for phylogenetic analysis: A comparison of five methods. Systematic Botany 32: 302–309.

GARDNER, S., P. SIDISUNTHORN & V. ANUSARNSUNTHORN. 2000. A field guide to forest trees of Northern Thailand. Kobfai Publishing Project, Bangkok.

GOLOBOFF, P.A. 1999a. Analyzing large data sets in reasonable times: solutions for composite optima. Cladistics 15: 415–428.

GOLOBOFF, P.A. 1999b. NONA (NO NAME) ver. 2. Published by the author, Tucumán, Argentina.

GOLOBOFF, P.A. & D. POL. 2005. Parsimony and Bayesian phylogenetics. In: V.A. Albert (ed.), Parsimony, phylogeny and genomics: 145–159. Oxford University Press, Oxford.

GOLOBOFF, P.A., J.S. FARRIS & K.C. NIXON. 2003a. T.N.T. Tree analysis using New Technology. Program and documentation, available at http://www.zmuc.dk/public/phylogeny/tnt.

GOLOBOFF, P.A., C.I. MATTONI & A.S. QUINTEROS. 2006. Continuous characters analyzed as such. Cladistics 22: 589–601.

GOLOBOFF, P.A., J.S. FARRIS, M. KÄLLERSJÖ, B. OXELMAN, M.J. RAMÍREZ & C.A. SZUMIK. 2003b. Improvements to resampling measures of group support. Cladistics 19: 324–332.

GOVAERTS, R., D.G. FRODIN & A. RADCLIFFE-SMITH. 2000. World checklist and bibliography of Euphorbiaceae (and Pandaceae). Royal Botanic Gardens, Kew.

GREEN, P.S. 1986. New combinations in *Baloghia* and *Codiaeum* (Euphorbiaceae). Kew Bull. 41: 1026.

GRIERSON, A.J.C. & D.G. LONG. 1987. Flora of Bhutan 1, 3. Royal Botanic Garden, Edinburgh.

GUINDON, S. & O. GASCUEL. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Syst. Biol. 52: 696–704.

HAEGENS, R.M.A.P. 2000. Taxonomy, phylogeny, and biogeography of *Baccaurea*, *Distichirhops*, and *Nothobaccaurea* (Euphorbiaceae). Blumea Suppl. 12.

HALL, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleid Acids Symp. Ser. 41: 95–98.

HENDERSON, C.P. & I.R. HANCOCK. 1988. A guide to the useful plants of Solomon Islands. Ministry of Agriculture and Lands, Honaira.

HENNIG, W. 1966. Phylogenetic systematics. University of Illinois Press, Urbana.

HEY, J. 2001. The mind of the species problem. Trends Ecol. Evol. 16: 326–329.

HEY, J., R.S. WAPLES, M.L. ARNOLD, R.K. BUTLIN & R.G. HARRISON. 2003. Understanding and confronting species uncertainty in biology and conservation. Trends Ecol. Evol. 18: 597–603.

HILLIS, D.M. & J.J. BULL. 1993. An empirical-test of bootstrapping as a method for assessing confidence in phylogenetic analysis. Syst. Biol. 42: 182–192.

HIPP, A.L., J.C. HALL & K.J. SYTSMA. 2004. Congruence versus phylogenetic accuracy: revisiting the Incongruence Lenght Difference Test. Syst. Biol. 53: 81–89.

HOLDER, M. & P.O. LEWIS. 2003. Phylogeny estimation: traditional and Bayesian approaches. Nat. Rev. Genet. 4: 275–284.

HOLLAND, B. & V. MOULTON. 2003. Consensus networks: a method for visualising incompatibilities in collections of trees. In: G. Benson & R. Page (eds.), Algorithms in bioinformatics, WABI 2003. Springer-Verlag, Berlin, Germany.

HOLLAND, B.R., F. DELSUC & V. MOULTON. 2005. Visualizing conflicting evolutionary hypotheses in large collections of trees: Using consensus networks to study the origins of placentals and hexapods. Syst. Biol. 54: 66–76.

HOOKER, J.D. 1887. Flora of British India 5. L. Reeve & Co, London.

HORIUCHI, J. & H. TAKIMOTO. 2001. Plant mega-fossils from the Late Early to Early Middle Miocene Asawaka formation at Inuboe pass, Ibaraki prefecture, Kanto district, Japan. Bull. Ibaraki Nat. Mus. 4: 1–32.

HUELSENBECK, J.P. & F. RONQUIST. 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17: 754–755.

HUSON, D.H. & D. BRYANT. 2006. Application of phylogenetic networks in evolutionary studies. Mol. Biol. Evol. 23: 254–267.

JANSA, S.A., S.M. GOODMAN & P.K. TUCKER. 1999. Molecular phylogeny and biogeography of the native rodents of Madagascar (Muridae: Nesomyinae): A test of the single-origin hypothesis. Cladistics 15: 253–270.

JUDD, W.S., C.S. CAMPBELL, E.A. KELLOGG & P.F. STEVENS. 1999. Plant systematics: a phylogenetic approach. Sinauer Associates, Sunderland.

KANJILAL, U.N., P.C. KANJILAL, R.N. DE & A. DAS. 1940. Flora of Assam. Prabasi Press, Calcutta.

KATHRIARACHCHI, H., P. HOFFMANN, R. SAMUEL, K.J. WURDACK & M.W. CHASE. 2005. Molecular phylogenetics of Phyllanthaceae inferred from five genes (plastid *atpB*, *matK*, 3 ‘ *ndhF*, *rbcL*, and nuclear *PHYC*). Mol. Phylog. Evol. 36: 112–134.

KATHRIARACHCHI, H., R. SAMUEL, P. HOFFMANN, J. MLINAREC, K.J. WURDACK, H. RALIMANANA, T.F. STUESSY & M.W. CHASE. 2006. Phylogenetics of tribe Phyllantheae (Phyllanthaceae; Euphorbiaceae sensu lato) based on nrITS and plastid *matK* DNA sequence data. Amer. J. Bot. 93: 637–655.

KELCHNER, S.A. 2000. The evolution of non-coding chloroplast DNA and its application in plant systematics. Ann. Missouri Bot. Gard. 87: 482–498.

KEßLER, P.J.A. 2000. Secondary forest trees of Kalimantan, Indonesia. A manual to 300 selected species. MOFEC-Tropenbos-Kalimantan Project, Balikpapan.

KITAMURA, S., S. SUZUKI, T. YOMOTO, P. CHUAILUA, K. PLONGMAI, P. POONSWAD, N. NOMA, T. MARUHASHI & C. SUCKASAM. 2005. A botanical invertory of a tropical seasonal forest in Khao Yai National Park, Thailand: implications for fruit-frugivore interactions. Biodiversity and Conservation 14: 1241–1262.

KITCHING, I.J., P.L. FOREY, C.J. HUMPHRIES & D.M. WILLIAMS. 1998. Cladistics. The theory and practice of parsimony analysis. Oxford University Press, Oxford.

KIU, H.S., S.M. HWANG & Y.T. CHANG. 1996. Euphorbiaceae 2. Flora Reipubl. Popularis Sin. 44, 2. Science Press, Beijing.

KLOTZSCH F. & A. GARCKE. 1862. Die Botanischen Ergebnisse der Reise seiner Königl. Hoheit des Prinzen Waldemar von Preussen. R. Decker, Berlin.

KLUGE, A.G. 1989. A concern for evidence and a phylogenetic hypothesis of relationships among Epicrates (Boidae, Serpenes). Syst. Zool. 38: 7–25.

KULJU, K.K.M & P.C. VAN WELZEN. 2005. Revision of the genus *Cleidion* (Euphorbiaceae) in Malesia. Blumea 50: 197–219.

KULJU, K.K.M. & P.C. VAN WELZEN. Proposal to conserve the name Mallotus against Trewia (Euphorbiaceae s.s.). Submitted to Taxon.

KULJU, K.K.M., S.E.C. SIERRA & P.C. VAN WELZEN. 2007. Re-shaping *Mallotus* [part 2]: Inclusion of *Neotrewia*, *Octospermum* and *Trewia* in *Mallotus* s.s. (Euphorbiaceae s.s.). Blumea 52: 115–136.

KULJU, K.K.M., S.E.C. SIERRA, S.G.A. DRAISMA, R. SAMUEL & P.C. VAN WELZEN. In press. Molecular phylogeny of *Macaranga*, *Mallotus*, and related genera (Euphorbiaceae s.s.): insights from plastid and nuclear DNA sequence data. Amer. J. Bot.

KURZ, S. 1877. Forest flora of British Burma 2. Goverment printing, Calcutta.

LEHTONEN, S. 2006. Phylogenetics of *Echinodorus* (Alismataceae) based on morphological data. Bot. J. Linn. Soc. 150: 291–305.

LINNAEUS, C. 1753. Species plantarum 2. Laurentii Salvii, Stockholm.

MADDISON, D.R. & W.P. MADDISON. 2001. MacClade 4: Analysis of phylogeny and character evolution. Version 4.08. Sinauer Associates, Sunderland, Massachusetts.

MAYDEN, R.I. 1997. A hierarchy of species concepts: the denouement in the saga of the species

problem. In: M.F. Claridge, A.H. Dawah & M.R. Wilson (eds.), Species: the units of biodiversity: 381–424. Chapman & Hall, London.
McLOUGHLIN, S. 2001. The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. Aust. J. Bot. 49: 271–300.
McNEILL, J., F.R. BARRIE, H.M. BURDET, V. DEMOULIN, D.L. HAWKSWORTH, K. MARHOLD, D.H. NICOLSON, J. PRADO, P.C. SILVA, J.E. SKOG, J.H. WIERSEMA & N.J. TURLAND (eds.) 2006. International Code of Botanical Nomenclature (Vienna Code) adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. Regnum Veg. 146.
McPHERSON, G. 1995. On *Mallotus* and *Deuteromallotus* (Euphorbiaceae) in Madagascar. Bull. Mus. Nation. Hist. Nat., B, Adansonia 17: 169–173.
McPHERSON, G. & C. TIREL. 1987. Flore de la Nouvelle-Calédonie et Dépendances 14. Euphorbiacées 1. Muséum National d'Histoire Naturelle, Paris.
MEEUSE, A.D.J. 1990. The Euphorbiaceae auct. plur., an unnatural taxon. Eburon, Delft.
MEIJER, W. 1967. Family Euphorbiaceae. Bot. News Bull. Forest Dept., Sabah 7: 52.
MERRILL, E.D. 1906. The flora of the Lamao forest reserve. Philipp. J. Sci. 1, suppl. 1: 1–141.
MERRILL, E.D. 1912. Notes on Philippine Euphorbiaceae. Philipp. J. Sci., C. Bot. 7: 379–410.
MERRILL, E.D. 1914. Notes on Philippine Euphorbiaceae, II. Philipp. J. Sci., C. Bot. 9: 461–493.
MERRILL, E.D. 1917. An interpretation of Rumphius's Herbarium Amboinense. Bureau of Printing, Manila.
MERRILL, E.D. 1922. New or noteworthy Philippine plants, XVII. Philipp. J. Sci. 20: 367–476.
MERRILL, E.D. 1923. An enumeration of Philippine flowering plants 2. Bureau of Printing, Manila.
MERRILL, E.D. 1926. Additions to our knowledge of the Philippine flora, III. Philipp. J. Sci. 30: 389–430.
MIQUEL, F.A.W. 1859. Flora van Nederlandsch Indië 1, 2. F. Fleischer, Leipzig.
MÜLLER [ARGOVIENSIS], J. 1865. Euphorbiaceae. Linnaea 34: 1–224.
MÜLLER [ARGOVIENSIS], J. 1866. Euphorbiaceae. In: A. de Candolle (ed.), Prodromus systematis universalis regni vegetabilis 15, 2: 189–1260. V. Masson & Filii, Paris.
MÜLLER, K. 2004. PRAP-computation of Bremer support for large data sets. Mol. Phylog. Evol. 31: 780–782.
MÜLLER, K. 2005a. SeqState: primer design and sequence statistics for phylogenetic DNA datasets. Appl. Bioinformatics 4: 65–69.
MÜLLER, K. 2005b. The efficiency of different search strategies in estimating parsimony jackknife, bootstrap, and Bremer support. BMC Evol. Biol. 5.
MÜLLER, K. 2006. Incorporating information from length-mutational events into phylogenetic analysis. Mol. Phylog. Evol. 38: 667–676.
NECKER, N.J. 1790. Elementa botanica 2. Neowedae, Rhenum.
NICOLSON, D.H., C.R. SURESH & K.S. MANILAL. 1988. An interpretation of van Rheede's Hortus Malabaricus. Regnum Veg. 119.
NIXON, K.C. 1999. The Parsimony Ratchet, a new method for rapid parsimony analysis. Cladistics 15: 407–414.
NOWICKE, J.W. & M. TAKAHASHI. 2002. Pollen morphology, exine structure and systematics of Acalyphoideae (Euphorbiaceae), Part 4. Rev. Palaeob. Palyn. 121: 231–336.
NOWICKE, J.W., M. TAKAHASHI & G.L. WEBSTER. 1998. Pollen morphology, exine structure and systematics of Acalyphoideae (Euphorbiaceae), Part 1. Rev. Palaeob. Palyn. 102: 115–152.
NOWICKE, J.W., M. TAKAHASHI & G.L. WEBSTER. 1999. Pollen morphology, exine structure and systematics of Acalyphoideae (Euphorbiaceae), Part 2. Rev. Palaeob. Palyn. 105: 1–62.
NYLANDER, J.A.A. 2004. MrModeltest v2. Program distributed by the author. Evolutionary Biology Center, Uppsala University.
OLSEN, K.M. & B.A. SCHAAL. 1999. Evidence on the origin of cassava: phylogeography of *Maninot esculenta*. Proc. Natl. Acad. Sci. U.S.A. 96: 5586–5591.
PAX, F. 1896. Euphorbiaceae. In: A. Engler & K. Prantl (eds.), Die natürlichen Pflanzenfamilien 3, 5: 1–119. W. Engelmann, Leipzig.
PAX, F. & K. HOFFMANN. 1914. Euphorbiaceae – Acalypheae – Mercurialinae. In: A. Engler (ed.),

Das Pflanzenreich IV.147.vii. W. Engelmann, Leipzig & Berlin.
PAX, F. & K. HOFFMANN. 1919. Euphorbiaceae – Additamentum VI. In: A. Engler (ed.), Das Pflanzenreich IV.147.xiv. Engelmann, Leipzig.
PAX, F. & K. HOFFMANN. 1931. Euphorbiaceae. In: A. Engler & H. Harms (eds.), Die natürlichen Pflanzenfamilien ed. 2, 19C: 11–233. W. Engelmann, Leipzig.
PHILCOX, D. 1997. Euphorbiaceae. In: M.D. Dassanayake (ed.), A revised handbook to the flora of Ceylon. A.A. Balkema, Rotterdam.
PICKETT, K.M. & C.P. RANDLE. 2005. Strange bayes indeed: uniform topological priors imply non-uniform clade priors. Mol. Phylog. Evol. 34: 203–211.
PIMENTEL, R. & R. RIGGINS. 1987. The nature of cladistic data. Cladistics 3: 201–209.
POSADA, D. & T.R. BUCKLEY. 2004. Model selection and model averaging in phylogenetics: Advantages of akaike information criterion and Bayesian approaches over likelihood ratio tests. Syst. Biol. 53: 793–808.
PRAIN, D. 1963. Bengal plants 2. Botanical Survey of India, Calcutta.
PRESL, C. 1845. Botanische Bemerkungen. Abh. Königl. Böhm. Ges. Wiss. 5, 3: 431–584.
PRIMACK, R.B. & H.S. LEE. 1991. Population dynamics of pioneer (Macaranga) trees and understorey (Mallotus) TREES (Euphorbiaceae) in primary and selectively logged Bornean rain forests. J. Trop. Ecol. 7: 439–458.
QUANDT, D., K. MÜLLER & S. HUTTUNEN. 2003. Characterisation of the chloroplast DNA *psbT-H* region and the influence of dyad symmetrical elements on phylogenetic reconstructions. Plant Biol. 5: 400–410.
QUISUMBING, E. 1978. Medicinal plant of the Philippines. Katha Publishing Co., Quezon City.
QUISUMBING, E. & E.D. MERRILL. 1928. New Philippine plants. Philipp. J. Sci. 37: 133–212.
RADCLIFFE-SMITH, A. 1987. Segregate families from the Euphorbiaceae. Bot. J. Linn. Soc. 94: 47–66.
RADCLIFFE-SMITH, A. 1996. Euphorbiaceae. In: G.V. Pope (ed), Flora Zambesiaca 9(4). Royal Botanic Gardens, Kew, UK.
RADCLIFFE-SMITH, A. 2001. Genera Euphorbiacearum. Royal Botanic Gardens, Kew, UK.
RADCLIFFE-SMITH, A. & R.H.A. GOVAERTS. 1997. New names and new combinations in the Euphorbiaceae–Acalyphoïdeae. Kew Bull. 52: 477–481.
RAE, T.C. 1998. The logical basis for the use of continuous characters in phylogenetic systematics. Cladistics 14: 221–228.
RANNALA, B. & Z.H. YANG. 1996. Probability distribution of molecular evolutionary trees: A new method of phylogenetic inference. J. Mol. Evol. 43: 304–311.
RAUBESON, L.A. & R.K. JANSEN. 2005. Chloroplast genomes of plants. In: R.J. Henry (ed.), Diversity en evolution of plants - Genotypic and phenotypic variation in higher plants: 44–68. CABI Publishing, Wallingford.
RAVEN, P.H. & D.I. AXELROD. 1974. Angiosperm biogeography and past continental movements. Ann. Missouri Bot. Gard. 61: 539–673.
RAXWORTHY, C.J., M.R.J. FORSTNER & R.A. NUSSBAUM. 2002. Chameleon radiation by oceanic dispersal. Nature 415: 784–787.
REICHENBACH, H.G. & H. ZOLLINGER. 1856. Rottlera. In: H. Zollinger (ed.), Über die Rottlera-Arten. Linnaea 28: 299–332.
RENNER, S.S. 2004. Multiple Miocene Melastomataceae dispersal between Madagascar, Africa and India. Philos. Trans. R. Soc. Lond., B, Biol. Sci. 359: 1485–1494.
RIDLEY, H.N. 1910. Symbiosis of ants and plants. Ann. Bot. 24: 457–483.
RITTERSHAUSEN, P. 1892. Anatomisch-systematische Untersuchung von Blatt und Axe der Acalypheen. Höfling, München.
RONQUIST, F. & J.P. HUELSENBECK. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574.
ROTH, A.G. 1821. Novae plantarum species praesertim Indiae Orientalis. H. Vogleri, Halberstadii.
SAMUEL, R., H. KATHRIARACHCHI, P. HOFFMANN, M.H.J. BARFUSS, K.J. WURDACK, C.C. DAVIS & M.W. CHASE. 2005. Molecular phylogenetics of Phyllanthaceae: Evidence from plastid *matK* and nuclear *PHYC* sequences. Amer. J. Bot. 92: 132–141.

SAMUEL, R., W. GUTERMANN, T.F. STUESSY, C.F. RUAS, H.W. LACK, K. TREMETSBERGER, S. TALAVERA, B. HERMANOWSKI & F. EHRENDORFER. 2006. Molecular phylogenetics reveals *Leontodon* (Asteraceae, Lactuceae) to be diphyletic. Amer. J. Bot. 93: 1193–1205.

SANG, T., D.J. CRAWFORD & T.F. STUESSY. 1995. Documentation of reticulate evolution in peonies (*Peonia*) using Internal Transcribed Spacer sequences of nuclear ribosomal DNA - Implications for biogeography and concerted evolution. Proc. Natl. Acad. Sci. U.S.A. 92: 6813–6817.

SATO, S. & A. SAKAI. 2005. Birds as dispersers of *Mallotus japonicus* in a coniferous plantation. Jpn. J. Ornithol. 54: 23–28.

SAVOLAINEN, V., M.F. FAY, D.C. ALBACH, A. BACKLUND, M. VAN DER BANK, M.C. CAMERON, S.A. JOHNSSON, M.D. LLEDO, J.-C. PINTAUD, M. POWELL, M.C. SHEAHAN, D.E. SOLTIS, P.S. SOLTIS, P. WESTON, W.M. WHITTEN, K.J. WURDACK & M.W. CHASE. 2000. Phylogeny of eudicots: a nearly complete familiar analysis based on *rbcL* gene sequences. Kew Bull. 55: 257–309.

SCHATZ, G.E. 1996. Malagasy/Indo-Australo-Malesian phytogeographic connections. In: W.R. Lourenco (ed.), Biogeography of Madagascar: 73–83. ORSTOM editions, Paris.

SCHOLS, P., C. D'HONDT, K. GEUTEN, V. MERCKX, S. JANSSENS & E. SMETS. 2004. MorphoCode: coding quantitative data for phylogenetic analysis. Phyloinformatics 4: 1–4.

SCHUMANN, K. 1895. Plantae Bamlerianae. Notizbl. Königl. Bot. Gart. Berlin 2: 44–57.

SCHUMANN, K. & K. LAUTERBACH. 1905. Nachträge zur Flora der Deutschen Schutzgebiete in der Südsee. Gebrüder Borntraeger, Leipzig.

SIERRA, S.E.C. & P.C. VAN WELZEN. 2005. A taxonomic revision of *Mallotus* section *Mallotus* (Euphorbiaceae) in Malesia. Blumea 50: 249–274.

SIERRA, S.E.C. & P.C. VAN WELZEN. 2006. *Mallotus kongkandae* (Euphorbiaceae): New record for China. Blumea 51: 365–366.

SIERRA, S.E.C., P.C. VAN WELZEN & J.W.F. SLIK. 2005. A taxonomic revision of *Mallotus* sections *Philippinenses* (former section *Rottlera* - Euphorbiaceae) in Malesia and Thailand. Blumea 50: 221–248.

SIERRA, S.E.C., M. APARICIO, K.K.M. KULJU, Ž. FIŠER, P.C. VAN WELZEN & R.W.J.M. VAN DER HAM. 2006. Re-shaping *Mallotus* [part 1]: Expanded circumscription and revision of the genus *Cordemoya* (Euphorbiaceae) Blumea 51: 519–540.

SIERRA, S.E.C., M. APARICIO, M.J.H. GEBRAAD, K.K.M. KULJU & P.C. VAN WELZEN. 2007. The morphological range in *Mallotus* (Euphorbiaceae) and a taxonomic revision of its section *Rottleropsis* (including *Axenfeldia*) in Malesia, Thailand and Africa. Blumea 52: 21–113.

SIMMONS, M.P. & H. OCHOTERENA. 2000. Gaps as characters in sequence-based phylogenetic analyses. Syst. Biol. 49: 369–381.

SIMMONS, M.P., K.M. PICKETT & M. MIYA. 2004. How meaningful are Bayesian support values? Mol. Biol. Evol. 21: 188–199.

SLIK, J.W.F. 2005. Assessing tropical lowland forest disturbance using plant morphological and ecological attributes. For. Ecol. Manag. 205: 241–250.

SLIK, J.W.F. & P.C. VAN WELZEN. 2001a. A phylogeny of *Mallotus* (Euphorbiaceae) based on morphology: Indications for a pioneer origin of *Macaranga*. Syst. Bot. 26: 786–796.

SLIK, J.W.F. & P.C. VAN WELZEN. 2001b. A taxonomic revision of *Mallotus* sections *Hancea* and *Stylanthus* (Euphorbiaceae). Blumea 46: 3–66.

SLIK, J.W.F., PRIYONO & P.C. VAN WELZEN. 2000. Key to the *Macaranga* and *Mallotus* species (Euphorbiaceae) of East Kalimantan, Indonesia. Gard. Bull. Singapore 52: 11–87.

SLIK, J.W.F., P.J.A. KEßLER & P.C. VAN WELZEN. 2003a. *Macaranga* and *Mallotus* species (Euphorbiaceae) as indicators for disturbance in the mixed lowland dipterocarp forest of East Kalimantan (Indonesia) Ecol. Ind. 2: 311–324.

SLIK, J.W.F., A.D. POULSEN, P.S. ASHTON, C.H. CANNON, K.A.O. EICHHORN, K. KARTAWINATA, I. LANNIARI, H. NAGAMASU, M. NAKAGAWA, M.G.L. VAN NIEUWSTADT, J. PAYNE, PURWANINGSIH, A. SARIDAN, K. SIDIYASA, R.W. VERBURG, C.O. WEBB & P. WILKIE. 2003b. A floristic analysis of the lowland dipterocarp forests of Borneo. J. Biogeogr. 30: 1517–1531.

SMITH, J.E. 1819. *Trewia*. In: A. Rees (ed.), The cyclopaedia; or universal dictionary of arts, sciences and literature 36. Longman et al., London.

SMITH, J.J. 1910. Bijdrage 12 tot de kennis der boomsoorten op Java. Euphorbiaceae. Meded. Dept.

Landb. Ned.-Indië 10: 9–637.
SOLTIS, D.E. & P.S. SOLTIS. 1998. Choosing an approach and an appropriate gene for phylogenetic analysis. In: D.E. Soltis, P.S. Soltis & J.J. Doyle (eds.), Molecular systematics of plants II. DNA sequencing. Kluwer Academic Publishers, Norwell.
SOLTIS, D.E., P.S. SOLTIS, M.E. MORT, M.W. CHASE, V. SAVOLAINEN, S.B. HOOT & C.M. MORTON. 1998. Inferring complex phylogenies using parsimony: an empirical approach using three large DNA data sets for angiosperms. Syst. Biol. 47: 32–42.
SOLTIS, D.E., A.E. SENTERS, M.J. ZANIS, S. KIM, J.D. THOMPSON, P.S. SOLTIS, L.P.R. DE CRAENE, P.K. ENDRESS & J.S. FARRIS. 2003. Gunnerales are sister to other core eudicots: Implications for the evolution of pentamery. Amer. J. Bot. 90: 461–470.
SOLTIS, D.E., P.S. SOLTIS, M.W. CHASE, M.E. MORT, D.C. ALBACH, M. ZANIS, V. SAVOLAINEN, W.H. HAHN, S.B. HOOT, M.F. FAY, M. AXTELL, S.M. SWENSEN, L.M. PRINCE, W.J. KRESS, K.C. NIXON & J.S. FARRIS. 2000. Angiosperm phylogeny inferred from 18S rDNA, *rbcL*, and *atpB* sequences. Bot. J. Linn. Soc. 133: 381–461.
SORENG, R.J. & J. DAVIS, I. 1998. Phylogenetics and character evolution in the grass family (Poaceae): Simultaneous analysis of morphological and chloroplast DNA restriction site character sets. Botanical Review 64: 1–85.
SPRENGEL, C. 1826. Systema vegatabilium 3. J.C. Dieterich, Göttingen.
STEUDEL, E.T. 1841. Nomenclator botanicus 2, 2. J.G. Cottae, Stuttgart & Tübingen.
STEVENS, P.F. 1991. Character states, morphological variation, and phylogentic analysis: a review. Systematic Botany 16: 533–583.
STOKES, J. 1812. A botanical materia medica 4. J. Johnson & Co., London.
STRAND, A.E., J. LEEBENS-MACK & B.G. MILLIGAN. 1997. Nuclear DNA-based markers for plant evolutionary biology. Mol. Ecol. 6: 113–118.
SULLIVAN, J. 1996. Combining data with different distributions of among-site rate variation. Syst. Biol. 45: 375–380.
SUSILA RANI, S.R.M. & N.P. BALAKRISHNAN. 1998. Notes on *Trewia nudiflora* L. (Euphorbiaceae) and its varieties. J. Econ. Tax. Bot. 22: 345–352.
SUZUKI, Y., G.V. GLAZKO & M. NEI. 2002. Overcredibility of molecular phylogenies obtained by Bayesian phylogenetics. Proc. Natl. Acad. Sci. U.S.A. 99: 16138–16143.
SWART J.J. 1942. A monograph of the genus *Protium* and some allied genera (Burseraceae). Rec. Trav. Bot. Néerl. 39: 211–446.
SWOFFORD, D.L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.
SWOFFORD, D.L., G.J. OLSEN, P.J. WADDELL & D.M. HILLIS. 1996. Phylogenetic Inference. In: D.M. Hillis, C. Moritz & B. Mable (eds.), Molecular Systematics: 407–514. Sinauer Associates, Sunderland, Massachusetts.
TABERLET, P., L. GIELLY, G. PAUTOU & J. BOUVET. 1991. Universal primers for amplification of 3 noncoding regions of chloroplast DNA. Plant Mol. Biol. 17: 1105–1109.
TAKAHASHI, M., J.W. NOWICKE, G.L. WEBSTER, S.S. ORLI & S. YANKOWSKI. 2000. Pollen morphology, exine structure, and systematics of Acalyphoideae (Euphorbiaceae), Part 3. Rev. Palaeob. Palyn. 110: 1–66.
TALBOT, W.A. 1911. Forest flora of the Bombay presidency and Sind 2. Government Photozincographic Department, Poona.
THIELE, K. 1993. The holy grail of the perfect character: the cladistic treatment of morphometric data. Cladistics 9: 275–304.
THOMPSON, J.D., D.G. HIGGINS & T.J. GIBSON. 1994. Clustal-W - Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res. 22: 4673–4680.
TOKUOKA, T. & H. TOBE. 2006. Phylogenetic analyses of Malpighiales using plastid and nuclear DNA sequences, with particular reference to the embryology of Euphorbiaceae sens. str. J. Plant. Res. 119: 599–616.
TOKUOKA, T. 2007. Molecular phylogenetic analysis of Euphorbiaceae sensu stricto based on plastid and nuclear DNA sequences and ovule and seed character evolution. J. Plant Res.

TROUP, R.S. 1921. The silviculture of Indian trees 3. Clarendon Press, Oxford.
TURCZANINOW, N. 1843. Decas generum plantarum hucusque non descriptorum. Bull. Soc. Imp. Nat. Moscou 16: 51–62.
VAN DE VEN, E.A. & R.W.J.M. VAN DER HAM. 2006. Pollen of *Melodinus* (Apocynaceae): Monads and tetrads. Grana 45: 1–8.
VAN RHEEDE, H. 1678. Hortus Indicus Malabaricus 1. J. van Someren & J. van Dyck, Amsterdam.
VAN STEENIS, C.G.G.J. 1957. Specific and infraspecific delimitation, Flora Malesiana I, Vol. 5: clxvii–ccxxxiv.
VAN WELZEN, P.C. 1998. Revisions and phylogenies of Malesian Euphorbiaceae: Subtribe Lasiococcinae (*Homonoia*, *Lasiococca*, *Spathiostemon*) and *Clonostylis*, *Ricinus*, and *Wetria*. Blumea 43: 131–164.
VAN WELZEN, P.C. & K. CHAYAMARIT. 2001. Two new *Mallotus* and two new *Sauropus* species (Euphorbiaceae) endemic to Thailand. Kew Bull. 56: 649–656.
VAN WELZEN, P.C. & K. PHATTARAHIRANKANOK. 2001. *Mallotus kongkandae* (Euphorbiaceae), a new species from Thailand. Blumea 46: 67–69.
VAN WELZEN, P.C. & S.E.C. SIERRA. 2006. The *Mallotus wrayi* complex (Euphorbiaceae). Blumea 51: 373–388.
VAN WELZEN, P.C., R.W.J.M. VAN DER HAM & K.K.M. KULJU. 2004. *Mallotus glomerulatus* (Euphorbiaceae), a new species: its description, pollen, and phylogenetic position. Thai Forest Bulletin 32: 173–178.
VAN WELZEN, P.C., S.E.C. SIERRA, J.W.F. SLIK & S.M. BOLLENDORFF. In press. *Mallotus*. In: P.C. Van Welzen & K. Chayamarit (eds.), Flora of Thailand 8(2). The Forest Herbarium, Bangkok.
VAN WELZEN, P.C., S.E.C. SIERRA, M.J.H. GEBRAAD & K.K.M. KULJU. 2006. The distinction of five *Mallotus* species formerly in section *Hancea* (Euphorbiaceae). Blumea 51: 367–372.
VIDAL Y SOLER, S. 1886. Revision de plantas vasculares Filipinas. M. Perez, Manila.
VU VAN DUNG (ed.). 1996. Vietnam forest trees. Agricultural Publishing House, Hanoi.
WATT, G. 1893. A dictionary of the economic products of India 6, 4. W.H. Allen & Co., London.
WEBSTER, G.L. 1975. Conspectus of a new classification of the Euphorbiaceae. Taxon 24: 593–601.
WEBSTER, G.L. 1987. The sage of the spurges: a review of classification and relationships in the Euphorbiales. Bot. J. Linn. Soc. 94: 3–46.
WEBSTER, G.L. 1994a. Classification of the Euphorbiaceae. Ann. Missouri Bot. Gard. 81: 3–32.
WEBSTER, G.L. 1994b. Synopsis of the genera and suprageneric taxa of Euphorbiaceae. Ann. Missouri Bot. Gard. 81: 33–144.
WENDEL, J.F. & J.J. DOYLE. 1998. Phylogenetic incongruence: window into genome history and molecular evolution. In: D.E. Soltis, P.S. Soltis & J.J. Doyle (eds.), Molecular systematics of plants II: DNA sequencing. Kluwer Academic Publishers, Boston.
WHITE, T.J., T. BRUNS, S. LEE & J. TAYLOR. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: M.A. Innis, D.H. Gelfand, J.J. Sninsky & T.J. White (eds.), PCR Protocols. A Guide to Methods and Applications.: 315–322. Academic Press, San Diego.
WHITMORE, T.C. 1965. Studies in *Macaranga*, an easy genus of Malayan wayside trees. Malayan Nat. J. 20: 89–99.
WHITMORE, T.C. 1966. Guide to the forests of the British Solomon Islands. Oxford University Press, London.
WHITMORE, T.C. 1969. Studies in *Macaranga*, III. First thoughts on species evolution in Malayan *Macaranga*. Biol. J. Linn. Soc. Lond. 1: 223–231.
WHITMORE, T.C. 1973. Tree Flora of Malaya 2. Longman, London.
WHITMORE, T.C. 1980. Studies in *Macaranga*, VIII. *Macaranga* in New Guinea and the Bismarck Archipelago. Kew Bull. 34: 599–606.
WHITMORE, T.C. In press. Prodromus of *Macaranga* (Euphorbiaceae) [edited by M.J.E. Coode and P. Hoffmann]. Royal Botanic Gardens, Kew, UK.
WIENS, J.J. 1998. Combining data sets with different phylogenetic histories. Syst. Biol. 47: 568–581.

WIENS, J.J. 2001. Character analysis in morphological phylogenetics: Problems and solutions. Syst. Biol. 50: 688–699.

WIENS, J.J. 2003. Missing data, incomplete taxa, and phylogenetic accuracy. Syst. Biol. 52: 528–538.

WIERINGA, J.J. 1999. Monopetalanthus exit. A systematic study of *Aphanocalyx*, *Bikinia*, *Icuria*, *Michelsonia* and *Tetraberlinia* (Leguminosae, Caesalpinioideae). Wageningen Agricultural University Papers 99: 1–320.

WIGHT, R. 1852. Icones plantarum Indiae Orientalis. J.B. Pharoah, Madras.

WILLDENOW, C.D. 1797. *Schradera* et *Rottlera*, bina nova plantarum genera. Gött. J. Naturwiss. 1: 1–9.

WILLDENOW, C.D. 1806. Species Plantarum ed. 4, 2. G.C. Nauk, Berlin.

WOLFE, K.H., P.M. SHARP & W.H. LI. 1989. Rates of Synonymous Substitution in Plant Nuclear Genes. J. Mol. Evol. 29: 208–211.

WOLFE, K.H., W.H. LI & P.M. SHARP. 1987. Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. Proc. Natl. Acad. Sci. U.S.A. 84: 9054–9058.

WURDACK, K.J. 2002. Molecular systematics and evolution of Euphorbiaceae sensu lato. Ph.D. dissertation, University of North Carolina, Chapel Hill, North Carolina.

WURDACK, K.J., P. HOFFMANN & M.W. CHASE. 2005. Molecular phylogenetic analysis of uniovulate Euphorbiaceae (Euphorbiaceae sensu stricto) using plastid *rbcL* and *trnL-F* DNA sequences. Amer. J. Bot. 92: 1397–1420.

WURDACK, K.J., P. HOFFMANN, R. SAMUEL, A. DE BRUIJN, M. VAN DER BANK & M.W. CHASE. 2004. Molecular phylogenetic analysis of Phyllanthaceae (Phyllanthoideae Pro Parte, Euphorbiaceae sensu lato) using plastid *rbcL* DNA sequences. Amer. J. Bot. 91: 1882–1900.

YODER, A.D., J.A. IRWIN & B.A. PAYSEUR. 2001. Failure of the ILD to determine data combinability for slow loris phylogeny. Syst. Biol. 50: 408–424.

YUAN, Y.-M., S. WOHLHAUSER, M. MÖLLER, J. KLACKENBERG, M.W. CALLMANDER & P. KÜPFER. 2005. Phylogeny and biogeography of *Exacum* (Gentianaceae): A disjunctive distribution in the Indian Ocean basin resulted from long-distance dispersal and extentive radiation. Syst. Biol. 54: 21–34.

ZOHARY, M. 1973. Geobotanical foundations of the Middle East. Gustav Fischer Verlag, Stuttgart.

ZOLLINGER, H. 1856. Über die *Rottlera*-Arten. Linnaea 28: 299–332.