



**Universiteit
Leiden**
The Netherlands

Spontaneous breathing and respiratory support of preterm infants at birth

Pas, A.B. de

Citation

Pas, A. B. de. (2009, March 5). *Spontaneous breathing and respiratory support of preterm infants at birth*. Retrieved from <https://hdl.handle.net/1887/13595>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13595>

Note: To cite this publication please use the final published version (if applicable).

Establishing functional residual capacity at birth: the effect of sustained inflation and positive end expiratory pressure in a preterm rabbit model

Arjan B te Pas^{*1,2}, Melissa Siew³, Megan J Wallace³, Marcus J Kitchen⁴, Andreas Fouras⁶, Robert A Lewis⁵, Naoto Yagi⁷, Kentaro Uesugi⁷, Susan Donath⁸, Peter G Davis¹, Colin J Morley¹, Stuart B Hooper³

¹ Neonatal Services, Royal Women's Hospital, Melbourne, Victoria 3052, Australia

² Division of Neonatology, Department of Pediatrics, Leiden University Medical Center, Leiden, 2300 RC, the Netherlands

³ Department of Physiology, Monash University, Melbourne, Victoria 3800, Australia

⁴ School of Physics, Monash University, Melbourne, Victoria 3800, Australia

⁵ Monash Centre for Synchrotron Science, Monash University, Melbourne, Victoria 3800, Australia

⁶ Division of Biological Engineering, Monash University, Melbourne, Victoria 3800, Australia

⁷ SPring-8/JASRI, Sayo, Hyogo 679-5198, Japan

⁸ Murdoch Children's Research Institute, Melbourne, Victoria 3052, Australia

Abstract

The effect of a 20 second sustained inflation (SI) and positive end-expiratory pressure (PEEP) on functional residual capacity (FRC) formation at birth were investigated. Preterm rabbit pups (28 days) were randomized at birth into 4 groups (n=6 for each): 1) SI, PEEP 5cmH₂O, 2) no SI, PEEP 5cmH₂O, 3) no SI, no PEEP, 4) SI, no PEEP. FRC and tidal volume (Vt) were measured by plethysmography and uniformity of lung aeration by phase contrast X-ray imaging. Ventilation with a SI and PEEP uniformly aerated the lung and Vt and FRC were recruited by the first tidal inflation. Ventilation without a SI, with PEEP, gradually recruited Vt and FRC with each inflation but aeration was not uniform. Ventilation without a SI or PEEP, gradually recruited Vt, but no FRC. Ventilation with a SI, without PEEP, uniformly aerated the lung and recruited Vt but no FRC. FRC was greater with SI (p=0.006) during the first minute, but was larger with PEEP than without PEEP throughout the first 7 minutes (p<0.0005). Effects of PEEP and SI were additive. In ventilated preterm rabbits at birth, combining a SI and PEEP improved FRC formation and uniformity of lung aeration, but PEEP had the greatest influence.

Introduction

There is currently little specific evidence about the appropriate respiratory support for very preterm infants immediately after birth (1). Ventilatory strategies in the delivery room may play an important role in the development or prevention of lung injury (2-6) as inappropriate ventilatory support during resuscitation can cause lung damage (2, 3, 7-11).

Adequate gas exchange requires the rapid creation and maintenance of a functional residual capacity (FRC). Many preterm infants need positive pressure ventilation to aerate their lungs and overcome the high surface tension and frictional forces caused by liquid-filled airways. Initiating respiratory support with a sustained inflation (SI) may improve FRC development by allowing an appropriate time-constant for the air/liquid interface to move into the distal airways. Although clinical studies have shown lower intubation rates when a SI is given (4, 6, 12), there is only one study evaluating the physiology of a SI in establishing an FRC in humans (13).

Once the lung is aerated and airway liquid is cleared, the low compliance of the immature lung results in a very short expiratory time constant and the lung easily loses FRC. However, the creation and maintenance of an FRC during resuscitation of preterm infants is assisted by a positive end-expiratory pressure (PEEP) which reduces distal airway collapse at end expiration (14, 15). While PEEP is always used during ventilation in the neonatal unit, international resuscitation guidelines provide few recommendations on the use of PEEP immediately after birth (1, 16).

We have used simultaneous phase contrast X-ray imaging and plethysmography to examine the effect of PEEP and a 20 sec SI on the spatial pattern of lung aeration and FRC recruitment in a preterm rabbit model. We hypothesized that the combination of a 20 sec SI followed by ventilation with PEEP would produce a larger FRC than either a 20 sec SI or PEEP alone or neither PEEP nor a SI.

Methods

Animal experiments

All animal procedures were approved by the SPring-8 Animal Care and Monash University's School of Biomedical Science's Animal Ethics Committees. Studies were conducted in experimental hutch 3 of beam line 20B2, in the Biomedical Imaging Centre at the SPring-8 synchrotron in Japan.

Experiments were performed on 6 litters of New Zealand white rabbits using 35 pups delivered at 28 days gestation (term 32 days; 28 days correlates approximately with human

infants of 26-28 weeks of gestation). The does were anesthetized (Propofol; 12 mg/kg bolus, 100 mg/kg/h i.v. infusion, then isoflurane 1-2%), intubated and ventilated. The pups were sequentially delivered by hysterotomy, leaving the fetal membranes over the nose and mouth to prevent air entering the lungs. The pups were sedated (Nembutal 0.1 mg intraperitoneal), a topical anesthetic (5% xylocaine) was applied and an endotracheal tube (18 gauge plastic catheter) inserted via tracheostomy; the endotracheal tube was blocked to prevent air from entering the lung before imaging commenced. The fetal membranes were then removed, the umbilical cord ligated and cut and the pup was placed in a water-filled plethysmograph (see below). The pups were then ventilated for 7 minutes while they were simultaneously imaged. After the experiment, pups were killed by anesthetic over-dose.

Plethysmography

Lung gas volume changes were measured using water-based plethysmography (17). The pup was placed in the plethysmograph (head out), which consisted of a sealed water-filled (37°C) perspex chamber open to the atmosphere via a water column. The pup's head was located outside the chamber and a rubber diaphragm formed a water-tight seal around the pup's neck. The increase in lung volume caused by ventilation displaced water from the chamber into the water column and the resulting increase in pressure was recorded electronically using a computerised data acquisition system (Powerlab, ADInstruments Sydney, Australia). The plethysmograph was calibrated before each experiment by measuring the pressure change caused by injecting a known volume of water (1 mL) into the sealed chamber. We recorded FRC, initial volume, tidal volume (V_t), airway pressure and gas flow.

Mechanical ventilation

A purpose-built small animal ventilator was used to ventilate pups with air in a time cycled, pressure limited mode (positive pressure ventilation, PPV). Ventilation started with a PIP of 35 cmH₂O and an inspiration time of 1 or 20 seconds, depending on the group (see below). After the first inflation, pups were ventilated with a fixed inspiration and expiration time (rate 24/min, 1:1.5 inspiration: expiration ratio), starting with a peak inspiratory pressure of 35 cm H₂O which was then adjusted (increased or decreased by 2 cmH₂O per minute) to generate a tidal volume of about 10 ml/kg, PEEP was set at either 0 or 5 cm H₂O.

Protocol

The pups were randomly allocated to one of the four ventilation groups:

Group 1: Sustained inflation for 20 s followed by PPV with PEEP 5 cm H₂O (20 SI + PEEP).

Group 2: PPV with PEEP 5 cm H₂O (No SI + PEEP).

Group 3: PPV with PEEP 0 cm H₂O (No SI + no PEEP).

Group 4: Sustained inflation for 20 s followed by PPV with PEEP 0 cm H₂O (20 SI + no PEEP).

Phase contrast X-ray imaging

Phase contrast X-ray imaging was used to image the effect of a SI and PEEP on the rate and spatial pattern of lung aeration. Details describing the imaging procedures have been reported previously (18, 19). Briefly, the X-ray energy used was 24 keV, the pups were located 2.0 m upstream of the detector (Hamamatsu, C4742-95HR) and a short exposure time (50 msec) and a relatively long inflation time (1 sec) were used to minimise motion blur. Image acquisition was synchronised with, and triggered by, the ventilator (onset of inflation) which triggered a sequence of images at 300 msec intervals. A pre-object shutter prevented radiation exposure between image acquisitions and the timing of each image acquisition was recorded digitally in unison with the plethysmograph recording.

The uniformity of lung aeration was measured by dividing the images into quadrants and measuring the air volume in each quadrant (at end-expiration or during the SI) using a phase retrieval analysis (17). This analysis has recently been validated (17) and uses extra-thoracic landmarks to track pup movement between consecutive images to ensure that the relative amount of lung in each quadrant does not change over time. However, as the amount of lung (thus potential air volume) varies between quadrants, the air volume of each quadrant at each time point was expressed as a percentage of the maximum volume achieved in that quadrant following complete lung aeration. The uniformity of lung aeration for each pup was then measured by determining the coefficient of variation in relative air volume between all four quadrants.

Analysis

FRC was defined as the volume of air remaining in the lung at end expiration. V_t recruitment refers to the gradual increase in V_t with time for a given pressure gradient.

At 20 s, 1, 3 and 7 min we calculated the average FRC over four consecutive inflations. For each group we used the area under the FRC time curve (FRC–time integral) to calculate the time-related increase in FRC between 0 and 1 minute and during the total ventilation period (0 - 7 min). FRC at start of ventilation was assumed to be zero. One minute was used as this is a critical time point during neonatal resuscitation when decisions are made about modes of respiratory support (1). To compare the increase in FRC-time integral (area under curve) between groups in the two time periods and to see which variable (SI or PEEP) influenced FRC most, or if there was interaction between the two variables, we performed a linear regression analysis and adjusted for the possible clustering effect of several pups having the same mother (24 pups, 6 different mothers). Significance was defined as $p < 0.05$. Statistical analyses were performed using Stata software (Stata 10 for Windows).

Results

Animal data

Twenty-seven of 35 pups were delivered alive and studied, of which 3 were excluded due to equipment malfunctions (n=2) and endotracheal tube obstruction (n=1). None of the pups developed a pneumothorax, as determined by observation of the images. The average (SD) weight was: Group 1 (20 SI + PEEP) 37.9 ± 2.3 g, group 2 (no SI + PEEP) 34.4 ± 2.4 g, group 3 (no SI + no PEEP) 37.3 ± 1.5 g, group 4 (20 SI + no PEEP) 39.9 ± 11.3 g. Each group contained 6 pups.

The lung gas volume changes with ventilation onset were very different between groups (figure 1). In group 1 (20 SI + PEEP), lung gas volumes rapidly increased and, although




Figure 1. Representative examples of recordings from each group. Change in lung gas volume from birth in anaesthetised ventilated preterm rabbit pups using plethysmography. With PEEP (top two panels) an end-expiratory gas volume (FRC) was rapidly formed whereas in the absence of PEEP a significant FRC was not formed (bottom two panels).

A = SI + PEEP, B = no SI + PEEP, C = no SI, no PEEP, D = SI, no PEEP.

different regions aerated at different rates, by the end of the SI the lung was uniformly aerated (figures 1, 2 & 3). Indeed, the coefficient of variation in relative air volumes across the lung (figure 3) decreased from 27.9 (12.7) % at 2.5 secs to 0.7 (0.6) % at 20 secs. As a result, following a SI (figures 1a and 1d), the Vt did not increase with time and the Vt was fully recruited (figure 4) and neither the Vt nor FRC increased further with subsequent ventilation (table 1; figures 1 and 4).

In group 2 (no SI + PEEP), the Vt gradually increased with each inflation (figure 4) as did the FRC (figure 1, table 1), although the FRC (table 1) and uniformity of lung aeration (figures 2 & 3) was markedly less than in group 1 (20 SI + PEEP) at 20 secs. Although the coefficient of variation in relative air volumes across the lung (figure 3) decreased from 63.0 (38.3) % at 2.5 secs to 27.6 (12.6) % at 20 secs in group 2 it remained significantly greater than in group 1 pups throughout this period.

In group 3 (no SI + no PEEP), the Vt gradually increased as in group 2 (figure 4), but a significant FRC failed to develop (figures 1-4, table 1).

In group 4 (20 SI + no PEEP), the lung gas volume rapidly increased during the SI as in group 1. However, after the SI the lungs were unable to retain an FRC (figures 1, 2, table 1), although the tidal volume was fully recruited (figure 4).

FRC time-integral: Zero to one minute

The FRC time-integral from 0 – 1 minute after birth (figure 5a) was larger when a 20 s SI was used compared to no SI (β coefficient 108; $p = 0.006$). Compared to when no PEEP was used, the increase in the FRC-time integral was always larger when PEEP was used (β coefficient 413; $p = 0.003$). There was no interaction between PEEP and the 20 sec SI (p value for interaction = 0.5), indicating that the effects of PEEP and SI on the FRC-time integral were additive (figure 5a).

FRC time-integral: Zero to seven minutes

The FRC time-integral from 0 – 7 minutes after birth (figure 5b) tended to be larger when a 20 s SI was used compared to no SI (β coefficient 1120; $p = 0.06$), although the differ-

Table 1. Mean (SD) FRC in time of the groups. (FRC= Functional residual capacity)

Group	FRC (ml/kg)			
	20 seconds	1 minute	3 minutes	7 minutes
20 SI + PEEP	10.8 (3.5)	10.8 (3.3)	11.2 (3.5)	11.7 (3.4)
No SI + PEEP	7.5 (5.7)	9.3 (6.1)	8.8 (6.6)	8.4 (7.3)
no SI - no PEEP	0.9 (1.0)	-0.3 (1.4)	-1.2 (2.9)	-0.9 (2.5)
20 SI - no PEEP	2.5 (2.3)	1.5 (2.1)	2.2 (1.9)	2.3 (1.3)




Figure 2. Representative examples of phase contrast X-ray images from each group. Images were acquired at end-expiration (FRC) 20 s after birth in anaesthetised preterm rabbit pups in which an initial sustained inflation (SI) of 20 sec or not was given and ventilated in the presence/absence of positive end-expiratory pressure (PEEP) (5 cmH₂O); A = SI +PEEP, B = no SI +PEEP, C = no SI, no PEEP, D = SI, no PEEP.




Figure 3. Uniformity of lung aeration was assessed by measuring the coefficient of variation in air volume across 4 different regions of the lung. Pups were ventilated either with a 20 second sustained inflation (SI; Group 1, open circles) or without a SI but with 5cmH₂O positive end-expiratory pressure (PEEP) (Group 2; closed circles); measurements were made at either end-expiration (no SI) or during the SI. Aeration across the lung was significantly more variable (less uniform) without a SI, compared with ventilation with a SI. After 15 sec of a SI, the lung was uniformly aerated as indicated by little variability in relative air volume between lung regions. Asterisks indicate values that are significantly different between groups for the same time point. Values are mean \pm SEM.




Figure 4. Effect of a 20 s sustained inflation (SI) and 5 cm H₂O PEEP on tidal volume recruitment after birth. In absence of a SI, tidal volume was recruited gradually with each subsequent inflation, whereas a 20 s SI fully recruited tidal volume. Values are mean \pm SEM.

—●— SI+PEEP, —○— no SI+PEEP, —▼— no SI & no PEEP, —△— SI & no PEEP

ence was not quite statistically significant. Compared to no PEEP, the FRC-time integral was much larger when PEEP was used (β coefficient 3808; $p < 0.0005$). There was no interaction between PEEP and the 20 sec SI on the FRC-time integral (p value for interaction = 0.8), again indicating that the effects of PEEP and SI were additive (figure 5b).




Figure 5a, b. FRC area under curve (median (IQR)) in anaesthetised ventilated preterm rabbit pups in the presence/absence of 5 cm H₂O PEEP, with/without a 20 s sustained inflation (SI) from 0 - 1 minute (a) and 0-7 minutes (b). A = SI + PEEP, B = no SI + PEEP, C = no SI, no PEEP, D = SI, no PEEP.

Discussion

This study has shown that with sedated, ventilated, preterm rabbit pups, in the first 7 minutes after birth, a larger FRC was created and maintained when a 20 s SI was followed by ventilation with 5 cm H₂O PEEP compared with either no SI or no PEEP. PEEP had a larger effect on the formation of FRC than a SI inflation. However, the effects of SI and PEEP were additive, making it possible to create and maintain a FRC as early as the first inflation, if it is sustained. In absence of PEEP, the lungs failed to retain an FRC, irrespective of whether a SI was used, although a SI produced uniform lung aeration before onset of tidal ventilation. This emphasises the importance of using PEEP during ventilation at birth in very preterm infants and indicates that a combination of the two techniques (PEEP + SI) may be better, particularly in infants with very immature lungs. Indeed, a SI can promote uniform lung aeration from the first inflation whereas the addition of PEEP helps to maintain FRC in the lung.

The FRC levels required by preterm infants to establish adequate gas exchange at birth are unclear. However, the FRC we measured in preterm rabbit pups exposed to a 20 s SI with PEEP was comparable to the average FRC measured at birth in spontaneously breathing term rabbit pups (10-15 ml/kg) (20). Vyas and Milner measured FRCs of 5-6 ml/kg in the first few breaths of spontaneously breathing term infants (21-23). A similar FRC was measured in asphyxiated infants during resuscitation with a 5 sec inflation (13) whereas no FRC was created without an initial long inflation (24). In both studies no PEEP was used which may have affected the development of FRC.

After birth, uniform lung aeration and the retention of an appropriate FRC are required to facilitate effective gas exchange and improve an infant's oxygenation and heart rate. We have shown that a combined 20 s SI and PEEP produced a larger FRC during the first minutes after birth. This could explain why intubation rates in very preterm infants are lower in the delivery room when a SI was given, followed by continuous positive airway pressure (4, 6, 12).

Healthy, term infants rapidly create an FRC during their first breaths by exerting large intra-thoracic sub-atmospheric pressures (25, 26). However, preterm infants often have problems clearing lung liquid and need help to overcome high frictional forces caused by airway liquid. As a result, either a large pressure gradient (inflating pressure) or a greater duration of inflation is required to move the air/liquid interface into the distal airways. For any given inflating pressure, if the duration of inflation is too short, the air/liquid interface will not reach the gas exchange regions. As application of high airway pressures may be injurious, a long initial inflation may be preferential, allowing time for air to enter alveoli (13). Indeed, the first breaths of spontaneously breathing infants are characterized by a long deep breath followed by a prolonged expiratory phase (27, 28). The phase contrast X-ray imaging analysis

clearly demonstrated that, at the end of a 20 sec SI, the lung was uniformly aerated, with very little variation (<1%; figure 3) in relative air volumes across the lung. As a result, when tidal ventilation commenced, ventilation appeared to occur uniformly throughout the lung. In the absence of a SI, lung aeration was gradual and regionalized, restricting tidal ventilation to small regions of the lung initially. This is likely to be injurious, particularly if high pressures and short inspiratory times are used, as these regions will be the most compliant and prone to over-expansion.

Following lung liquid clearance, a surfactant deficient immature lung has a low compliance and tends to collapse at end-expiration when PEEP is not used. We found that, although the affects of a SI and PEEP were additive, PEEP had a much greater influence on FRC formation than a SI. Indeed, independent of whether a SI was given or not, no FRC was created if PEEP was not used. This is consistent with previous data suggesting that, during resuscitation at birth, PEEP is necessary in immature lungs to establish and maintain FRC (14, 15) and improve oxygenation (29, 30). No randomized trials have evaluated the role of PEEP or no PEEP during resuscitation in the delivery room (31, 32).

In some pups ventilated without PEEP, the FRC decreased slightly below zero (table 1), which occurred after an initial transient increase. Although the reason for this is unclear, it was not caused by lung aeration before the onset of ventilation and is likely to be an experimental artefact. Although the weight of water on the chest wall may have slightly increased deflation forces on the lung, particularly in the absence of PEEP, it can not explain a decrease in FRC below zero in these pups. Another possible explanation is a small upward displacement of the pups within the plethysmograph following ventilation onset, although this could not be confirmed from the images.

There was much individual variation in FRC within a group, especially in pups receiving PEEP but no SI (Group 2; table 1); this is also reflected by the high variability in relative air volumes across the lung in this group (figure 3). The variability between pups may be caused by the natural variation in maturity at 28 days which is also influenced by the time of day the experiments were done. It is interesting that the variability in FRC was significantly lower in the SI group compared with no SI (coefficient of variation = 30.8 (1.2) % vs 75.9 (7.5) %; $p < 0.001$), despite similar levels of expected variability in pup maturation (table 1). This is most likely due to the greater uniformity and degree of lung aeration achieved before the onset of tidal ventilation (figure 3). It is possible that a 20 sec SI at 35 cm H₂O was enough to overcome all the different factors that contribute to the time constant for lung aeration, leading to less variation in FRC. This indicates that a SI may be able to achieve a consistently higher FRC when used with PEEP and that a SI may have major benefits in very immature lungs, but is of less benefit in more mature lungs. Indeed, we observed this difference in two very small (~25-30g) pups with very stiff lungs. The pup ventilated with PEEP alone, failed to

increase its FRC to a level seen in larger pups (see movie SIPEEP-3 on disc). However, ventilation with a SI + PEEP in the other pup caused aeration of the apical lobes before the onset of tidal ventilation), resulting in the recruitment of a larger FRC. (movie SIPEEP-4 on disc).

Although our model is not directly applicable to preterm infants, our study was designed to specifically assess the affects of SI and PEEP on the formation of FRC in preterm lungs from birth. Preterm pups were all born by caesarean section so that we could image and record the first inflation and antenatal steroids were not administered to maximise differences between groups. Spontaneous breathing was inhibited to assess the affects of PEEP and SI. Pups were ventilated in an upright position in water to simultaneously image and measure lung gas volumes. Although it would not be possible to perform such experiments in very preterm infants, we would expect similar results. Indeed, similar studies were done with preterm rabbits to initially demonstrate the efficacy of surfactant treatment (33, 34).

The inflating pressure (35 cm H₂O) and length of SI used were based on earlier observations in preterm pups. As phase contrast X-ray imaging is ideal for detecting pneumothoraces, we can confirm that no pups developed pneumothoraces in the first 7 minutes. The plethysmography data also demonstrated that a SI did not cause lung over-expansion as the gas volumes achieved were not different from the volumes achieved at end-inspiration during ventilation with PEEP and a V_t of ~10ml/kg. However, as most preterm infants breathe at birth and the lung may be partially aerated, the use of fixed inflating pressures is problematic as the starting gas volume is unknown. Similarly, the application of a sustained fixed inflation pressure in the presence of breathing may be inadvisable and could lead to considerable lung over-expansion (2). It is likely that different inflation pressures and times, with a maximum limit placed on the volume inhaled, may be preferential in preterm infants.

In conclusion, we have shown that both a 20 s SI and 5 cm H₂O of PEEP promote uniform lung aeration and create a much larger FRC at birth than ventilation without these techniques. A 20 s initial SI resulted in uniform lung aeration before the onset of tidal ventilation, allowing subsequent tidal ventilation to be uniformly distributed throughout the lung. PEEP was essential to maintain FRC both with and without a SI. The effects of PEEP and SI were additive for the creation of FRC. Following a SI the variability in FRC during treatment with PEEP was greatly reduced. It is possible that combining these two techniques during stabilisation of preterm infants would increase the chance of establishing effective breathing, promote uniform ventilation and reduce the need for intubation in the delivery room.

Reference List

1. 2006 The International Liaison Committee on Resuscitation (ILCOR) consensus on science with treatment recommendations for pediatric and neonatal patients: neonatal resuscitation. *Pediatrics* 117:e978-e988
2. Bjorklund LJ, Ingimarsson J, Curstedt T, John J, Robertson B, Werner O, Vilstrup CT 1997 Manual ventilation with a few large breaths at birth compromises the therapeutic effect of subsequent surfactant replacement in immature lambs. *Pediatr Res* 42:348-355
3. Wada K, Jobe AH, Ikegami M 1997 Tidal volume effects on surfactant treatment responses with the initiation of ventilation in preterm lambs. *J Appl Physiol* 83:1054-1061
4. Lindner W, Vossbeck S, Hummler H, Pohlandt F 1999 Delivery room management of extremely low birth weight infants: spontaneous breathing or intubation? *Pediatrics* 103:961-967
5. Aly H, Massaro AN, Patel K, El Mohandes AA 2005 Is it safer to intubate premature infants in the delivery room? *Pediatrics* 115:1660-1665
6. te Pas AB, Walther FJ 2007 A randomized, controlled trial of delivery-room respiratory management in very preterm infants. *Pediatrics* 120:322-329
7. Jobe AH, Kramer BW, Moss TJ, Newnham JP, Ikegami M 2002 Decreased indicators of lung injury with continuous positive expiratory pressure in preterm lambs. *Pediatr Res* 52:387-392
8. Thomson MA, Yoder BA, Winter VT, Martin H, Catland D, Siler-Khodr TM, Coalson JJ 2004 Treatment of immature baboons for 28 days with early nasal continuous positive airway pressure. *Am J Respir Crit Care Med* 169:1054-1062
9. Bjorklund LJ, Ingimarsson J, Curstedt T, Larsson A, Robertson B, Werner O 2001 Lung recruitment at birth does not improve lung function in immature lambs receiving surfactant. *Acta Anaesthesiol Scand* 45:986-993
10. Ingimarsson J, Bjorklund LJ, Curstedt T, Gudmundsson S, Larsson A, Robertson B, Werner O 2004 Incomplete protection by prophylactic surfactant against the adverse effects of large lung inflations at birth in immature lambs. *Intensive Care Med* 30:1446-1453
11. Ikegami M, Kallapur S, Michna J, Jobe AH 2000 Lung injury and surfactant metabolism after hyperventilation of premature lambs. *Pediatr Res* 47:398-404
12. Lindner W, Hogel J, Pohlandt F 2005 Sustained pressure-controlled inflation or intermittent mandatory ventilation in preterm infants in the delivery room? A randomized, controlled trial on initial respiratory support via nasopharyngeal tube. *Acta Paediatr* 94:303-309
13. Vyas H, Milner AD, Hopkin IE, Boon AW 1981 Physiologic responses to prolonged and slow-rise inflation in the resuscitation of the asphyxiated newborn infant. *J Pediatr* 99:635-639
14. Upton CJ, Milner AD 1991 Endotracheal resuscitation of neonates using a rebreathing bag. *Arch Dis Child* 66:39-42
15. Naik AS, Kallapur SG, Bachurski CJ, Jobe AH, Michna J, Kramer BW, Ikegami M 2001 Effects of ventilation with different positive end-expiratory pressures on cytokine expression in the preterm lamb lung. *Am J Respir Crit Care Med* 164:494-498
16. Morley C 2007 New Australian Neonatal Resuscitation Guidelines. *J Paediatr Child Health* 43:6-8
17. Kitchen MJ, Lewis RA, Morgan MJ, Wallace MJ, Siew ML, Siu KK, Habib A, Fouras A, Yagi N, Uesugi K, Hooper SB 2008 Dynamic measures of regional lung air volume using phase contrast x-ray imaging. *Phys Med Biol* 53:6065-6077
18. Hooper SB, Kitchen MJ, Wallace MJ, Yagi N, Uesugi K, Morgan MJ, Hall C, Siu KK, Williams IM, Siew M, Irvine SC, Pavlov K, Lewis RA 2007 Imaging lung aeration and lung liquid clearance at birth. *FASEB J* 21:3329-37
19. Lewis RA, Yagi N, Kitchen MJ, Morgan MJ, Paganin D, Siu KK, Pavlov K, Williams I, Uesugi K, Wallace MJ, Hall CJ, Whitley J, Hooper SB 2005 Dynamic imaging of the lungs using x-ray phase contrast. *Phys Med Biol* 50:5031-5040

20. Lachmann B, Grossmann G, Nilsson R, Robertson B 1979 Lung mechanics during spontaneous ventilation in premature and fullterm rabbit neonates. *Respir Physiol* 38:283-302
21. Vyas H, Field D, Milner AD, Hopkin IE 1986 Determinants of the first inspiratory volume and functional residual capacity at birth. *Pediatr Pulmonol* 2:189-193
22. Vyas H, Milner AD, Hopkins IE 1981 Intrathoracic pressure and volume changes during the spontaneous onset of respiration in babies born by cesarean section and by vaginal delivery. *J Pediatr* 99:787-791
23. Milner AD, Saunders RA, Hopkin IE 1978 Is air trapping important in the maintenance of the functional residual capacity in the hours after birth? *Early Hum Dev* 2:97-105
24. Boon AW, Milner AD, Hopkin IE 1979 Lung expansion, tidal exchange, and formation of the functional residual capacity during resuscitation of asphyxiated neonates. *J Pediatr* 95:1031-1036
25. Milner AD, Saunders RA 1977 Pressure and volume changes during the first breath of human neonates. *Arch Dis Child* 52:918-924
26. Saunders RA, Milner AD 1978 Pulmonary pressure/volume relationships during the last phase of delivery and the first postnatal breaths in human subjects. *J Pediatr* 93:667-673
27. Karlberg P, Cherry RB, Escardo FE, Koch G 1962 pulmonary ventilation and mechanics of breathing in the first minutes of life, including the onset of respiration. *Acta Paediatr Scand* 51:121-136
28. Mortola JP, Fisher JT, Smith JB, Fox GS, Weeks S, Willis D 1982 Onset of respiration in infants delivered by cesarean section. *J Appl Physiol* 52:716-724
29. Probyn ME, Hooper SB, Dargaville PA, McCallion N, Crossley K, Harding R, Morley CJ 2004 Positive End Expiratory Pressure during Resuscitation of Premature Lambs Rapidly Improves Blood Gases without Adversely Affecting Arterial Pressure. *Pediatr Res* 56:198-204
30. Probyn ME, Hooper SB, Dargaville PA, McCallion N, Harding R, Morley CJ 2005 Effects of tidal volume and positive end-expiratory pressure during resuscitation of very premature lambs. *Acta Paediatr* 94:1764-1770
31. O'donnell C, Davis P, Morley C 2004 Positive end-expiratory pressure for resuscitation of newborn infants at birth. *Cochrane Database Syst Rev* CD004341
32. Finer NN, Carlo WA, Duara S, Fanaroff AA, Donovan EF, Wright LL, Kandefer S, Poole WK 2004 Delivery room continuous positive airway pressure/positive end-expiratory pressure in extremely low birth weight infants: a feasibility trial. *Pediatrics* 114:651-657
33. Enhorning G, Robertson B 1972 Lung expansion in the premature rabbit fetus after tracheal deposition of surfactant. *Pediatrics* 50:58-66
34. Enhorning G, Grossman G, Robertson B 1973 Effect of tracheal deposition of surfactant on air expansion of lungs--study on premature rabbit fetuses. *Arch Dis Child* 48:162

