



Universiteit
Leiden
The Netherlands

The psychological impact of breast reconstruction after prophylactic or therapeutic mastectomy for breast cancer

Gopie, J.P.

Citation

Gopie, J. P. (2013, January 9). *The psychological impact of breast reconstruction after prophylactic or therapeutic mastectomy for breast cancer*. Retrieved from <https://hdl.handle.net/1887/21092>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/21092>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/21092> holds various files of this Leiden University dissertation.

Author: Gopie, Jessica Premdee

Title: The psychological impact of breast reconstruction after prophylactic or therapeutic mastectomy for breast cancer

Issue Date: 2013-01-09

CHAPTER 1 INTRODUCTION



1. Breast cancer

In the Netherlands breast cancer is the most common type of cancer in women with a lifetime risk of 1 out of 8. Every year there are more than 13,000 new cases mostly occurring in women older than 50 years of age [1]. Therefore, since 1989 women aged between 50 and 75 years old have been invited to participate biannually in a national breast cancer screening program using mammography which has proven to be effective in the early detection of breast cancer and the reduction of mortality [2-4]. Several treatment options are available for breast cancer, such as surgery, chemotherapy, radiotherapy and hormonal therapy. Despite the introduction of breast conserving therapy with or without neo-adjuvant treatment, mastectomy is still performed in about 46% of all cases [5], while more recent results have shown that long-term survival rates are similar after breast conserving therapy including radiotherapy [6].

About 5% to 10% of all breast cancer cases are caused by a genetic predisposition [7]. Women from families with a BRCA1/2 mutation have a significantly increased cumulative lifetime risk for developing breast cancer (39-85%) as well as ovarian cancer (10-63%) [8-11]. These women are offered a specific breast surveillance program including an annual MRI-scan, a mammography and clinical breast examination. Women at high risk for breast cancer based on their family history are also offered a specific breast surveillance program, as suggested by the Dutch Breast Cancer Guidelines 2012 [12;13]. As breast cancer screening programs may detect breast cancer at an early stage but cannot prevent it, prophylactic mastectomy may also be discussed with these high risk patients as an alternative option to reduce the risk for developing breast cancer [14-16].

1.1. Therapeutic mastectomy and breast reconstruction

Mastectomy can have a significant impact a woman's body image and self-identity, including feelings of abnormality, depressive symptoms, loss of wholeness, and mourning for the lost breast [17-21]. Breast reconstruction (BR) can significantly improve patient satisfaction and body image after mastectomy [22-27]. However, general psychological outcomes do not differ significantly between women with and without BR after mastectomy [28-32]. Recently, it has been found that BR is associated with decreased breast cancer mortality compared to women undergoing mastectomy only, but this is more likely to be explained by socioeconomic factors and access to health care than to oncologic factors [33;34].

The number of patients receiving BR after mastectomy ranges from 10% to 40% but is still increasing. However, there is a great ethnic and geographic variation, for instance non-white women and patients from deprived or rural areas are less likely to undergo BR [35]. In the Netherlands, the estimated uptake of BR after therapeutic mastectomy is about 20%, indicating the majority of patients does not receive BR [36].

This thesis focuses on the impact of two different types of BR after mastectomy: the most commonly performed BR method using silicone implants and the most advanced BR type using abdominal tissue; the Deep Inferior Epigastric artery Perforator (DIEP) flap (Figure 1). There was very little data on psychosocial outcomes regarding DIEP flap BR available at the beginning of this study [37-41]. Therefore, it was investigated what possible benefits

and disadvantages this BR method would bring patients, in comparison to implant BR. The effect of demographic and other clinical variables on the psychosocial outcomes after BR was also investigated.

The following paragraphs provide an overview of therapeutic and prophylactic mastectomy, the types and timing of BR, psychosocial aspects related to the decision to undergo BR and the psychosocial outcomes after BR. In the final paragraph, the aims and research questions of this thesis are described.

1.2. Prophylactic mastectomy and breast reconstruction

Prophylactic mastectomy (PM) is considered to be an effective method for decreasing the breast cancer risk of high risk women significantly with a risk reduction of more than 90% [14-16;42;43], but there is insufficient data to support an improvement in survival rates after PM [44-46]. The combination of the cancer risk, fear of cancer and increasing possibilities for BR has popularized this risk-reducing surgery. More than ten years ago 55% of the Dutch unaffected high risk patients chose bilateral PM [43], but more recently this percentage has decreased to 33% [47]. The reasons for a declining uptake of bilateral PM may be related to recent study results describing the risk for adverse effects on body image and sexuality [48-59]. In addition, more advanced diagnostic imaging techniques have become available, providing earlier detection of breast cancer development [60-65] and it is likely that this improved diagnosis may dissuade doctors and patients from choosing bilateral PM.

This said, the uptake of bilateral PM in the Netherlands is still high compared to the mean international uptake of 18% [47;66] and only the United States of America (36%) and the United Kingdom (40%) have a higher uptake rate. The uptake is lowest in Poland (3%), Israel (4%) and Norway (5%). Regarding contralateral PM in high risk women with a history of breast cancer, the uptake of 53% in the Netherlands is also high, compared to the average international uptake of 27% [46;67]. The large variance in uptake may reflect cultural differences in which both patients and clinicians may play a role.

Currently, BRCA1/2 mutation carriers are advised to undergo prophylactic bilateral salpingo-oophorectomy (PBSO) to reduce the risk for ovarian cancer as well as the prevalence of breast cancer and to improve the survival rate in premenopausal BRCA1/2 mutation carriers [44;68-71]. BRCA1 mutation carriers are recommended to undergo PBSO between 35 to 40 years of age, whereas for BRCA2 mutation carriers this is advised between the age of 40 and 45 years as recommended by the Netherlands Foundation for the Detection of Hereditary Tumours (STOET) [72]. However, PBSO may have a profound effect on psychological wellbeing as it induces the pre-menopause, which may result in adverse psychological, somatic and sexual consequences, including vasomotor symptoms, vaginal dryness and decreased libido [73-75].

1.3. Types of breast reconstruction

In general, three types of BR are available: with silicone implants, autologous tissue, or a combination of both. BR, using a silicone implant is the most commonly performed [76;77] and is usually preceded by the insertion of a tissue expander during the first stage of the

reconstruction, which is gradually inflated to expand the breast skin and muscle [78], and after a few months is replaced with the definite implant.

Several autologous BR techniques exist, commonly including skin, fat and muscle tissue from a distant donor site, such as the latissimus dorsi myocutaneous (LD) flap from the back [79]. The transverse rectus abdominis myocutaneous (TRAM) flap includes skin, fat and muscle from the lower abdomen which is transferred to the chest wall [80], and the TRAM flap is either a pedicled or a free flap requiring a smaller proportion of the abdominal muscle. Other autologous BR techniques include the gluteal artery perforator (GAP) flap and the transverse myocutaneous gracilis (TMG) flap, which consist of skin and fat from the buttock and skin, fat and muscle from the medial thigh area [81;82], respectively. Currently, the most popular autologous BR method is the Deep Inferior Epigastric artery Perforator (DIEP) flap, using abdominal skin and fat, but leaving the abdominal muscles intact, which reduces the chance of abdominal muscle weakness or hernia [83]. However, it requires microsurgical expertise and for the reconstructive surgeon to have specialist knowledge, and this expertise is only available in specialized centers, resulting in long patient waiting lists and higher costs. In general, higher satisfaction rates and a better body image have been reported after autologous BR compared to implant BR, but more prospective studies are required to be conclusive [40;41;49;84-87].

1.4. Timing of breast reconstruction

BR can be performed either immediately, at the same time with mastectomy or delayed; months or years after mastectomy. A randomized controlled trial (RCT) investigating the specific impact of immediate versus delayed reconstruction found that three months postoperatively immediate BR was associated with better psychosocial outcomes compared to delayed or no BR [24]. As randomization to the type or timing of BR is generally seen as unethical with regard to withholding information and not observing patient preferences, no other RCT has been performed in this field since 1983. Cross-sectional and cohort studies have demonstrated that aesthetic outcome is generally better following immediate BR as typically the skin envelope and occasionally the nipple can be preserved [84;88;89]. Women with immediate BR may have less distress and a better body image, more self-esteem and may feel more sexually attractive as well [90]. However, with regard to the type of BR, more positive outcomes on body image and sexual satisfaction were found after delayed autologous BR [40;49;84;87]. The timing of BR is of great significance here, as women with delayed BR have lived for a period with only one or no breasts and therefore have a greater opportunity to experience improvement, as their body image and sexual satisfaction may be at the lowest point before they undergo BR. Furthermore, after delayed BR the complication risk has been found to be lower compared to immediate BR [88].

Generally, the timing of BR does not have an impact on the incidence of recurrent breast cancer and women undergoing immediate BR generally experience no delay in adjuvant treatment compared to women without undergoing BR [33;91-93].

2. Psychological aspects of Breast Reconstruction

2.1. Decisional aspects of breast reconstruction

The decision for the type of BR can be influenced by different factors such as surgeon's preferences and expertise, treatment characteristics (therapeutic or prophylactic indication for mastectomy, timing of BR), and patient characteristics [51;77;94-99]. Some patients are limited in their reconstructive options by clinical and logistic circumstances, such as having insufficient autologous tissue or being restricted by long waiting lists [96]. However, some demographic conditions facilitate the access to BR: being younger, Caucasian, higher educated, wealthier and more often married or in a relationship [31;96;98-100].

Women generally choose BR because they feel too young to live without breasts, want to avoid wearing an external prosthesis, and wish to feel feminine and self-confident [96;98;99]. Common reasons not to choose BR are: feeling reconstruction is not essential for physical and emotional well-being, uncertainty about the procedure and not wanting anything unnatural in the body [96;98].

A lack of information and unrealistic expectations regarding the outcome of BR are also generally associated with regret or low satisfaction with the end result of BR [21;49;95;101-104]. Therefore, it is extremely important to inform patients as comprehensively as possible about the options regarding the types of BR and possible outcomes. Currently, there is no standardized information provided regarding BR options for patients, and, one study demonstrated that autologous BR was chosen by active information seekers and patients who referred themselves to a reconstructive surgeon [105]. This suggests that less independent decision-makers are less aware of the different reconstructive options and as a consequence, due to the increased risk of not selecting the optimal option may be less satisfied with their decision.

2.2. Complications after breast reconstruction

Implant BR can be performed either as a one-stage procedure (direct insertion of prosthesis) or a two-stage procedure (insertion of tissue expander followed by replacement with a definite implant). Presently, the two-stage procedure is generally performed. This procedure may take months as tissue expansion requires several months before the expander can be replaced with the implant. The overall short and long-term complication rates after implant BR vary from 18% to 51% [106-110]. Complications like infections, seroma and hematoma occur in 15% to 39% of the cases, of which 3% to 20% result in implant removal in the short-term [107;109].

Regarding autologous BR, free perforator flaps are preferred as there is less interference with muscle tissue and therefore donor site complications are minimized [76]. The most preferred free perforator flap for BR is the DIEP flap. The overall short and long-term complication rates after autologous BR vary between 32% and 43% [108;111]. Common immediate complications following DIEP flap BR are hematoma, seroma and partial flap necrosis occurring in about 2%, 5% and 15%, respectively, and in experienced hands total flap loss occurs in about 2% of DIEP flap reconstructions [112-114].

The chance for general failure of BR depends on the type of prosthesis used, the reconstruction method, and on risk factors for wound healing problems such as obesity, smoking, hypertension and pre- or postoperative radiotherapy [107;115-117]. Postoperative radiotherapy increases the risk of major complications, in particular after implant BR [106;116]. Postoperative complications can be an important indicator of dissatisfaction with reconstruction [118;119].

2.3. General quality of life

In general, the overall quality of life does not appear to differ between women with BR and women with mastectomy only [27;29;31;120-126]. One study found that women with BR even had a poorer wellbeing compared to women with mastectomy only, for which age might be a contributing factor, as the women with BR were significantly younger [30]. The general quality of life in these studies usually includes parameters such as distress or mood disturbances and overall wellbeing. It is more likely that other psychosocial aspects are affected, such as the altering body image and consequent effects on intimacy and the relationship, which is underlined by in depth-interview studies regarding the personal experiences of patients [54;56;97;101;127-130]. However, it remains relevant to correct for the overall quality of life when investigating these outcomes as they highly correlate with each other [131-134].

2.4. Body image

Conflicting results regarding body image after BR have been reported, with most studies having found no substantial differences between women who had received BR and women who had undergone mastectomy only [29;31;121;135-139]. Other studies reported better body image in women who had received BR [22;24;26;27;122;140;141]. For high risk women who are treated with PM and BR, body image is commonly negatively affected [48-50;57;58]. An explanation for this could be that losing the own healthy breasts without having had breast cancer, and directly exchanging them for reconstructed breasts might reduce satisfaction with appearance, self-esteem and self-consciousness.

The conflicting results regarding body image after BR could also be explained by methodological issues, as most studies had a retrospective design and could therefore not compare baseline scores with postoperative scores. It is also necessary to further investigate predictors of the patient evaluation of body image [142], therefore prospective studies are needed on the predictors of body image.

2.5. Sexual satisfaction and the partner relationship

Sexuality can be complex after breast cancer, as sexual changes can be caused by clinical factors, such as the influence of hormonal and chemotherapy with consequent menopause. A range of physical changes, including dyspareunia, fatigue, vaginal dryness, loss or decreased sexual desire or pleasure and numbness of the breasts, have been reported as a consequence of breast cancer treatment [121;143-150]. Of the BRCA1/2 mutation carriers 35% to 74% opts for PBSO to considerably lower the risk of ovarian cancer [47;151], but this

can, however, induce menopause and cause sexual side-effects, such as vaginal dryness, greater discomfort, and less satisfaction with sexual activities, although most complaints tend to decrease over time [152]. Psychosocial changes regarding sexuality include a fear of loss of fertility, negative body image, loss of femininity and attractiveness, depression and anxiety [143;153-155].

The quality of the partner relationship appears to be the strongest predictor of sexual satisfaction, sexual functioning and sexual desire compared to the physical or chemical changes due to cancer treatment [146;156;157]. If women can discuss their sexual practices with their partner, they are more able to manage changes in their sexual relationship [158]. After BR, sexual functioning has not generally been found to differ between breast cancer patients who had mastectomy only or with BR [22;24;29;31;121;122;125;137-140;159]. However, in high risk women who had PM, adverse sexual effects have been found in a substantial proportion of the patients [48-56].

The specific impact of BR on the partner relationship has received relatively little attention world-wide. A Franco-British study regarding patients' and partners' satisfaction with BR, showed satisfaction was related to the level of preoperative information, type of reconstruction and cosmetic result [160]. This suggests that it is important that further research should be conducted on the impact of BR on partner relationship satisfaction.

2.6. Patient regrets

About 18% of breast cancer survivors regret having had BR [161]. Regrets after BR are associated with a lack of information and unrealistic expectations regarding the outcome, as described previously [21;49;95;101-104]. In addition, women who are more concerned about the future or who had problems in communicating with their doctors are more likely to experience regret five years after cancer treatment [161]. Unaffected high risk women who underwent bilateral PM are less likely to report feelings of regret, which suggests that the relief from the reduced breast cancer risk outweighs the negative side effects of bilateral PM, such as pain, discomfort and adverse sexual effects [49;50;54;162].

3. Aims and outline of the present thesis

The Leiden – Rotterdam collaboration on breast cancer and the psychological issues of BRCA mutation carriers has started over 15 years ago [49;58;95;163-183]. Distress surrounding presymptomatic testing for BRCA1/2 was investigated, and high post-test anxiety was detected in 20% of the mutation carriers and in 35% of their partners [171]. Mutation carriers opting for PM had significantly higher distress levels than those opting for surveillance. In addition, adverse effects in body image and intimacy were observed in women who underwent PM combined with BR [49;163;169;175]. However, distress appeared to decline over time [58;164;169]. More recent studies of Den Heijer et al. showed that low self-esteem and feelings of stigmatization, increased distress and that communication regarding cancer within the family was associated with a reduced sense of vulnerability and less distress [167;168].

The current study aimed to expand this field of research and to explore patient motivation for BR and the specific psychological impact of this surgery on psychological distress, body image, sexuality and the partner relationship. This could increase the body of evidence to support a tailor-made psychosocial care-program for women dealing with either mastectomy after breast cancer or with prophylactic mastectomy.

3.1. Aims and research questions

In our study patients with implant and DIEP flap BR were included as our general aim was to prospectively investigate the psychosocial impact of BR after mastectomy comparing patients with implant and DIEP flap BR. At first, the decision-making process before surgery was explored, to investigate factors or thoughts related to unrealistic expectations. Secondly, our aim was to identify possible demographic, clinical and psychological risk factors that are related to a decreased psychosocial functioning.

The research questions were:

- What are the motives to opt for either implant or DIEP flap BR? (Chapter 2)
- To what extent do women make an independent decision regarding the type of BR? (Chapter 3)
- What is the short-term impact of complications after BR? (Chapter 4)
- What is the long-term impact of complications after BR? (Chapter 5)
- What is the impact of delayed BR on body image and the sexual and partner relationship satisfaction? (Chapter 6)
- What is the impact of prophylactic mastectomy with immediate BR on body image and the sexual and partner relationship satisfaction? (Chapter 7)

3.2. Study procedure

The study was a multi-center prospective follow-up study and the participants were women who opted for post mastectomy reconstruction after breast cancer (n=152) and healthy women who chose prophylactic mastectomy with BR because of an increased hereditary risk for developing breast cancer (n=50, Figure 2). Reconstructions were either implant or DIEP flap based, and were immediate or delayed. Exclusion criteria were:

- A BR in the past
- A detection of recurrence or metastasis of breast cancer during the study period, and
- Not being able to understand and speak the Dutch language sufficiently.

Patients were approached between December 2007 and May 2010 at the Leiden University Medical Center, Erasmus University Medical Center Rotterdam, Daniel den Hoed Cancer Center Rotterdam, Haga Teaching Hospital The Hague, Rijnland Hospital Leiderdorp, the Lange Land Hospital Zoetermeer, Hospital Walcheren, Admiraal de Ruyter Hospital Goes and at the Hospital Zorgsaam Terneuzen. Ethics approval was obtained from all participating hospitals..

Before their operation, an invitation letter explaining the procedure and purpose of the study, an informed consent, and a prepaid envelope were sent to all women on the BR

waiting lists of the participating hospitals. If patients did not respond within two weeks, a reminder was sent. Patients who returned their completed consent form received a self-report questionnaire that they had to complete before their operation. They were requested to complete the same questionnaire post-surgery, after 6 months and at the end of the BR procedure (21 months). In addition, questions regarding distress were sent at 1 month and 12 months post-surgery as well and patients were called by telephone to ask for the occurrence of complications and pain symptoms (Figure 3). After surgery, patient-reported clinical characteristics (including complications) were checked in the medical records.

The questionnaires included demographic information (e.g. age, having a partner or children, educational level), clinical data (e.g. indication for mastectomy, body mass index) and psychological assessments regarding general quality of life, anxiety, depression, cancer-specific distress, body image, sexual satisfaction and partner relationship satisfaction. At the final measurement patient satisfaction regarding the aesthetic result and complications experienced during the whole BR process were also requested. The content and psychometric properties of the self-report questionnaires are described in detail in the relevant chapters.

References

1. <http://www.iknl.nl>. Integraal Kankercentrum Nederland. 2012
2. Otto SJ, Fracheboud J, Verbeek AL, Boer R et al. Mammography Screening and Risk of Breast Cancer Death: A Population-Based Case-Control Study. *Cancer Epidemiology Biomarkers & Prevention* 2012; 21:66-73.
3. Holland R, Rijken H and Hendriks J. The dutch population-based mammography screening: 30-year experience. *Breast Care* 2007; 2:12-18.
4. Otto SJ, Fracheboud J, Looman CWN, Broeders MJM et al. Initiation of population-based mammography screening in Dutch municipalities and effect on breast-cancer mortality: a systematic review. *Lancet* 2003; 361:1411-1417.
5. Struikmans H, Aarts MJ, Jobsen JJ, Koning CCE et al. An increased utilisation rate and better compliance to guidelines for primary radiotherapy for breast cancer from 1997 till 2008: A population-based study in The Netherlands. *Radiotherapy and Oncology* 2011; 100:320-325.
6. Litiere S, Werutsky G, Fentiman IS, Rutgers E et al. Breast conserving therapy versus mastectomy for stage I-II breast cancer: 20 year follow-up of the EORTC 10801 phase 3 randomised trial. *Lancet Oncology* 2012; 13:412-419.
7. Lux MP, Fasching PA and Beckmann MW. Hereditary breast and ovarian cancer: review and future perspectives. *Journal of Molecular Medicine-Imm* 2006; 84:16-28.
8. King MC, Marks JH and Mandell JB. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* 2003; 302:643-646.
9. Ford D, Easton DF, Stratton M, Narod S et al. Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. *American Journal of Human Genetics* 1998; 62:676-689.
10. Antoniou A, Pharoah PDP, Narod S, Risch HA et al. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case series unselected

- for family history: A combined analysis of 22 studies. *American Journal of Human Genetics* 2003; 72:1117-1130.
11. Chen SN and Parmigiani G. Meta-analysis of BRCA1 and BRCA2 penetrance. *Journal of Clinical Oncology* 2007; 25:1329-1333.
 12. <http://www.oncoline.nl/mammacarcinoom>. Dutch Breast Cancer Guidelines. 2012
 13. Gadzicki D, Evans DG, Harris H, Julian-Reynier C et al. Genetic testing for familial/hereditary breast cancer-comparison of guidelines and recommendations from the UK, France, the Netherlands and Germany. *Journal of community genetics* 2011; 2:53-69.
 14. Hartmann LC, Sellers TA, Schaid DJ, Frank TS et al. Efficacy of bilateral prophylactic mastectomy in BRCA1 and BRCA2 gene mutation carriers. *Journal of the National Cancer Institute* 2001; 93:1633-1637.
 15. Hartmann LC, Schaid DJ, Woods JE, Crotty TP et al. Efficacy of bilateral prophylactic mastectomy in women with a family history of breast cancer. *New England Journal of Medicine* 1999; 340:77-84.
 16. Rebbeck TR, Friebel T, Lynch HT, Neuhausen SL et al. Bilateral Prophylactic Mastectomy Reduces Breast Cancer Risk in BRCA1 and BRCA2 Mutation Carriers: The PROSE Study Group. *Journal of Clinical Oncology* 2004; 22:1055-1062.
 17. Bjorkman B, Arner S and Hyden LC. Phantom Breast and Other Syndromes After Mastectomy: Eight Breast Cancer Patients Describe Their Experiences Over Time: A 2-Year Follow-up Study. *Journal of Pain* 2008; 9:1018-1025.
 18. Cromptvoets S. Comfort, control, or conformity: women who choose breast reconstruction following mastectomy. *Health Care for Women International* 2006; 27:75-93.
 19. Hill O and White K. Exploring women's experiences of TRAM flap breast reconstruction after mastectomy for breast cancer. *Oncology Nursing Forum* 2008; 35:81-88.
 20. Piot-Ziegler C, Sassi ML, Raffoul W and Delaloye JF. Mastectomy, body deconstruction, and impact on identity: A qualitative study. *British Journal of Health Psychology* 2010; 15:479-510.
 21. Rolnick SJ, Altschuler A, Nekhlyudov L, Elmore JG et al. What women wish they knew before prophylactic mastectomy. *Cancer Nursing* 2007; 30:285-291.
 22. Al-Ghazal SK, Fallowfield L and Blamey RW. Comparison of psychological aspects and patient satisfaction following breast conserving surgery, simple mastectomy and breast reconstruction. *European Journal of Cancer* 2000; 36:1938-1943.
 23. Cordeiro PG. Breast reconstruction after surgery for breast cancer. *New England Journal of Medicine* 2008; 359:1590-1601.
 24. Dean C, Chetty U and Forrest APM. Effects of Immediate Breast Reconstruction on Psycho-Social Morbidity After Mastectomy. *Lancet* 1983; 1:459-462.
 25. Elder EE, Brandberg Y, Bjorklund T, Rylander R et al. Quality of life and patient satisfaction in breast cancer patients after immediate breast reconstruction: a prospective study. *Breast* 2005; 14:201-208.

26. Nicholson RM, Leinster S and Sassoon EM. A comparison of the cosmetic and psychological outcome of breast reconstruction, breast conserving surgery and mastectomy without reconstruction. *Breast* 2007; 16:396-410.
27. Pusic A, Thompson TA, Kerrigan CL, Sargeant R et al. Surgical options for early-stage breast cancer: Factors associated with patient choice and postoperative quality of life. *Plastic and Reconstructive Surgery* 1999; 104:1325-1333.
28. Collins KK, Perez M, Schootman M, Aft R and Jeffe DB. Effects of breast cancer surgery and surgical side effects on body image over time. *Breast Cancer Research and Treatment* 2011; 126:263-264.
29. Harcourt DM, Rumsey NJ, Ambler NR, Cawthorn SJ et al. The psychological effect of mastectomy with or without breast reconstruction: A prospective, multicenter study. *Plastic and Reconstructive Surgery* 2003; 111:1060-1068.
30. Nissen MJ, Swenson KK, Ritz LJ, Farrell JB et al. Quality of life after breast carcinoma surgery - A comparison of three surgical procedures. *Cancer* 2001; 91:1238-1246.
31. Rowland JH, Desmond KA, Meyerowitz BE, Belin TR et al. Role of breast reconstructive surgery in physical and emotional outcomes among breast cancer survivors. *Journal of the National Cancer Institute* 2000; 92:1422-1429.
32. Parker PA, Youssef A, Walker S, Basen-Engquist K et al. Short-term and long-term psychosocial adjustment and quality of life in women undergoing different surgical procedures for breast cancer. *Annals of Surgical Oncology* 2007; 14:3078-3089.
33. Bezuhly M, Temple C, Sigurdson LJ, Davis RB et al. Immediate Postmastectomy Reconstruction Is Associated With Improved Breast Cancer-Specific Survival Evidence and New Challenges From the Surveillance, Epidemiology, and End Results Database. *Cancer* 2009; 115:4648-4654.
34. Agarwal J, Agarwal S, Pappas L and Neumayer L. A Population-Based Study of Breast Cancer-Specific Survival Following Mastectomy and Immediate or Early-Delayed Breast Reconstruction. *Breast Journal* 2012; 18:226-232.
35. Nelson JA, Nelson P, Tchou J, Serletti JM and Wu LC. The ethnic divide in breast reconstruction: A review of the current literature and directions for future research. *Cancer Treatment Reviews* 2012; 38:362-367.
36. Dutch Medical Registration. Kiwa Prismant BV. 2008
37. Lindegren A, Halle M, Skogh ACD and Edsander-Nord A. Postmastectomy Breast Reconstruction in the Irradiated Breast: A Comparative Study of DIEP and Latissimus Dorsi Flap Outcome. *Plastic and Reconstructive Surgery* 2012; 130:10-18.
38. Zhong T, McCarthy C, Min S, Zhang J et al. Patient satisfaction and health-related quality of life after autologous tissue breast reconstruction. *Cancer* 2012; 118:1701-1709.
39. Haekens CM, Enajat M, Keymeulen K and Van der Hulst RRWJ. Self-esteem and patients' satisfaction after deep inferior epigastric perforator flap breast reconstruction. *Plastic surgical nursing : official journal of the American Society of Plastic and Reconstructive Surgical Nurses* 2011; 31:160-166.

40. Damen THC, Timman R, Kunst EH, Gopie JP et al. High satisfaction rates in women after DIEP flap breast reconstruction. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2010; 63:93-100.
41. Yueh JH, Slavin SA, Adesiyun T, Nyame TT et al. Patient Satisfaction in Postmastectomy Breast Reconstruction: A Comparative Evaluation of DIEP, TRAM, Latissimus Flap, and Implant Techniques. *Plastic and Reconstructive Surgery* 2010; 125:1585-1595.
42. Heemskerk-Gerritsen BAM, Brekelmans CTM, Menke-Pluymers MBE, van Geel AN et al. Prophylactic mastectomy in BRCA1/2 mutation carriers and women at risk of hereditary breast cancer: Long-term experiences at the rotterdam family cancer clinic. *Annals of Surgical Oncology* 2007; 14:3335-3344.
43. Meijers-Heijboer H, van Geel B, van Putten WLJ, Henzen-Logmans SC et al. Breast cancer after prophylactic bilateral mastectomy in women with a BRCA1 or BRCA2 mutation. *New England Journal of Medicine* 2001; 345:159-164.
44. Domchek SM, Friebel TM, Singer CF, Evans DG et al. Association of Risk-Reducing Surgery in BRCA1 or BRCA2 Mutation Carriers With Cancer Risk and Mortality. *Jama-Journal of the American Medical Association* 2010; 304:967-975.
45. Heemskerk-Gerritsen BAM, Hooning MJ, Jager A, Tilanus-Linthorst MMA et al. Is risk-reducing mastectomy in BRCA1/2 mutation carriers with a history of unilateral breast cancer beneficial with respect to distant disease free survival and overall survival? *Ejc Supplements* 2010; 8:206.
46. van Sprundel TC, Schmidt MK, Rookus MA, Brohet R et al. Risk reduction of contralateral breast cancer and survival after contralateral prophylactic mastectomy in BRCA1 or BRCA2 mutation carriers. *Br J Cancer* 2005; 93:287-292.
47. Metcalfe KA, Birenbaum-Carmeli D, Lubinski J, Gronwald J et al. International variation in rates of uptake of preventive options in BRCA1 and BRCA2 mutation carriers. *International Journal of Cancer* 2008; 122:2017-2022.
48. Brandberg Y, Sandelin K, Erikson S, Jurell G et al. Psychological reactions, quality of life, and body image after bilateral prophylactic mastectomy in women at high risk for breast cancer: A prospective 1-year follow-up study. *Journal of Clinical Oncology* 2008; 26:3943-3949.
49. Bresser PJC, Seynaeve C, Van Gool AR, Brekelmans CT et al. Satisfaction with prophylactic mastectomy and breast reconstruction in genetically predisposed women. *Plastic and Reconstructive Surgery* 2006; 117:1675-1682.
50. Gahm J, Wickman M and Brandberg Y. Bilateral prophylactic mastectomy in women with inherited risk of breast cancer - Prevalence of pain and discomfort, impact on sexuality, quality of life and feelings of regret two years after surgery. *Breast* 2010; 19:462-469.
51. Eldor L and Spiegel A. Breast Reconstruction after Bilateral Prophylactic Mastectomy in Women at High Risk for Breast Cancer. *Breast Journal* 2009; 15:S81-S89.
52. Frost MH, Schaid DJ, Sellers TA, Slezak JM et al. Long-term satisfaction and psychological and social function following bilateral prophylactic mastectomy. *Jama-Journal of the American Medical Association* 2000; 284:319-324.

53. Geiger AM, Nekhlyudov L, Herrinton LJ, Rolnick SJ et al. Quality of life after bilateral prophylactic mastectomy. *Annals of Surgical Oncology* 2007; 14:686-694.
54. Hatcher MB and Fallowfield LJ. A qualitative study looking at the psychosocial implications of bilateral prophylactic mastectomy. *Breast* 2003; 12:1-9.
55. McGaughey A. Body image after bilateral prophylactic mastectomy: an integrative literature review. *J Midwifery Womens Health* 2006; 51:e45-e49.
56. Payne DK, Biggs C, Tran KN, Borgen PI and Massie MJ. Women's regrets after bilateral prophylactic mastectomy. *Annals of Surgical Oncology* 2000; 7:150-154.
57. Matloff ET, Barnett RE and Bober SL. Unraveling the Next Chapter Sexual Development, Body Image, and Sexual Functioning in Female BRCA Carriers. *Cancer Journal* 2009; 15:15-18.
58. van Oostrom I, Meijers-Heijboer H, Ladder LN, Duivenvoorden HJ et al. Long-term psychological impact of carrying a BRCA1/2 mutation and prophylactic surgery: A 5-year follow-up study. *Journal of Clinical Oncology* 2003; 21:3867-3874.
59. Rijnsburger AJ, Obdeijn IM, Kaas R, Tilanus-Linthorst MMA et al. BRCA1-Associated Breast Cancers Present Differently From BRCA2-Associated and Familial Cases: Long-Term Follow-Up of the Dutch MRISC Screening Study. *Journal of Clinical Oncology* 2010; 28:5265-5273.
60. Kriege M, Brekelmans CTM, Boetes C, Besnard PE et al. Efficacy of MRI and mammography for breast-cancer screening in women with a familial or genetic predisposition. *New England Journal of Medicine* 2004; 351:427-437.
61. Leach MO, Boggis CRM, Dixon AK, Easton DF et al. Screening with magnetic resonance imaging and mammography of a UK population at high familial risk of breast cancer: a prospective multicentre cohort study (MARIBS). *Lancet* 2005; 365:1769-1778.
62. Sardanelli F, Podo F, D'Agnolo G, Verdecchia A et al. Multicenter comparative multimodality surveillance of women at genetic-familial high risk for breast cancer (HIBCRIT study): Interim results. *Radiology* 2007; 242:698-715.
63. Warner E, Plewes DB, Hill KA, Causer PA et al. Surveillance of BRCA1 and BRCA2 mutation carriers with magnetic resonance imaging, ultrasound, mammography, and clinical breast examination. *Jama-Journal of the American Medical Association* 2004; 292:1317-1325.
64. Weinstein SP, Localio A, Conant EF, Rosen M et al. Multimodality Screening of High-Risk Women: A Prospective Cohort Study. *Journal of Clinical Oncology* 2009; 27:6124-6128.
65. Hagen AI, Kvistad KA, Maehle L, Holmen MM et al. Sensitivity of MRI versus conventional screening in the diagnosis of BRCA-associated breast cancer in a national prospective series. *Breast* 2007; 16:367-374.
66. Evans DGR, Lalloo F, Ashcroft L, Shenton A et al. Uptake of Risk-Reducing Surgery in Unaffected Women at High Risk of Breast and Ovarian Cancer Is Risk, Age, and Time Dependent. *Cancer Epidemiology Biomarkers & Prevention* 2009; 18:2318-2324.

67. Metcalfe KA, Lubinski J, Ghadirian P, Lynch H et al. Predictors of contralateral prophylactic mastectomy in women with a BRCA1 or BRCA2 mutation: The hereditary breast cancer clinical study group. *Journal of Clinical Oncology* 2008; 26:1093-1097.
68. Kauff ND, Satagopan JM, Robson ME, Scheuer L et al. Risk-reducing salpingo-oophorectomy in women with a BRCA1 or BRCA2 mutation. *New England Journal of Medicine* 2002; 346:1609-1615.
69. Finch A, Beiner M, Lubinski J, Lynch HT et al. Salpingo-oophorectomy and the risk of ovarian, fallopian tube, and peritoneal cancers in women with a BRCA1 or BRCA2 mutation. *Jama-Journal of the American Medical Association* 2006; 296:185-192.
70. Hermsen BBJ, van Diest PJ, Berkhof J, Menko FH et al. Low prevalence of (pre) malignant lesions in the breast and high prevalence in the ovary and fallopian tube in women at hereditary high risk of breast and ovarian cancer. *International Journal of Cancer* 2006; 119:1412-1418.
71. Brekelmans CTM, Tilanus-Linthorst MMA, Seynaeve C, Van der Ouweland A et al. Tumour characteristics, survival and prognostic factors of hereditary breast cancer from BRCA2-, BRCA1- and non-BRCA1/2 families as compared to sporadic breast cancer cases. *European Journal of Cancer* 2007; 43:867-876.
72. <http://www.stoet.nl/uploads/richtlijnenboekje.pdf>. Netherlands Foundation for the Detection of Hereditary Tumours (STOET) and the Dutch Society of Clinical Geneticists (VKGN). 2012
73. Benshushan A, Rojansky N, Chaviv M, Arbel-Alon S et al. Climacteric symptoms in women undergoing risk-reducing bilateral salpingo-oophorectomy. *Climacteric* 2009; 12:404-409.
74. Madalinska JB, Hollenstein J, Bleiker E, van Beurden M et al. Quality-of-life effects of prophylactic salpingo-oophorectomy versus gynecologic screening among women at increased risk of hereditary ovarian cancer. *Journal of Clinical Oncology* 2005; 23:6890-6898.
75. Finch A and Narod SA. Quality of life and health status after prophylactic salpingo-oophorectomy in women who carry a BRCA mutation: A review. *Maturitas* 2011; 70:261-265.
76. Petit JY, Rietjens M, Lohsiriwat V, Rey P et al. Update on Breast Reconstruction Techniques and Indications. *World Journal of Surgery* 2012; 36:1486-1497.
77. Rosson GD, Magarakis M, Shridharani SM, Stapleton SM et al. A Review of the Surgical Management of Breast Cancer: Plastic Reconstructive Techniques and Timing Implications. *Annals of Surgical Oncology* 2010; 17:1890-1900.
78. Singh N, Reaven NL and Funk SE. Immediate 1-stage vs. tissue expander postmastectomy implant breast reconstructions: A retrospective real-world comparison over 18 months. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2012; 65:917-923.
79. Maxwell GP, Tansini,Iginio and the Origin of the Latissimus Dorsi Musculocutaneous Flap. *Plastic and Reconstructive Surgery* 1980; 65:686-692.

80. Hartrampf CR, Schefflan M and Black PW. Breast Reconstruction with a Transverse Abdominal Island Flap. *Plastic and Reconstructive Surgery* 1982; 69.
81. Allen RJ and Tucker C. Superior Gluteal Artery Perforator Free-Flap for Breast Reconstruction. *Plastic and Reconstructive Surgery* 1995; 95:1207-1212.
82. Higgins JP, Orlando GS and Blondeel PN. Ischial pressure sore reconstruction using an inferior gluteal artery perforator (IGAP) flap. *British Journal of Plastic Surgery* 2002; 55:83-85.
83. Allen RJ and Treece P. Deep Inferior Epigastric Perforator Flap for Breast Reconstruction. *Annals of Plastic Surgery* 1994; 32:32-38.
84. Atisha D, Alderman AK, Lowery JC, Kuhn LE et al. Prospective analysis of long-term psychosocial outcomes in breast reconstruction - Two-year postoperative results from the Michigan Breast Reconstruction Outcomes Study. *Annals of Surgery* 2008; 247:1019-1028.
85. Christensen BO, Overgaard J, Kettner LO and Damsgaard TE. Long-term evaluation of postmastectomy breast reconstruction. *Acta Oncologica* 2011; 50:1053-1061.
86. Tonseth KA, Hokland BM, Tindholdt TT, Abyholm FE and Stavem K. Quality of life, patient satisfaction and cosmetic outcome after breast reconstruction using DIEP flap or expandable breast implant. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2008; 61:1188-1194.
87. Wilkins EG, Cederna PS, Lowery JC, Davis JA et al. Prospective analysis of psychosocial outcomes in breast reconstruction: One-year postoperative results from the Michigan Breast Reconstruction Outcome Study. *Plastic and Reconstructive Surgery* 2000; 106:1014-1025.
88. Chevray PM. Timing of breast reconstruction: Immediate versus delayed. *Cancer Journal* 2008; 14:223-229.
89. Roth RS, Lowery JC, Davis J and Wilkins EG. Quality of life and affective distress in women seeking immediate versus delayed breast reconstruction after mastectomy for breast cancer. *Plastic and Reconstructive Surgery* 2005; 116:993-1002.
90. Al-Ghazal SK, Sully L, Fallowfield L and Blamey RW. The psychological impact of immediate rather than delayed breast reconstruction. *European Journal of Surgical Oncology* 2000; 26:17-19.
91. Holmich LR, Durning M, Henriksen TF, Krag C et al. Delayed breast reconstruction with implants after invasive breast cancer does not impair prognosis. *Annals of Plastic Surgery* 2008; 61:11-18.
92. Gouy S, Rouzier R, Missana MC, Atallah D et al. Immediate reconstruction after neoadjuvant chemotherapy: Effect on adjuvant treatment starting and survival. *Annals of Surgical Oncology* 2005; 12:161-166.
93. Wilson CR, Brown IM, Weiller-Mithoff E, George WD and Doughty JC. Immediate breast reconstruction does not lead to a delay in the delivery of adjuvant chemotherapy. *Ejso* 2004; 30:624-627.

94. Alderman AK, Hawley ST, Waljee J, Morrow M and Katz SJ. Correlates of referral practices of general surgeons to plastic surgeons for mastectomy reconstruction. *Cancer* 2007; 109:1715-1720.
95. Contant CME, van Wersch AMEA, Wiggers T, Wai RTJ and van Geel AN. Motivations, satisfaction, and information of immediate breast reconstruction following mastectomy. *Patient Education and Counseling* 2000; 40:201-208.
96. Handel N, Silverstein MJ, Waisman E and Waisman JR. Reasons Why Mastectomy Patients do Not Have Breast Reconstruction. *Plastic and Reconstructive Surgery* 1990; 86:1118-1122.
97. Lee CN, Hultman CS and Sepucha K. What Are Patients' Goals and Concerns About Breast Reconstruction After Mastectomy? *Annals of Plastic Surgery* 2010; 64:567-569.
98. Reaby LL. Reasons why women who have mastectomy decide to have or not to have breast reconstruction. *Plastic and Reconstructive Surgery* 1998; 101:1810-1818.
99. Shameem H, Yip CH and Fong E. Immediate Breast Reconstruction after Mastectomy - Why do Women Choose this Option? *Asian Pacific J Cancer Prev* 2008; 9:409-412.
100. Fallbjork U, Karlsson S, Salander P and Rasmussen BH. Differences between women who have and have not undergone breast reconstruction after mastectomy due to breast cancer. *Acta Oncologica* 2010; 49:174-179.
101. Abu-Nab Z and Grunfeld EA. Satisfaction with outcome and attitudes towards scarring among women undergoing breast reconstructive surgery. *Patient Education and Counseling* 2007; 66:243-249.
102. Brandberg Y, Malm M, Rutqvist LE, Jonsson E and Blomqvist L. A prospective randomised study (named SVEA) of three methods of delayed breast reconstruction. Study design, patients' preoperative problems and expectations. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery* 1999; 33:209-216.
103. Sheehan J, Sherman KA, Lam T and Boyages J. Association of information satisfaction, psychological distress and monitoring coping style with post-decision regret following breast reconstruction. *Psycho-Oncology* 2007; 16:342-351.
104. Snell L, McCarthy C, Klassen A, Cano S et al. Clarifying the Expectations of Patients Undergoing Implant Breast Reconstruction: A Qualitative Study. *Plastic and Reconstructive Surgery* 2010; 126:1825-1830.
105. Matros E, Yueh JH, Bar-Meir ED, Slavin SA et al. Sociodemographics, referral patterns, and internet use for decision-making in microsurgical breast reconstruction. *Plast Reconstr Surg* 2010; 125:1087-1094.
106. Berry T, Brooks S, Sydow N, Djohan R et al. Complication rates of radiation on tissue expander and autologous tissue breast reconstruction. *Ann Surg Oncol* 2010; 17:202-210.
107. McCarthy CM, Mehrara BJ, Riedel E, Davidge K et al. Predicting complications following expander/implant breast reconstruction: An outcomes analysis based on preoperative clinical risk. *Plastic and Reconstructive Surgery* 2008; 121:1886-1892.

108. Sullivan SR, Fletcher DRD, Isom CD and Isik FF. True incidence of all complications following immediate and delayed breast reconstruction. *Plastic and Reconstructive Surgery* 2008; 122:19-28.
109. Woerdeman LAE, Hage JJ, Hofland MMI and Rutgers EJT. A prospective assessment of surgical risk factors in 400 cases of skin-sparing mastectomy and immediate breast reconstruction with implants to establish selection criteria. *Plastic and Reconstructive Surgery* 2007; 119:455-463.
110. Delgado JF, Garcia-Guilarte RF, Palazuelo MR, Mendez JIS and Perez CC. Immediate Breast Reconstruction with Direct, Anatomic, Gel-Cohesive, Extra-Projection Prosthesis: 400 Cases. *Plastic and Reconstructive Surgery* 2010; 125:1599-1605.
111. Hofer SOP, Damen THC, Mureau MAM, Rakhorst HA and Roche NA. A critical review of perioperative complications in 175 free deep inferior epigastric perforator flap breast reconstructions. *Annals of Plastic Surgery* 2007; 59:137-142.
112. Chen CM, Halvorson EG, Disa JJ, McCarthy C et al. Immediate postoperative complications in DIEP versus Free/Muscle-Sparing TRAM flaps. *Plastic and Reconstructive Surgery* 2007; 120:1477-1482.
113. Blondeel PN. One hundred free DIEP flap breast reconstructions: a personal experience. *British Journal of Plastic Surgery* 1999; 52:104-111.
114. Hamdi M, Weiler-Mithoff EM and Webster MHC. Deep inferior epigastric perforator flap in breast reconstruction: Experience with the first 50 flaps. *Plastic and Reconstructive Surgery* 1999; 103:86-95.
115. Reefy S, Patani N, Anderson A, Burgoyne G et al. Oncological outcome and patient satisfaction with skin-sparing mastectomy and immediate breast reconstruction: a prospective observational study. *Bmc Cancer* 2010; 10:171-183.
116. Christante D, Pommier SJ, Diggs BS, Samuelson BT et al. Using Complications Associated With Postmastectomy Radiation and Immediate Breast Reconstruction to Improve Surgical Decision Making. *Archives of Surgery* 2010; 145:873-878.
117. Chew BK, Yip C and Malyon AD. Becker expander implants: Truly a long term single stage reconstruction? *Journal of Plastic Reconstructive and Aesthetic Surgery* 2010; 63:1300-1304.
118. Andrade WN, Baxter N and Semple JL. Clinical determinants of patient satisfaction with breast reconstruction. *Plast Reconstr Surg* 2001; 107:46-54.
119. Metcalfe KA, Esplen MJ, Goel V and Narod SA. Predictors of quality of life in women with a bilateral prophylactic mastectomy. *Breast J* 2005; 11:65-69.
120. Arora NK, Gustafson DH, Hawkins RP, McTavish F et al. Impact of surgery and chemotherapy on the quality of life of younger women with breast carcinoma - A prospective study. *Cancer* 2001; 92:1288-1298.
121. Avis NE, Crawford S and Manuel J. Psychosocial problems among younger women with breast cancer. *Psycho-Oncology* 2004; 13:295-308.
122. Janz NK, Mujahid M, Lantz PM, Fagerlin A et al. Population-based study of the relationship of treatment and sociodemographics on quality of life for early stage breast cancer. *Quality of Life Research* 2005; 14:1467-1479.

123. Lantz PM, Janz NK, Fagerlin A, Schwartz K et al. Satisfaction with surgery outcomes and the decision process in a population-based sample of women with breast cancer. *Health Services Research* 2005; 40:745-767.
124. Nano MT, Gill PG, Kollias J, Bochner MA et al. Psychological impact and cosmetic outcome of surgical breast cancer strategies. *Anz Journal of Surgery* 2005; 75:940-947.
125. Rubino C, Figus A, Loretto L and Sechi G. Post-mastectomy reconstruction: a comparative analysis on psychosocial and psychopathological outcomes. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2007; 60:509-518.
126. Veiga DF, Neto MS, Ferreira LM, Garcia EB et al. Quality of life outcomes after pedicled TRAM flap delayed breast reconstruction. *British Journal of Plastic Surgery* 2004; 57:252-257.
127. Fallbjork U, Frejeus E and Rasmussen BH. A preliminary study into women's experiences of undergoing reconstructive surgery after breast cancer. *European journal of oncology nursing : the official journal of European Oncology Nursing Society* 2012; 16:220-226.
128. Fang SY, Balneaves LG and Shu BC. "A Struggle Between Vanity and Life" The Experience of Receiving Breast Reconstruction in Women of Taiwan. *Cancer Nursing* 2010; 33:E1-E11.
129. McQuirter M, Castiglia LL, Loiselle CG and Wong N. Decision-Making Process of Women Carrying a BRCA1 or BRCA2 Mutation Who Have Chosen Prophylactic Mastectomy. *Oncology Nursing Forum* 2010; 37:313-320.
130. Wasteson E, Sandelin K, Brandberg Y, Wickman M and Arver B. High satisfaction rate ten years after bilateral prophylactic mastectomy - a longitudinal study. *European Journal of Cancer Care* 2011; 20:508-513.
131. Al-Ghazal SK, Fallowfield L and Blamey RW. Does cosmetic outcome from treatment of primary breast cancer influence psychosocial morbidity? *European Journal of Surgical Oncology* 1999; 25:571-573.
132. Helms RL, O'Hea EL and Corso M. Body image issues in women with breast cancer. *Psychol Health Med* 2008; 13:313-325.
133. Schover LR, Yetman RJ, Tuason LJ, Meisler E et al. Partial Mastectomy and Breast Reconstruction - A Comparison of Their Effects on Psychosocial Adjustment, Body-Image, and Sexuality. *Cancer* 1995; 75:54-64.
134. Zimmermann T, Scott JL and Heinrichs N. Individual and dyadic predictors of body image in women with breast cancer. *Psycho-Oncology* 2010; 19:1061-1068.
135. Mock V. Body-Image in Women Treated for Breast-Cancer. *Nursing Research* 1993; 42:153-157.
136. Noyan MA, Sertoz OO, Elbi H, Kayar R and Yilmaz R. Variables affecting patient satisfaction in breast surgery: A cross-sectional sample of Turkish women with breast cancer. *International Journal of Psychiatry in Medicine* 2006; 36:299-313.

137. Wellisch DK, Silverstein M, Hoffman R, Handel N et al. Psychosocial Outcomes of Breast-Cancer Therapies - Lumpectomy Versus Mastectomy. *Psychosomatics* 1989; 30:365-373.
138. Yurek D, Farrar W and Andersen BL. Breast cancer surgery: Comparing surgical groups and determining individual differences in postoperative sexuality and body change stress. *Journal of Consulting and Clinical Psychology* 2000; 68:697-709.
139. Zweifler M, Rodriguez E, Reilly J, Lewis T and Glasberg SB. Breast reconstruction among inner city women with breast carcinoma. *Annals of Plastic Surgery* 2001; 47:53-59.
140. Fung KW, Lau Y, Fielding R, Or A and Yip AWC. The impact of mastectomy, breast-conserving treatment and immediate breast reconstructions on the quality of life of Chinese women. *Australian and New Zealand Journal of Surgery* 2001; 71:202-206.
141. Nano MT, Gill PG, Kollias J, Bochner MA et al. Qualitative assessment of breast reconstruction in a specialist breast unit. *Anz Journal of Surgery* 2005; 75:445-453.
142. Beesley H, Ullmer H, Holcombe C and Salmon P. How patients evaluate breast reconstruction after mastectomy, and why their evaluation often differs from that of their clinicians. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2012; 65:1064-1071.
143. Wilmoth MC. The aftermath of breast cancer: An altered sexual self. *Cancer Nursing* 2001; 24:278-286.
144. Fobair P, Stewart SL, Chang SB, D'Onofrio C et al. Body image and sexual problems in young women with breast cancer. *Psycho-Oncology* 2006; 15:579-594.
145. Meyerowitz BE, Desmond KA, Rowland JH, Wyatt GE and Ganz PA. Sexuality following breast cancer. *Journal of Sex & Marital Therapy* 1999; 25:237-250.
146. Speer JJ, Hillenberg B, Sugrue DP, Blacker C et al. Study of sexual functioning determinants in breast cancer survivors. *Breast Journal* 2005; 11:440-447.
147. Ganz PA, Rowland JH, Desmond K, Meyerowitz BE and Wyatt GE. Life after breast cancer: Understanding women's health-related quality of life and sexual functioning. *Journal of Clinical Oncology* 1998; 16:501-514.
148. Ganz PA, Desmond KA, Belin TR, Meyerowitz BE and Rowland JH. Predictors of sexual health in women after a breast cancer diagnosis. *Journal of Clinical Oncology* 1999; 17:2371-2380.
149. Ganz PA, Greendale GA, Petersen L, Kahn B and Bower JE. Breast cancer in younger women: Reproductive and late health effects of treatment. *Journal of Clinical Oncology* 2003; 21:4184-4193.
150. Knobf MT. The menopausal symptom experience in young mid-life women with breast cancer. *Cancer Nursing* 2001; 24:201-211.
151. Finch A, Beiner M, Lubinski J, Lynch HT et al. Salpingo-oophorectomy and the risk of ovarian, fallopian tube, and peritoneal cancers in women with a BRCA1 or BRCA2 mutation. *Jama-Journal of the American Medical Association* 2006; 296:185-192.

152. Fang CY, Cherry C, Devarajan K, Li TY et al. A prospective study of quality of life among women undergoing risk-reducing salpingo-oophorectomy versus gynecologic screening for ovarian cancer. *Gynecologic Oncology* 2009; 112:594-600.
153. Archibald S, Lemieux S, Byers ES, Tamlyn K and Worth J. Chemically-induced menopause and the sexual functioning of breast cancer survivors. *Women & Therapy* 2006; 29:83-106.
154. Bertero C and Wilmoth MC. Breast cancer diagnosis and its treatment affecting the self - A meta-synthesis. *Cancer Nursing* 2007; 30:194-202.
155. Garrusi B and Faezee H. How do Iranian women with breast cancer conceptualize sex and body image? *Sexuality and Disability* 2008; 26:159-165.
156. Zee B, Huang C, Mak S, Wong J et al. Factors related to sexual health in Chinese women with breast cancer in Hong Kong. *Asia-Pacific Journal of Clinical Oncology* 2008; 4:218-226.
157. Alder J, Zanetti R, Wight E, Urech C et al. Sexual dysfunction after premenopausal Stage I and II breast cancer: Do androgens play a role? *Journal of Sexual Medicine* 2008; 5:1898-1906.
158. Gilbert E, Ussher JM and Perz J. Renegotiating Sexuality and Intimacy in the Context of Cancer: The Experiences of Carers. *Archives of Sexual Behavior* 2010; 39:998-1009.
159. Reaby LL and Hort LK. Postmastectomy Attitudes in Women Who Wear External Breast Prostheses Compared to Those Who Have Undergone Breast Reconstructions. *Journal of Behavioral Medicine* 1995; 18:55-67.
160. Guyomard V, Leinster S, Wilkinson M, Servant JM and Pereira J. A Franco-British patients' and partners' satisfaction audit of breast reconstruction. *Journal of Plastic Reconstructive and Aesthetic Surgery* 2009; 62:782-789.
161. Fernandes-Taylor S and Bloom JR. Post-treatment regret among young breast cancer survivors. *Psycho-Oncology* 2011; 20:506-516.
162. Lloyd SM, Watson M, Oaker G, Sacks N et al. Understanding the experience of prophylactic bilateral mastectomy: A qualitative study of ten women. *Psycho-Oncology* 2000; 9:473-485.
163. Bresser PJC, Seynaeve C, Van Gool AR, Brekelmans CTM et al. Long-term satisfaction with bilateral prophylactic mastectomy and immediate breast reconstruction in genetically predisposed women. *Breast Cancer Research and Treatment* 2003; 82:32.
164. Bresser PJC, Seynaeve C, Van Gool AR, Niermeijer MF et al. The course of distress in women at increased risk of breast and ovarian cancer due to an (identified) genetic susceptibility who opt for prophylactic mastectomy and/or salpingo-oophorectomy. *European Journal of Cancer* 2007; 43:95-103.
165. Bresser PJC, Van Gool AR, Seynaeve C, Duivenvoorden HJ et al. Who is prone to high levels of distress after prophylactic mastectomy and/or salpingo-ovariectomy? *Annals of Oncology* 2007; 18:1641-1645.
166. den Heijer M, Vanheusden K, Seynaeve C, Duivenvoorden HJ et al. Distress in partners of high-risk women undergoing breast cancer surveillance. *Journal of Mens Health* 2010; 7:413-419.

167. den Heijer M, Seynaeve C, Vanheusden K, Duivenvoorden HJ et al. The contribution of self-esteem and self-concept in psychological distress in women at risk of hereditary breast cancer. *Psycho-Oncology* 2011; 20:1170-1175.
168. den Heijer M, Seynaeve C, Vanheusden K, Duivenvoorden HJ et al. Psychological distress in women at risk for hereditary breast cancer: the role of family communication and perceived social support. *Psycho-Oncology* 2011; 20:1317-1323.
169. den Heijer M, Seynaeve C, Timman R, Duivenvoorden HJ et al. Body image and psychological distress after prophylactic mastectomy and breast reconstruction in genetically predisposed women: A prospective long-term follow-up study. *European Journal of Cancer* 2012; 48:1263-1268.
170. den Heijer M, Vos J, Seynaeve C, Vanheusden K et al. The impact of social and personal resources on psychological distress in women at risk for hereditary breast cancer. *Psycho-Oncology* 2012; 21:153-160.
171. Lodder L, Frets PG, Trijsburg RW, Meijers-Heijboer EJ et al. Psychological impact of receiving a BRCA1/BRCA2 test result. *American Journal of Medical Genetics* 2001; 98:15-24.
172. Lodder L, Frets PG, Trijsburg RW, Klijn JGM et al. Attitudes and distress levels in women at risk to carry a BRCA1/BRCA2 gene mutation who decline genetic testing. *American Journal of Medical Genetics Part A* 2003; 119A:266-272.
173. Lodder LN, Frets PG, Trijsburg RW, Meijers-Heijboer EJ et al. Presymptomatic testing for BRCA1 and BRCA2: how distressing are the pre-test weeks? *Journal of Medical Genetics* 1999; 36:906-913.
174. Lodder LN, Frets PG, Trijsburg RW, Meijers-Heijboer EJ et al. Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the BRCA1/BRCA2 gene mutation in The Netherlands. *Journal of Medical Genetics* 2000; 37:883-884.
175. Lodder LN, Frets PG, Trijsburg RW, Meijers-Heijboer EJ et al. One year follow-up of women opting for presymptomatic testing for BRCA1 and BRCA2: Emotional impact of the test outcome and decisions on risk management (Surveillance or prophylactic surgery). *Breast Cancer Research and Treatment* 2002; 73:97-112.
176. van Dijk S, Otten W, Zoetewij MW, Timmermans DRM et al. Genetic counselling and the intention to undergo prophylactic mastectomy: Effects of a breast cancer risk assessment. *British Journal of Cancer* 2003; 88:1675-1681.
177. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Brocker-Vriens AHJT et al. Experience of parental cancer in childhood is a risk factor for psychological distress during genetic cancer susceptibility testing. *Annals of Oncology* 2006; 17:1090-1095.
178. van Oostrom I, Meijers-Heijboer H, Duiven-Voorden H, Brocker-Vriens AHJT et al. Open family communication and the well-being of individuals opting for genetic cancer susceptibility testing. *Psycho-Oncology* 2006; 15:1039.
179. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Brocker-Vriens AHJT et al. Comparison of individuals opting for BRCA1/2 or HNPCC genetic susceptibility testing with regard to coping, illness perceptions, illness experiences, family system

- characteristics and hereditary cancer distress. *Patient Education and Counseling* 2007; 65:58-68.
180. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Brocker-Vriends AHJT et al. Family system characteristics and psychological adjustment to cancer susceptibility genetic testing: a prospective study. *Clinical Genetics* 2007; 71:35-42.
181. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Brocker-Vriends AHJT et al. A prospective study of the impact of genetic susceptibility testing for BRCA1/2 or HNPCC on family relationships. *Psycho-Oncology* 2007; 16:320-328.
182. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Brocker-Vriends AHJT et al. Prognostic factors for hereditary cancer distress six months after BRCA1/2 or HNPCC genetic susceptibility testing. *European Journal of Cancer* 2007; 43:71-77.
183. van Oostrom I, Meijers-Heijboer H, Duivenvoorden HJ, Broecker-Vriends AHJT et al. The common sense model of self-regulation and psychological adjustment to predictive genetic testing: a prospective study. *Psycho-Oncology* 2007; 16:1121-1129.

Figure 1. Implant (A) and DIEP flap (B) breast reconstruction

Figure 2. Patient inflow




Figure 3. Timing of measurements

