



Universiteit
Leiden
The Netherlands

Impact and benefits of early hearing screening

Korver, A.M.H.

Citation

Korver, A. M. H. (2010, October 8). *Impact and benefits of early hearing screening*. Retrieved from <https://hdl.handle.net/1887/16023>

Version: Corrected Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/16023>

Note: To cite this publication please use the final published version (if applicable).

REFERENCES

- (1) Wessex Universal Neonatal Hearing Screening Trial Group. Controlled trial of universal neonatal screening for early identification of permanent childhood hearing impairment. *Lancet* 1998;352:1957-1964.
- (2) Watkin PM, Baldwin M. Confirmation of deafness in infancy. *Arch Dis Child* 1999;81:380-389.
- (3) Kennedy CR. Controlled trial of universal neonatal screening for early identification of permanent childhood hearing impairment: coverage, positive predictive value, effect on mothers and incremental yield. Wessex Universal Neonatal Screening Trial Group. *Acta Paediatr Suppl* 1999;88:73-75.
- (4) Fortnum H, Davis A. Epidemiology of permanent childhood hearing impairment in Trent Region, 1985-1993. *Br J Audiol* 1997;31:409-446.
- (5) Fortnum HM, Summerfield AQ, Marshall DH, Davis AC, Bamford JM. Prevalence of permanent childhood hearing impairment in the United Kingdom and implications for universal neonatal hearing screening: questionnaire based ascertainment study. *BMJ* 2001;323:536-540.
- (6) Morton CC, Nance WE. Newborn hearing screening—a silent revolution. *N Engl J Med* 2006;354:2151-2164.
- (7) Konigsmark BW, Gorlin RJ. *Genetic and metabolic deafness*. W.B. Saunders Co., 1976.
- (8) Fraser GR. *The causes of profound deafness in childhood*. John Hopkins University Press, 1976.
- (9) Korver AM, Meuwese-Jongejeugd J, Briare JJ, Frijns JH, Oudesluys-Murphy AM. [Permanent hearing loss in the prelingual phase in children with a non-aberrant neonatal hearing screening result]. *Ned Tijdschr Geneeskd* 2008;152:426-430.
- (10) Nance WE, Lim BG, Dodson KM. Importance of congenital cytomegalovirus infections as a cause for pre-lingual hearing loss. *J Clin Virol* 2006;35:221-225.
- (11) Malm G, Engman ML. Congenital cytomegalovirus infections. *Semin Fetal Neonatal Med* 2007;12:154-159.
- (12) Dollard SC, Grosse SD, Ross DS. New estimates of the prevalence of neurological and sensory sequelae and mortality associated with congenital cytomegalovirus infection. *Rev Med Virol* 2007;17:355-363.
- (13) Ross SA, Boppana SB. Congenital cytomegalovirus infection: outcome and diagnosis. *Semin Pediatr Infect Dis* 2005;16:44-49.
- (14) Conyn-van Spaendonck MA, Veldhuijzen IK, Suijkerbuijk AW, Hirasing RA. [Significant decline of the number of invasive Haemophilus influenzae infections in the first 4 years after introduction of vaccination against H. influenzae type B in children]. *Ned Tijdschr Geneeskd* 2000;144:1069-1073.

- (15) Year 2007 position statement: Principles and guidelines for early hearing detection and intervention programs. *Pediatrics* 2007;120:898-921.
- (16) Theunissen EJ. [Screening of hearing in children up to 18 months of age]. *Ned Tijdschr Geneesk* 2000;144:589-593.
- (17) Kemper AR, Downs SM. A cost-effectiveness analysis of newborn hearing screening strategies. *Arch Pediatr Adolesc Med* 2000;154:484-488.
- (18) Keren R, Helfand M, Homer C, McPhillips H, Lieu TA. Projected cost-effectiveness of statewide universal newborn hearing screening. *Pediatrics* 2002;110:855-864.
- (19) Yoshinaga-Itano C, Apuzzo ML. Identification of hearing loss after age 18 months is not early enough. *Am Ann Deaf* 1998;143:380-387.
- (20) Oudesluys-Murphy AM, Harlaar J. Neonatal hearing screening with an automated auditory brainstem response screener in the infant's home. *Acta Paediatr* 1997;86:651-655.
- (21) Yoshinaga-Itano C, Sedey AL, Coulter DK, Mehl AL. Language of early- and later-identified children with hearing loss. *Pediatrics* 1998;102:1161-1171.
- (22) Krishnan LA. Universal newborn hearing screening follow-up: a university clinic perspective. *Am J Audiol* 2009;18:89-98.
- (23) Uus K, Bamford J, Taylor R. An analysis of the costs of implementing the National Newborn Hearing Screening Programme in England. *J Med Screen* 2006;13:14-19.
- (24) Bischof HJ. Behavioral and neuronal aspects of developmental sensitive periods. *Neuroreport* 2007;18:461-465.
- (25) Sharma A, Nash AA, Dorman M. Cortical development, plasticity and re-organization in children with cochlear implants. *J Commun Disord* 2009;42:272-279.
- (26) Zanten GAv, Zanten. Oto-akoestische emissies. In: Meuwese-Jongejouw J, Straaten HLM, eds. *Werkboek Neonatale gehoorscreening*; 2008.
- (27) van Straaten HL, Hille ET, Kok JH, Verkerk PH. Implementation of a nation-wide automated auditory brainstem response hearing screening programme in neonatal intensive care units. *Acta Paediatr* 2003;92:332-338.
- (28) Yoshinaga-Itano C. Early intervention after universal neonatal hearing screening: impact on outcomes. *Ment Retard Dev Disabil Res Rev* 2003;9:252-266.
- (29) Yoshinaga-Itano C. Benefits of early intervention for children with hearing loss. *Otolaryngol Clin North Am* 1999;32:1089-1102.
- (30) Yoshinaga-Itano C, Coulter D, Thomson V. The Colorado Newborn Hearing Screening Project: effects on speech and language development for children with hearing loss. *J Perinatol* 2000;20:S132-S137.

- (31) Kennedy CR, McCann DC, Campbell MJ et al. Language ability after early detection of permanent childhood hearing impairment. *N Engl J Med* 2006;354:2131-2141.
- (32) Moeller MP. Early intervention and language development in children who are deaf and hard of hearing. *Pediatrics* 2000;106:E43.
- (33) Wake M, Hughes EK, Poulakis Z, Collins C, Rickards FW. Outcomes of children with mild-profound congenital hearing loss at 7 to 8 years: a population study. *Ear Hear* 2004;25:1-8.
- (34) Wake M, Poulakis Z, Hughes EK, Carey-Sargeant C, Rickards FW. Hearing impairment: a population study of age at diagnosis, severity, and language outcomes at 7-8 years. *Arch Dis Child* 2005;90:238-244.
- (35) Nicholas JG, Geers AE. Effects of early auditory experience on the spoken language of deaf children at 3 years of age. *Ear Hear* 2006;27:286-98.
- (36) Wolff R, Hommerich J, Riemsma R, Antes G, Lange S, Kleijnen J. Hearing screening in newborns: systematic review of accuracy, effectiveness, and effects of interventions after screening. *Arch Dis Child* 2010;95:130-135.
- (37) Thompson DC, McPhillips H, Davis RL, Lieu TL, Homer CJ, Helfand M. Universal newborn hearing screening: summary of evidence. *JAMA* 2001;286:2000-2010.
- (38) Vandebroucke JP. When are observational studies as credible as randomised trials? *Lancet* 2004;363:1728-1731.
- (39) Martens EP, Pestman WR, de BA, Belitser SV, Klungel OH. Instrumental variables: application and limitations. *Epidemiology* 2006;17:260-267.
- (40) Hertogh EM, Schuit AJ, Peeters PH, Monninkhof EM. Noncompliance in lifestyle intervention studies: the instrumental variable method provides insight into the bias. *J Clin Epidemiol* 2010.
- (41) Rassen JA, Brookhart MA, Glynn RJ, Mittleman MA, Schneeweiss S. Instrumental variables I: instrumental variables exploit natural variation in nonexperimental data to estimate causal relationships. *J Clin Epidemiol* 2009;62:1226-1232.
- (42) Morton CC, Nance WE. Newborn hearing screening—a silent revolution. *N Engl J Med* 2006;354:2151-2164.
- (43) Year 2007 position statement: Principles and guidelines for early hearing detection and intervention programs. *Pediatrics* 2007;120:898-921.
- (44) Binda S, Caroppo S, Dido P et al. Modification of CMV DNA detection from dried blood spots for diagnosing congenital CMV infection. *J Clin Virol* 2004;30:276-279.
- (45) Korver AM, de Vries JJ, konings S et al. DECIBEL study: Congenital cytomegalovirus infection in young children with permanent bilateral hearing impairment in the Netherlands. *J Clin Virol* 2009;46 Suppl 4:S27-S31.

- (46) Juni P, Witschi A, Bloch R, Egger M. The hazards of scoring the quality of clinical trials for meta-analysis. *JAMA* 1999;282:1054-1060.
- (47) Stroup DF, Berlin JA, Morton SC et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA* 2000;283:2008-2012.
- (48) Kelsell DP, Dunlop J, Stevens HP et al. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997;387:80-83.
- (49) Admiraal RJ, Huygen PL. Changes in the aetiology of hearing impairment in deaf-blind pupils and deaf infant pupils at an institute for the deaf. *Int J Pediatr Otorhinolaryngol* 2000;55:133-142.
- (50) Bajaj Y, Sirimanna T, Albert DM et al. Causes of deafness in British Bangladeshi children: a prevalence twice that of the UK population cannot be accounted for by consanguinity alone. *Clin Otolaryngol* 2009;34:113-119.
- (51) Billings KR, Kenna MA. Causes of pediatric sensorineural hearing loss: yesterday and today. *Arch Otolaryngol Head Neck Surg* 1999;125:517-521.
- (52) Dietz A, Lopponen T, Valtonen H, Hyvarinen A, Lopponen H. Prevalence and etiology of congenital or early acquired hearing impairment in Eastern Finland. 2009.
- (53) Egeli E, Cicekci G, Silan F et al. Etiology of deafness at the Yeditepe School for the deaf in Istanbul. *Int J Pediatr Otorhinolaryngol* 2003;67:467-471.
- (54) Fortnum HM, Marshall DH, Summerfield AQ. Epidemiology of the UK population of hearing-impaired children, including characteristics of those with and without cochlear implants-audiology, aetiology, comorbidity and affluence. *Int J Audiol* 2002;41:170-179.
- (55) Nekahm D, Weichbold V, Welzl-Muller K. Epidemiology of permanent childhood hearing impairment in the Tyrol, 1980-94. *Scand Audiol* 2001;30:197-202.
- (56) Uus K, Davis AC. Epidemiology of permanent childhood hearing impairment in Estonia, 1985-1990. *Audiology* 2000;39:192-197.
- (57) Fowler KB, Pass RF. Risk factors for congenital cytomegalovirus infection in the offspring of young women: exposure to young children and recent onset of sexual activity. *Pediatrics* 2006;118:e286-e292.
- (58) Fowler KB, McCollister FP, Dahle AJ, Boppana S, Britt WJ, Pass RF. Progressive and fluctuating sensorineural hearing loss in children with asymptomatic congenital cytomegalovirus infection. *J Pediatr* 1997;130:624-630.
- (59) Fowler KB, Dahle AJ, Boppana SB, Pass RF. Newborn hearing screening: will children with hearing loss caused by congenital cytomegalovirus infection be missed? *J Pediatr* 1999;135:60-64.

- (60) Barbi M, Binda S, Primache V et al. Cytomegalovirus DNA detection in Guthrie cards: a powerful tool for diagnosing congenital infection. *J Clin Virol* 2000;17:159-165.
- (61) de Vries JJ, Claas EC, Kroes AC, Vossen AC. Evaluation of DNA extraction methods for dried blood spots in the diagnosis of congenital cytomegalovirus infection. *J Clin Virol* 2009;46 Suppl 4:S37-S42.
- (62) Kalpoe JS, Kroes AC, de J et al. Validation of clinical application of cytomegalovirus plasma DNA load measurement and definition of treatment criteria by analysis of correlation to antigen detection. *J Clin Microbiol* 2004;42:1498-1504.
- (63) Barbi M, Binda S, Primache V, Luraschi C, Corbetta C. Diagnosis of congenital cytomegalovirus infection by detection of viral DNA in dried blood spots. *Clin Diagn Virol* 1996;6:27-32.
- (64) Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol* 1993;46:1417-1432.
- (65) Kenneson A, Cannon MJ. Review and meta-analysis of the epidemiology of congenital cytomegalovirus (CMV) infection. *Rev Med Virol* 2007;17:253-276.
- (66) Yamamoto AY, Mussi-Pinhata MM, Pinto PC, Figueiredo LT, Jorge SM. Usefulness of blood and urine samples collected on filter paper in detecting cytomegalovirus by the polymerase chain reaction technique. *J Virol Methods* 2001;97:159-164.
- (67) Atkinson C, Walter S, Sharland M et al. Use of stored dried blood spots for retrospective diagnosis of congenital CMV. *J Med Virol* 2009;81:1394-1398.
- (68) Statistics Netherlands. <http://statline.cbs.nl/StatWeb/publication/?DM=SLEN&PA=37259eng&D1=1,3,8-9,14,16,21-22&D2=0&D3=0&D4=43-45&LA=EN&HDR=G3,T&STB=G1,G2&VW=T>.
- (69) Staras SA, Dollard SC, Radford KW, Flanders WD, Pass RF, Cannon MJ. Seroprevalence of cytomegalovirus infection in the United States, 1988-1994. *Clin Infect Dis* 2006;43:1143-1151.
- (70) Grosse SD, Ross DS, Dollard SC. Congenital cytomegalovirus (CMV) infection as a cause of permanent bilateral hearing loss: a quantitative assessment. *J Clin Virol* 2008;41:57-62.
- (71) Ogawa H, Suzutani T, Baba Y et al. Etiology of severe sensorineural hearing loss in children: independent impact of congenital cytomegalovirus infection and GJB2 mutations. *J Infect Dis* 2007;195:782-788.
- (72) Barbi M, Binda S, Caroppo S, Ambrosetti U, Corbetta C, Sergi P. A wider role for congenital cytomegalovirus infection in sensorineural hearing loss. *Pediatr Infect Dis J* 2003;22:39-42.

- (73) Harris S, Ahlfors K, Ivarsson S, Lernmark B, Svanberg L. Congenital cytomegalovirus infection and sensorineural hearing loss. *Ear Hear* 1984;5:352-355.
- (74) Foulon I, Naessens A, Foulon W, Casteels A, Gordts F. A 10-year prospective study of sensorineural hearing loss in children with congenital cytomegalovirus infection. *J Pediatr* 2008;153:84-88.
- (75) Stehel EK, Shoup AG, Owen KE et al. Newborn hearing screening and detection of congenital cytomegalovirus infection. *Pediatrics* 2008;121:970-975.
- (76) Hicks T, Fowler K, Richardson M, Dahle A, Adams L, Pass R. Congenital cytomegalovirus infection and neonatal auditory screening. *J Pediatr* 1993;123:779-782.
- (77) Conboy TJ, Pass RF, Stagno S et al. Early clinical manifestations and intellectual outcome in children with symptomatic congenital cytomegalovirus infection. *J Pediatr* 1987;111:343-348.
- (78) Casteels A, Naessens A, Gordts F, De CL, Bougatet A, Foulon W. Neonatal screening for congenital cytomegalovirus infections. *J Perinat Med* 1999;27:116-121.
- (79) Cannon MJ, Davis KF. Washing our hands of the congenital cytomegalovirus disease epidemic. *BMC Public Health* 2005;5:70.
- (80) Picone O, Vauloup-Fellous C, Cordier AG et al. A 2-year study on cytomegalovirus infection during pregnancy in a French hospital. *BJOG* 2009;116:818-823.
- (81) Harvey J, Dennis CL. Hygiene interventions for prevention of cytomegalovirus infection among childbearing women: systematic review. *J Adv Nurs* 2008;63:440-450.
- (82) Jeon J, Victor M, Adler SP et al. Knowledge and awareness of congenital cytomegalovirus among women. *Infect Dis Obstet Gynecol* 2006;2006:80383.
- (83) Center for Disease Control. Morbidity and Mortality Weekly Report: Knowledge and practices of obstetricians and gynecologists regarding cytomegalovirus infection during pregnancy-United States 2007. Volume 57, number 3. 2008.
- (84) Read JS, Cannon MJ, Stanberry LR, Schuval S. Prevention of mother-to-child transmission of viral infections. *Curr Probl Pediatr Adolesc Health Care* 2008;38:274-297.
- (85) Biernacki P, Waldorf D. Snowball sampling: Problems and techniques of chain referral sampling. *Social Methods Research* 1981;2:141-63.
- (86) Kimberlin DW, Lin CY, Sanchez PJ et al. Effect of ganciclovir therapy on hearing in symptomatic congenital cytomegalovirus disease involving the central nervous system: a randomized, controlled trial. *J Pediatr* 2003;143:16-25.
- (87) Sininger YS, Grimes A, Christensen E. Auditory Development in Early Amplified Children: Factors Influencing Auditory-Based Communication Outcomes in Children With Hearing Loss. *Ear Hear* 2010; 31: 166-185.

- (88) Nelson HD, Bougatsos C, Nygren P. Universal newborn hearing screening: systematic review to update the 2001 US Preventive Services Task Force Recommendation. *Pediatrics* 2008;122:e266-e276.
- (89) Yoshinaga-Itano C. Early intervention after universal neonatal hearing screening: impact on outcomes. *Ment Retard Dev Disabil Res Rev* 2003;9:252-266.
- (90) Calderon C, Naidu S. Further support of the benefits of early identification and intervention with children with hearing loss [abstract]Calderon C, Naidu S. *Volta Rev* 2000;53:84
- (91) Moeller MP, Tomblin JB, Yoshinaga-Itano C, Connor CM, Jerger S. Current state of knowledge: language and literacy of children with hearing impairment. *Ear Hear* 2007;28:740-753.
- (92) Kauffman-de Boer MA, Uilenburg N, Schuitema T, Vinks E, Van den Brink G, van der PK et al. Landelijke Implementatie Neonatale Gehoorscreening. 2006.
- (93) Straaten HLM, Dommelen van P, Verkerk PH. Jaarverslag neonatale gehoorscreening in de Neonatale Intensive Care Units 2007.
- (94) Kennedy C, McCann D, Campbell MJ, Kimm L, Thornton R. Universal newborn screening for permanent childhood hearing impairment: an 8-year follow-up of a controlled trial. *Lancet* 2005;366:660-662.
- (95) Hille ET, Elbertse L, Gravenhorst JB, Brand R, Verloove-Vanhorick SP. Nonresponse bias in a follow-up study of 19-year-old adolescents born as preterm infants. *Pediatrics* 2005;116:e662-e666.
- (96) Mason CA, Gaffney M, Green DR, Grosse SD. Measures of follow-up in early hearing detection and intervention programs: a need for standardization. *Am J Audiol* 2008;17:60-67.
- (97) Yoshinaga-Itano C, Coulter D, Thomson V. Developmental outcomes of children with hearing loss born in Colorado hospitals with and without universal newborn hearing screening programs. *Semin Neonatol* 2001;6:521-529.
- (98) Yoshinaga-Itano C. From Screening to Early Identification and Intervention: Discovering Predictors to Successful Outcomes for Children With Significant Hearing Loss. *J Deaf Stud Deaf Educ* 2003;8:11-30.
- (99) Sininger YS, Martinez A, Eisenberg L, Christensen E, Grimes A, Hu J. Newborn hearing screening speeds diagnosis and access to intervention by 20-25 months. *J Am Acad Audiol* 2009;20:49-57.
- (100) Uus K, Bamford J. Effectiveness of population-based newborn hearing screening in England: ages of interventions and profile of cases. *Pediatrics* 2006;117:e887-e893.

- (101) Young A, Gascon-Ramos M, Campbell M, Bamford J. The design and validation of a parent-report questionnaire for assessing the characteristics and quality of early intervention over time. *J Deaf Stud Deaf Educ* 2009;14:422-435.
- (102) Ireton H, Glascoe FP. Assessing children's development using parents' reports. The Child Development Inventory. *Clin Pediatr (Phila)* 1995;34:248-255.
- (103) Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol* 1993;46:1417-1432.
- (104) Zink I, Lejaegere M. NCDI-3, *aanpassing en hernormering van MacArthur CDI*. 9789033466632 ed. Acco, 2007.
- (105) Varni JW, Seid M, Kurtin PS. PedsQL 4.0: reliability and validity of the Pediatric Quality of Life Inventory version 4.0 generic core scales in healthy and patient populations. *Med Care* 2001;39:800-812.
- (106) Engelen V, Haentjens MM, Detmar SB, Koopman HM, Grootenhuis MA. Health related quality of life of Dutch children: psychometric properties of the PedsQL in the Netherlands. *BMC Pediatr* 2009;9:68.
- (107) Korver AMH, Konings S, Meuwese-Jongejeugd J et al. Newborn hearing screening: who is unidentified? 2010.
- (108) Moeller MP. Current state of knowledge: psychosocial development in children with hearing impairment. *Ear Hear* 2007;28:729-739.
- (109) Wolff R, Hommerich J, Riemsma R, Antes G, Lange S, Kleijnen J. Hearing Screening in Newborns. Systematic Review of Accuracy, Effectiveness, and Effects of Interventions after Screening. *Arch Dis Child* 2009.
- (110) Yoshinaga-Itano C, Coulter D, Thomson V. Developmental outcomes of children with hearing loss born in Colorado hospitals with and without universal newborn hearing screening programs. *Semin Neonatol* 2001;6:521-529.
- (111) Fellinger J, Holzinger D, Sattel H, Laucht M. Mental health and quality of life in deaf pupils. *Eur Child Adolesc Psychiatry* 2008;17:414-423.

ABBREVIATIONS

A-ABR	Automated Auditory Brainstem Response
AC	Audiology Center (Speech and Hearing Center)
CANG	Central Administration program Neonatal Hearing screening
CBS	Statistics Netherlands
CDI	Child Development Inventory
CMV	Cytomegalovirus
dB	Decibel
DECIBEL-study	Developmental Evaluation of Children: Impact and Benefits of Early hearing screening strategies Leiden
DHS	Distraction Hearing Screening
DBS	Dried blood spot
ELQ	Expressive Language Quotient
GJB2	Gap Junction protein Beta 2
IQR	Interquartile range
JCIH	Joint Committee on Infant Hearing
GDQ	General Development Quotient
LCQ	Language Comprehension Quotient
MCDI	Minnesota Child Development Inventory
MLU	Mean Length of longest Utterance
NHS	Newborn Hearing Screening
NICU	Neonatal Intensive Care Unit
NSDSK	Dutch Foundation for the Deaf and Hard of Hearing Child
PedsQL™ 4.0	Parental Quality of Life Inventory™ 4.0
PCHI	Permanent Childhood Hearing Impairment
PCR	Polymerase Chain Reaction
QoL	Quality of Life
RIVM	National Institute for Public Health and the Environment
SNHL	Sensorineural hearing loss
SNP	Single Nucleotide Polymorphism
YHC	Youth Health Care organization

MEMBERS OF THE DECIBEL- COLLABORATIVE STUDY GROUP

INITIATORS LEIDEN UNIVERSITY MEDICAL CENTER

H.M. Oudesluys-Murphy, Department of Paediatrics
J.H.M. Frijns, Department of Oto-rhinolaryngology
F.W. Dekker, Department of Clinical Epidemiology

EXECUTIVE AND WRITING COMMITTEE LEIDEN UNIVERSITY MEDICAL CENTER

A.M.H. Korver, Department of Paediatrics
S. Konings, Department of Oto-rhinolaryngology
H.M. Oudesluys-Murphy, Department of Paediatrics
J.H.M. Frijns, Department of Oto-rhinolaryngology
F.W. Dekker, Department of Clinical Epidemiology

OTHER WRITING PARTICIPANTS

126

C.C. Wever and W. Beers, Department of Oto-rhinolaryngology, LUMC
J.J.C de Vries and A.C.T.M. Vossen, Department of Medical Microbiology, LUMC
J.W. de Jong, medical student
S.G. Kant, Department of Clinical Genetics, LUMC
H.L.M. van Straaten, Isala Clinics, Zwolle
N.N. Uilenburg, Dutch Foundation for the Deaf and Hard of Hearing Child
J. Meuwese-Jongejeugd, National Institute for Public Health and Environment, Bilthoven
R.J.C. Admiraal, Department of Oto-rhinolaryngology, Radboud University Nijmegen Medical Centre
H.P.M. Kunst, Department of Oto-rhinolaryngology, Radboud University Nijmegen Medical Centre

OTHER PARTNERS

B. Elvers and G. Loeber, National Institute for Public Health and Environment
C. Rieffe, Faculty of Social Sciences, Developmental Psychology, Leiden University
M.H. Ens-Dokkum, Royal Dutch Kentalis, Zoetermeer
Medical students: L. Seekles, A. Thijssen, A. Lievense, M. van Egdom-van der Wind,
S.C.P.M. Theunissen, S.H. Mooij and C.S. Schouten

PARTICIPATING SPEECH AND HEARING CENTERS

Academic Medical Center: M.J. Maré
Audiology Center Eindhoven: C.T. ter Huurne
Audiology Center Holland Noord: P. Brienesse
Audiology Center Tilburg and Breda: F. Jorritsma and M. van Beurden
Erasmus Medical Center -Sophia Children's Hospital: A. Goedegebuure
Hoensbroeck Audiology Center: D. Pans
Leiden University Medical Center: W. Soede
Pento Audiology Centers: G. Rijpma, W. Graef, D. Linschoten, J. Kuijper, N.J. Hof and
G.J. Koldewijn
Radboud University Nijmegen Medical Centre: R.J.C. Admiraal, C.W.R.J. Cremers and
H.P.M. Kunst
Royal Dutch Kentalis: J. Dijkhuizen, M. Scharloo and D. Hoeben
Royal Auris Group: M. de Leeuw
University Medical Center Utrecht, Wilhelmina Children's Hospital: G.A. van Zanten
University Medical Center Groningen: F. Coster and P. van Dijk
VU University Medical Center: S.T. Goverts

CURRICULUM VITAE

130

ANNA MARIA HELENA (MARLEEN) KORVER
25 september 1979, Echteld

1991-1997

Voortgezet Wetenschappelijk Onderwijs
Christelijk Lyceum Veenendaal

1997-1999

Gezondheidswetenschappen
Beleid en Management van de Gezondheidszorg
Universiteit Maastricht

1999-2004

Geneeskunde, Universiteit Maastricht
Arts-examen Cum Laude

2004

Arts-assistent in de zorg voor
verstandelijk gehandicapten
's Heerenloo Midden Nederland

2005-2007

Arts-assistent niet in opleiding
Kindergeneeskunde
Gelderse Vallei Ziekenhuis Ede
Juliana Kinderziekenhuis Den Haag
Leids Universitair Medisch Centrum

2007- 2010

Wetenschappelijk onderzoek
Sociale Pediatrie
Leids Universitair Medisch Centrum

Per oktober 2010

Kinderarts in opleiding
Leids Universitair Medisch Centrum
(opleider: Prof. Dr. H. A. Delemarre-van de Waal)

LIST OF PUBLICATIONS

134

Korver, A.M.H., De Vries, J.J.C., Konings S., De Jong, J.W., Dekker F.W. Vossen, A.C.T.M., Frijns, J.H.M., Oudesluys-Murphy A.M. and the DECIBEL collaborative study Group. DECIBEL study: Congenital cytomegalovirus infection in young children with permanent bilateral hearing impairment in the Netherlands. *J Clin Virol.* 2009;46 Suppl S27-S31.

Korver, A.M.H., De Vries, J.J.C., De Jong, J.W., Dekker F.W., Vossen, A.C.T.M. and Oudesluys-Murphy A.M. Awareness concerning congenital cytomegalovirus among doctors in the Netherlands. *J Clin Virol.* 2009;46 Suppl S11-S15.

Korver, A.M.H., Meeuwese, J., Briaire, J.J., Frijns, J.H.M. en Oudesluys-Murphy, A.M. Pre-linguaal gehoorverlies na een gunstige screeningsuitslag. In Werkboek "Neonatale gehoorschreeing", VU University Press November 2008.

Diepen van, M.M.L., **Korver, A.M.H.**, Ens-Dokkum, M.H. en Oudesluys-Murphy, A.M. Etiologie van verworven gehoorverlies. In Werkboek Neonatale gehoorschreeing, VU University Press November 2008.

Korver, A.M.H., Meeuwese, J., Briaire, J.J., Frijns, J.H.M. en Oudesluys-Murphy, A.M. Permanent gehoorverlies in de prelinguale fase bij kinderen met een niet-afwijkende uitslag bij de neonatale gehoorschreeing. *Ned. Tijdschr Geneeskd.* 2008; 152; 425-30.

Korver, A.M.H., Freund, M.W., Haas, F. and Strengers, J.L.M. Pentalogy of Cantrell : Successful Early Correction. *Pediatr Cardiol.* 2008 Jan; 29(1):146-149.

Peek, F.A.W., Briaire, J.J., Mylanus, E.A.M., **Korver, A.M.H.**, Mulder, J.J.S., Snik, A.F.M., Oudesluys-Murphy, A.M. en Frijns, J.H.M. Ingezonden reactie op artikel van Merkus P, Furth van AM, Suer M, Doofheid na bacteriële meningitis bij jonge kinderen: ingrijpen vóór de cochlea is geoblitereerd *Ned Tijdschr Geneeskd.* 2007;151(22) 1209-1213. *Ned Tijdschr Geneeskd.* 2007 Okt 6;151(40): 2239-2240

Korver, A.M.H. en Moons, P. Diagnose in beeld: Een meisje met hemorragische bullae. *Ned Tijdschr Geneeskd.* 2007; 151; 1403

Korver, A.M.H., Walther F.J. en de Beaufort, A.J. Ernstige complicaties van een navelvenelin. *Ned Tijdschr Geneeskd.* 2007; 151; 2219-23.

