



Universiteit
Leiden
The Netherlands

Circadian timekeeping: from basic clock function to implications for health

Lucassen, Eliane Alinda

Citation

Lucassen, E. A. (2016, March 31). *Circadian timekeeping: from basic clock function to implications for health*. Retrieved from <https://hdl.handle.net/1887/38651>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/38651>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/38651> holds various files of this Leiden University dissertation

Author: Lucassen, Eliane Alinda

Title: Circadian timekeeping: from basic clock function to implications for health

Issue Date: 2016-03-31



CHAPTER 2

Interacting epidemics? Sleep curtailment, insulin resistance, and obesity

Eliane A. Lucassen,^{1,2} Kristina I. Rother,³ and Giovanni Cizza⁴

Ann N Y Acad Sci 2012, 1264: 110-34.

1. Clinical Center, National Institutes of Health, Bethesda, United States.
2. Laboratory of Neurophysiology, Department of Molecular Cell Biology, Leiden University Medical Center, Leiden, the Netherlands.
3. Section on Pediatric Diabetes and Metabolism, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, United States.
4. Section on Neuroendocrinology of Obesity, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, United States.

Abstract

In the last 50 years, the average self-reported sleep duration in the United States has decreased by 1.5–2 hrs in parallel with an increasing prevalence of obesity and diabetes. Epidemiological studies and meta-analyses report a strong relationship between short or disturbed sleep, obesity, and abnormalities in glucose metabolism. This relationship is likely to be bidirectional and causal in nature, but many aspects remain to be elucidated. Sleep and the internal circadian clock influence a host of endocrine parameters. Sleep curtailment in humans alters multiple metabolic pathways, leading to more insulin resistance, possibly decreased energy expenditure, increased appetite, and immunological changes. On the other hand, psychological, endocrine, and anatomical abnormalities in individuals with obesity and/or diabetes can interfere with sleep duration and quality, thus creating a vicious cycle. In this review, we address mechanisms linking sleep with metabolism, highlight the need for studies conducted in real-life settings, and explore therapeutic interventions to improve sleep, with a potential beneficial effect on obesity and its comorbidities.



Introduction

According to the National Health and Nutrition Examination Survey (NHANES) from 2009 to 2010, 33% of adult Americans were overweight (body mass index (BMI) of 25.0–29.9 kg/m²), and an even larger proportion (36%) were obese (BMI \geq 30.0 kg/m²).¹ The report contained good and bad news: for more than a decade, the age-adjusted prevalence of obesity had not changed in women overall, but there were marked ethnic, gender, and socioeconomic differences. Mexican American and Black women and men continued to show a significant rise in the prevalence of obesity. These figures are reflected in important obesity-related complications, such as hypertension, hyperlipidemia, and type 2 diabetes, all leading causes of cardiovascular disease.² Type 2 diabetes affects 8.3% of the U.S. population (27% of U.S. residents aged 65 yrs and older), whereas 35% of U.S. adults have fasting glucose and glycosylated hemoglobin (HbA1c) levels in the prediabetic range.³ Thus, it is of great importance to characterize contributing etiological factors in order to effectively counteract the obesity and diabetes epidemics. The exact mechanisms explaining the relationship between short sleep, obesity, and diabetes remain unclear.

Accumulating evidence has pointed to an association between sleep curtailment and obesity,^{4,5} type 2 diabetes,^{6–9} and possibly mortality (see Ref. 10 for a review). In the last 50 years, self-reported sleep duration in the United States has decreased by 1.5–2 hrs due to lifestyle changes in our society.¹¹ Worldwide, secular trends in adult sleep duration may be more variable. A recent review article summarizing 12 studies from 15 countries from 1960 to 2000 did not find a consistent decrease in self-reported adult sleep duration: self-reported sleep duration had increased in seven countries, decreased in six, and results were inconsistent for the United States and Sweden.¹² Imprecision in assessing sleep duration, as well as socioeconomic and demographic factors may play a role in this heterogeneity.

In this paper, we will first present the epidemiologic evidence for these relationships; we will then focus on sleep physiology and its role in metabolic functions. Then, we will critically evaluate the studies of sleep manipulation on glucose metabolism and energy expenditure (EE). We will highlight endocrine parameters that may influence these processes, including thyroid hormones, growth hormone (GH), cortisol, sympathovagal balance, and the immune system. We will also delineate the relationships between short sleep and some of the hormones that control appetite, including leptin, ghrelin, orexin, and neuropeptide Y (NPY). In addition, we will present evidence for the reverse effect: of obesity/diabetes on sleep. Finally, we will describe emerging evidence for a role of melatonin in the sleep–metabolism relationship and its therapeutic potential in metabolic disturbances.

Epidemiologic evidence for an association among sleep duration, obesity, and diabetes

A meta-analysis of 45 cross-sectional studies, including 604,509 adults and 30,002 children, confirmed the relationship between short sleep (generally fewer than five hrs per night in

adults and fewer than 10 hrs per night in children) and obesity, and quantified the associated risk (OR 1.55; 1.43–1.68; OR 1.89; 1.46–2.43, for adults and children, respectively).¹³ A similar OR of 1.58 (1.26–1.98) was found in children in another meta-analysis.¹⁴ Most of the included studies used actigraphy or self-reported sleep duration. A polysomnography study, not included in these meta-analyses, of 2,700 men above the age of 65 yrs revealed an inverse relationship between slow wave sleep (SWS) duration and BMI or waist circumference.¹⁵ Another study reported that obese adults (BMI 41 ± 1 kg/m²) without sleep apnea slept 88 min fewer than lean subjects.¹⁶ The relationship between short sleep and BMI was also confirmed by actigraphy in 612 participants (35–50 yrs old).¹⁷

In principle, causality is best addressed in prospective randomized controlled studies. This ideal approach may not be practical in this case: studies of sleep extension or curtailment cannot be blinded, and participants may not behave according to the allocation group (i.e. control subjects changing sleep duration or, *vice versa*, intervention subjects not changing sleep duration). For these reasons, the existing prospective studies have all been observational in nature. Short sleepers experience greater weight increases over time according to most,^{4,18–22} but not all studies.^{17,23,24} However, average weight gain was modest and of relative clinical meaning. Short sleepers (fewer than five to six hours) gained 2 kg more in a 6-year study,¹⁹ and 0.4 kg more in another 16-year study.¹⁸ In addition, subjects had a 35% and 31% greater chance of gaining more than 5 kg in 6 yrs, and gaining more than 15 kg in 16 yrs, respectively.^{18,19} Thus, there is a large variability among studies in weight gain associated with short sleep for unknown reasons: the magnitude of this effect may be gender dependent or depend on other yet-to-be identified genetic and environmental factors. For example, short sleep may affect metabolism differently in natural “short sleepers” vs. chronically sleep-deprived individuals. A large study including 35,247 subjects found that lean men, but not lean women, who slept fewer than five hrs had increased odds of becoming overweight within one year (OR 1.91; 1.36–2.67).²⁵ In contrast, in a cohort of 3,576 elderly subjects, women, but not men, who slept fewer than five hours, had a higher risk of gaining 5 kg in two yrs (OR 3.41; 1.31–8.69).²⁰ Interestingly, in this study, BMI was also increased in women sleeping more than eight hrs. A similar U-shaped association has been reported in other studies.^{19,25} Collectively, these studies indicate that self-reported short or long sleep is associated with increased BMI. The amount of weight gain over time, however, varied greatly, ranging from less than 1 kg to several kg, and age may be one of the parameters modulating this association.

Many mechanisms may be responsible for the frequently observed insulin resistance in obese individuals, including increased release of free fatty acids, leptin, and TNF- α from adipose tissue.²⁶ We suggest that short sleep (fewer than five to six hours) may now qualify as an additional clinical factor in the development of insulin resistance and diabetes, as shown in several cross-sectional studies.^{27–29} A recent study of 174,542 middle-aged subjects

showed that short sleepers had an OR of 1.46 (1.31–1.63) for developing diabetes after 3–10 yrs of follow-up.³⁰ A meta-analysis of 10 prospective reported a pooled OR of 1.28 (1.03–1.60) for diabetes in short sleepers.³¹ Interestingly, this correlation was only present in men (OR 2.07; 1.16–3.72). Both long sleep (OR 1.48; 1.13–1.96) and decreased sleep efficiency also resulted in greater chances of developing diabetes.³¹ Difficulty in falling asleep resulted in a greater incidence of diabetes after 4.2–14.8 yrs of follow up.^{32–34} In smaller studies, the association between sleep duration and diabetes is more variable.^{7–9,35} In addition, this variability may be real and differ on the basis of ethnic group and socioeconomic status: a study found increased risk of developing diabetes only in Whites and Hispanics who slept fewer than seven hrs.³⁶ Besides sleep-duration, poor sleep quality relates to insulin resistance. Frequency of nocturnal awakenings, as measured by actigraphy, correlated positively with HbA1c levels in patients with type 2 diabetes.³⁷ Furthermore, sleep disordered breathing was related to worse blood glucose control.³⁸ Taken together, there is solid epidemiologic evidence for an interconnection between sleep duration (too short or too long), insulin resistance, and obesity.



Physiological changes of metabolism during normal sleep

Sleep architecture, the structure and pattern of sleep, is best described by a combined approach that uses electroencephalographic, electromyographic, and other criteria (**Fig. 1**).³⁹ Many metabolic parameters display diurnal rhythms, which are the resultant of both homeostatic and circadian mechanisms. From a homeostatic point of view, sleep pressure increases with sustained wakefulness. Circadian rhythms are endogenously generated in the suprachiasmatic nucleus of the hypothalamus by feedback loops of transcripts of a group of “core clock” genes, including the CLOCK gene.^{40,41} Timekeeping signals are passed on from the suprachiasmatic nucleus to oscillators in peripheral tissues also containing core clock genes. Endogenous rhythms, in turn, are synchronized to the outside world by various environmental clues, mostly by light via the retina. During sleep, energy requirements are lower as metabolic needs for processes such as breathing, gut motility, heart rate and muscular activity, are decreased.⁴² Indeed, resting EE, as determined by indirect calorimetry, was lower in healthy volunteers during sleep compared to wakefulness.⁴³

During sleep glucose levels remain stable or fall only minimally in spite of fasting, mostly because of decreased energy needs.⁴⁴ In contrast, if subjects are kept awake in bed while fasting for a comparable number of hours, glucose levels do fall, approximately 15 mg/dl.⁴⁵ During the first part of the night, when most slow wave sleep (SWS) occurs, glucose production is diminished (**Fig. 2**).⁴⁴ Brain glucose metabolism, as measured by PET, decreases by 30–40%.^{46,47} During the second half of the night, glucose consumption increases due to increased REM sleep requiring more energy.^{46,47}




Fig. 1. Representative sleep architecture during eight hrs of uninterrupted sleep (adapted from Ref. 39). Sleep can be divided into rapid eye movement (REM) and nonrapid eye movement (NREM) sleep. On average, adults spend 75–80% in NREM sleep, which can be further subdivided into stages 1 to 4.

Stages 3 and 4 are often referred to as slow-wave sleep (SWS) because of the appearance of well-defined waves of 0.5–4.0 Hz. In REM sleep, a phasic and a tonic phase can be distinguished. Skeletal muscles are atonic or hypotonic during REM sleep (except for the diaphragm, extraocular, and sphincter muscles), but bursts of muscle activity can occur during the phasic phase. Furthermore, dreams mostly take place during REM sleep and are more complex during this stage. Eye movement is only present during the phasic phase. Throughout a normal night of sleep, there are three to five sleep cycles, from NREM (stage 1 to 4) to REM, each cycle taking 90–120 minutes. Humans display less SWS and more REM sleep toward the end of the night.

EE and glucose metabolism are modulated by a number of hormones. Thyroid hormones are important determinants of EE; cortisol and GH are powerful modulators of circadian rhythms of glucose metabolism. The sympathovagal system and adiponectin influence both EE and glucose metabolism. Thyroid stimulating hormone (TSH) levels rise at night, mostly due to increased hypothalamic secretion of thyroid releasing hormone (TRH) (**Fig. 2**).⁴⁸

GH secretion is under dual hypothalamic control: somatostatin is inhibitory and growth hormone releasing hormone (GHRH) is stimulatory. Somatostatin and GHRH are secreted in alternation, thus generating the typical pulsatile pattern of GH secretion. GH typically peaks at sleep onset during SWS (**Fig. 2**).⁴⁹ GH levels were lower in SWS-deprived individuals⁵⁰ and higher in individuals with pharmacologically induced SWS.⁵¹ In addition, REM sleep deprivation did not affect plasma GH levels.⁵² Intravenous GH administration decreases SWS, suggesting a modulatory role for GH on SWS.^{53,54} The GH peak at sleep onset has insulin lowering effects.⁵⁵ GHRH injection increased glucose levels at awakening by almost 50%⁵⁶ and GH administration rapidly decreased muscular glucose uptake.^{55,57}

Cortisol concentrations reach their zenith in the morning, experience a gradual fall during the day, which is briefly interrupted by meals, and have their nadir around 3 am (**Fig. 2**).⁴⁸ Cortisol impairs insulin sensitivity with a latency of four to six hrs.^{56,58,59} Plasma insulin levels also display a modest diurnal variation with a 10% excursion, a nadir between midnight and 6 am and a peak between noon and 6 pm.⁶⁰ Adiponectin is a hormone secreted by the adipocytes in an inverse fashion compared to fat mass.⁶¹ Its role in energy homeostasis is unclear: increases, decreases, or no changes in energy balance are derived from peripheral adiponectin administration in rodents.⁶² Adiponectin knockout mice develop insulin resistance and increased TNF- α levels when fed a high fat diet. In humans, plasma adiponectin levels correlated negatively with insulin resistance, and higher levels correlated with future weight gain in 1,063 women.^{63,64}




Fig. 2. Mean levels of metabolic parameters and sleep stages in lean young men ($n = 8$ for TSH, cortisol, GH, melatonin, SWS, and REM sleep;⁴⁸ $n = 14$ for insulin, glucose, total and acetylated ghrelin;⁷⁵ $n = 8$ for catecholamines⁷³); in 23 lean women for leptin and adiponectin;⁶⁹ and in 25 individuals (13 females, 12 males) for IL-6 and TNF- α .⁸³ Ghrelin levels are indicated as a percent of mean 24 hrs. values (1027 pg/mL for total ghrelin; 80 pg/mL for acetylated ghrelin); leptin levels are shown as percent change from levels at 08:00. Dark bars below each plot indicate bedtimes. Sleep was monitored polysomnographically during all measurements. Striped bars indicate meal times. Modified with permission from Refs. 48, 69, 73, 76, and 83.

Individuals with type 2 diabetes typically have low adiponectin levels, even when adjusted for body weight.⁶⁵ In contrast, patients with type 1 diabetes have higher than expected adiponectin levels: an observation that is poorly understood.⁶⁶ Sleep duration was inversely related to serum adiponectin in 109 lean Japanese males.⁶⁷ Adiponectin has anti-inflammatory properties: *in vitro* exposure to adiponectin decreases macrophage activation and TNF- α production. Finally, adiponectin displays some diurnal variability, with lower levels at night (**Fig. 2**).^{68,69}

During sleep, vagal tone increases.⁷⁰ Sympathovagal balance is lowest during REM sleep, when the locus coeruleus, the main brain source of catecholamine biosynthesis, is silent.⁷¹ Plasma norepinephrine (NE) and epinephrine (EPI) are produced by the sympathetic system: EPI is secreted exclusively from the adrenal medulla, whereas NE can also be released from postganglionic sympathetic nerves.⁷² In a small study of eight lean males, plasma catecholamines were consistently lower during the night with no differences among sleep stages (**Fig. 2**).⁷³ Heart rate and mean blood pressure were lower during NREM sleep, compared to during awakenings and REM sleep.⁷⁰

Several hormones involved in appetite regulation display circadian rhythms. Leptin is secreted by the white adipose tissue in a highly pulsatile fashion.⁷⁴ Leptin reflects fat stores: its levels increase after meals and during the night and are associated with decreased appetite (**Fig. 2**).⁷⁴ Ghrelin, produced in the stomach, stimulates appetite and circulates mainly in a nonacetylated form.⁷⁵ Acetylated ghrelin, the active form, is necessary for most of its endocrine actions, including GH release, appetite stimulation, and gastric emptying. Total and acetylated ghrelin levels rise with fasting and during sleep (**Fig. 2**).⁷⁶ Leptin and ghrelin exert their effects in the arcuate nucleus of the hypothalamus, through anorexigenic proopiomelanocortin/cocaine- and amphetamine-regulated transcript (POMC/CART) and orexigenic neuropeptide Y/agouti-related protein (NPY/AgRP) neurons. Leptin activates POMC/CART neurons, which induces hunger and inhibits NPY/AgRP neurons,^{77,78} which induces satiety. Ghrelin has the opposite effect on the arcuate nucleus, stimulating appetite, and prolonging postprandial glucose responses, while stimulating GH release.⁷⁹ Insulin, while peripherally lowering blood glucose and stimulating appetite, inhibits appetite centrally in a leptin-like manner.⁷⁸

Orexin A and B, two neuropeptides released by lateral hypothalamic neurons, stimulate appetite. Orexin neurons are active during waking and quiescent during sleep; consistently, orexin levels in the CSF are maximal at the end of the waking period.^{80,81} The orexin system is activated by ghrelin and inhibited by leptin and glucose. Projections from orexin neurons in the lateral hypothalamus activate NPY neurons in the arcuate nucleus.⁸⁰ Melatonin is a sleep-promoting hormone that is produced by the pineal gland. Melatonin is exclusively secreted during the dark, as light inhibits its production (**Fig. 2**).

Interleukin 6 (IL-6), interleukin 1 β (IL-1 β), TNF- α , and C-reactive protein (CRP) are pro-inflammatory factors. IL-6 displays a circadian rhythm, with a first peak around 2 am and a second peak around 5 am (**Fig. 2**).^{82,83} IL-1 β decreases during the night and reaches its nadir at 8 am in the morning.⁸² TNF- α also displays a circadian rhythm, peaking close to the awakening (6 am) and reaching a nadir around 3 pm (**Fig. 2**). IL-6 and TNF- α stimulate secretion of cortisol and, in turn, cortisol inhibits their secretion.⁸⁴ Reports on diurnal variability of CRP are few and inconsistent: some report no diurnal variance,⁸⁵ whereas others find increased morning levels.⁸⁶ In 10 healthy, lean men, white blood cell counts peaked around 11 pm, decreased throughout the night and reached a nadir at 8 am.⁸²

The relationships here described between hormone secretion and sleep stages are necessarily descriptive. In theory, as new pharmacological tools aimed at manipulating hormonal status (i.e., selective hormone agonists/antagonists) and/or sleep become available for investigation, we may be able to better understand the numerous causal links interconnecting these phenomena. In summary, many metabolic parameters involved in EE regulation, glucose metabolism, and appetite control display diurnal rhythms, reflecting the different metabolic needs during the sleep and wake states. In the next section, we will report the effect of sleep deprivation on these metabolic parameters.

Effects of sleep restriction on metabolic parameters

The relationship between sleep and metabolism has been extensively studied in human subjects under varying conditions, such as total sleep deprivation,^{87–100} partial sleep deprivation,^{101–122} and sleep fragmentation (**Table 1**).^{123–125} Partial sleep deprivation affects SWS less than other sleep stages. Most of the experiments on the effects of acute sleep deprivation were conducted in healthy, lean volunteers. Food intake and physical activity were often, but not always, strictly controlled (**Table 1**). It is important to note that experimentally induced sleep alterations may not be representative of changes occurring in clinical condition of chronic sleep deprivation. In addition, important differences exist between acute and chronic sleep deprivation. We will describe effects of sleep curtailment on important metabolic parameters that may contribute to insulin resistance and obesity, including effects on appetite and EE (**Fig. 3**).



Table 1: Results of 37 human experimental studies assessing influence of sleep curtailment on metabolism.

Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Benedict et al., 2011 [97] Randomized cross-over study Controlled, post-intervention <i>Ad libitum</i> Bed rest	men ($n = 14$) 22.6 \pm 0.8 23.9 \pm 0.5 kg/m ² 6 wks regular bedtimes*	1: 1 night of 8h bedtime*** 2: 1 night of TSD \geq 4 wks between conditions	Indirect calorimetry pre- and postbreakfast; morning VAS hunger ratings 24 hours, every 1.5–3 hours	\uparrow postprandial glucose (8%) \downarrow RMR (5%) and postprandial (20%) metabolic rate \uparrow cortisol (7%) ¹ \uparrow NE (12%) ¹ \sim food intake; \uparrow hunger (60%) ¹ \uparrow morning ghrelin (11%) \sim leptin
Born et al., 1997 [79] Randomized cross-over study Standardized meals Synchronous inpatient activities	men ($n = 10$) 24.7 (21-29) NA NA	1: 1 night of 8h bedtimes*** then 1 night of <i>ad libitum</i> sleep*** 2: 24h TSD*** \geq 10 days between conditions	LPS-stimulated whole blood and ELISA (TNF- α ; IL-1 β ; IL-6) 51h, every 3h	\sim IL-6 \sim IL-1 β ; \uparrow nocturnal whole blood \sim TNF- α \uparrow white blood cells (8%) ¹
Brondel et al., 2010 [110] Randomized cross-over study <i>Ad libitum</i> Free physical activity**	men ($n = 12$) 22 \pm 3 22.3 \pm 1.83 kg/m ² 2 days sleep/diet diaries*	1: 1 night of 4h bedtime*** and 1 night free bedtime at home*** 2: 1 night of 8h bedtime*** and 1 night of free bedtimes at home*** Conditions \geq 5 days apart	Actigraphy; 12 daytime VAS hunger ratings	\sim EE; \uparrow physical activity (13%) \uparrow hunger; \uparrow food intake (22%)
Buxton et al., 2010 [111] Consecutive experimental phases Controlled isocaloric (finishing meal required) Controlled sedentary living	men ($n = 20$) 26.8 \pm 5.2 23.3 \pm 3.1 kg/m ² \geq 5 days of 10hr bedtimes***	1: 3 nights of 10h bedtimes*** 2: 7 nights of 5h bedtimes***	ivGTT penultimate day; clamp last day; salivary sampling 15:00-21:00 last 2 days, every hour, 24h urine sampling last 2 days; indirect calorimetry	\uparrow insulin resistance (20% by ivGTT; 11% by clamp) \sim fasting RMR \uparrow salivary cortisol (51%) \uparrow urinary NE (18%) and EPI (22%) ¹
Clore et al., 1989 [41] Comparative study Similar evening meal Bed rest	5 females, 12 males ($n = 17$) 25 \pm 1 24.1 \pm 0.6 kg/m ² NA	1. Group 1 ($n = 11$): 1 night of 8h30min bedtime 2. Group 2 ($n = 6$): 1 night of TSD	Continuous ³ -3H glucose infusion	\downarrow nocturnal <u>fall</u> in glucose utilization \downarrow nocturnal <u>fall</u> in glucose production

Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM Prestudy conditions	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Donga et al., 2010 [112] Randomized cross-over study NA NA	3 females, 4 males ($n = 9$) 44 \pm 7 23.5 \pm 0.9 kg/m ² Type 1 diabetes, HbA1c \leq 8.5%, 1 week of regular bedtimes ^{**}	1: 3 nights of 8h30min bedtimes ^{***} 2: 3 nights of 4h bedtimes ^{***} Conditions \geq 3 wks apart	Hyperinsulinemic euglycemic clamp	\uparrow insulin resistance (21%)
Dzaja et al., 2004 [92] Randomized cross-over study 1,800 kcal/24h Bed rest	men ($n = 10$) 28 \pm 3.1 24.0 \pm 2.9 kg/m ² 1 week of regular bedtimes [*]	1: 1 night of 8h bedtime ^{***} 2: 1 night of TSD Conditions \geq 2 wks apart	24h, every hour	\downarrow GH peak (71%) ² \sim mean cortisol levels \downarrow nocturnal total ghrelin increase (39%) ²
Frey et al., 2007 [93] Consecutive phases study Hourly controlled intake Bed rest	9 females, 10 males ($n = 19$) 28.1 \pm 8.6 18.5-24.5 kg/m ² 3wks of regular bedtimes ^{**}	1: 3 nights of 8h bedtimes ^{***} 2: 1 night of TSD	Saliva 36h, every hour; hsELISA (IL-6; IL-1 β ; CRP) 40h, every 30min	\sim saliva cortisol; \downarrow at 13:00 (53%) ² and 20:00 (17%) ² \downarrow daytime IL-6 \uparrow IL-1 β \downarrow morning hsCRP
González-Ortiz et al., 2000 [87] Consecutive experimental phases Controlled isocaloric Free inpatient movement	7 females, 7 males ($n = 14$) 21.0 \pm 2.1 22.1 \pm 1.3 kg/m ² 3 days diet diary	1: baseline "normal sleep" 2: 1 night of TSD	Insulin suppression test with octeotide Once, morning	\uparrow insulin resistance (18%) ¹ \sim morning cortisol
Haack et al., 2007 [105] Comparative study Controlled isocaloric Free inpatient movement	6 females, 12 males ($n = 18$) 27.3 \pm 5.8 23.1 \pm 3.3 <30 kg/m ² \geq 10 days regular bedtimes [*]	($n = 8$) 1: 10 nights of 8h bedtimes ^{**} ($n = 10$) 2: 10 nights of 4h bedtimes ^{**}	24h urine 24h blood, every 4h (high sensitivity ELISA IL-6 and hsCRP)	\sim food intake \uparrow IL-6 (62%) ¹ \sim TNF-receptor 1 \uparrow hsCRP (102%; $P = 0.11$) ¹
Hursel et al., 2011 [122] Randomized cross-over Controlled isocaloric Sedentary activity	men ($n = 15$) 23.7 \pm 3.5 24.1 \pm 1.9 kg/m ² 2 days controlled feeding	1: 2 nights of 8h bedtimes ^{***} 2: 2 nights of sleep fragmentation hourly 2-min-long alarms ^{***} Conditions \geq 2 wks apart	Indirect calorimetry	\sim total EE \uparrow physical activity (8%) ¹ \uparrow RQ (3%) ¹
Irwin et al., 2006 [104] Consecutive phases study NA No exercise	13 females, 17 males ($n = 30$) 37.6 \pm 9.8 <30 kg/m ² 2 wks regular bedtimes [*]	1: 3 nights of 8h bedtimes ^{***} 2: 1 night of 4h bedtime ^{***}	Flow cytometry; real time PCR ($n = 10$); high-density oligonucleotide array ($n = 5$) Daytime (08:00-23:00), every 3-4h	\uparrow IL-6 monocyte expression (344%) ¹ ; \uparrow IL-6 mRNA (250%) ¹ \uparrow TNF- α monocyte expression (87%) ¹ ; \uparrow TNF- α mRNA (100%) ¹



Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM Prestudy conditions	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Jung et al., 2011 [40] Consecutive experimental phases Controlled isocaloric Sedentary	2 females, 5 males ($n = 7$) 22.4 \pm 4.8 22.9 \pm 2.4 kg/m ² 1 week 8h sleep ^{***} ; 3 days diet	1: 2 nights of 8h bedtimes ^{***} 2: 40h of TSD	Indirect calorimetry	\uparrow total EE (7%); \uparrow nocturnal EE (32%); \sim daytime EE \sim RQ
Kuhn et al., 1969 [84] Consecutive phases study Balanced diet NA	human subjects ($n = 28$) 20 – 30 NA NA	1: 4-5 days “control period” 2: 72-126h of TSD 3: 3 days “control period”	OGTT Once daily	\downarrow glucose tolerance after 3–4 days \uparrow 17-OH corticosteroids on day 2 and 3, then return to baseline \uparrow urinary catecholamines
Mullington et al., 2003 [91] Consecutive phases study Controlled isocaloric (finishing not required) Sedentary	men ($n = 10$) 27.2 26.1 \pm 1.9 kg/m ² 1 week of regular bedtimes ^{**}	1. 3 nights 8h bedtimes ^{***} 2. 3 nights of TSD	120 hours, every 90 min	\sim leptin mesor, \downarrow leptin circadian amplitude
Nedeltcheva et al., 2009 [106] Kessler et al., 2010 [113] Randomized crossover study <i>Ad libitum</i> Sedentary	5 females, 6 males ($n = 11$) 39 \pm 5 26.5 \pm 1.5 kg/m ² NA	1: 14 nights of 8.5h bedtimes ^{***} 2: 14 nights of 5.5h bedtimes ^{***} Conditions \geq 3 months apart	OGTT; ivGTT 24h, every 15-30min	\downarrow insulin sensitivity (18%) ¹ \downarrow TSH (7%); especially in females) ¹ \downarrow free T4 (8%) ¹ \sim GH; \downarrow during first 4hours of bedtime period in males (31%) ¹ \sim cortisol; \downarrow and 44min later acrophase (10%) ² ; 71min later nadir \uparrow EPI (21%) ¹ \sim mean NE; \uparrow nocturnal (24%) ¹
Nedeltcheva et al., 2010 [115] Randomized cross-over study Controlled restricted diet (90% RMR) Sedentary	3 females, 7 males ($n = 10$) 41 \pm 5 27.4 \pm 2 kg/m ² NA	1: 14 nights of 8.5h bedtimes ^{***} 2: 14 nights of 5.5h bedtimes ^{***} Conditions \geq 3 months apart	Indirect calorimetry; VAS hunger rating scale before meal/at 22:30 24h, every hour	\sim total energy expenditure; \downarrow RMR (8%) ¹ ; \uparrow fasting and postprandial RQ (4%) ¹ \sim thyroid hormones \sim GH \sim cortisol \downarrow EPI (12%) ¹ ; \sim NE \sim food intake; \uparrow hunger \sim total weight loss (3 kg), \downarrow fat weight loss (55%), \uparrow fat-free weight loss (60%) \uparrow mean acylated ghrelin; (7%) ¹ \sim leptin

Authors, Year [Reference] Study design	Gender/Sample Size Age ± SEM (yrs) Mean weight ± SEM	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Omisade et al., 2010 [114] Consecutive phases study Controlled Free inpatient movement	women (<i>n</i> = 15) 21.6 ± 2.2 24.5 ± 8 kg/m ² 1 week regular bedtimes and diet ^{*,**}	1: 1 night of 10h bedtime ^{**} 2: 1 night of 3h bedtime ^{**}	VAS hunger ratings; salivary cortisol daytime, every 1-2h; salivary leptin morning and afternoon	↓ median morning cortisol (19%) ¹ ; slower decline; ↑ afternoon/evening (44%) ¹ ~ hunger ↑ morning leptin (8%) ¹
Parker et al., 1987 [85] Consecutive phases study Controlled diet Sedentary	men (<i>n</i> = 4) 21-30 NA NA	1: 1 night of 8h bedtime ^{**} 2: 2 nights of TSD	72h, every 30min	↑ TSH; ↑ amplitude, longer peak, later nadir/acrophase
Pejovic et al., 2010 [96] Consecutive phases study <i>Ad libitum</i> Ambulatory	11 females, 10 males (<i>n</i> = 21) 24.1 ± 3.1 24.1 ± 2.6 kg/m ² 2 wks regular bedtimes ^{**}	1: 4 nights of 8h bedtimes ^{***} 2: 1 night of TSD, half 2h afternoon nap	Hunger scale (0-10) 24h, every 30 min	~ cortisol ~ adiponectin ~ hunger ↑ leptin (14%) ¹
Schmid et al., 2007 [94] Randomized cross-over study None Bed rest	men (<i>n</i> = 10) 25.3 ± 1.4 23.8 ± 0.5 kg/m ² Light dinner before arrival in the lab (21:00)	1: 1 night of 8h bedtime ^{**} 2: 1 night of TSD Conditions ≥ 2 wks apart	Hypoglycemic clamp Two morning samples	↓ glucagon levels (16%) ¹ ; ↑ response to hypoglycemia ↓ cortisol levels (16%) ¹ ; ~ ACTH ~ catecholamines ↑ hunger
Schmid et al., 2008 [95] Randomized cross-over study None Bed rest	men (<i>n</i> = 9) 24.2 ± 1.0 23.8 ± 0.6 kg/m ² NA	1: 1 night of 7h30min bedtime ^{***} 2: 1 night of 5h bedtime ^{***} 3: 1 night of TSD Conditions ≥ 2 wks apart	Hunger graded 0-9 Two morning samples	↑ hunger (129%) ¹ ~ leptin ↑ total ghrelin (22%)
Schmid et al., 2009 [107] Schmid et al., 2011 [116] Randomized cross-over study <i>Ad libitum</i> Free outpatient movement	men (<i>n</i> = 15) 27.1 ± 1.3 22.9 ± 0.3 kg/m ² 4 wks regular bedtimes [*]	1: 2 nights of 8h bedtimes ^{***} 2: 2 nights of 4h bedtimes ^{***} Conditions ≥ 6 wks apart	Actigraphy, buffet on Day 2, ELISA (IL-6) 08:00-23:00 (every hour)	↑ insulin (40%) and glucose (11%) peak response to breakfast ↓ physical activity (13%) ¹ ~ cortisol; ~ ACTH ~ total energy intake; ↑ fat intake (23%) ¹ ~ hunger; ~ intake ~ leptin; ~ ghrelin ~ IL-6
Schmid et al., 2009 [108] Randomized cross-over study None Bed rest	men (<i>n</i> = 10) 25.3 ± 1.4 23.8 ± 0.5 kg/m ² 2 wks regular bedtimes [*]	1: 1 night of 7h bedtime ^{**} 2: 1 night of 4.5h bedtime ^{**} Conditions ≥ 2 wks apart	Hypoglycemic clamp Two morning samples	~ insulin resistance ↓ glucagon (8%) ² ~ GH ↓ ACTH (44%) ¹ ; ↓ cortisol (44%) ¹ ~ EPI; ~ NE



Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Shearer et al., 2001 [90] Comparative study Isocaloric controlled No exercise	men ($n = 42$) 28.7 (21-47) NA 2 wks regular bedtimes**	($n = 21$) 1: 3 nights of 8h bedtimes, then 4 nights of 2h bedtimes 2: 3 nights of 8h bedtimes, 4 nights of TSD	5 days, every 6h (ELISA)	No changes in condition 1 ; in condition 2 : \uparrow IL-6 \sim TNF- α \uparrow sTNF- α receptor I \sim sTNF- α receptor II
Simpson et al., 2010a [117] Consecutive phases study <i>Ad libitum</i> Free inpatient movement	67 females, 69 males ($n = 136$) 30.4, 22-45 24.7, 17.7-32.6 kg/m ² 1 week regular bedtimes***	1: 2 nights of 10h bedtimes 2: 5 nights of 4h bedtimes	Once, morning (10:30-12:00)	\uparrow leptin (33%) ¹
Simpson et al., 2010b [118] Consecutive phases study <i>Ad libitum</i> Free inpatient movement	33 females, 41 males ($n = 74$) 29.9, 22-45 24.6, 17.7-33.1 kg/m ² 1 week regular bedtimes***	1: 2 nights of 10h bedtimes*** 2: 5 nights of 4h bedtimes***	Once, morning (10:30-12:00)	\sim adiponectin in men; \uparrow in African American women; \downarrow in Caucasian women
Spiegel et al., 1999 [98] Spiegel et al., 2000 [99] Spiegel et al., 2003 [100] Spiegel et al., 2004a [101]	men ($n = 11$) 22 \pm 1 23.4 \pm 0.5 kg/m ² ≥ 1 week of 8h bedtimes, diet***	1: 6 nights of 4h bedtimes*** 2: 6 nights of 12h bedtimes***	ivGTT; 24h heart rate variability; salivary sampling 15:00-bedtime, every 30min; blood sampling 24h, every 30min	\downarrow glucose clearance (30%) \downarrow TSH (26%), \uparrow fT4 (7%) ¹ \sim GH, different profile \uparrow afternoon saliva/plasma cortisol (25%) ¹ \uparrow sympathovagal balance (17%) ¹ \downarrow leptin (19%); \downarrow and 2.5h earlier acrophase (26%); \downarrow amplitude (20%) \downarrow melatonin; later onset; \downarrow acrophase
Spiegel et al., 2004b [102] Randomized cross-over study Controlled Free outpatient movement	men ($n = 12$) 22 \pm 2 23.6 \pm 2.0 ≥ 1 week of 8h bedtimes	1: 2 nights of 10h bedtimes*** 2: 2 nights of 4h bedtimes*** Conditions ≥ 6 wks apart	VAS hunger rating scale 08:00-21:00, every 20min	\uparrow hunger (23%; 33-45% for sweets/salty foods) \downarrow leptin (18%) \uparrow ghrelin (28%)
Stamatakis et al., 2010 [121] Consecutive phases <i>Ad libitum</i> NA	2 females, 9 males ($n = 11$) 23.2 (18 – 29) 24.3 \pm 0.9 kg/m ² 3 days diet and >7h bedtimes***	1: 1 night of unfragmented sleep*** 2: 2 nights of fragmented sleep (~ 31.4 auditory/mechanical stimuli per h)***	ivGTT; heart rate variability; enzyme-linked immunosorbent assay techniques (IL-6; hsCRP) 2 morning samples (08:00 and 16:00)	\downarrow insulin sensitivity (25%) \uparrow morning cortisol (13%) \sim adiponectin \uparrow sympathetic tone (17%) \sim leptin \sim IL-6; \sim hsCRP

Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
St-Onge et al., 2011 [119] Randomized cross-over study Controlled isocaloric, self-selected on Day 5 Free inpatient movement	15 females; 15 males ($n = 30$) 33.9 \pm 4.3; 36.6 \pm 5.6 23.0 \pm 1.1; 24.1 \pm 1.1 kg/m ² 2 wks regular bedtimes ^{**}	1: 5 nights of 9h bedtimes ^{***} 2: 5 nights of 4h bedtimes ^{***} Conditions \geq 4 wks apart	Double labeled water (EE); indirect calorimetry (RMR); actigraphy; VAS hunger scales	\sim RMR; \sim EE; \sim physical activity \sim hunger; \uparrow food intake (12%) ¹ ; especially \uparrow saturated fat intake in females (62%)
Tasali et al., 2008 [120] Randomized cross-over study Controlled Sedentary activities	4 females, 5 males ($n = 9$) 20 – 31 64.2 kg; 19-24 kg/m ² NA 1 week regular bedtimes ^{**}	1: 2 nights of undisturbed sleep ^{***} 2: 3 nights of disturbed sleep by acoustic stimuli during SWS (\sim 33 micro-arousals/h) ^{***} Conditions \geq 4 wks apart	ivGTT; heart rate variability 24h, every 20min	\downarrow insulin sensitivity (25%) \sim cortisol; \sim daytime; \sim nocturnal \uparrow sympathovagal balance (14%)
Thomas et al., 2000 [88] Consecutive experimental phases NA NA	men ($n = 17$) 24.7 \pm 2.8 NA \geq 7 days of 7h45min bedtimes ^{**}	1: 3 nights of 7h45min bedtimes ^{***} 2: 85h of TSD (only results from 24h shown)	18-FDG and PET, every 24h	\downarrow global cerebral glucose metabolic rate (8%), especially the thalamus, prefrontal and posterior parietal cortices
Van Cauter et al., 1991 [42] Consecutive phases study Only breakfast on study day Bed rest	men ($n = 8$) 22 - 27 22.7 \pm 0.7 kg/m ² 1 week 23:00-07:00 bedtimes [*]	1: 2 nights of 8h bedtimes ^{***} 2: 28h of TSD	Continuous glucose infusion during phase 2 24 hours, every 30 min	\downarrow nocturnal GH pulses (70%) ¹ \sim cortisol, \downarrow acrophase (4%) ¹ ; 1h earlier nadir
VanHelder et al., 1993 [86] Consecutive phases study Isocaloric controlled Sedentary and exercise	men ($n = 10$) 22 \pm 3 74.5 \pm 11.8 kg 1 night of 7h bedtimes ^{***}	60h of TSD followed by 7h of recovery sleep ^{***} 1: sedentary activities 2: daily exercise Conditions \geq 10 days apart	OGTT at 10h, 60h and after recovery sleep	\uparrow insulin response (21% in group 1)
Van Leeuwen et al., 2009 [109] Consecutive phases study Isocaloric controlled Free outpatient movement	men ($n = 13$) 23.1 \pm 2.5 NA 2 wks regular bedtimes [*]	1: 2 nights of 8h bedtimes ^{***} 2: 5 nights of 4h bedtimes ^{***}	Flow cytometry, real time PCR (1 morning sample), salivary cortisol (10 times a day)	\sim cortisol \uparrow IL-6 (63%) \uparrow IL-1 β (37%) \sim TNF- α \uparrow hsCRP (45%)



Authors, Year [Reference] Study design	Gender/Sample Size Age \pm SEM (yrs) Mean weight \pm SEM	Experimental sleep protocol	Techniques used Blood sampling (time, frequency)	Main findings (sleep deprived vs. longer sleep)
Vgontas et al., 2004 [103] Consecutive phases study Usual diet Free outpatient movement (sedentary during sampling)	13 females, 12 males ($n = 25$) 25.2 \pm 3.8 23.8 \pm 2.3 kg/m ² 2 wks regular bedtimes**	1: 4 nights of 8h bedtime*** 2: 8 nights of 6h bedtimes***	24h, every 30min (ELISA TNF- α and IL-6)	~ cortisol; \downarrow and 2h earlier peak \uparrow IL-6 \uparrow TNF- α in males; ~ in females

Sleep curtailment was verified by *questionnaires/diaries **by actigraphy or ***by polysomnography. Percentage of change in metabolic parameters was given in the article or calculated^{1,2}: from ¹absolute levels in the text/tables or from ²graph values of the original article. Metabolic measurements are in plasma/serum unless otherwise specified. Bed rest was always required during the sleep deprived state. *Abbreviations:* ~ similar; TSD: total sleep deprivation; VAS: visual analogue scale; RMR: resting metabolic rate; IL-6: interleukin 6; IL-1 β : interleukin 1 β ; TNF- α : tumor necrosis factor- α ; EE: energy expenditure; ivGTT: intravenous glucose tolerance test; RQ: respiratory quotient; OGTT: oral glucose tolerance test; REM: rapid-eye movement; GH: growth hormone; ACTH: adrenocorticotrophic hormone; FDG: fludeoxyglucose; PET: position emission tomography.

Effects on glucose metabolism

In the course of the night, glucose supply to the brain must remain adequate in the setting of slightly declining plasma glucose levels. Infusion of [3H3]-labeled glucose in healthy, lean subjects revealed decreased glucose usage during the night, whereas in subjects kept awake glucose usage was increased (**Table 1**).⁴⁴ Total and partial sleep deprivation reduced glucose tolerance in lean individuals. This has been documented by intravenous glucose tolerance tests (ivGTT),^{101,109,114,123,124} oral glucose tolerance tests (OGTT),^{87,89,109} insulin suppression tests with octreotide,⁹⁰ as well as hypoglycemic and euglycemic hypoinsulinemic clamps (**Table 1**).^{97,114} Impaired glucose metabolism was not only shown in healthy study participants, but also in individuals with type 1 diabetes: after a four-hr night in adults with type 1 diabetes, glucose infusion rate decreased by 21%, as assessed by hyperinsulinemic euglycemic clamp (**Table 1**).¹¹⁵ In addition to altered peripheral glucose metabolism during total sleep deprivation, cerebral glucose uptake decreased, especially in the prefrontal and posterior parietal cortex and in the thalamus of healthy volunteers (**Table 1**).⁹¹ Interestingly, most studies found no compensatory rise in plasma insulin secondary to increased insulin resistance after short sleep,^{109,114} fragmented sleep,¹²³ or total sleep deprivation (**Table 1**).⁹⁰ However, increased insulin secretion in response to ivGTTs was found after 60 hrs of sleep deprivation⁸⁹ and two nights of sleep fragmentation.¹²⁴ Thus, almost all studies find decreased glucose tolerance and a dampened compensatory insulin response after sleep curtailment.



Effects on EE

Two groups found that one night of total sleep deprivation increased EE during the night, whereas the following day, resting and postprandial EE were reduced by 5% and 20%, respectively, overall leading to a positive energy balance (**Fig. 3** and **Table 1**).^{43,100} Similarly, Schmid et al. reported that volunteers had decreased spontaneous physical activity after a night of short sleep compared to a night with restful sleep (**Table 1**).¹¹⁰ Fragmenting sleep for two consecutive nights lead to an increase in the respiratory quotient (RQ), indicating a shift from fat toward carbohydrate oxidation, without affecting total EE (**Table 1**).¹²⁵ High RQ predicts weight gain and insulin resistance.¹²⁶ Two weeks of partial sleep deprivation (5.5 hrs) decreased resting metabolic rate and increased RQ in a randomized, cross-over study (**Table 1**).¹⁰⁹ In contrast, subjects with chronic insomnia had increased metabolic rate.¹²⁷ We have recently shown that poor sleep quality was associated with higher REE, a higher RQ, and an activation of the stress system in obese subjects with short sleep duration.¹²⁸ In summary, EE may be differentially affected by chronic vs. acute sleep curtailment and may also depend on differences in reasons of sleep curtailment (experimentally induced short sleep vs. chronic insomnia).




Fig. 3. A simplified schematic representation of putative pathways of sleep curtailment leading to obesity and diabetes, via endocrine mechanisms that stimulate appetite, decrease energy expenditure, and increase insulin resistance. The direction of change (an increase or decrease) of each mechanism due to sleep loss is displayed; increased sympathovagal balance could stimulate EE, depicted by +, while a decreased EE would contribute to obesity. For readability, the relationships between decreases in leptin levels associated with IL-6 and TNF- α production are not displayed. Cortisol and proinflammatory cytokines display a positive bidirectional relationship. Increased insulin levels stimulate orexin secretion and orexin arcuate neurons, while decreasing the activity of anorexic arcuate neurons, leading to a further increase in appetite. Adiponectin influences energy homeostasis, possibly via modulating appetite through arcuate neurons and modulating EE.

Additional endocrine mechanisms linking sleep, glucose metabolism, and energy expenditure

As mentioned above, thyroid hormones, GH, cortisol, adipokines (e.g., adiponectin), and sympathovagal balance modulate glucose metabolism and energy homeostasis (**Fig. 3**). Acute total sleep deprivation increased TSH and free T4 levels,⁸⁸ whereas partial sleep deprivation of 5.5 hrs nights for two wks or six 4-hr nights decreased TSH and T4 levels, suggesting a dose-related effect (**Table 1**).^{104,116} Decreased peripheral thyroid hormone levels are likely secondary to suppression of the nocturnal TSH peak, possibly secondary to decreased drive from TRH neurons in the PVN.¹¹⁶ Sleep-deprived rodents show reduced levels of hypothalamic TRH mRNA in the PVN of the hypothalamus.¹²⁹ Of note, a different stressor such as acute immobilization also decreases TRH mRNA hypothalamic levels and plasma TSH levels in rodents.¹³⁰

Insulin resistance can result from increases in insulin counter-regulatory hormones, including GH and cortisol (**Fig. 3**). One night of total sleep deprivation markedly decreased the GH peak that usually occurs around sleep onset (**Table 1**).^{45,95} Partial sleep deprivation did not affect 24-hr levels of GH in lean or obese individuals,^{101,109,118} but six 4-hr nights changed GH nocturnal secretory patterns: sleep deprivation was associated with a biphasic pattern compared to the usual single large GH peak at sleep onset.¹⁰² This extended exposure to GH may decrease glucose uptake in muscles, thus contributing to insulin resistance. However, the threshold of GH levels contributing to insulin resistance is unknown. In contrast to the previous study, decreased GH levels during the first four hrs of sleep were reported after a two-week partial sleep-deprivation period, in association with more SWS.^{109,118} GH is usually secreted during SWS, thus this unexpected finding may be due to a dissociation of GH secretion from SWS. In summary, the effects of sleep deprivation on GH plasma levels are variable.

Partial and total sleep deprivation resulted in elevated salivary and plasma cortisol the following afternoon and evening (**Table 1**),^{100,101,109,114,117} while morning cortisol levels may be unchanged or even decreased.^{90,97,117} As awakening is usually associated with a cortisol surge, this finding probably reflects the lack of the awakening response. In addition, a delay and/or a reduction in the acrophase of cortisol have been reported after partial and total sleep deprivation.^{95,102,109} In a sleep-deprived state, cortisol levels decreased more slowly after reaching their acrophase, possibly indicating decreased sensitivity to the negative feedback effects on the hypothalamic–pituitary–adrenal axis. Mean 24-hr levels of plasma cortisol often remain unchanged by sleep deprivation,^{43,95,96,99,106,109,123} but the circadian cortisol-pattern changes, as described above.

Adiponectin levels were similar after partial and total sleep deprivation and sleep fragmentation (**Table 1**).^{99,121,124} However, subgroup analysis revealed that effects of sleep restriction on adiponectin levels may be gender and race dependent: adiponectin decreased

in Caucasian women and increased in African American women, while adiponectin levels in males did not change after short sleep (**Table 1**).¹²¹ Besides influencing energy homeostasis, decreased adiponectin levels worsen insulin resistance.

Sympathovagal balance, defined as the ratio between the activity of the sympathetic and the parasympathetic nervous system, can be indirectly assessed with heart rate variability analysis. Since the activity of this system is very diverse, on the basis of the different anatomic branches, it is unclear to which extent perturbations in heart-rate variability may be predictive of changes in basal metabolic rate and insulin sensitivity. Furthermore, heart rate, EPI and NE levels are markers of sympathetic activity. Acute sleep deprivation is associated with increased sympathetic activity, decreased parasympathetic tone and therefore, with increased sympathovagal balance (**Table 1**).^{101,109,123,124} Total EE is usually decreased in acute sleep deprivation; thus, other mechanisms, such as decreased thyroid hormones or adiponectin levels, must prevail over increased sympathovagal tone (**Fig. 3**). Increased sympathovagal balance also adds to insulin resistance (**Fig. 3**).¹³¹ Furthermore, activation of the sympathetic nervous system and inhibition of the parasympathetic nerves decreases insulin secretion; thus, changes in the autonomic nervous system could explain the absence of a compensatory insulin response after sleep curtailment.¹³²

In summary, sleep loss impairs glucose metabolism and may decrease EE. This is mediated by multiple factors, such as decreased thyroid hormones and, possibly, adiponectin levels; increased sympathovagal balance; and altered patterns of cortisol secretion, while the role of GH is still unclear (**Fig. 3**).

Effects on appetite

It is not clear to what extent obesity in subjects with short sleep is caused by increased energy intake vs. decreased EE (**Fig. 3** and **Table 1**).¹³³ If food is provided ad libitum, partial or total sleep deprivation may not change hunger ratings, while likely increasing food intake.^{99,117,120} On the other hand, when food intake is controlled, sleep deprivation usually increases appetite. Self-reported sleep quality was inversely related to appetite in 53 first degree, normal weight relatives of subjects with diabetes.¹³⁴ After two 4-hr nights, there was a 24% increase in daytime appetite ratings;¹⁰⁵ in addition, sleep deprivation specifically increased appetite for sweet and salty foods. The effects of sleep deprivation on appetite may not be immediate: one 4-hr night of sleep deprivation was associated with increases in food intake 36 hrs later, while this parameter was unaffected immediately following sleep deprivation.¹¹³ In addition, the amount of sleep deprivation needed to increase hunger ratings may be variable. One night of total, but not partial, sleep deprivation, significantly increased hunger ratings in normal weight males.^{98,113} No differences in total food intake were observed after two 4-hr, 15 min vs. 8-hr, 15-min nights, but subjects ate relatively more fat when sleeping less.¹¹⁹ In a randomized, cross-over, five-day study of 4 hrs vs. 9 hrs



of sleep, with a controlled diet on days 1–4 vs. an ad libitum diet on day five women, but not men, exhibited increased total caloric intake and a preference for fatty foods, especially if saturated, suggesting that women may be more susceptible to the orexic effects of sleep deprivation than men.¹²²

Chronotype, the individual attribute determining morning or evening preference, modulates metabolism. Obese subjects carrying the CLOCK 3111TC/CC polymorphism, a variant of one of the core clock genes associated with being an “owl” or an evening person, sleep 20 minutes less.¹³⁵ In addition, these subjects had a higher intake of transfatty acids and proteins, and consumed relatively later in the day, while having the same total energy intake of noncarriers. Subjects carrying the CLOCK 3111TC/CC genotype are also more resistant to weight loss: they lost about 2 kg less while on a hypocaloric diet for 30 wks.¹³⁵

Leptin and ghrelin inhibit and stimulate appetite, respectively (**Fig. 3**). Following partial sleep deprivation, mean daytime values of leptin were approximately 18% lower compared to the rested state.^{104,105} In addition, the circadian amplitude of leptin was dampened and the acrophase was decreased and phase advanced by two hrs. A decreased circadian amplitude was also observed after three nights of total sleep deprivation.⁹⁴ In a large cohort study of 1,024 subjects, shorter habitual sleep duration as assessed by sleep diaries related to lower morning values of leptin.⁵ Some studies found no association between short sleep and leptin levels.^{94,98,100,119,136} On the contrary, morning leptin levels were found to be increased after five nights of four hours,¹²⁰ after one night of three hours¹¹⁷ and after one night of total sleep deprivation.⁹⁹ Short sleep duration, as assessed by one night of polysomnography in 561 subjects, also related to higher morning leptin levels.¹³⁷ It is difficult to explain the inconsistent results of these different studies. Of note, studies that found decreased leptin after short sleep controlled food intake, whereas when sleep-deprived subjects were allowed to eat ad libitum, leptin levels were unchanged or even increased.

In obese subjects, leptin levels are elevated but several of its actions are impaired, a concept referred to as leptin resistance. In a cross-over study design, no association among habitual sleep duration, measured actigraphically, and morning leptin levels was observed in obese subjects.¹³⁸ Mean 24-hr leptin levels in obese subjects on a controlled diet were also unchanged after 14 nights of 5.5 hrs vs. 8.5 hrs.¹¹⁸

The physiological increase in total ghrelin levels in the early night was blunted when individuals were kept awake.⁹⁵ A similar phenomenon is reported in chronic insomniacs.¹³⁵ However, morning total ghrelin levels were increased after one night of sleep deprivation^{95,98} and in individuals with shorter habitual sleep durations.⁵ Mean daytime total ghrelin levels were higher after two nights of 4-hr sleep.¹⁰⁵ CLOCK 3111TC/CC carriers, who display an evening preference and sleep 20 min shorter, also had increased morning ghrelin levels.¹³⁴ Mean 24-hr levels of the metabolically active acetylated ghrelin were higher in obese subjects after two wks of 5.5-hr nights.¹¹⁸ Overall, sleep curtailment is consistently associated with higher daytime ghrelin.

The effect of sleep deprivation on the orexin system in normal subjects is not well studied. However, it is known that patients with narcolepsy have abnormally low orexin-A CSF levels, which correlate both with body weight and sleep abnormality.¹³⁹ Total sleep deprivation increased orexin levels in the CSF of squirrel monkeys, and increased cFos expression in orexin neurons in rats.^{81,140} It is challenging to assess the activity of the orexin system in humans. Central determination requires collection of CSF, an invasive technique. Determination of orexin in plasma is hampered by the lack of adequate assays, in terms of sensitivity and specificity. Hyperactivity of the orexin system could contribute to increased food intake during sleep deprivation. In summary, appetite and food intake are often increased following sleep curtailment, possibly via decreased leptin, increased ghrelin and a hyperactive orexin system (**Fig. 3**).



Effects on the immune system

Enhanced levels of circulating leucocytes, proinflammatory cytokines (e.g., IL-6, TNF- α , IL-1 β) and CRP have been associated with an increased risk of cardiovascular disease and type 2 diabetes in long-term human and experimental studies.^{141–144} Increased plasma levels of IL-6 were reported after partial and total sleep deprivation in most cases,^{93,106,107,108,112} although two studies only reported a loss of circadian rhythms (**Table 1**).^{82,96} IL-1 β was elevated after partial,^{106,112} and total sleep deprivation,⁹⁶ but one study did not find any changes after total sleep deprivation (**Table 1**).⁸² TNF- α levels are most often,^{82,93,112} but not always,¹⁰⁷ unchanged after partial sleep restriction. One study only found increases in men, but not in women, after partial sleep deprivation (**Table 1**).¹⁰⁶ Another study reported higher levels of soluble TNF- α receptor I, through which TNF- α exerts its sleep promoting function.⁹³ Effects of sleep curtailment on CRP levels are inconsistent: they have been reported lower,⁹⁶ unchanged,¹²⁴ or higher.^{108,112} Prolonged total sleep deprivation has been associated with leucocytosis (**Table 1**).⁸² Thus, differences in study outcomes may be influenced by individual or gender differences. In addition, sensitivity and specificity of commercial cytokine assays are highly variable. Highly sensitive analytical methods, such as mass spectrometry, may be needed for accurate cytokine measurements.¹⁴⁵ We conclude that short sleep promotes a proinflammatory state, which, in turn, exerts its negative consequences on insulin resistance.

The other side of the coin: metabolic dysfunction influencing sleep

Most of the studies of sleep deprivation have been performed in healthy, lean subjects and therefore could not address the effect of sleep deprivation on subjects with obesity and diabetes. Obesity and diabetes are often associated with increased stress or differences in life style that could negatively influence sleep (**Fig. 3**).¹⁴⁶ Depression and obesity may occur together as well: obesity may negatively affect mood and both pharmacologic antidepressant therapy and major depression itself are associated with weight gain.¹⁴⁷ We

recently reported that obese subjects often suffer from sleep-disordered breathing, which induces frequent microawakenings and loss of SWS.¹⁴⁸ Similar effects of BMI on sleep have also been observed in youth. Decreased total sleep time and less sleep efficiency were observed in severely obese adolescents, of which 74% displayed sleep apnea (16.5 yrs; BMI $60.3 \pm 2.1 \text{ kg/m}^2$).¹⁴⁹ Changes in sleep architecture in obesity have been reported: severely obese adolescents (12–15 yrs; BMI 50.9; $39.8\text{--}63.0 \text{ kg/m}^2$, apnea index 14 episodes/hr) have a decrease in REM sleep.¹⁵⁰

Symptoms associated with diabetes, such as thirst, nocturia, extreme glucose excursions, and mood alterations, may also interfere with sleep, independent of obesity. To this point, adults with type 1 diabetes, who are typically not obese, report a poorer sleep quality than their matched controls without diabetes.¹⁵¹ These awakenings were associated with a rapid decline of glucose levels, but not with the absolute values of hypoglycemia.¹⁵² Furthermore, studies in animals suggest that diet composition may directly influence sleep: mice fed a high-fat diet display an attenuated rhythm in the circadian expression of core clock genes.⁴⁰ Of note, augmented amplitude of the peripheral and central circadian rhythms favors wakefulness during the day and sleep at night, whereas an attenuated circadian amplitude disrupts the sleep–wake cycle.¹⁵³ Likewise, studies in rodents demonstrate that physical activity at the congruent time (i.e. nocturnal for mice) increases the central circadian amplitude, suggesting that active behavior during the day and sleep at night in humans may be beneficial for high-amplitude circadian rhythms.¹⁵⁴ Thus, sleep and circadian rhythm are bidirectionally connected (**Fig. 3**). Leptin influences sleep architecture in normal fed rodents, increasing SWS by 13% and decreasing REM by 30%.¹⁵⁵ Furthermore, *ob/ob* mice that lack leptin and *db/db* mice that lack the leptin receptor, both display sleep disturbances that are reverted by leptin replacement.

In summary, metabolic dysfunction, altered hormone levels, and sleep abnormalities probably all influence each other.^{156,157} In summary, sleep deprivation and obesity may potentiate each other in a vicious reverberating circuit; short sleep may induce weight gain, via the mechanisms outlined in this paper, whereas obesity, in turn, via sleep apnea and other symptoms, may disrupt sleep. Observational studies, especially if cross-sectional, may not be well suited to disentangle these aggregate, causative factors. This underlies the need for behavioral, life-style intervention studies.

Possible therapeutic role of melatonin

In nonequatorial areas, the duration, intensity, and spectral quality of natural light vary across seasons.¹⁵⁸ For example, during a summer in England, there was relatively more blue light vs. green and red light exposure, in addition to increased light intensity and duration.¹⁵⁸ These are important signals for seasonal rhythms, transmitted to the body via the duration of nocturnal melatonin secretion. As a consequence of artificial lighting and

other environmental modifications, seasonality is greatly attenuated in modern humans. Furthermore, because sleep is usually the only time during which humans do not experience light exposure, shorter sleep duration will result in longer light exposure. Light suppresses the pineal production of melatonin in a dose related fashion. White artificial light greater than 200 lux completely suppressed nocturnal plasma melatonin production, dim room light (106 lux) reduced melatonin by 88% and light intensities lower than 80 lux did not affect melatonin levels.¹⁵⁹ As the night progresses, the intensity of continuous artificial light needed to suppress circulating melatonin increases.¹⁶⁰ Of note, light of shorter wavelength, such as the light that is typically emitted by electronic devices such as televisions and computers, induces relatively more melatonin suppression and likely leads to greater sleep disruption.¹⁶⁰

Urinary 24-hr melatonin concentrations decreased by more than 50% when subjects slept five vs. eight hours, while being continuously exposed to white light (700 lux) during wake hours.¹⁶¹ Diminished levels of melatonin were likely due to three additional hrs of light exposure. Likewise, mean 24-hr plasma melatonin concentrations were decreased after 6 nights of 4 hrs vs. 12 hrs bedtimes.¹⁰³ The onset of melatonin secretion was delayed, due to four hrs longer light exposure before bedtime (300 lux) and the acrophase was reduced in sleep-deprived subjects.

Recent studies suggest that melatonin may have a physiological role in modulating metabolism. For example, removal of the pineal gland decreased the responsiveness of several insulin-dependent hepatic kinases during the dark phase in rats.¹⁶² This effect was reversed by nocturnal supplementation of melatonin in the drinking water. Chronic melatonin supplementation in drinking water also reduced body weight in rodents.^{163,164} Patients with metabolic syndrome have disturbances in melatonin production, characterized by an alteration of the night melatonin-insulin ratio of unclear clinical significance.¹⁶⁵ In a large genome-wide association studies conducted in individuals of European origin, certain genetic variants of the melatonin receptor 1B were associated with an increased risk of type 2 diabetes.¹⁶⁶⁻¹⁶⁸ However, the most common receptor variant did not influence the relationship between sleep abnormalities and type 2 diabetes.¹⁶⁶ Interestingly, the melatonin receptor 1B is present on pancreatic β -cells, and melatonin has an inhibitory effect on insulin secretion. Thus, it was feared that treatment with melatonin in subjects with diabetes would increase blood glucose levels. On the other hand, since type 2 diabetes is associated with reduced melatonin levels toward the end of the night;¹⁶⁹ thus, there may be a physiological rationale for melatonin supplementation at bedtime. Few studies have addressed these questions. Sleep efficiency improved after five months of treatment with 2 mg prolonged-release melatonin administered two hrs before bedtime, while total sleep time was unaffected in 26 subjects with type 2 diabetes.¹⁷⁰ HbA1c levels also improved (8.47% vs. 9.13% at baseline). Lack of a control group prevented understanding whether



improved HbA1c levels were in part due to other factors than melatonin administration, such as better pharmacological and diet compliance due to study participation. In lean subjects, melatonin administration at night for 2 months was associated with decreased LDL levels and improved oxidative status.¹⁷¹ Another study found a positive correlation between plasma melatonin levels at 2 am and HDL levels in 36 women treated with 1 mg melatonin for one month.¹⁷² As melatonin promotes sleep, the decrease in obesity and improvement in insulin resistance could be due to longer and better sleep *per se*, but it could also relate to other melatonin-dependent pathways yet to be identified (**Fig. 3**).

Discussion and future directions

Epidemiological and laboratory data prove the existence of a relationship between short or disturbed sleep and adverse metabolic outcomes. In general, these effects are small in size, but certain specific populations may be more susceptible than others. For example, age plays a clear role as the relationship between short sleep and obesity is stronger in children. Sleep duration and sleep architecture change with age in healthy individuals, with SWS becoming less common with age.¹⁷³ It is also well known that sex and reproductive hormones affect sleep regulatory mechanisms¹⁷⁴ and that sleep disorder affects women differently from men.¹⁷⁵ It is unfortunate that experimental studies, in both humans and in rodents, are mostly conducted in the male gender, thus unduly discriminating the female gender.

Similarly, sleep deprivation is more common in minorities but the effects of socioeconomic status *per se* vs. ethnicity remain to be determined. Individual susceptibility within these categories varies greatly: identification of contributing factors will greatly help prevention and treatment and be one of the tenets of individualized medicine.¹⁷⁶ Few negative effects are expected to be observed in individuals with voluntary sleep curtailment (natural short sleepers), in contrast to individuals with short sleep due to social or work-related pressure or chronic insomnia. Current epidemiologic studies often lack an objective measurement of sleep duration, such as actigraphy or repeated sleep diaries. Thus, prospective studies with objective sleep duration measurements are warranted in different subgroups to further explore this relationship.

In addition, the causality of the relationship between sleep and metabolic disorders remains unclear, even in prospective studies. Obese/diabetic individuals may be sleeping less than lean subjects because of differences in life style or stress; their sleep could be interrupted more often by symptoms of disturbed glucose levels or sleep apnea; evidence arises that possibly circulating hormones affect the brain inducing worsened sleep (**Fig. 3**). As already mentioned, causality can be addressed best in long-term studies where “prescribed” sleep duration is randomized, but this may present with practical difficulties

and significant challenges, including the fact that the biological need for optimal sleep duration varies and cannot be estimated at an individual level. Sleep is regulated by circadian and homeostatic mechanisms. As far as homeostatic mechanisms are concerned, it has been the matter of a long scientific debate whether during wakefulness, single or rather multiple metabolites accumulate, progressively increasing the pressure to sleep during the day. Classic experiments of parabiosis seem to support the notion of endogenous “sleep promoting” substances. Recently, the importance of adenosinergic neurotransmission in the regulation of sleep, especially non-REM sleep, has been underlined.¹⁷⁷ Nevertheless, it is certain that there is a large individual variability in sleep needs. Sensitive and specific biomarkers of sleep deprivation are needed for various reasons, including clinical and medico-legal reasons.¹⁷⁸

The effect of sleep extension in subsets of short sleepers on metabolism can be addressed in randomized controlled trials. Currently, we are conducting such a study in short-sleeping obese individuals.¹⁷⁹ Investigating the effects of sleep extension in other subgroups of varying age, BMI, and ethnicity will also be of interest. The effect of metabolic disturbances on sleep, such as hyperinsulinemia or increased circulating levels of proinflammatory cytokines, may be empirically addressed in polysomnographical studies. As it is difficult to artificially mimic the internal milieu of obese/diabetic individuals, effects of metabolism may be modeled in animal studies using parabiosis, connecting circulations of lean and obese animals and comparing sleep characteristics.

As indicated by studies of sleep deprivation, the mechanisms connecting short sleep and obesity/ insulin resistance are probably mediated by three pathways: increases in appetite, decreases in EE, and influences on glucose metabolism (**Fig. 3**). Endocrine mechanisms influencing these pathways are complex and interconnected. The existing studies of sleep deprivation have generated variable results; small sample size, differences in the control of food-intake and physical activity levels, and inherent individual differences may have all contributed to this variability. In addition, these studies only addressed the effects of acute sleep deprivation and were not performed in real-life situations. We suggest that future studies should focus on the effects of sleep deprivation in specific patient populations, including obese/insulin-resistant individuals and chronic insomniacs vs. natural short sleepers, and should mimic real-life conditions.

Most studies find worsened glucose tolerance after sleep curtailment, which may be beneficial from a teleological perspective, at least in the short term. When our ancestors remained awake during the night, they were probably experiencing a threat (e.g., absence of food, presence of predators), so inducing a state in which glucose availability was advantageous. Likewise, increases in appetite result in more food intake. Together with reduced EE, this will further increase blood glucose availability. In modern societies, however, prolonged exposure to such a state may result in obesity and type 2 diabetes.



At this point, there is a general consensus that short sleep has negative effects on health, but the amount of sleep curtailment causing metabolic disturbances is unclear, and specific groups at greater risk are yet to be identified. Whether melatonin would benefit obese/diabetic subjects, or a subset of these individuals, remains to be proven in rigorous randomized clinical trials. Supplementation of melatonin has been shown to improve sleep efficiency, and HbA1c levels in patients with type 2 diabetes.¹⁷⁰ Lean subjects may also profit from melatonin administration as well, as it improved lipid profiles.^{171,172} Thus, melatonin appears to be a promising supplement in battling the obesity/diabetes epidemics, but controlled studies need to confirm its positive effect on metabolism. Furthermore, the long-term effects and side effects of chronic melatonin supplementation need to be examined in different subgroups.

Overall, improving sleep duration and quality is a potential tool to counteract the epidemics of obesity and diabetes. However, before sleep extension advice is translated to the clinic, more research is needed.

Acknowledgments

This study was supported by the National Institutes of Health (NIH), Intramural Research Program: National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Eliane A. Lucassen was the recipient of a Fulbright Scholarship and received scholarships from VSBfonds and the Leiden University Fund.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Flegal KM, Carroll MD, Kit BK, Ogden CL (2012) Prevalence and trends in obesity in the distribution of body mass index among US adults, 1999–2010. *JAMA* 307:491–497.
2. Korner J, Woods SC, Woodworth KA (2009) Regulation of energy homeostasis and health consequences in obesity. *Am J Med* 122:S12–S18.
3. Centers for Disease Control and Prevention (2011) National diabetes fact sheet: national estimates and general information on diabetes and prediabetes in the United States, 2011. Available from: http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2011.pdf.
4. Hasler G, Buysse DJ, Klaghofer R, et al. (2004) The association between short sleep duration and obesity in young adults: a 13-Year prospective study. *Sleep* 27:661–666.
5. Taheri S, Lin L, Austin D, Young T, Mignot E (2004) Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased Body Mass Index (BMI). *Sleep* 27:A146–A147.
6. Ayas N, White DP, Al-Delaimy WK, et al. (2003) A prospective study of sleep duration and coronary heart disease in women. *Arch Intern Med* 163:205–209.
7. Björkelund C, Bondyr-Carlsson D, Lapidus L, et al. (2005) Sleep disturbances in midlife unrelated to 32-year diabetes incidence: the prospective population study of women in Gothenburg. *Diabetes Care* 28:2739–2744.
8. Yaggi HK, Araujo AB, McKinlay JB (2006) Sleep duration as a risk factor for the development of type 2 diabetes. *Diabetes Care* 29:657–661.
9. Gangwisch JE, Heymsfield SB, Boden-Albala B, et al. (2007) Sleep duration as a risk factor for diabetes incidence in a large US sample. *Sleep* 30:1667–1673.
10. Gallicchio L, Kalesan B (2009) Sleep duration and mortality: a systematic review and meta-analysis. *J Sleep Res* 18:148–158.
11. National Sleep Foundation (2005) Sleep in America Poll. National Sleep Foundation. Washington DC.
12. Bin YS, Marshall NS, Glozier N (2012) Secular trends in adult sleep duration: A systematic review. *Sleep Med Rev* 16:223–230.
13. Cappuccio FP, Taggart FM, Kandala NB, et al. (2008) Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 31:619–626.
14. Chen X, Beydoun MA, Wang Y (2008) Is sleep duration associated with childhood obesity? A systematic review and meta-analysis. *Obesity* 16:265–274.
15. Rao MN, Blackwell T, Redline S, et al. (2009) Association between sleep architecture and measures of body composition. *Sleep* 32:483–490.
16. Rasmussen MH, Wildschiodt G, Juul A, Hilsted J (2008) Polysomnographic sleep, growth hormone, insulin like growth factor-I axis, leptin, and weight loss. *Obesity* 16:1516–1521.
17. Lauderdale DS, Knutson KL, Rathouz PJ, et al. (2009) Cross-sectional and longitudinal associations between objectively measured sleep duration and body mass index: the CARDIA Sleep Study. *Am J Epidemiol* 170:805–813.
18. Patel SR, Malhotra A, White DP, Gottlieb DJ, Hu FB (2006) Association between reduced sleep and weight gain in women. *Am J Epidemiol* 164:947–954.
19. Chaput JP, Després JP, Bouchard C, Tremblay A (2008) The association between sleep duration and weight gain in adults: a 6-year prospective study from the Quebec Family Study. *Sleep* 31:517–523.
20. López-García E, Faubel R, León-Muñoz L, et al. (2008) Sleep duration, general and abdominal obesity, and weight change among the older adult population of Spain. *Am J Clin Nutr* 87:310–316.
21. Gunderson EP, Rifas-Shiman SL, Oken E, et al. (2008) Association of fewer hours of sleep at 6 months postpartum with substantial weight retention at 1 year postpartum. *Am J Epidemiol* 167:178–187.
22. Hairston KG, Bryer-Ash M, Norris JM, et al. (2010) Sleep duration and five-year abdominal fat accumulation in a minority cohort: the IRAS family study. *Sleep* 33:289–295.
23. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB (2005) Inadequate sleep as a risk factor for obesity: analyses of the NHANES I. *Sleep* 28:1289–1296.
24. Stranges S, Cappuccio FP, Kandala NB, et al. (2008) Cross-sectional versus prospective associations of sleep duration with changes in relative weight and body fat distribution: the Whitehall II Study. *Am J Epidemiol* 167:321–329.
25. Watanabe M, Kikuchi H, Tanaka K, Takahashi M (2010) Association of short sleep duration with weight gain and obesity at 1-year follow-up: a large-scale prospective study. *Sleep* 33:161–167.



26. Boden G, Shulman GI (2002) Free fatty acids in obesity and type 2 diabetes: defining their role in the development of insulin resistance and beta-cell dysfunction. *Eur J Clin Invest* 32:14–23.
27. Knutson KL, Ryden AM, Mander BA, van Cauter E (2006) Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus. *Arch Intern Med* 166:1768–1774.
28. Choi KM, Lee JS, Park HS, et al. (2008) Relationship between sleep duration and the metabolic syndrome: Korean National Health and Nutrition Survey 2001. *Int J Obes* 32:1091–1097.
29. Tuomilehto H, Peltonen M, Partinen M, et al. (2008) Sleep duration is associated with an increased risk for the prevalence of type 2 diabetes in middle-aged women – The FIND2D survey. *Sleep Med* 9:221–227.
30. Xu Q, Song Y, Hollenbeck A, et al. (2010) Day napping and short night sleeping are associated with higher risk of diabetes in older adults. *Diabetes Care* 33:78–83.
31. Cappuccio FP, D’Elia L, Strazzullo P, Miller MA (2010) Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care* 33:414–420.
32. Kawakami N, Takatsuka N, Shimizu H (2004) Sleep disturbance and onset of type 2 diabetes. *Diabetes Care* 27:282–283.
33. Nilsson PM, Rööst M, Engström G, Hedblad B, Berglund G (2004) Incidence of diabetes in middle-aged men is related to sleep disturbances. *Diabetes Care* 27:2464–2469.
34. Chaput JP, Després JP, Bouchard C, Astrup A, Tremblay A (2009) Sleep duration as a risk factor for the development of type 2 diabetes or impaired glucose tolerance: analyses of the Quebec Family Study. *Sleep Med* 10:919–924.
35. Meisinger C, Heier M, Loewel H (2005) Sleep disturbance as a predictor of type 2 diabetes mellitus in men and women from the general population. *Diabetologia* 48:235–241.
36. Beihl DA, Liese AD, Haffner SM (2009) Sleep duration as a risk factor for incident type 2 diabetes in a multiethnic cohort. *Ann Epidemiol* 19:351–357.
37. Trento M, Broglio F, Riganti F, et al. (2008) Sleep abnormalities in type 2 diabetes may be associated with glycemic control. *Acta Diabetol* 45:225–229.
38. Aronsohn RS, Whitmore H, van Cauter E, Tasali E (2010) Impact of untreated obstructive sleep apnea on glucose control in type 2 diabetes. *Am J Res Crit Care Med* 181:507–513.
39. Neubauer DN (1999) Sleep problems in the Elderly. *Am Fam Physician* 59:2551–2558.
40. Kohsaka A, Laposky AD, Ramsey KM, et al. (2007) High-fat diet disrupts behavioral and molecular circadian rhythms in mice. *Cell Metab* 6:414–421
41. Asher G, Schibler U (2011). Crosstalk between components of circadian and metabolic cycles in mammals. *Cell Metab* 13: 125–137.
42. Cizza G, Requena M, Galli G, de Jonge L (2011) Chronic sleep deprivation and seasonality: implications for the obesity epidemic. *J Endocrinol Invest* 34:793–800.
43. Jung CM, Melanson EL, Frydelhall EJ, et al. (2011) Energy expenditure during sleep, sleep deprivation and sleep following sleep deprivation in adult humans. *J Physiol* 589:235–244.
44. Clore JN, Nestler JE, Blackard WG (1989) Sleep associated fall in glucose disposal and hepatic glucose output in normal humans. Putative signaling mechanism linking peripheral and hepatic events. *Diabetes* 38:285–290.
45. Van Cauter E, Blackman JD, Roland D, et al. (1991) Modulation of glucose regulation and insulin secretion by circadian rhythmicity and sleep. *J Clin Invest* 88:934–942.
46. Boyle PJ, Nagy RJ, O’Connor AM, et al. (1994) Adaptation in brain glucose uptake following recurrent hypoglycemia. *Proc Natl Acad Sci USA* 91:9352–9356.
47. Maquet P (1997) Positron emission tomography studies of sleep and sleep disorders. *J Neurol* 244:S23–S28.
48. Van Coevorden A, Mockel J, Laurent E, et al. (1991) Neuroendocrine rhythms and sleep in aging men. *Am J Physiol* 260:E651–E661.
49. Holli RW, Hartman ML, Veldhuis JD, Taylor WM, Thorner MO (1991) Thirty second sampling of plasma growth hormone in man: correlation with sleep stages. *J Clin Endocrinol Metab* 72:854–861.
50. Karacan I, Rosenbloom AL, Williams RL, Finley WW, Hirsch CJ (1971) Slow wave sleep deprivation in relation to plasma growth hormone concentration. *Behav Neuropsychiatry* 2:11–14.
51. Van Cauter E, Plat L, Scharf MB, et al. (1997) Simultaneous stimulation of slow-wave sleep and growth hormone secretion by gamma-hydroxybutyrate in normal young men. *J Clin Invest* 100:745–753.
52. Othmer E, Daughaday V, Guze S (1969) Effects of 24 hour REM deprivation on serum growth hormone levels in humans. *Electroencephalogr Clin Neurophysiol* 27:685.
53. Mendelson WB, Slater S, Gold P, Gillin JC (1980) The effect of growth hormone administration on human sleep: a dose-response study. *Biol Psychiatry* 15:613–618.

54. Kern W, Halder R, al-Reda S, et al. (1993) Systemic growth hormone does not affect human sleep. *J Clin Endocrinol Metab* 76:1428–1432.
55. Møller N, Jorgensen JOL, Schmitz O, et al. (1990) Effects of a growth hormone pulse on total and forearm substrate fluxes in humans. *Am J Physiol Endocrinol Metab* 258: E86–E91.
56. Byrne M, Sturis J, Blackman JD, Polonsky KS, van Cauter E (1992) Decreased glucose tolerance during sleep may be partially mediated by GH. Program of the 74th Annual Meeting of The Endocrine Society, San Antonio, TX (Abstract 40).
57. Møller N, Butler PC, Antsiferov MA, Alberti KGMM (1989) Effects of growth hormone on insulin sensitivity and forearm metabolism in normal man. *Diabetologia* 32:105–110.
58. McMahon M, Gerich J, Rizza R (1988) Effects of glucocorticoids on carbohydrate metabolism. *Diabetes Metab Rev* 4:17–30.
59. Dinneen S, Alzaid A, Miles J, Rizza R (1993) Metabolic effects of the nocturnal rise in cortisol on carbohydrate metabolism in normal humans. *J Clin Invest* 92:2283–2290.
60. Boden G, Ruiz J, Ubrain J., Chen X (1996) Evidence for a circadian rhythm of insulin secretion. *Am J Physiol* 271:E246–E252.
61. Stanley K, Wynne K, McGowan B, Bloom S (2005) Hormonal regulation of food intake. *Physiol Rev* 85:1131–1158.
62. Dridi S, Taouis M (2009) Adiponectin and energy homeostasis: consensus and controversy. *J Nutr Biochem* 20:831–839.
63. Hivert MF, Sullivan LM, Fox CS, et al. (2008) Associations of adiponectin, resistin, and tumor necrosis factor alpha with insulin resistance. *J Clin Endocrinol Metab* 93:3165–3172.
64. Hivert MF, Sun Q, Shrader P, et al. (2011) Higher adiponectin levels predict greater weight gain in healthy women in the Nurses' Health Study. *Obesity* 19: 409–415.
65. Lilja M, Rolandsson O, Norberg M, Soderberg S (2012) The impact of leptin and adiponectin on incident type 2 diabetes is modified by sex and insulin resistance. *Metab Syndr Relat Disord* 10:143–151.
66. Maahs DM, Ogden LG, Snell-Bergeon JK, et al. (2007) Determinants of serum adiponectin in persons with and without type 1 diabetes. *Am J Epidemiol* 166:731–740.
67. Kontani K, Sakane N, Saiga K, et al. (2007) Serum adiponectin levels and lifestyle factors in Japanese men. *Heart Vessels* 22:291–296.
68. Scheer FA, Chan JL, Fargnoli J, et al. (2010) Day/night variations of high-molecular-weight adiponectin and lipocalin-2 in healthy men studied under fed and fasted conditions. *Diabetologia* 53:2401–2405.
69. Cizza G, Nguyen VT, Eskandari F, et al. (2010) Low 24 hour adiponectin and high nocturnal leptin concentrations in a case-control study of community-dwelling premenopausal women with major depressive disorder: the Premenopausal, Osteopenia/Osteoporosis, Women, Alendronate, Depression (POWER) study. *J Clin Psychiatry* 71:1079–1087.
70. Somers VK, Dyken ME, Mark AL, Abboud FM (1993) Sympathetic-nerve activity during sleep in normal subjects. *N Engl J Med* 328:303–307.
71. Aston-Jones G, Chen S, Zhu Y, Oshinsky ML (2001) A neural circuit for circadian regulation of arousal. *Nat Neurosci* 4:732–738.
72. Hjemdahl P (1993) Plasma catecholamines—analytical challenges and physiological limitations. *Baillieres Clin Endocrinol Metab* 7:307–353.
73. Linsell CR, Lightman SL, Mullen PE, Brown MJ, Causon RC (1985) Circadian rhythms of epinephrine and norepinephrine in man. *J Clin Endocrinol Metab* 60:1210–1205.
74. Licinio J, Mantzoros C, Negrao AB, et al. (1997) Human leptin levels are pulsatile and inversely related to pituitary adrenal function. *Nat Med* 3:575–579.
75. Badman MK, Flier JS (2005) The gut and energy balance: visceral allies in the obesity wars. *Science* 307:1909–1914.
76. Spiegel K, Tasali E, Leproult R, Scherberg N, van Cauter E (2011) Twenty-four-hour profiles of acylated and total ghrelin: relationship with glucose levels and impact of time of day and sleep. *J Clin Endocrinol Metab* 96:486–493.
77. Elias CF, Aschkenasi C, Lee C, et al. (1999) Leptin differentially regulates NPY and POMC neurons projecting to the lateral hypothalamic area. *Neuron* 23:775–786.
78. Morton GJ, Schwartz MW (2011) Leptin and the central nervous system control of glucose metabolism. *Physiol Rev* 91:389–411.



79. Hataya Y, Akamizu T, Takaya K, et al. (2001) A low dose of ghrelin stimulates growth hormone (GH) release synergistically with GH-releasing hormone in humans. *J Clin Endocrinol Metab* 86:4552.
80. Van den Top M, Lee K, Whyment AD, Blanks AM, Spanswick D (2004) Orexigen-sensitive NPY/AgRP pacemaker neurons in the hypothalamic arcuate nucleus. *Nat Neurosci* 7: 493–494.
81. Zeitzer JM, Buckmaster CL, Lyons DM, Mignot E (2007) Increasing length of wakefulness and modulation of hypocretin-1 in the wake-consolidated squirrel monkey. *Am J Physiol Regul Integr Comp Physiol* 293:R1736–R1742.
82. Born J, Lange T, Hansen K, et al. (1997) Effects of sleep and circadian rhythm on human circulating immune cells. *J Immunol* 158:4454–4464.
83. Vgontzas AN, Zoumakis M, Papanicolaou DA (2002) Chronic insomnia is associated with a shift of interleukin-6 and tumor necrosis factor secretion from nighttime to daytime. *Metabolism* 51:887–892.
84. Tsigos C, Papanicolaou DA, Defensor R, et al. (1997) Dose effects of recombinant human interleukin-6 on pituitary hormone secretion and energy expenditure. *Neuroendocrinology* 66:54–62.
85. Meier-Ewert HK, Ridker PM, Rifai N, et al. (2001) Absence of diurnal variation of C-reactive protein concentration in healthy human subjects. *Clin Chem* 47:426–430.
86. Koc M, Karaarslan O, Abali G, Batur MK (2010) Variations in high-sensitivity C-reactive protein levels over 24 hours in patients with stable coronary artery disease. *Tex Heart Inst J* 37 42–48.
87. Kuhn E, Brodan V, Brodanova M, Rysanek K. 1969. Metabolic reflection of sleep deprivation. *Acta Nerv Super* 11:165–174.
88. Parker DC, Rossman LG, Pekary AE, Hershman JM (1987) Effect of 64-hour sleep deprivation on the circadian waveform of thyrotropin (TSH): further evidence of sleep related inhibition of TSH release. *J Clin Endocrinol Metab* 64:157–161.
89. VanHelder T, Symons JD, Radomski MW (1993) Effects of sleep deprivation and exercise on glucose tolerance. *Aviat Space Environ Med* 64:487–492.
90. González-Ortiz M, Martínez-Abundis E, Balcázar-Muñoz BR, Pascoe-González S (2000) Effect of sleep deprivation on insulin sensitivity and cortisol concentration in healthy subjects. *Diabetes Nutr Metab* 13:80–83.
91. Thomas M, Sing H, Belenky G, et al. (2000) Neural basis of alertness and cognitive performance impairments during sleepiness: I. Effects of 24 h of sleep deprivation on waking human regional brain activity. *J Sleep Res* 9:335–352.
92. Garfinkel D, Zorin M, Wainstein J, et al. (2011) Efficacy and safety of prolonged-release melatonin in insomnia patients with diabetes: a randomized, double-blind, crossover study. *Diabetes Metab Syndr Obes* 4:307–313.
93. Shearer WT, Reuben JM, Mullington JM, et al. (2001) Soluble TNF- α receptor 1 and IL-6 plasma levels in humans subjected to the sleep deprivation model of spaceflight. *J Allergy Clin Immunol* 107:165–170.
94. Mullington JM, Chan JL, Dongen HP, et al. (2003) Sleep loss reduces diurnal rhythm amplitude of leptin in healthy men. *J Neuroendocrinol* 15:851–854.
95. Dzaja A, Dalal MA, Himmerich H, et al. (2004) Sleep enhances nocturnal plasma ghrelin levels in healthy subjects. *Am J Physiol Endocrinol Metab* 286:E963–E967.
96. Frey DJ, Fleshner M, Wright KP (2007) The effects of 40 hours of total sleep deprivation on inflammatory markers in healthy young adults. *Brain Behav Immun* 21:1050–1057.
97. Schmid SM, Hallschmid M, Jauch-Chara K, et al. (2007) Sleep loss alters basal metabolic hormone secretion and modulates the dynamic counterregulatory response to hypoglycemia. *J Clin Endocrinol Metab* 92:3044–3051.
98. Schmid SM, Hallschmid M, Jauchchara K, Born J, Schultes B (2008) A single night of sleep deprivation increases ghrelin levels and feelings of hunger in normal weight healthy men. *J Sleep Res* 17:331–334.
99. Pejovic S, Vgontzas AN, Basta M, et al. (2010) Leptin and hunger levels in young healthy adults after one night of sleep loss. *J Sleep Res* 19:552–558.
100. Benedict C, Hallschmid M, Lassen A, et al. (2011) Acute sleep deprivation reduces energy expenditure in healthy men. *Am J Clin Nutr* 93:1229–1236.
101. Spiegel K, Leproult R, Van Cauter E (1999) Impact of sleep debt on metabolic and endocrine function. *Lancet* 354:1435–1439.
102. Spiegel K, Leproult R, Colechia EF, et al. (2000) Adaptation of the 24h growth hormone profile to a state of sleep debt. *Am J Psychol* 279:R874–R883.
103. Spiegel K, Leproult R, van Cauter E (2003) Impact of sleep debt on physiological rhythms. *Rev Neurol* 159:6S11–6S20.
104. Spiegel K, Tasali E, Penev P, Van Cauter E. (2004a). Brief communication: sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med*. 141:846–850.

105. Spiegel K, Leproult R, L'Hermite-Baleriaux M, et al. (2004b). Leptin levels are dependent on sleep duration: relationships with sympathovagal balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin Endocrinol Metab* 89:5762–5771.
106. Vgontzas AN, Zoumakis E, Bixler EO, et al. (2004) Adverse effects of modest sleep restriction on sleepiness, performance, and inflammatory cytokines. *J Clin Endocrinol Metab* 89:2119–2126.
107. Irwin MR, Wang M, Campomayor CO, et al. (2006) Sleep deprivation and activation of morning levels of cellular and genomic markers of inflammation. *Arch Intern Med* 166:1756–1762.
108. Haack M, Sanchez E, Mullington JM (2007) Elevated inflammatory markers in response to prolonged sleep restriction are associated with increased pain experience in healthy volunteers. *Sleep* 30:1145–1152.
109. Nedeltcheva AV, Kessler L, Imperial J, Penev PD (2009) Exposure to recurrent sleep restriction in the setting of high caloric intake and physical inactivity results in increased insulin resistance and reduced glucose tolerance. *J Clin Endocrinol Metab* 94:3242–3250.
110. Schmid SM, Hallschmid M, Jauch-Chara K, et al. (2009) Short-term sleep loss decreases physical activity under free-living conditions but does not increase food intake under time-deprived laboratory conditions in healthy men. *Am J Clin Nutr* 90: 1476–1482.
111. Schmid SM, Jauch-Chara K, Hallschmid M, Schultes B (2009) Mild sleep restriction acutely reduces plasma glucagon levels in healthy men. *J Clin Endocrinol Metab* 94:5169–5173.
112. van Leeuwen WM, Lehto M, Karisola P, et al. (2009) Sleep restriction increases the risk of developing cardiovascular diseases by augmenting proinflammatory responses through IL-17 and CRP. *PLoS One* 4:e4589.
113. Brondel L, Romer MA, Nougues PM, Touyarou P, Davenne D (2010) Acute partial sleep deprivation increases food intake in healthy men. *Am J Clin Nutr* 91:1550–1559.
114. Buxton OM, Marcelli E (2010) Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States. *Soc Sci Med* 71:1027–1036.
115. Donga E, van Dijk M, van Dijk JG, et al. (2010) Partial sleep restriction decreases insulin sensitivity in type 1 diabetes. *Diabetes Care* 33:1573–1577.
116. Kessler L, Nedeltcheva A, Imperial J, Penev PD (2010) Changes in serum TSH and free T4 during human sleep restriction. *Sleep* 33:1115–1118.
117. Omisade A, Buxton OM, Rusak B (2010) Impact of acute sleep restriction on cortisol and leptin levels in young women. *Physiol Behav* 99:651–656.
118. Nedeltcheva AV, Kilkus JM, Imperial J, Schoeller DA, Penev PD (2010) Insufficient sleep undermines dietary efforts to reduce adiposity. *Ann Intern Med* 153:435–441.
119. Schmid SM, Hallschmid M, Jauch-Chara K, et al. (2011) Disturbed gluco-regulatory response to food intake after moderate sleep restriction. *Sleep* 34:371–377.
120. Simpson NS, Banks S, Dinges DF (2010) Sleep restriction is associated with increased morning plasma leptin concentrations, especially in women. *Biol Res Nurs* 12:47–53.
121. Simpson NS, Banks S, Arroyo S, Dinges DF (2010) Effects of sleep restriction on adiponectin levels in healthy men and women. *Physiol Behav* 101:693–698.
122. St-Onge M, Roberts AL, Chen J, et al. (2011) Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr* 94:410–416.
123. Tasali E, Leproult R, Ehrmann DA, Van Cauter E (2008) Slow-wave sleep and the risk of type 2 diabetes in humans. *Proc Natl Acad Sci USA* 105:1044–1049.
124. Stamatakis KA, Punjabi NM (2010) Effects of sleep fragmentation on glucose metabolism in normal subjects. *Chest* 137:95–101.
125. Hursel R, Rutters F, Gonnissen HK, Martens EA, Westertep-Plantenga MS (2011) Effects of sleep fragmentation in healthy men on energy expenditure, substrate oxidation, physical activity, and exhaustion measured over 48 h in a respiratory chamber. *Am J Clin Nutr* 94:804–808.
126. Ravussin E (1995) Metabolic differences and the development of obesity. *Metabolism* 44:12–14.
127. Bonnet MH, Arand DL (1995) 24-hour metabolic rate in insomniacs and matched normal sleepers. *Sleep* 18:581–588.
128. De Jonge L, Zhao X, Mattingly MS, et al. (2012) Poor sleep quality and sleep apnea are associated with higher energy expenditure in obese individuals with short sleep duration. *J Clin Endocrinol Metab* 97:2881–9.
129. Everson CA, Nowak Jr TS (2002) Hypothalamic thyrotropin-releasing hormone mRNA responses to hypothyroxinemia induced by sleep deprivation. *Am J Physiol Endocrinol Metab* 283:E85–E93.
130. Cizza G, Brazy LS, Esclapes ME, et al. (1996) Age and gender influence basal and stress-modulated hypothalamic pituitary-thyroidal function in Fischer 344/N rats. *Neuroendocrinology* 64:440–448.



131. Deibert DC, DeFronzo RA (1980) Epinephrine-induced insulin resistance in man. *J Clin Invest* 65:717–721.
132. Avogaro A, Toffolo G, Valerio A, Cobelli C (1996) Epinephrine exerts opposite effects on peripheral glucose disposal and glucose stimulated insulin secretion. A stable label intravenous glucose tolerance test minimal model study. *Diabetes* 45:1373–1378.
133. Cizza G, Rother KI (2012) Beyond fast food and slow motion: weighty contributors to the obesity epidemic. *J Endocrinol Invest* 35:236–242.
134. Kilkus JM, Booth JN, Bromley LE, et al. (2012) Sleep and eating behavior in adults at risk for type 2 diabetes. *Obesity* 20: 112–117.
135. Garaulet M, Moreno C, Smith CE, et al. (2011) Ghrelin, sleep reduction and evening preference: relationships to CLOCK3111 T/C SNP and weight loss. *PLoS One* 6:e17435.
136. Motivala SJ, Tomiyama J, Ziegler M, Khandrika S, Irwin MR (2009) Nocturnal levels of ghrelin and leptin and sleep in chronic insomnia. *Psychoneuroendocrinology* 34:540–545.
137. Hayes AL, Xy F, Babineau D, Patel SR (2011) Sleep duration and circulatory adipokine levels. *Sleep* 34:147–152.
138. Knutson KL, Galli G, Zhao X, Mattingly M, Cizza G (2011) No association between leptin levels and sleep duration or quality in obese adults. *Obesity* 19:2433–2435.
139. Nakamura M, Kanbayashi T, Sugiura T, Inoue Y (2011) Relationship between clinical characteristics of narcolepsy and CSF orexin-A levels. *J Sleep Res* 20:45–49.
140. Estabrooke IV, McCarthy MT, Ko, E et al. (2001) Fos expression in orexin neurons varies with behavioral state. *J Neurosci* 21:1656–1662.
141. Hotamisligil GS, Budavari A, Murray D, Spiegelman BM (1994) Reduced tyrosine kinase activity of the insulin receptor in obesity-diabetes. Central role of tumor necrosis factor- α . *J Clin Invest* 94:1543–1549.
142. Kristiansen OP, Mandrup-Poulsen T (2005) Interleukin-6 and diabetes: the good, the bad, or the indifferent? *Diabetes* 54:S114–S124.
143. Rother KI (2007) Diabetes treatment – bridging the divide. *New Eng J Med* 356:1499–1501.
144. Nieto-Vazquez I, Fernández-Veledo S, Krämer DK, et al. (2008) Insulin resistance associated to obesity: the link TNF α . *Arch Physiol Biochem* 114:183–194.
145. Cizza G, Marques AH, Eskandari F, et al. (2008) Elevated neuroimmune biomarkers in sweat patches and plasma of premenopausal women with major depressive disorder in remission: the POWER study. *Biol Psychiatry* 64:907–911.
146. Siervo M, Wells JC, Cizza G (2009) The contribution of psychosocial stress to the obesity epidemic: an evolutionary approach. *Horm. Metab Res* 41:261–270.
147. Patten SB, Williams JV, Lavorato DH, Khaled S, Bulloch AG (2011) Weight gain in relation to major depression and antidepressant medication use. *J Affect Disord* 134:288–293.
148. Knutson KL, Zhao X, Mattingly M, Galli G, Cizza G (2012) Predictors of sleep-disordered breathing in obese adults who are chronic short sleepers. *Sleep Med* 13:484–489.
149. Kalra M, Manna M, Fitz K, et al. (1998) Effect of surgical weight loss on sleep architecture in adolescents with severe obesity. *Pediatrics* 101:61–67.
150. Willi SM, Oexmann MJ, Wright NM, Collop NA, Lindon Key L (2008) The effects of a high-protein, low-fat, ketogenic diet on adolescents with morbid obesity: body composition, blood chemistries, and sleep abnormalities. *Obes Surg* 18:675–679.
151. van Dijk M, Donga E, van Dijk JG, et al. (2011) Disturbed subjective sleep characteristics in adult patients with long-standing type 1 diabetes mellitus. *Diabetologia* 54:1967–1976.
152. Pillar G, Schusheim G, Weiss R, et al. (2003) Interactions between hypoglycemia and sleep architecture in children with type 1 diabetes mellitus. *J Pediatrics* 142:163–168.
153. Pandi-Perumal SR, Moscovitch A, Srinivasan V, et al. (2009) Bidirectional communication between sleep and circadian rhythms and its implications for depression: lessons from agomelatine. *Prog Neurobiol* 88:264–271.
154. van Oosterhout F, Lucassen EA, Houben T, et al. (2012) Amplitude of the SCN clock enhanced by the behavioral activity rhythm. *PLoS One* 2012 7:e39693.
155. Sinton CM, Fitch TE, Gershenfeld HK (1999) The effects of leptin on REM sleep and slow wave delta in rats are reversed by food deprivation. *J Sleep Res* 8:197–203.
156. Laposky AD, Shelton J, Bass J, et al. (2006) Altered sleep regulation in leptin-deficient mice. *Am J Physiol Regul Integr Comp Physiol* 290:R894–R903.
157. Laposky AD, Bradley MA, Williams DL, Bass J, Turek FW (2008) Sleep-wake regulation is altered in leptin resistance (db/db) genetically obese and diabetic mice. *Am J Physiol Regul Integr Comp Physiol* 295:R2059–R2066.

158. Thorne HC, Jones KH, Peters SP, Archer SN, Dijk DJ (2009) Daily and seasonal variation in the spectral composition of light exposure in humans. *Chronobiol Int* 26:854–866.
159. Zeitzer JM, Dijk DJ, Kronauer R, Brown E, Czeisler C (2000) Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting suppression. *J Physiol* 526:695–702.
160. Gooley JJ, Rajaratnam SM, Brainard GC, et al. (2010) Spectral responses of the human circadian system depend on the irradiance and duration of exposure to light. *Sci Trans Med* 2:31ra33.
161. Minami M, Takahashi H, Sasaki T, et al. (2012) The effect of sleep restriction and psychological load on the diurnal metabolic changes in tryptamine-related compounds in human urine. *Environ Health Prev Med* 17:87–97.
162. Nogueira TC, Lellis-Santos C, Jesus DS, et al. (2011) Absence of melatonin induces night-time hepatic insulin resistance and increased gluconeogenesis due to stimulation of nocturnal unfolded protein response. *Endocrinology* 152:1253–1263.
163. Wolden-Hanson T, Mitton DR, McCants RL, et al. (2000) Daily melatonin administration to middle-aged rats suppresses body weight, intra-abdominal adiposity, and plasma leptin and insulin dependent of food intake and total body fat. *Endocrinology* 141:487–497.
164. Cardinali DP, Pagano ES, Scacchi Bernasconi PA, Reynoso R, Scacchi P (2011) Disrupted chronobiology of sleep and cytoprotection in obesity: possible therapeutic value of melatonin. *Neuro Endocrinol Lett* 32:588–606.
165. Robeva R, Kirilov G, Tomova A, Kumanov P (2008) Melatonin-insulin interactions in patients with metabolic syndrome. *J Pineal Res* 44:52–56.
166. Olsson L, Pettersen E, Ahlborn A, et al. (2011) No effect by the common gene variant rs10830963 of the melatonin receptor 1B on the association between sleep disturbances and type 2 diabetes: results from the Nord-Trøndelag Health Study. *Diabetologia* 54:1375–1378.
167. Prokopenko I, Langenberg C, Florez JC, et al. (2009) Variant in MTNR1B influence fasting glucose levels. *Nat Genet* 41:77–81.
168. Bonnefond A, Clément N, Fawcett K, et al. (2012) Rare MTNR1B variants impairing melatonin receptor 1B function contribute to type 2 diabetes. *Nat Genet* 44:297–301.
169. Peschke E, Frese T, Chankiewicz E, et al. (2006) Diabetic Goto Kakizaki rats as well as type 2 diabetic patients show a decreased diurnal serum melatonin level and an increased pancreatic melatonin-receptor status. *J Pineal Res* 40:135–143.
170. Garfinkel C, Zorin M, Wainstein J, et al. (2011) Efficacy and safety of prolonged-release melatonin in insomnia patients with diabetes: a randomized, double-blind, crossover study. *Diabetes Metab Syndr Obes* 4:307–313.
171. Kozirog M, Poliwczak AR, Duchnowicz P, et al. (2011) Melatonin treatment improves blood pressure, lipid profile, and parameters of oxidative stress in patients with metabolic syndrome. *J Pineal Res* 50:261–266.
172. Tamura H, Nakamura Y, Narimatsu A, et al. (2008) Melatonin treatment in peri- and postmenopausal women elevates serum high density lipoprotein cholesterol levels without influencing total cholesterol. *J Pineal Res* 45:101–105.
173. Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV (2004) Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep* 27:1255–1273.
174. Paul KN, Turek FW, Kryger MH (2008) Influence of sex on sleep regulatory mechanisms. *J Womens Health* 17:1201–1208.
175. Phillips BA, Collop NA, Drake C, et al. Sleep disorders and medical conditions in women (2008) Proceedings of the Women & Sleep Workshop, National Sleep Foundation, Washington, DC, March 5–6, 2007. *J Womens Health* 17:1191–1199.
176. Ershow AG (2009) Environmental influences on development of type 2 diabetes and obesity: challenges in personalizing prevention and management. *J Diabetes Sci Technol* 2:727–734.
177. Landolt HP (2008) Sleep homeostasis: a role for adenosine in humans? *Biochem Pharmacol* 75:2070–2079.
178. Czeisler CA (2011) Impact of sleepiness and sleep deficiency on public health-utility of biomarkers. *J Clin Sleep Med* 7(5 Suppl):S6–S8.
179. Cizza G, Marincola P, Mattingly M, et al. (2010) Treatment of obesity with extension of sleep duration: a randomized, prospective, controlled trial. *Clinical Trials* 7:274–285.



