

Genetically modified (GM) corn in the Philippines : Ecological impacts on agroecosystems, effects on the economic status and farmers' experiences

Mabutol-Afidchao, M.B.

Citation

Mabutol-Afidchao, M. B. (2013, November 20). *Genetically modified (GM) corn in the Philippines : Ecological impacts on agroecosystems, effects on the economic status and farmers' experiences*. Retrieved from https://hdl.handle.net/1887/22273

Version:	Corrected Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/22273

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/22273</u> holds various files of this Leiden University dissertation

Author: Mabutol-Afidchao, Miladis B.

Title: Genetically modified (GM) corn in the Philippines : ecological impacts on agroecosystems, effects on the economic status and farmers' experiences **Issue Date:** 2013-11-20



Analyzing the farm level economic impact of GM corn in the Philippines

Miladis M. Afidchao, C.J.M. Musters, Ada Wossink, Orlando F. Balderama and Geert R. de Snoo

Submitted to: Netherlands Journal of Agricultural Science (NJAS)

Abstract

This paper analyses the farm economic viability of GM corn as compared to non-GM corn isohybrids in the Philippines. Data was collected from 114 farmers in Isabela province including non-GM, *Bt*, *Bt*HT and HT corn farmers. Results of univariate analysis showed that non-GM corn was not statistically different from GM *Bt*, *Bt*HT and HT corn in terms of production output (PO), net income (NI), production-cost ratio (M) and return on investment (RoI). Multivariate econometric analysis for the agronomic input variables showed a higher RoI at P=0.05 for *Bt* corn as the only difference between seed types. Next, pest occurrence and severity variables were included in the regression to address endogeneity and the Blinder-Oaxaca-decomposition method was used to further investigate differences between growers of *Bt*HT corn and non-GM corn into an endowment and a coefficient effect. The BO analysis showed that *Bt*HT corn has a negative impact on RoI as revealed by the negative signs of the overall mean gap and the characteristics and coefficient components. Contrary to RoI, the overall mean gap indicated that adopting *Bt*HT corn could potentially increase non-GM growers' income mainly from better control of corn borer pest even though mean levels of corn borer occurrence are lower for non-GM growers.

Introduction

The adoption of GM corn, cotton and soybean improves yields and reduces pesticides usage (e.g., Klotz-Ingram *et al.*, 1999; Thirtle *et al.*, 2003; Huesing and English, 2004; Gianessi, 2005). A recent meta-analysis by Finger *et al.*, (2011) of 203 publications on GM corn and cotton provides evidence that these crops lead, on average, to a higher economic performance than conventional crops. Other studies have confirmed these higher averages for specific countries.

In the US for example, the observed overall adoption of glyphosate-resistant crops translated to an annual saving of \$1.2 billion for US farmers in 2001 (Gianessi, 2005). Edgerton *et al.* (2012) estimate that *Bt* corn enabled US farmers to harvest an additional 8.4 million ton of corn in 2010, i.e., an additional average yield benefit of 0.51 ± 0.95 t/ha. In addition, there are important non-pecuniary benefits when adopting GM corn such as increased management flexibility and convenience, savings on machinery use, and human health benefits from reduced handling and use of pesticides (Marra and Piggot, 2006; Brookes and Barfoot, 2009).

Similar estimations have been made for developing countries. For the year 2007, the farm income gains in these countries were estimated at \$302 and \$41 million for *Bt* and HT corn, respectively (Brookes and Barfoot, 2009). The yield increasing effects of GM corn is considered of most importance for developing countries. This is because yield gaps (the difference between farmers' field results and the genetic potential of a crop) tend to be larger in developing countries (Edgerton *et al.*, 2012). Thus, the most obvious pecuniary benefit is increase in yield (Finger *et al.*, 2011; Raney, 2006; Qaim & Zilberman, 2003). In addition there would also be non-pecuniary benefits but no studies have been undertaken to value these for farmers in developing countries (Raney, 2006; Brookes and Barfoot, 2009).

Contrariwise. GM corn and the results reported in studies such as those listed above remain a matter of great controversy. The first issue is with data collection. Yield data for the hybrids might be from breeding programs (field trials) and not from actual production fields. Yield and pesticide usage data from field/farm surveys also has limitations. The main problem with surveys is that neither the early adopters nor the fields chosen for the GM crop are randomly selected leading to a selection bias and this, again, makes a comparison with the non-GM crop problematic (Stone, 2011). Second, studies often provide a partial analysis of yield levels, returns and cost for pest control whereas it is the change in the gross margin which is decisive for farmers' income (Wossink and Denaux, 2006). A further critique is that average figures are misleading and that the performance of GM crops is variable, socio-economically differentiated, and contingent on a range of agronomic and institutional factors (Raney, 2006; Smale et al., 2009; Glover, 2010; Mutuc et al., 2011). There is particularly a need for further evidence on the experience by small, resource-poor farmers. GM crop technology is seen as being capable of benefiting these farmers but this is conditional on institutional settings. For example, the perspective that planting GM seeds would improve the life of poor farmers has been challenged as they have to buy new seeds every season and this makes them dependent on seed suppliers. Finally, many studies build on cross-section data, so that longer term effects have not been analysed (Marvier et al., 2007; Krisna and Qaim, 2012).

Against this background, the present paper evaluates the farm level economic impact of pesticide producing (*Bt*), herbicide tolerant (HT) and stacked gene (*Bt*HT) corn in the Philippines. Asian countries have been slow in the uptake of GM crops that are grown for food and feed and the Philippines is the first and so far only country in Asia to have approved the commercial cultivation of GM corn. After *Bt* corn was first commercialized in the Philippines in 2003, there was a dramatic increase in its adoption. Corn production increased tremendously because yield and farm income levels with *Bt* corn were significantly higher (Yorobe and Quicoy, 2006; Anonymous, 2011). By 2010, GM corn was grown on over a quarter million hectares by 270,000 small-scale, resource-poor Filipino farmers (James, 2010).

In Isabela province, the focus of this paper, yield of *Bt* corn per ha was reported to exceed yield of conventional corn by up to 33% in the 2003-2005 seasons. In 2008-2009, *Bt* and *Bt*HT corn yields surpassed conventional corn by 4-5% and by 13-22%, respectively (Gonzales *et al.*, 2009). Previous studies on the social and economic impact of GM corn in the Philippines (Gonzales *et al.*, 2009; Yorobe and Quicoy, 2006) reported that increased yield and income were the driving factors for the high level of GM corn adoption in the country. Mutuc *et al.* (2011) confirmed the yield enhancing effect of *Bt* corn under poor weather conditions. Yet, a recent study (Afidchao *et al.*, Chapter 6) found striking evidence of negative farmer perceptions with regard to the statement if GM corn could improve their present economic status.

These conflicting findings motivated us to conduct a more in-depth study of the economics of GM corn hybrids in the Philippines. We focus at the farm level rather than at the national level or field level and at the variability across farms/farmers. We take explicitly into account that GM corn seed is substantially higher in price and hard to afford by a resource poor farmer. This price can be up to 84% higher than for non GM-corn depending on the type and number of transgenic traits included in the seed. Thus, to deal with the farm economic issues we seek to know and answer the research question: Is GM corn more economically viable and worth the investment than non-GM corn at the farm level? We investigate farm level differences by corn variety in expenditure for agricultural inputs (labour, seed, and fertilizer costs), gross and net return, production-cost ratio and return on investments. Econometric analyses were done to evaluate if and how agronomic variables (i.e. labour costs, agricultural inputs, corn types and farm area) affect production cost, total return, net income, production-cost ratio and return on investment.

This paper further contributes to the literature by employing the Blinder-Oaxaca decomposition method (Blinder, 1973; Oaxaca, 1973) to decompose the observed differences in economic performance between GM adopters and non-adopters into two components, namely a characteristics effect and a coefficients effect. This decomposition technique is widely used in labour economic applications to study mean outcome differences between groups. For example, the technique is often used to analyse wage gaps by gender or race. More recently it is also used in other areas (Park and Lohr, 2010; Tárrega *et al.*, 2010; Wu *et al.*, 2012). The counterfactual exercise answers the question, what would happen to the GM adopters if their distribution of characteristics was as for the non-GM adopters but if they maintained the returns to their characteristics? A comparison of the counterfactual and estimated performance distribution for the GM group and the non-GM group yields the part of the performance difference that

is attributable to differences in covariates (farm and farmer endowments). The remainder of performance difference is then attributable to differences in returns to covariates. To the best of our knowledge, no other study has employed this decomposition technique to investigate the GM-economic impact nexus.

Material and methods

Area description: GM Corn and the family farm in the Philippines

The Philippines has a total of 9.6 million hectares (32%) agricultural land area of which 51% and 44% are arable and permanent croplands, respectively (Anonymous, 2011). There are ~1.8 million corn farmers in the country and 60% of these cultivate yellow corn. Mostly, these farmers are categorized as small, semi-subsistence farmers with a farm area of less than 4 hectares (Gerpacio *et al.*, 2004). All corn in the country is grown on rainfed non-irrigated land. The cornfields of these small farmers are mostly situated in marginal places. In contrast, most of the large-scale plantations of yellow corn are found in well-situated lowland or upland areas.

Small-scale farmers and their families perform the major agricultural activities such as seeding, harvesting and weeding. These households plant one corn variety, sometimes intercropped with tobacco, fruits (pineapple) and vegetables. Post-harvest activities include de-husking, shelling and grain drying which is done manually by both family and hired labour. Harvested corn is sun-dried immediately after harvest (Gerpacio *et al.*, 2004). This is accomplished on drying pavements at home or on the barangay multipurpose pavements but mostly along paved or asphalted national highways and provincial roads notably in the case study region of Cagayan Valley. The small-scale farmers are dependent on trader-financiers for full-season input financing because they lack the necessary capital. Farmer's payback their loans with a certain interest (~7-15%) either in cash or in corn product upon harvest. The trader-financier decides on the terms of condition of the payback agreement. For large-scale farmers that have large cornfields (cornfield size of more than 3 hectares) hired labour and mechanized farming are common practices.

Among the sixteen regions in the Philippines, the Cagayan Valley region ranks first in terms of corn production. Isabela province in the Cagayan Valley region was chosen as the case study area for the farm level economic assessment. In this province, farm demonstrations showcased the advantages of using GM corn including both its pecuniary and non-pecuniary benefits. One of the non-pecuniary benefits of GM corn, especially of *Bt*HT (insecticide plus herbicide tolerant) corn is that less labour inputs are required for weed management. With proper spraying of herbicide, the weed problem can be reduced or totally controlled. Since GM corn seeds cost are higher than the available commercial iso-hybrid corn in the market, high income and large-scale farmers were the first adopters of this technology. More recently small scale and poor farmers have also adopted the technology. However many poor farmers cannot afford to buy herbicides and still resort to manual weeding in *Bt*HT corn employing the labour force of the (extended) family on a cooperative basis.

Survey

The survey was conducted from October to December 2010 to obtain data for the wet growing season. In order to select our respondents within the group of general farmers who were best able to give us the first-hand information we needed, we applied a purposive sampling technique. Purposive sampling was accomplished of 114 corn farmers in the province of which 42, 8, 44 and 20 were non-GM, *Bt*, *Bt*HT and HT corn adopters, respectively (Table 1). Ninety-percent of the respondents were classified as small scale farmers with farm sizes of not more than 3 ha. Only 10% of the respondents were large scale farmers with farm sizes of 4 to 8 ha.

A self-structured questionnaire was used during the face-to-face interview of the respondents who were from 10 municipalities and 33 villages of the province. The questionnaire was structured to obtain information on respondents' farming background and on costs and returns, i.e. labour cost, input cost and other expenses. The labour cost encompasses the labour service fee for man machine day, man animal day and man day entailed during land preparation and cultivation practices (ploughing, harrowing, furrowing, off-barring and hilling-up), chemical application (fertilizer application and spraying of insecticide and herbicide) and pre- and post-harvesting practices (seed planting, harvesting, threshing, hauling and drying) for the 2010 wet growing season. The service fees for man day include both paid labour (hired labour) and non-paid labour (labour by family members). The corresponding wage per farming practice (e.g. harvesting, spraying) employed was calculated by multiplying the number of labourers to the existing standard service fee given per labourer per day (e.g., harvesting cost=10 persons [paid and unpaid labourers] x \$4.65 per man day). Input cost covers the payment for the seeds, fertilizers, pesticides and other expenses entailed from land preparation to post harvest.

Prior to employing statistical analyses of the data, the total cost of production (TCP), gross income (GI) or production output (PO), net income (NI), production-cost ratio (M) and return on investment (RoI) were computed in US\$ per hectare. Table 2 reports the summation of all the expenses entailed throughout the production to harvest period that was obtained as TCP. The PO refers to the total yield in kg of the 2010 wet season multiplied by the prevailing prize of corn grain per kilogram. The NI was calculated by subtracting TCP from PO. The production-cost ratio (M) was computed as the quotient of the production output and the total cost production per hectare (M=PO/TCP). Finally, the RoI was calculated by subtracting the net income to the product of interest rate paid on loans (IR) and the total cost of production (TCP) i.e. RoI = NI-(IR x TCP).

Univariate and multivariate analysis

A univariate analysis was first employed to evaluate differences on the respondents' information, farming background and production cost and to deal with the single response variables (i.e. corn types, agronomic inputs). A Holm-Bonferroni post hoc test (Quinn and Keough, 2007) was used to assess significant differences of the responses between GM and non-GM adopters.

While the means for production cost, total return, net income, production-cost ratio and return on investment provide the realistic farm economic result of the corn types under farm conditions, a comparison of these means by seed type would be misleading. A correct comparison needs to account for the fact that it is not just (a) the corn type that differs but at the same time (b) many other agronomic inputs and (c) farm characteristics as well. This confounds the impact of seed type on the economic results.

Multivariate analysis was used to evaluate how production output (PO), net income (NI), costproduction ratio (M) and return on investment (RoI) by seed type are directly or indirectly affected by other agronomic input variables. For comparison of the individual response variable between corn types the following conventional production function specification was estimated:

 $y_{i} = \alpha + \beta_{n} x_{ni} + \varepsilon_{i}$ (1)

where y_i denotes the response variables (i.e., the natural logs of PO, NI, M and RoI) in US\$ ha⁻¹ of farm i; α is the intercept and x_{ni} is a vector of the natural logs of the explanatory variables 1..., n of farm i, including labour cost in US\$ ha⁻¹, agricultural input cost (fertilizer, seeds or pesticides) in US\$ ha⁻¹, area planted, corn type, and ε_i is the error term with the usual classical properties. The estimated model was formulated following the Cobb-Douglas production function approach of Yorobe and Quicoy (2006) which is linear in the natural logs of the variables.

Starting from the full model for each of the response variable, stepwise regression analyses were performed through gradual elimination of those variables with insignificant p-values. The final model retains the variables with significant p-values. This enables evaluation which agronomic input variables have influence on the response variables tested in this study. The tables present only the results from the final model obtained after the series of stepwise regression analyses. All econometric analyses were performed using R stat. version 2.12.2.

Blinder-Oaxaca decomposition between GM and non-GM corn

The agronomic production function in eqn (1) above covers only part of the heterogeneity among the farmers that is expected to affect their input and output decisions. To proxy farmers' individual production environment a common approach is to include addition variables in the production function. Particularly important in this context is that GM seed and pesticides are applied in response to pest problems. This can give rise to endogeneity of pesticide use decisions and seed type selection and thus inconsistent parameter estimates. Following Mutuc *et al.* (2010) we included a pest occurrence and a severity variable in the production function to eliminate this potential bias.

Next, to the extended equations the Blinder-Oaxaca decomposition technique was applied to further investigate the mean differences in the response variables between GM and non-GM corn farmers. We assumed that GM corn has advantages compared to non-GM corn in terms of the responses because farmers will not shift to GM corn otherwise. Thus we expect that the non-GM corn have a lower mean of responses as compared to GM *Bt*HT corn.

For the decomposition, the extended equation is estimated separately for two groups of farmers (by seed type):

$$\overline{y}_{GM} = \overline{x}_{GM} \hat{\beta}_{GM} + \hat{\alpha}_{GM} \quad \text{for } n_1 \text{ obs}$$

$$\overline{y}_{nonGM} = \overline{x}_{nonGM} \hat{\beta}_{nonGM} + \hat{\alpha}_{nonGM} \quad \text{for } n_2 \text{ obs}$$
(2)
(3)

Recall that residuals sum to zero in eqs (2) and (3). Next, the mean gap in performance between the two groups of parcels, \overline{y}_{GM} - \overline{y}_{nonGM} , is split into two parts:

$$\overline{y}_{\rm GM} - \overline{y}_{\rm nonGM} = \underbrace{\left(\overline{x}_{\rm GM} - \overline{x}_{\rm nonGM}\right)\hat{\beta}_{\rm GM}}_{(4)} + \underbrace{\overline{x}_{\rm nonGM}\left(\hat{\beta}_{\rm GM} - \hat{\beta}_{\rm nonGM} + \hat{\alpha}_{\rm GM} - \hat{\alpha}_{\rm nonGM}\right)}_{(4)}$$

Mean gap = Characteristics effect + Coefficients effect

where $\bar{x}_{\rm GM}$ and $\bar{x}_{\rm nonGM}$ refer to the means of the explanatory variables, and α and β are the intercept and the coefficient estimates on the explanatory variables for the two samples, respectively. The eqn (4) follows the proposed decomposition formulation of Neumark (1988). Subtracting and adding $\bar{x}_{\rm nonGM} \hat{\beta}_{GM}$ to the right hand side of eqn. (4) and rearrangement gives the decomposition in the characteristics and coefficients effect. An alternative and equally valid formulation in eqn (4) multiplies differences in mean observables characteristics by difference in non-GM coefficient estimates and multiplies differences in coefficient estimates by GM mean observable characteristics.

In eqn (4), the first term of the right-hand side is the part of the performance differential 'explained' by group differences in the predictors, i.e. the part of the gap attributed to differences in observed individual characteristics. The second term is attributable to differences in returns to co-variates, this is the unexplained "coefficient" part. It is important to recognize that this second term includes also all potential effects of differences in unobserved variables. In our case, it is the part of the gap that is due to different returns to the field characteristics and input levels. This second part answers the question if the growers non-GM corn were to switch to GM corn overnight but nothing else observable changed (i.e. the field/farmers' characteristics remained the same) would this lead to better results? A further detailed decomposition examines the percentage contribution of each individual explanatory variable to the total raw differential between the two samples to assess the comparative impact.

A decomposition of the mean gap as discussed above is only useful if the two compared equations are significantly different. Thus, first a Chow test for the difference between eqns. (2) and (3) is required; the null hypothesis is that the parameters of the two equations are equal, meaning that all the independent variables have uniform effects for both subgroups. The formula of the Chow test is:

$$F = \frac{\left(RSS_{\text{pooled}} - \sum RSS_{j}\right)}{\sum RSS_{j}/(n_{1}+n_{2}-2k-2)}$$
(5)

Chapter 5

Table 1. Respondents' information and farming background. Similar superscript letters represent no differences between corn à P<0.05 after post-hoc analysis using

Bonferroni test													
Respondents' Information	non-G	М (n=42)		St (n	8	B	革	(n=44)		HT (n=20)	-	Sig
	Mean	1+	sd	Mean	1+	sd	Mean ±	ŝ	<u>u</u>	Mean	+ sd		
Age	49.29	1+	8.97	43.38	1+	8.90	46.84 <u>+</u>		12.49	49.50	<u>+</u> 12.34	0.939	0.424 ^{ns}
Household size	6.43	1+	3.12	5.00	1+	0.54	5.70 ±	••	2.46	6.30	<u>+</u> 2.64	0.980	0.405 ^{ns}
Years residing in the area	41.95	1+	16.88	34.63	1+	18.10	41.00 <u>+</u>		15.89	43.35	<u>+</u> 16.91	0.560	0.643 ^{ns}
Highest educational attainment ¹	3.93	۱+	1.96	4.75	1+	1.91	4.00 <u>+</u>		1.95	4.70	± 2.20	1.040	0.378 ^{ns}
Cornfield information					1+		1+				I+		
Area of farm devoted to the new variety ²	1.33	1+	1.36	1.56	1+	0.82	2.06 ±		1.43	1.85	± 1.44	2.104	0.104 ^{ns}
Weeds problem ³	0.71	1+	0.46	0.75	1+	0.46	0.59 ±	••	0.50	0.60	<u>+</u> 0.50	0.657	0.580 ^{ns}
Concerns on weeds pest ⁴	2.61ª	1+	0.67	3.50 ^b	1+	0.93	2.91 ^{ab} ±		0.94	2.95 ^{ab}	± 0.97	2.794	0.044*
Asian corn borer (ACB) problem ³	1.00 ª	1+	0.00	0.71 ^b	1+	0.49	0.82 ^{ab} ±	••	0.39	0.89 ^{ab}	<u>+</u> 0.32	3.431	0.020*
Concerns on ACB pest ⁴	3.12	1+	0.83	2.75	1+	1.39	3.07 ±		0.77	2.95	± 0.83	0.517	0.671 ^{ns}
Severity of ACB damage ⁵	3.72	I+	1.02	3.13	1+	1.55	3.42 <u>+</u>		1.03	3.40	± 1.23	0.968	0.411 ^{ns}
¹ Scale: 1-No schooling; 2- Elementary level; 3-	Elemen	tary	graduate	; 4- High	Sch	ool Level; 5	-High School į	grad	uate;				

²Scale: in hectare 1-No schooling; 2- Elementary level; 3- Elementary graduate;

Scale: Yes-1, No-0

4 moderately 1- Highly unconcern

Highly concern;
 Highly severe; 4-

4- severe; 3-tely concern; 3- concern; 2-unconcern; 1- Highly uncc
 moderately severe; 2- negligible; 1- Highly negligible

Results

Respondent's information and farming background

As shown in Table 1, the mean age for the farmers' respondents, of which 25% are female, ranged from 43 to 50. Almost all (94%) of the respondents are married with mean household size of 5-6 members. Respondents have been living in their respective municipality for 35 to 43 years. Most of them reached high school or had a high school diploma or 10th grade. Almost all farmers (98%) in the sample practice mono-cropping. Respondents do not differ significantly in any of these characteristics by corn type.

where RSS_{pooled} is the residual sum of squares (RSS) in the pooled regression, ΣRSS_i is the sum of

the RSS from the two subgroup regressions, k is the number of predictor variables in the model

and n_1 and n_2 are the number of observations in the subgroups (Otineno, 2009). The Chow test

statistic follows an *F*-distribution with k+1 and n_1+n_2-2k-2 degrees of freedom.

On Asian Corn Borer (ACB) infestation, respondents vary in responses on the occurrence, concerns on damage and severity of ACB infestations (Table 1). All farmers encountered weed problem but their level of concern varies. Further analyses, revealed large differences between non-GM and Bt farmers' responses about: a) concerns on weeds pest and; b) the Asian corn borer (ACB) problem (Table 1). The non-GM respondents were less concerned about weeds pest in their farms than the Bt respondents. Likewise, a difference was noted between non-GM and Bt farmers' responses about the existence of the Asian corn borer (ACB) problem in their fields. All non-GM respondents confirmed that they have encountered the ACB problem whilst only part of the *Bt* farmers did encounter the ACB problem in their fields.

GM vs. non-GM corn: Production Cost

The total cost of production (TPC) was obtained by summing up the overall cost entailed by farmers per corn type in one hectare corn production (Table 2). This includes all cost components (labour and agricultural inputs) entailed from pre-harvesting to post harvesting activities. Table 2 showed that non-GM corn had significantly lower mean total cost of production than the total cost of production incurred when using GM corn hybrids.

Univariate analyses showed that the total input cost differed between GM and non-GM corn (Table 2). Agricultural input cost between GM corn types, i.e. Bt vs. BtHT vs. HT, did not differ but all these GM corn types differed from non-GM corn. This corresponds to the big difference in seed cost between GM and non-GM corn. Seed prize of non-GM corn was statistically lower than GM corn. Seeds costs of all the GM corn types were more than 60% higher than non-GM corn. The cost incurred by non-GM farmers for pesticide use was statistically similar to that by GM farmers (Table 2.2). Total labour cost per hectare of production showed no difference between corn types.

ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine pesos). Similar superscript letters repre fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p<0.01, * = c0.05, (*) = <0.10)	esent no	
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine pesos). Similar superscript lett Ferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p<0.01, * = <0.05, (*) = <0.10)	ers repre	
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine pesos). Similar supersci fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p<0.011, ** = c0.05, (*) =	ipt lette	0.10)
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine pesos). Similar. Ferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p<0.01, * = <0.05	supersci	> = (*) ,
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine pesos). Freences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p<0.01, *	Similar	= <0.05
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Philippine fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001, ** = p	pesos).	<0.01, *
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$:42.50 Pl fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** = p<0.001	ilippine	d = **
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha at 1US\$: fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P values: *** =	42.50 Ph	p<0.001
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ per ha a fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test. P value	t 1US\$:	S: *** =
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are in US\$ fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferroni test.	per ha a	P value
ble 2. Cost of production (labour, agricultural inputs, other expenses. Values are fferences between corn varieties at P<0.05 after post-hoc analysis using Bonferr	e in US\$	oni test.
ble 2. Cost of production (labour, agricultural inputs, other expenses. V fferences between corn varieties at P<0.05 after post-hoc analysis using	alues are	Bonferr
ble 2 . Cost of production (labour, agricultural inputs, other expe Ferences between corn varieties at P<0.05 after post-hoc analy:	enses. Va	sis using
ble 2. Cost of production (labour, agricultural inputs, otl fferences between corn varieties at P<0.05 after post-hc	her expe	oc analys
ble 2. Cost of production (labour, agricultural in fferences between corn varieties at P<0.05 after	puts, otł	post-hc
ble 2. Cost of production (labour, agricu fferences between corn varieties at P<0.	ltural in	05 after
ble 2 . Cost of production (labou fferences between corn varieties	r, agricu	s at P<0.
ble 2 . Cost of production fferences between corn	uodal) ר	varieties
ble 2 . Cost of pr fferences betwee	oductio	en corn
ble 2 . Co fferences	sst of pr	s betwei
	ble 2 . Cc	ference:

		-47		D+ [n-0]						LT (n-20)			(01-0)	on los a
Production cost			- + 2	10-11 10								-	-value	h-value
Total Bundunction Cont	Mean For 70b	+	5d 120.60	Mean 660 03	4	sd 20.67	Mean 522 20a	4	sd 110.01	Mean	4	sd 11F OF	600.0	***0000
		- -	00.0CT	76.600		10.01	cc.cc0		+C.011	21.000	-ı ·		700.0	00000
1.Total labour cost	208.60	+1	52.12	231.90	+1	61.90	228.36	+1	47.58	220.17	+1	39.61	1.336	0.267"
Ploughing	42.13	+1	20.91	38.09	+1	13.69	36.47	+1	16.82	37.41	+1	14.44	0.776	0.510 ^{ns}
Harrowing	8.64	+1	7.44	7.53	+1	11.77	9.85	+1	10.58	10.91	+1	11.96	0.362	0.780 ^{ns}
Furrowing	1.34	+1	6.84	2.94	+1	8.32	1.30	+1	3.48	1.59	+1	4.04	0.220	0.882 ^{ns}
Second harrowing	1.01	+1	6.54	2.94	+1	8.32	1.12	+1	3.64	0.00			0.664	0.576 ^{ns}
Basal application	4.91	+1	4.98	9.03	+1	4.93	5.50	+1	4.44	7.34	+1	6.52	2.165	0. 096(*)
Side dress	5.90	+1	4.96	6.62	+1	3.44	7.85	+1	5.29	6.55	+1	6.01	1.025	0.385 ^{ns}
Planting	25.04	+1	10.37	24.82	+1	7.41	29.39	+1	10.06	28.05	+1	15.14	1.264	0.291 ^{ns}
Off-baring	8.10	+1	7.97	8.59	+1	9.55	5.88	+1	8.46	3.79	+1	7.84	1.496	0.220 ns
Hilling-up	3.03	+1	5.89	4.12	+1	8.03	2.98	+1	6.71	1.76	+1	4.51	0.336	0.799 ^{ns}
Total spraying	8.49	+1	9.18	8.65	+1	7.48	8.94	+1	7.72	5.89	+1	4.31	0.736	0.533 ^{ns}
Insecticide Spraying	2.87	+1	3.97	2.35	+1	4.53	2.86	+	1.93	1.06	+1	2.70	066.0	0.400 ^{ns}
Herbicide spraying	5.17	+1	6.85	6.29	+1	6.00	5.60	+	5.76	4.84	+1	4.70	0.150	0.929 ^{ns}
Fungicide Spraying	0.45	+1	1.82	00.0			0.48	+	2.25	0.00			0.478	0.698 ^{ns}
Harvesting	30.84	+1	8.67	37.06	+1	14.38	33.88	+1	6.98	36.29	+1	10.60	2.344	0.077(*)
Thresher	35.30	+1	13.65	39.58	+1	16.52	39.84	+1	13.96	42.49	+1	16.16	1.338	0.266 ^{ns}
Hauling	14.23	+1	9.94	18.85	+1	7.67	19.46	+1	12.86	17.11	+1	7.76	1.770	0.157 ^{ns}
Drying	18.84	+1	10.36	20.32	+1	10.41	22.79	+1	10.57	19.48	+1	8.92	1.177	0.322 ^{ns}
2.Total input cost	262.84 ^b	+1	106.92	393.68ª	+1	79.84	369.10ª	+1	88.46	354.51°	+1	88.08	11.071	0.000***
Seed costs	90.01°	+1	32.46	152.35 ^a	+1	33.55	163.71^{a}	+1	32.11	168.00^{a}	+1	26.14	49.120	0.000***
Total Fertilizer	165.44	+1	88.71	238.38	+1	79.93	198.73	+1	81.08	181.86	+1	72.57	2.309	0.080(*)
Organic	126.86	+1	12.85	157.12	+1	29.79	148.13	+1	12.70	156.06	+1	18.39	0.821	0.485 ^{ns}
Inorganic	36.59	+1	85.09	81.26	+1	68.39	50.60	+1	77.41	34.89	+1	79.07	0.886	0.451 ^{ns}
Pesticides	7.39	+1	8.57	2.94	+1	8.32	6.67	+1	11.73	4.65	+1	8.71	0.678	0.567 ^{ns}
3. Others expenses	33.26	+1	51.79	44.35	+1	26.27	35.93	+1	27.10	28.45	+1	26.58	0.380	0.768 ^{ns}

GM vs. non-GM: Production and Income

The *Bt*HT and HT corns exhibited the highest gross income or production output (PO) as compared to *Bt* and non-GM corn hybrids (Table 3). *Bt*HT and HT corn out yielded non-GM corn by 8% and 7% but non-GM corn out yielded *Bt* corn by 1%. However, there was no statistical difference in PO between GM and non-GM corn.

The computed net income (NI) showed that *Bt*HT exhibited the highest NI followed in descending order by HT, non-GM and *Bt* corn hybrids (Table 3). *Bt*HT and HT corn net income were higher than non-GM corn by 7% and 5%, respectively. The NI of non-GM corn was higher to *Bt* corn by 15%. However, statistics shows that NI was not different between corn types.

The net-cost ratio (M) was computed by corn type. The lowest net-cost ratio was observed for *Bt* corn; yet, this did not differ statistically from other GM corn types and was found to be not significantly different from non-GM corn (Table 3). Finally, we measured the performance of each corn types under study in terms of return on investment (Rol). *Bt*, *Bt*HT and HTcorn had, respectively, 28%, 10% and 6% higher Rol than non-GM corn (Table 3). Yet, the efficiency as reflected from the computed Rol of non-GM corn was found to be statistically not different to the GM corn hybrids.

Table 3. Production output, net income, production-cost ratio and return on investment between corn typescategories using univariate analysis. (Values are in US\$ per ha at 1US\$:42.50 Philippine pesos).

	non-Gl	VI (r	n=42)	Bt	(n=8	3)	<i>Bt</i> HT	(n=	44)	HT (n=2	:0)	F- value	p- value
	Mean	±	sd	Mean	±	sd	Mean	±	sd	Mean	±	sd		
PO	1,103.98	±	539.36	1,071.84	±	455.98	1,299.17	±	372.12	1,272.12	±	442.09	1.671	0.177 ^{ns}
NI	612.28	±	489.98	436.12	±	456.77	687.54	±	345.17	684.21	±	410.63	0.940	0.424 ^{ns}
М	2.28	±	1.01	1.698	±	0.70	2.158	±	0.59	2.208	±	0.71	1.231	0.302 ns
Rol	503.23	±	341.66	885.64	±	676.05	618.39	±	417.93	572.16	±	424.53	2.046	0.112 ^{ns}

PO=Production Output NI=Net Income M=Production-cost ratio RoI=Return on Investment

Multivariate analysis

We applied production function analysis to PO, NI, M and Rol. Before the analysis, we first evaluated the residual plots (residual vs. fitted, normal Q-Q, scale-location and residual vs. leverage) for its normal distribution. Data that were non-normally distributed were ln(x+1) transformed. Data presented here are results of the minimal model per response variable obtained after series of stepwise regression analyses.

In explaining the variation in PO, costs of threshing, harvesting and plowing were found to have the largest effect. Among input cost, differences in seed cost seems to be important as expected from the summary statistics in Table 2. The R² value was estimated 0.53 for the final model used (Table 4 column A).

Table 4 (column B) shows the multi-agronomic variables that are affecting NI. Area planted and fungicide spraying had the highest impact on NI. The R² estimate values are 0.39 for the final model.

On M, variables such as area, fertilizer and labour costs for thresher showed great influence (Table 4 column C). The R² estimate values are 0.39 for the final model.

This analysis showed that *Bt* corn had a significantly higher Rol than non-GM corn (Table 4 column D), although the overall effect of corn type was not significant (table 3) whilst none of the other tested agronomic variables did have an effect (R^2 : 0.05).

Blinder-Oaxaca decomposition

Next the regression equations as above were extended with a pest occurrence and a severity variable to eliminate potential endogeneity bias. The equations for two response variables (i.e. Rol and NI) and *Bt*HT and non-GM corn were selected for the BO analysis on the basis of the results obtained after subjecting the extended regression models for all the response variables to a Chow test as shown in Table 6.

For the decomposition, the RoI and NI equations are estimated separately as discussed above. The regression results for return on investment (Table 7, column 4 and 5) show that among the assessed variables, corn borer occurrence and costs of labour, seeds and pesticides manifested significant negative effects on GM corn. It is interesting to note that together with farm size and fertilizer, corn borer severity showed positive effects on GM corn's RoI. For non-GM corn, the costs of seeds and pesticide have significant positive effects. All other variables including corn borer occurrence and severity show significant negative effects on non-GM corn's RoI.

The estimated models were then used to split the observed gap between corn types in two portions (Tables 7, last three columns). The sum in the bottom row of Table 7 shows that of the overall raw gap of -3.397 for Rol only 21% (-0.705) can be explained by differences in characteristics of the two samples. The remaining 79% (-2.692) can be attributed to the coefficient or unexplained effect. Notice that the gap is negative and thus the switch to GM corn would mean a drop in Rol for the farmers on average. The last two columns of Table 7 present the contribution of each explanatory variable to the explained and the unexplained component, respectively. In terms of the explained part, most important contributions to explaining the negative gap come from the seed cost (147%) followed by some distance by labour costs (20%). Notice that all the other characteristics reduce the gap (negative percentages).

For NI, the regression results in Table 8 show that among the assessed variables, seed cost, fertilizer cost and corn borer severity carry negative signs. These are variables which manifest negative effects on NI. Farm size, labour cost and pesticide cost have positive signs hence, exhibit significant positive effects on NI for both corn types. Further analysis shows that the two main parts of the mean gap (1.144) have opposite signs; we find a small negative characteristics effect (-23%) and a large positive coefficient effects (123%). In particular, among the explanatory variables of the negative characteristic components, seed cost has the largest percentage (112%) followed by fertilizer cost (61%). Except for farm size and labour cost, the remaining characteristics contribute to increasing the negative gap (positive percentages). Contrary to Rol, the overall gap indicates that adopting GM corn could potentially increase the growers' income. The results in the last two columns of Table 8 show that the mean income advantage from switching to *Bt*HT corn is mainly due to better control of corn borer pest.

Table 4. Estimates of agronomic variables identified to affect PO, NI, M and Rol ha⁻¹ employing series stepwise regression analyses. All data was natural log (In) transformed. P values: *** = p<0.001, ** = p<0.01, * = <0.05, (*) = <0.10); L_i = man labor cost ha⁻¹; I_i = agricultural input cost ha⁻¹; SE = standard error.

	(A) Production Output (PO) [r ² =0.526]	(B) Net Income (NI) [r²=0.391]	(C) Production- Cost Ratio (M) [r ² =0.386]	(D) Return on Investment (Rol) [r²=0.053]
	Estimate ± se	Estimate ± se	Estimate ± se	Estimate ± se
Intercept				
Corn types (C _i): Non-GM corn	3.698** ± 0.743	-1181.230 ± 297.630	1.029 ± 0.262	503.190 ^{***} ± 63.990
Contrast with intercept				
-Bt corn	-0.011 ± 0.144	-193.200 ± 133.710	-0.085 ± 0.086	382.560 [*] ± 159.970
-BtHT corn	0.036 ± 0.103	-15.450 ± 77.290	-0.018 ± 0.050	115.220 ± 89.460
-HT corn	-0.002 ± 0.121	-34.550 ± 96.260	-0.037 ±0.061	69.010 ± 112.660
Covariates				
Plowing cost (L _{1i})	-0.002** ± 0.050			
Furrowing cost (L _{2i})	-0.056* ± 0.047	-92.860* ± 41.630	-0.047* ± 0.026	
Second harrowing cost $(L_{_{3i}})$	-0.124* ± 0.058			
Insecticide spraying $(L_{_{4i}})$	$0.081^{*} \pm 0.031$			
Harvesting cost $(L_{_{5i}})$	0.112 ^{**} ± 0.118			
Thresher cost (L _{6i})	0.575*** ± 0.081	125.050 ^(*) ± 77.030	0.286 ^{***} ± 0.052	
Side dress cost (L _{7i})		-66.590 ^(*) ± 31.130	-0.048* ± 0.020	
Fungicide spraying (L _{8i})		504.290*** ± 82.010		
Seed cost (I _{1i})	0.143* ±0.125			
Fertilizer cost (I _{2i})			-0.178*** ± 0.041	
Area planted (A _{ri})	0.077** ± 0.024	83.940** ± 25.000	0.045** ±0.016	

 Table 5. Correlation values (upper) and p-values (lower) between corn agronomic variables ha⁻¹. (PO= production output/yield; NI = Net income; M= Cost-production ratio; Rol=

 Return on investment; TPC= Total Production Cost). P values: *** = p<0.001, ** = p<0.01, * = <0.05, (*) = <0.10), ns = not significant</td>

Parametric contration coefficient. Pearson's contration coefficient. Pearson's contration coefficient. Poince Poince Nin Nin No Fact inter Seat cost Co														
Po Ni Ni Rearron's correlation coefficient. Antibles Po Ni Ni Rearron's correlation coefficient. Poto Dev Dev <thdev< th=""> <</thdev<>		Weeds	-0.031	-0.016	-0.060	-0.109	0.074	0.052	0.084	0.003	0.150	0.209	0.069	
Parametericitation coefficient. Parametericitation coefficient. Variable. PO N No No TPC Labor Fartilizer Seed Pesticitie Area PO mm mo mo mo mo mot mod mod mode mo mode mode mode<		ACB	-0.025	-0.090	-0.067	-0.022	0.129	0.026	0.111	0.224	-0.091	0.083		0.468
Pariables Pariables Person's corretation coefficient Aniables PO NI M Rol TPC Labor Fertilizer Seed Position PO 2.2e-16 ^{···} 0.898 0.724 0.096 0.425 0.344 0.137 0.273 0.006 NI <.2.2e-16 ^{···} 0.883 0.724 0.096 0.425 0.344 0.138 0.017 NI <.2.2e-16 ^{···} 0.863 0.075 0.160 0.237 0.273 0.007 NI <.2.2e-16 ^{···} 0.863 0.074 0.014 0.014 0.138 0.017 NI <.2.2e-16 ^{···} 0.244 0.014 0.014 0.138 0.007 NI <.2.2e-16 ^{···} 0.244 0.014 0.137 0.075 NI <.2.2e-16 ^{···} 0.1044 0.014 0.137 0.075 NI <.2.2e-16 ^{···} 0.0926 ^{···} 0.241 ^{···} <th></th> <th>Area planted</th> <td>0.300</td> <td>0.287</td> <td>0.171</td> <td>0.000</td> <td>0.222</td> <td>0.061</td> <td>0.346</td> <td>0.310</td> <td>-0.089</td> <td></td> <td>0.378 ns</td> <td>0.025</td>		Area planted	0.300	0.287	0.171	0.000	0.222	0.061	0.346	0.310	-0.089		0.378 ns	0.025
Pariables Po NI Rearron's correlation coefficient Point Point NI Rol TPC Labor Fertilizer seed NI < 2.2e-16 ^{···} 0.398 0.724 0.096 0.425 0.344 0.237 0.273 NI < 2.2e-16 ^{···} < 0.898		Pesticide cost	0.006	-0.017	-0.077	-0.064	0.272	0.073	0.076	-0.089		0.349 ns	0.337 ^{ns}	0.110
Parson's correlation coefficient Variables PO Ni Moi Roi TPC Labor Fartilizer PO 2.22e-16 ^{···} 0.898 0.724 0.096 0.425 0.344 0.237 Ni <.2.2e-16 ^{···} 0.863 0.075 0.160 0.209 0.014 Ni <.2.2e-16 ^{···} 0.863 0.075 0.160 0.209 0.014 Ni <.2.2e-16 ^{···} 0.863 0.075 0.160 0.209 0.014 Ni <.2.2e-16 ^{···} 0.863 0.079 0.079 0.014 0.014 Ni <.2.2e-16 ^{···} 0.309 ^{···} 0.221 ^{···} 0.079 0.014 0.014 M <.2.2e-16 ^{···} 0.090 ^{···} 0.021 ^{···} 0.079 0.014 0.014 M 0.309 ^{···} 0.021 ^{···} 0.021 ^{···} 0.014 0.014 0.014 Fertilizer cost 0.001 ^{···} 0.022 ^{···} 0.926 ^{···} 0.926 ^{···} 0.411 ^{···} 0.411 ^{···}	L	Seed cost	0.273	0.138	-0.007	0.157	0.506	0.073	0.270		0.086 ^(*)	0.001**	0.016*	0.973
Variables PO Ni Moi TPC Dearson's correlatio Variables PO Ni M Noi TPC Labor PO \sim 0.898 0.724 0.096 0.425 0.344 Ni $<$ 2.2e-16 ^{···} \sim 0.863 0.075 0.160 0.209 M $<$ $<$ \sim 0.309 ^{···} $<$ 0.204 0.001 M $<$ $<$ $<$ $<$ $<$ $<$ M $<$ $<$ $<$ $<$ $<$ $<$ M $<$ $<$ $<$ $<$ $<$ $<$ $<$ M $<$ $<$ $<$ $<$ $<$ $<$ $<$ M $<$ $<$ $<$ $<$ M	n coefficient	Fertilizer cost	0.237	0.014	-0.116	0.025	0.795	0.188		0.004**	0.423 ^{ns}	0.065(*)	0.240 ns	0.374
Artiables PO NI Moi Roi TPC PO 22.2e-16 ^{···} 0.898 0.724 0.096 0.425 NI <.2.2e-16 ^{···} 0.863 0.075 0.160 0.425 M <.2.2e-16 ^{···} 0.888 0.724 0.096 0.425 M <.2.2e-16 ^{···} 0.271 ^{···} 0.075 0.160 0.014 M <.2.2e-16 ^{···} 0.271 ^{···} 0.075 0.160 M <.2.2e-16 ^{···} 0.090 ^{···} 0.014 0.014 M 0.014 0.014 Mol 0.014 0.014 Labor cost Fertilizer cost Fertilizer cost	's correlatio	Labor cost	0.344	0.209	0.001	600.0	0.579		0.045*	o.441 ^{ns}	0.437 ^{ns}	0.516 ^{ns}	0.784 ^{ns}	0.586
Variables PO NI M Rol PO 2.2e-16*** 0.898 0.724 0.096 NI < 2.2e-16***	Pearson	ТРС	0.425	0.160	-0.014	0.079		1.5e-11***	<2.2e-16***	9.6e-09***	0.324 ^{ns}	0.018*	0.172 ^{ns}	0.431
Variables PO NI M PO 0.898 0.724 NI < 2.2e-16 ^{***} 0.863 NI < 2.2e-16 ^{***} 0.863 M 0.309 ^{***} 0.039 ^{***} Pace 0.309 ^{***} 0.039 ^{***} Pace 0.000 ^{***} 0.099 ^{***} Pesticide cost 0.001 ^{***} 0.939 ^{***} Pesticide cost 0.927 ^{***} 0.069 ^{***} Area planted 0.001 ^{***} 0.000 ^{****} Meeds		Rol	0.096	0.075	0.104		0.401 ^{ns}	0.926 ^{ns}	0.794 ^{ns}	0.095(*)	0.497 ^{ns}	su 266.0	0.816 ^{ns}	0.248
Variables PO NI PO 0.898 NI NI <2.2e-16***		Σ	0.724	0.863		0.271 ^{ns}	0.035*	0.994 ns	0.002**	^{su} 6£6.0	0.414 ^{ns}	0.069(*)	0.476 ^{ns}	0.527
Variables PO PO PO NI < 2.2e-16***		IN	0.898		< 2.2e-16***	0.428 ^{ns}	0.090(*)	0.026*	0.884 ^{ns}	0.144 ^{ns}	0.857 ^{ns}	0.002**	0.341 ^{ns}	0.868
Variables Variables M M M M M M M M M M M M M		Ю		< 2.2e-16***	< 2.2e-16***	0.309 ^{ns}	2.48e-06***	0.000	0.011*	0.003**	0.952 ^{ns}	0.001**	0.788 ^{ns}	0.741
> salues -9		ariables	РО	ĪZ	Σ	Rol	трс	Labor cost	Fertilizer cost	Seed cost	Pesticide cost	Area planted	ACB	Weeds
		>		<u> </u>	<u> </u>	<u> </u>	sənli	ev -9	<u> </u>					

Table 6. Chow test outo	come for production output, net income, return on investment and cost-production ratio.
P values: ** = p<0.01,	ns = not significant

Response variable	df numerator	df denominator	RSS _{BtHT}	RSS _{non-GM}	ΣRSS	F	p-values
Production output	8	70	3.252	9.120	13.210	0.592599	0.955569 ns
Net Income	8	70	21.380	198.940	298.978	3.12391	0.000158**
Return on Investment	8	70	177.493	106.396	358.910	2.31229	0.003652**
Cost-Production Ratio	8	70	50.719	119.100	168.703	0.057502	1.000000 ns

RSS = residual sum of squares in the pooled regression; ΣRSS = sum of the RSS

Table 7. The Blinder-Oaxaca decomposition of the return on investment (ROI) of GM and non-GM corn types.P values: ** = p<0.01, * = <0.05, (*) = <0.10), ns = not significant</td>

Explanatory	Charact	eristics	Coeffi	icients	ROI GM evaluated at means	ROI non-GM evaluated at means	Mean Gap	Characteristics Effects (share in %)	Coefficients effects (share in %)
Variables	\overline{X}_{GM}	\overline{X}_{nonGM}	\hat{eta}_{GM}	${\hat eta}_{nonGM}$	$\hat{eta}_{GM}\overline{X}_{GM}$	$\hat{eta}_{nonGM}\overline{X}_{nonGM}$		$\left(\overline{X}_{GM} - \overline{X}_{nonGM}\right)\hat{\beta}_{GM}$	$\overline{X}_{nonGM}\left(\hat{m{eta}}_{GM}-\hat{m{eta}}_{nonGM} ight)$
Intercept	1.000	1.000	11.413	6.806	11.413	6.806	4.607		4.607 (-171%)
Farm size	2.182	1.500	0.042 ^{ns}	-0.341 ^{ns}	0.091	-0.512	0.602	0.028 (-4%)	0.574 (-21%)
Labor cost	228.341	208.571	-0.007 ^{ns}	-0.002 ^{ns}	-1.598	-0.417	-1.181	-0.138 (20%)	-1.043 (39%)
Seed cost	163.818	90.024	-0.014 ^{ns}	0.033**	-2.293	2.971	-5.264	-1.033 (147%)	-4.231 (157%)
Fertilizer cost	198.750	165.405	0.004 ^{ns}	-0.007(*)	0.795	-1.158	1.953	0.133 (-19%)	1.819 (-68%)
Pesticides	6.727	7.429	-0.287 ^{ns}	0.318 ^{ns}	-1.931	2.362	-4.293	0.201 (-29%)	-4.495 (167%)
Corn borer severity	3.659	3.439	0.127 ns	-0.730*	0.465	-2.510	2.975	0.028 (-4%)	2.947 (-109%)
Corn borer occurrence	1.091	1.119	-2.689(*)	-0.123 ns	-2.934	-0.138	-2.796	0.075 (-11%)	-2.871 (107%)
Sums					4.007	7.405	-3.397	-0.705 (100%)	-1.692 (100%)

Table 8.	The Blinder-Oaxaca decomposition of the Net Income (NI) of GM <i>Bt</i> HT and non-GM corn types.
P values	: (*) = <0.10), ns = not significant

Explanatory	Charac	teristics	Coeffi	cients	NI GM evaluated at means	NI non-GM evaluated a means	Mean Gap	Characteristics Effects (share in %)	Coefficients effects (share in %)
Variables	\overline{X}_{GM}	\overline{X}_{nonGM}	$\hat{eta}_{_{GM}}$	\hat{eta}_{nonGM}	$\hat{eta}_{_{GM}}\overline{X}_{_{GM}}$	$\hat{eta}_{nonGM}\overline{X}_{nonGM}$		$\left(\overline{X}_{GM} - \overline{X}_{nonGM}\right)\hat{\beta}_{GM}$	$\overline{X}_{nonGM}\left(\hat{\beta}_{GM}-\hat{\beta}_{nonGM}\right)$
Intercept	1.000	1.000	4.522	5.321	4.522	5.321	-0.799		-0.799 (-57%)
Farm size	2.182	1.500	0.113 ^{ns}	0.526 ns	0.247	0.788	-0.542	0.077 (-29%)	-0.619 (-44%)
Labor cost	228.341	208.571	0.010(*)	0.012 ns	2.215	2.461	-0.246	0.192 (-73%)	-0.438 (-31%)
Seed cost	163.818	90.024	-0.004 ^{ns}	-0.001 ns	-0.655	-0.108	-0.547	-0.295 (112%)	-0.252 (-18%)
Fertilizer cost	198.750	165.405	-0.005 ^{ns}	-0.003 ns	-0.954	-0.568	-0.386	-0.160 (61%)	-0.226 (-16%)
Pesticides	6.727	7.429	0.074 ^{ns}	0.013 ns	0.495	0.093	0.402	-0.052 (20%)	0.454 (32%)
Corn borer severity	3.659	3.439	-0.014 ^{ns}	-0.182 ns	-0.053	-0.627	0.574	-0.003 (1%)	0.577 (41%)
Corn borer occurrence	1.091	1.119	0.762 ^{ns}	-1.660 ns	0.831	-1.858	2.689	-0.021 (8%)	2.710 (193%)
Sums					6.648	5.504	1.144	-0.263 (100%)	1.407 (100%)

Discussion

GM corn effect on Cost

One of the most often highlighted reasons for non-GM corn adopters is the high cost of seed per hectare of corn production (Afidchao *et al.*, Chapter 6). This study once again shows that cost of seeds per hectare was far higher for GM corn than for the leading conventional corn hybrids available on the market. This is also one of the main factors influencing the high level of total production cost for GM corn (Tables 2).

Reduction of pesticides usage is one of the benefits that was promised to be achieved when using GM corn (Mutucet al., 2011; Brookes and Barfoot, 2009; Kleter et al., 2007; Wilson et al., 2005; Huang et al., 2003; Rice, 2003). Yet, our study showed that pesticide cost entailed in all corn types are statistically the same. Our result confirms results reported by Afidchao et al. (Chapter 6) where BtHT and HT farmers perceived no reduction in pesticides usage and exposure. This is likewise supported by the findings of Wossink and Denaux (2006) where efficiency of pest control cost between transgenic and conventional cotton found no statistical difference. Finally, the claim that pesticides usage can be reduced was not supported by our study as shown in Table 4. Although, it has been shown in US and Europe that GM corn reduce pesticide and its environmental footprints at 14% reduction rate (Brookes and Barfoot, 2006). Also, a savings of \$25-\$75/acre due to no insecticide is achieved with Bt corn (Rice, 2003). This reduction in pesticide usage observed in US was not manifested at the farm-scale level in Isabela province plausibly due to Bt farmers' fear and anticipation of yield loss by pests other than ACB. Hence insecticide spraying is usually done even with Bt seed and by HT farmers who opted to have manual weeding due to financial constraints (no money to buy herbicide).

GM corn effect on production, income and return on investment

In terms of yield or production output, our result for conventional and *Bt* corn was similar to the comparisons of yield in 2004-2005 and 2007-2008 in the Philippine provinces of General Santos City and Isabela, respectively where conventional corn was statistically higher than GM corn (Gonzales *et al.*, 2009).

*Bt*HT and HT corn produced on average higher yields, *Bt* corn lower than non-GM corn, but these differences were not statistically significant. This shows that GM corn has no straight forward overall advantages compared with non-GM corn. *Bt* corn may produce higher yields (Dilehay *et al.*, 2004; Stanger & Lauer, 2006; Qaim & Zilberman, 2003; Rice, 2003) but other additional points should be taken into account when assessing economic returns. As stated by Dilehay *et al.* (2004) and Stanger & Lauer (2006), *Bt* corn has higher grain moisture, lower test weight and higher harvest & seeds cost; these counterweigh increased yield and might result in adding no benefits when using GM corn. Ma and Subedi (2005) show that on the same maturity, non-*Bt* corn accumulates more nitrogen and leads to highest grain yield. In addition, low to moderate infestation of corn borer provides no advantage in using *Bt* corn. According to

Wolf and Vögeli (2009) using *Bt* corn, an increased yields of up to 15 percent can be obtained when infestation is severe to very severe but at low and moderate infestation conventional maize hybrids are superior when appropriately grade-selected.

The severe to moderately severe ACB infestation in the respondents' cornfields (Table 1) indicate that ACB is still a pest problem in Isabela province. Under high pressure of corn borer infestations, *Bt* corn should have yield advantages. However, in our study the production in GM corn did not exhibit significant yield advantages compared with non-GM corn (Table 3). In the same vein, the reduction of weeds incidence using herbicide tolerant (*Bt*HT and HT) corn varieties (Table 1) did not result in economic advantages compared with non-herbicide tolerant corn varieties (non-GM and *Bt* corn) (Table 3).

In our study, the Rol did not significantly differ among corn types thereby supporting the experimental data of Nolte and Young (2002). Nolte and Young (2002) found no differences between GM herbicide tolerant and conventional corn hybrids in terms of economic return in their 1999 field experiment. Although they have seen significant variations between these corn types in 2000 yet the grain yield effect was stronger than the corn type effect. However, in our study, econometrics showed that Rol could be positively influenced by corn types specifically; *Bt* corn had a significantly higher Rol than non-GM corn (Table 4).

The findings of our study on non-significant difference in mean PO, NI, M and Rol among corn types (Table 3) do not show more profits when using GM corn. In particular, our data did not affirm that *Bt* corn adoption could provide higher yield (Stanger & Lauer, 2006; Dilehay *et al.*, 2004) and higher profits (Qaim & Zilberman, 2003; Rice, 2003). Hence, this does not support the general concept that GM corn provides higher income than non-GM corn. Relatively, our study supplement the data of Baute *et al.* (2002) which refuted the notion that *Bt* corn hybrids in general are higher yielding compared to conventional corn.

Lastly, past studies (Yorobe and Quicoy, 2007 and Gonzales *et al.*, 2009) stated that the farmers that adopted GM corn found it profitable, i.e. the farmers with high risks of ACB, have adopted GM corn by now. Yet, in our present study we found that with moderately severe ACB infestation as observed by the respondents (see Table 1), GM corn did not manifest advantage in terms of profit. This means that further increase of GM corn is no longer profitable, although it might have been in the past.

Agronomic variables effect on PO, NI and M

Several variables could substantially affect PO as shown in Table 4. Labour cost, agricultural input cost and area planted are the influential variables on PO. On labour cost, plowing, harvesting and thresher are noted to greatly affect PO. Among the agricultural inputs, seed cost was shown to have great influence on PO. This may indicate that an increase in PO could require a high input of seeds. Lastly, area planted could as well influence PO. Increasing area planted results to a higher PO and this was supported by the positive and significant correlations of PO.

The relationship between PO and NI was strong and positive as shown in Table 5. Yet econometric shows different inputs have an effect on NI. For NI fungicide spraying is the most important input, followed by the area planted.

The agronomic variables like area, fertilizer and labour costs for furrowing, side dress fertilizer application and thresher are shown to influence M (Table 4). This demonstrates that an increase in area devoted to corn leads to an increase in production cost and production output. On the other hand, fertilizer cost that constitutes around 33 to 44% in the cost production depending on corn types (Gonzales *et al.*, 2009), showed a significant negative correlations to M. This directs us to the point that any increase in fertilizer inputs does not warrant higher production (correlation= -0.116; p-value: 0.002, table 5). Lastly, an increase in production or yield also entails an increase in thresher cost with positive significant correlations of 0.589 (p-value= 5.448e-12, table not shown).

Finally, the econometric analysis revealed that among the tested agronomic variables, area planted is the variable that has encompassing positive influence to PO, NI and M. This further mean that any increase in area of corn plantation may contribute to the increase in yield and income as well as production cost.

Blinder-Oaxaca decomposition of return on investment and net income

The RoI and NI as such showed insignificant differences between GM and non-GM corn types and suggest that GM corn does not show superior economic performance compared with non-GM corn (Table 3). Thus, the cultivation of an iso-hybrid non-GM corn seems to have comparable economic results in terms of RoI and NI based on this partial analysis. The application of BO analysis in this study served to check whether other characteristics (such as agricultural input, cost and cornfield pest history) that vary at the same time as seed type could explain some of the difference and thus might confound the overall assessment and determination of which corn type is worth investing. For GM *Bt*HT and non-GM corn growers, the two largest subsamples in this study, the Chow test revealed that there are indeed concomitant differences in the other underlying characteristics (Table 6).

The BO technique served to compare the contribution of independent variables RoI and NI between GM *Bt*HT and non-GM corn through the distinction of an observable characteristics effects and an unexplained coefficient effect. The coefficient component can have a different sign from the characteristics component and this can give insightful information in particular. If both components have the same sign, differences in RoI or NI are as expected. A situation of opposite signs and a substantial coefficient effect is often associated with discrimination in the sociological and labour economics literature in which the Blinder-Oaxaca decomposition is commonly applied.

The last two columns of Table 7 show that for RoI the sums of the two components have identical signs (negative). However for individual variables differences in signs do occur. For

both pesticide costs and for corn borer occurrence there is a negative impact on the Rol which is unexpected given the lower average for these variables for the non-GM sample. Finally, the intercept is responsible for most of the coefficients effects indicating the contribution of unobservable characteristics (such as physico-chemical characteristics of cornfields) to the difference in Rol.

In contrast, in case of NI, the sums of the two components shown in the last two columns of Table 8 do not have identical signs. The characteristics effect making up a small portion (23%) of the gap bears a negative sign. This indicates a negative effect on NI by the differences in *Bt*HT and non-GM farmers' observable characteristics which is mainly attributed to seed costs and costs of fertilizer inputs. However this is counteracted by the coefficients or unexplained component which has carry a positive sign and is mainly due to pesticide input, corn borer severity and occurrences. In general, this shows that *Bt*HT has disadvantages on NI based on observable characteristics yet, could provide economic advantage overall due to better pest control even for cornfields less heavily infested with corn borer pest and also due to savings on pesticide costs.

Conclusion

This study focused on small-scale farmers as they constitute the majority of corn farmers and are usually at the bottom in the economic production spectrum. They are likewise the most vulnerable groups easily malleable to be influenced with new introduced technologies that promise superior economic gains. The vast increment and wide-scale cultivation of GM corn in the Philippines is attributed to risk-averse farmers as well as driven by economic benefits offered by these novel varieties. While it is true that past studies showed the adoption of GM corn could increase yield and provide more profits to farmers, our study showed no difference in production output between corn varieties anymore. This study showed that GM corn adoption does no longer directly provide superior economic advantage against non-GM corn considering all the variables studied.

We found that the Blinder-Oaxaca decomposition technique usually used in racial and gender discrimination studies can as well be applied to agriculture economic related studies. Employing this technique allowed us to compare and identify variables with marked influences on the results of our study. Finally, this study can be undertaken on a larger scale to obtain more information on the economic benefits from GM corn technology overtime viz a viz its wide scale adoption in different economic settings and locations.

Acknowledgement

We would like to extend our appreciation and gratitude to the farmers for their cooperation and unselfishly sharing their first-hand information with us. In addition, we thanks to the following people: A. Domingo Sr., A.M. Vanyvan, R.G. Mabutol Jr., R.M. del Rosario, P.M. Afidchao, R.G. Salazar and W.B. Saliling. This study was supported by the Louwes scholarship program of Leiden University in the Netherlands.

References

- Afidchao, M.A., Musters, C.J.M., O.F. Balderama, G.R. Snoo de, Chapter 6. Farmers' adoption, perceptions and hurdles of GM corn in the Philippines.
- Anonymous, 2011. Country STAT Philippines. In: http://countrystat.bas.gov.ph/. Accessed on August 7, 2011.
- Baute, T.S., Sears, M.K., Schaafsma, A.W., 2002. Use of transgenic *Bacillus thuringiensis* Berliner corn hybrids to determine the direct economic impact of the European corn borer (Lepidoptera: Crambidae) on field corn in Eastern Canada. Journal of Economic Entomology 95 (1): 57-64.
- Blinder, A.S., 1973. Wage discrimination: reduced form and structural estimates. Journal of Human Resources 8(4): 436-455.
- Brookes, G., Barfoot, P., 2009. Global impact of biotech crops: Income and production effects 1996-2007. AgBioForum 12(2): 184-208.
- Brookes G., and P. Barfoot, 2006. Global impact of biotech crops: Socio-economic and environmental effects in the first ten years of commercial use. AgBioForum 9(3): 139-151.
- Dillehay, B.L., Roth, G.W., Calvin, D.D., Kratochvil, R.J., Kuldau, G.A., Hyde, J.A., 2004. Performance of *Bt* corn hybrids, their near isolines, and leading corn hybrids in Pennsylvania and Maryland. Agronomic Journal 96: 818-824.
- Edgerton, M.D. et al., 2012. Transgenic insect traits increase corn yield and yield stability. Nature biotechnology 30: 493-496.
- Finger, R., El Benni, N., Kaphengst, T., Evans, C., Herbert, S., Lehmann, B., Morse, S., Stupak, N., 2011. A Meta-analysis on farm-level costs and benefits of GM crops. Sustainability 3: 743-762.
- Gerpacio, R.V., Labios, J.D., Labios, R.V, and Diangkinay, E.I., 2004. Maize in the Philippines: Production Systems, Constraints and Research Priorities. CIMMYT, El Batan, Mexico.
- Gianessi, L.P., 2005. Economic and herbicide use impacts of glyphosate-resistant crops. Pest Management Science 61(3): 241–245.
- Glover, D., 2010. Is *Bt* Cotton a Pro-poor Technology: A Review and Critique of the Empirical Record. Journal of Agrarian Change 10(4): 482-509.
- Gonzales, L.A., Javier, E.Q., Ramirez, D.A., Cariño, F.A., Baria, A.R., 2009. Modern biotechnology and agriculture: A history of the commercialization of biotech maize in the Philippines. STRIVE Foundation. ISBN 978-971-91904-8-6.
- Huesing, J., English, L., 2004. The impact of *Bt* crops on the developing world. AgBioForum 7(1&2): 84-95.
- Huang, J., Hu, R., Pray, C., Qiao, F., Rozelle, S., 2003. Biotechnology as an alternative to chemical pesticides: a case study of *Bt* cotton in China. Agricultural Economics 29: 55-67.
- Kleter, G.A., Bhula, R., Bodnaruk, K., Carazo, E., Felsot, A.S., Harris, C.A., Katayama, A., Kuiper, H.A., Racke, K.D., Rubin, B., Shevah, Y., Stephenson, G.R., Tanaka, K., Unsworth, J., Wauchope, R.D., Wong, S.S., 2007. Altered pesticide use on transgenic crops and the associated general impact from an environmental perspective. Pest Management Science 63: 1107-1115.
- Klotz-Ingram, C., Jans, S., Fernandez-Cornejo, J., McBride, W., 1999. Farm-level production effects related to the adoption of genetically modified cotton for pest management. The Journal of Agrobiotechnology Management and Economics 2(3): 73-84.

- Krisna, V.V., Qaim, M., 2012. *Bt* Cotton and sustainability of pesticide reductions in India. Agricultural Systems 107: 47-55.
- James, C., 2010. ISAAA's Brief No. 42: Global Status of Commercialized Biotech/GM Crops.
- Ma, B.L., Subedi, K.D., 2005. Development, yield, grain moisture and nitrogen uptake of *Bt* corn hybrids and their conventional near-isolines. Field Crops Research 93(2-3): 199-211.
- Marra, M.C. and Piggot, N.E., 2006. The value of non-pecuniary characteristics of crop biotechnologies: A new look at the evidence. In R. Just, J. Alston and D. Ziberman (eds.), Regulating agricultural biotechnology: Economics and policy 145-178. New York: Springer-Verlag Publishers.
- Marvier, M., McCreedy, C., Regetz, J., Kareieva, P., 2007. A meta-analysis of effects of *Bt* cotton and maize on nontarget invertebrates. Science 316(5830): 1475-1477.
- Mutuc, M.E., Rejesus, R.M., Yorobe, J.M. Jr., 2011. Yields, Insecticide Productivity, and *Bt* Corn: Evidence form Damage Abatement Models in the Philippines. AgBioForum 14(2): 35-46.
- Nolte, A.N., Young, B.G., 2002. Efficacy and Economic Return on Investment for Conventional and Herbicide-Resistant Corn (*Zea mays*). Weed Technology 16 (2): 371-378.
- Neumark, D. 1988. Employers' Discriminatory Behavior and the Estimation of Wage Discrimination. Journal of Human Resources 23(3): 279-95.
- Oaxaca, R. 1973. Male–female wage differentials in urban labor markets. International Economic Review 14: 693-709.
- Otieno, D.J., Omiti, J., Nyanamba, T., McCullough, E., 2009. Application of Chow test to improve analysis of farmer participation in markets in Kenya. Paper presented. 27th Conference of the International Association of Agricultural Economists (IAAE), 16-22 August, Beijing, China.
- Park, T., Lohr, L., 2010. A Oaxaca-Blinder decomposition for count data models. Applied Economics. 17(5): 451-455.
- Qaim, M., Zilberman, D., 2003. Yield effects of genetically modified crops in developing countries. Science 299(5608): 900-902.
- Quinn, G.P., Keough, M.J., 2007. Experimental design and data analysis for biologists. Cambridge University Press, Cambridge, UK. ISBN 0521811287.
- Raney, T., 2006. Economic impact of transgenic crops in developing countries. Current Opinion in Biotechnology 17: 1-5.
- Rice, M.E., 2003. Transgenic rootworm corn: Assessing potential agronomic, economic, and environmental benefits Plant Health Progress.
- Smale, M., Zambrano, P., Gruère, G., Falck-Zepeda, J., Matuschke, I., Horna, D., Nagarajan, L.,
 Yerramareddy, I., Jones, H., 2009. Measuring the Economic Impacts of Transgenic Crops in
 Developing Agriculture during the First Decade. Approaches, Findings, and Future Directions.
 Food Policy Review 10. Washington DC: IFPRI.
- Stanger, T.F., Lauer, J.G., 2006. Optimum plant population of *Bt* and non-*Bt* corn in Wisconsin. Agronomic Journal. 98: 914-921.
- Stone, G.D., 2011. Field versus Farm in Warrangal: *Bt* Cotton, Higher Yields, and Larger Questions. World Development 39(3): 387-398.
- Tárrega, A., Bayarri, S., Carbonell, I., Izquierdo, L., 2010. Blinder–Oaxaca decomposition applied to sensory and preference data. Short Communication. Food Quality and Preference 21: 662-665.
- Thirtle, C., Beyers, L., Ismael, Y., Piesse, J., 2003. Can GM-technologies help the poor? The impact of *Bt* cotton in Makhathini Flats, KwaZulu-Natal. World Development 31(4): 717-732.

- Wilson, T.A., Rice, M.E., Tollefons, J.J., Pilcher, C.D., 2005. Transgenic corn for control of the European corn borer and corn rootworms: a survey of Midwestern farmer's practices and perceptions. Journal of Economic Entomology 98: 237-247.
- Wolf, D., Vögeli, G.A., 2009. Economic value of *Bt* maize is relative. Agrarforschung 16(1): 4-9.
- Wossink, A., Denaux, Z., 2006. Environmental and cost efficiency of pesticide use in transgenic and conventional cotton production. Agricultural Systems 90(1-3): 312-328.
- Wu, Y., Escalante, C.L., Gunter, L.F., Epperson, J., 2012. A decomposition approach to analysing racial and gender biases in farm service agency's lending decisions. Applied Economics 44(22): 2841-2850.
- Yorobe, J.M. Jr., Quicoy, C.B., 2006. Economic impact of *Bt* corn in the Philippines. The Philippine Agricultural Scientist 89(3): 258-267.