



Universiteit  
Leiden  
The Netherlands

## Genetic patterns of Black-tailed Godwit populations and their implications for conservation

Trimbos, K.B.

### Citation

Trimbos, K. B. (2013, November 5). *Genetic patterns of Black-tailed Godwit populations and their implications for conservation*. Retrieved from <https://hdl.handle.net/1887/22110>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/22110>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden




The handle <http://hdl.handle.net/1887/22110> holds various files of this Leiden University dissertation

**Author:** Trimbos, Krijn

**Title:** Genetic patterns of Black-tailed Godwit populations and their implications for conservation

**Issue Date:** 2013-11-05



## **Curriculum Vitae**



Krijn Baptist Trimbos was born in Leiden on the 6<sup>th</sup> of October in the year 1979. After successfully finishing his VWO at the Rijnlands Lyceum in Oegstgeest he started studying biology at Leiden University in 1999. He finished his Propadeutics in 2000 after which he started a specialization track in medical biology and after that in environmental biology. During this time he conducted on three internships, two were done at the Leiden University Medical Centre (LUMC) and one was conducted at the Institute of Environmental Sciences Leiden (CML). His internships involved studying acquired systemic B cell memory in response to Human Papilloma Virus infection (2004 with Prof van den Burg), acquired systemic T cell memory in the Balinese population in response to Human Papilloma Virus infection (2005 with Prof. Lex Peters as a supervisor) and differences in breeding success of the Skylark at conventional and organic arable farms (2006 with Steven Kragten as a supervisor). His biology doctorate was finalized in 2007 after a literature study on the complexity of the human genome.

In 2007 he was research assistant for a short while at the Institute of Environmental Sciences Leiden (CML) focusing mainly on the value offield margins for Agricultural biodiversity in Zeeland, with special attention for the presence of butterflies and birds. At the end of 2007 Krijn started his PhD on genetic patterns of Black-tailed Godwit populations and their implications for conservation at the Institute of Environmental Sciences Leiden (CML) resulting in this thesis. During his PhD he supervised two students during their internships, gave several scientific lectures, assisted in a number of scientific courses and attended several symposia. Prof. de Snoo and dr. Musters were his supervisors during this time. Since 2011 he works at Landschap Noord Holland as a project manager meadow bird protection. The work there involves the coordination, training and education of volunteer meadow bird protectors and writing and executing research projects involving meadow bird conservation.







## **Publications**





Trimbos JB, Mouw R, Ranke G, **Trimbos KB** and Zwinderman K (2002). *The Donati stitch revisited: favorable cosmetic results in randomized clinical trial*. Journal of Surgical Research, 107, 131-134.

Kragten S, **Trimbos KB** and de Snoo GR (2008). *Breeding Skylarks (Alauda arvensis) on organic and conventional arable farms in the Netherlands*. Agriculture, ecosystems and environment, 126, 163-167.

Verkuil YI, **Trimbos KB**, Haddrath O, Baker OJ and Piersma, T (2009). *Characterization of polymorphic microsatellite DNA markers in the black-tailed godwit (Limosa limosa: Aves)*. Molecular Ecology Resources, 9(5), 1415-1418.

**Trimbos KB**, Broekman J, Kentie R, Musters CJM and de Snoo G (2009). *Using eggshell membranes as a DNA source for population genetic research*. Journal of Ornithology, 150(4), 915-920.

**Trimbos KB**, Musters CJM, Verkuil Y, Kentie R, Piersma T and de Snoo GR (2011) *No evident spatial genetic structuring in the rapidly declining Dutch Black-tailed Godwit Limosa l. Limosa population*. Conservation Genetics, 12, 629-636.

Van der Velde M, Galama Y, Hooijmeijer J, **Trimbos K** and Piersma T (2011). *Black-tailed Godwit Limosa limosa forensics: who laid a strangely shaped depredated egg?* Limosa, 84, 155-161.

**Trimbos KB** and Musters CJM (2012). *Genetic differentiation between Dutch Black-tailed Godwit breeding sites re-examined using D statistics*. Wader study group bulletin, 119, 59-60.

**Trimbos KB**, Kentie R, van der Velde M, Hooijmeijer JCEW, Poley C, Musters CJM, de Snoo GR and Piersma T (2013). *Intronic variation on the CHD1-Z gene in Black-tailed Godwits Limosa limosa: correlations with fitness components revisited*. Ibis, 155, 508-517.

Kentie R, Hooijmeijer JCEW, **Trimbos KB**, Groen NM and Piersma T (2013). *Intensified Agricultural use of grasslands reduces growth and survival of precocial shorebird chicks*. Journal of Applied Ecology, 50, 243-251.

**Trimbos KB**, Doorenweerd C, Kraaijeveld K, Musters CJM, Groen NM, de Knijff P, Piersma T and de Snoo GR. *Patterns in nuclear and mitochondrial DNA reveal historical and recent isolation in the Black-tailed Godwit (Limosa limosa)*. Submitted PLOS One.



