



Universiteit
Leiden
The Netherlands

Genetic factors in human reproduction a trade off between procreation and longevity

Dunné, F.M. van

Citation

Dunné, F. M. van. (2006, October 18). *Genetic factors in human reproduction a trade off between procreation and longevity*. Retrieved from <https://hdl.handle.net/1887/8781>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/8781>

Note: To cite this publication please use the final published version (if applicable).

Factor V Leiden Mutation in Relation to Fecundity and Miscarriage in Women with Venous Thrombosis

F.M. van Dunné, C.J.M. Doggen, M. Heemskerk, F.R. Rosendaal,
F.M. Helmerhorst

ABSTRACT

BACKGROUND: Factor V Leiden mutation (Arg506Gln) increases the likelihood of venous thrombosis. Factor V Leiden mutation may also have a positive effect through facilitation of embryo implantation. This may manifest itself as a reduced time to pregnancy (increased fecundity) and fewer miscarriages in the first trimester.

METHODS: From March 1999 onwards, consecutive patients with a first venous thrombosis (VT) were recruited. The first 115 female VT patients with factor V Leiden and 230 age-matched female VT patients without factor V Leiden were included. All patients, unaware of their genotype, received a structured questionnaire.

RESULTS: 297 (86%) Women returned the questionnaire, 220 had been pregnant at least once. Time to first pregnancy was unaffected by carrier status: 58% factor V Leiden carriers reported a pregnancy within 3 months compared to 54% non-carriers. The miscarriage proportion was 14%, similar in both groups. First trimester miscarriage was less frequent among carriers (46%) than among non-carriers (95%) (RR 0.5, 95% CI: 0.3–0.9).

CONCLUSIONS: Factor V Leiden mutation may support embryo implantation, as factor V Leiden carriers had fewer miscarriages in the first trimester with a similar overall miscarriage rate. Miscarriage of embryos with poor viability may be postponed until the second trimester in factor V Leiden carriers. Fecundity was not influenced by factor V Leiden status.

INTRODUCTION

Factor V Leiden mutation (Arg506Gln) is present in 4 to 10% of people of Caucasian origin (Bertina *et al.*, 1994; Rees, 1996). The factor V Leiden mutation induces a hypercoagulable state which increases the risk of venous thrombosis seven-fold among heterozygous carriers and about eighty-fold among homozygous carriers compared to non-carriers (Rosendaal *et al.*, 1995). It has been suggested that factor V Leiden mutation may be associated with negative outcomes of reproduction such as (recurrent) abortion, pre-eclampsia, prematurity and small-for-gestational-age neonates. However, the available data are conflicting (Pauer *et al.*, 2003; Hundsdoerfer *et al.*, 2003; Morrison *et al.*, 2002; Rai *et al.*, 2001). Because of the high prevalence of this mutation in certain populations positive effects associated with factor V Leiden have been postulated, possibly through human reproduction. Women who carry the factor V Leiden mutation lose less blood in menstruation, have higher haemoglobin levels, and possibly a lower incidence of life threatening post-partum haemorrhage which could be an evolutionary advantage (Lindqvist *et al.*, 2001). Furthermore, a higher than expected prevalence of factor V Leiden mutation carriers was found in healthy pregnant women (9.2%) (De Groot *et al.*, 1999) compared to general population figures (3%) (Rosendaal *et al.*, 1995). A similar finding was reported in a recurrent miscarriage study where the prevalence of factor V Leiden was higher in women without a history of recurrent miscarriages (14%) compared to those with recurrent miscarriages (1.7%) (Dilley *et al.*, 2002). Facilitation of embryo implantation was suggested as a possible positive pathway for factor V Leiden (Majerus, 1994). In agreement with this hypothesis an improved implantation rate in factor V Leiden carriers compared to non-carriers was reported in intra-cytoplasmic sperm injection (ICSI) pregnancies. If either the mother or the fetus carried the factor V Leiden mutation the proportion of live births was 90% (9/10) for the first embryo transfer compared to 49% (45/92) in factor V Leiden negative pairs (Göpel *et al.*, 2001). A reduced time to pregnancy (increased fecundity) in spontaneous pregnancies of factor V Leiden carriers would be an indication of a protective effect of factor V Leiden.

To study the effect of factor V Leiden mutation on embryo implantation and human reproduction, we investigated the association of factor V Leiden mutation on fecundity and miscarriages and the trimester in which the miscarriages occurred in 297 women. To be able

to construct a large cohort of women with factor V Leiden, we used information from a large study on venous thrombosis. We also investigated the effect of factor V Leiden carriership on term birth rate and birth weight.

MATERIALS AND METHODS

The women described in this study were enrolled in the Multiple Environmental and Genetic Assessment of risk factors for venous thrombosis study (MEGA study). The primary aim of the MEGA study is to assess interaction between environmental and genetic risk factors for venous thrombosis. From March 1999 onwards, all consecutive patients who suffered a first deep-vein thrombosis or pulmonary embolism between the age of 18 and 70 were recruited from six anticoagulation clinics in The Netherlands. The anticoagulation clinics monitor the anticoagulant therapy of all patients in a well-defined geographical area, which allowed us to identify consecutive and unselected patients with venous thrombosis. All participants filled in a questionnaire on risk factors for venous thrombosis. A blood sample was drawn three months after discontinuation of anticoagulation. Subjects who were unable or declined to give a blood sample, were given the opportunity to give DNA by use of a buccal swab. All participants completed an informed consent form. The Leiden University Medical Center medical ethical committee approved of the study.

For the present analysis, the first 115 female thrombosis patients identified with factor V Leiden mutation were matched on age to 230 female thrombosis patients (controls). Besides the age matching and absence of factor V Leiden, controls were randomly selected from the study participants. All 345 patients, who were unaware of their genotype, were asked to fill in an additional structured questionnaire concerning their reproductive history. Questions consisted of age at first pregnancy attempt, the period of unprotected intercourse until the desired pregnancy occurred, number of pregnancies, and the duration of each pregnancy. When there had been no pregnancies, we enquired whether this was despite efforts to become pregnant (infertility). If there was no response to the initial questionnaire, a written reminder accompanied with an identical questionnaire was sent after three weeks. To increase response,

patients were contacted by telephone if they did not return the questionnaire after five additional weeks.

Time to pregnancy was defined as self-reported time between unprotected intercourse and the occurrence of pregnancy. Miscarriage ratio was calculated as the total number of miscarriages per number of pregnancies. First trimester miscarriage was defined as embryonic or fetal loss before the completion of 12 weeks gestation, second trimester miscarriage was defined as fetal loss from 13 to 24 weeks gestation, stillbirth was defined as a loss after 24 weeks gestation. A number of factors known to influence miscarriage or fecundity were included in the questionnaire, such as age at first pregnancy attempt, age at first birth, level of education (primary school, secondary school, college or university), smoking habits, and alcohol use. The body-mass index (BMI) was calculated (weight/height^2) from the weight (in kg) and height (in m) obtained at the time of thrombosis.

DNA was isolated from whole blood or buccal swabs. For the latter, three large cotton swabs in a total of 6 ml SDS-proteinase K solution (100 mM NaCl, 10 mM EDTA, 10 mM Tris-HCl, pH = 8.0, 0.5% SDS, 0.1 mg/ml proteinase K) were obtained from each patient. Upon arrival, the proteinase K concentration was raised to 0.2 mg/ml and the sample was incubated for 2 hrs at 65°C. Subsequently, the suspension was recovered by centrifugation. Potassium acetate was added to a final concentration of 1.6M. After 15 min incubation on ice, proteins were removed using chloroform/ isoamylalcohol (24:1) treatment. The DNA in the water-phase was subsequently ethanol precipitated. After centrifugation, the pellet was resuspended in 200 µl 10 mM Tris-HCl, 10 mM EDTA pH=8.0 and frozen at -20°C until further analysis. Assessment of the factor V Leiden mutation in DNA retrieved from the buccal swabs was performed identically to the method for DNA from whole blood, and determined by polymerase chain reaction (PCR) and *Mnl I* restriction digestion as described elsewhere (Bertina *et al.*, 1994).

Data are presented as simple counts and percentages. Relative risks were computed as the ratio of these counts, and 95% confidence intervals were based on a binomial distribution.

RESULTS

Two hundred and ninety-seven (86%) women returned a completed questionnaire, 89 % of the factor V Leiden carriers and 84% of the non-carriers. Of the 48 non-responders, 4 women were deceased, 6 were lost to follow-up and for 21 women, the reason for not returning the questionnaire was unknown. Seventeen questionnaires were sent back blank, for reasons of lack of motivation (4 women), 3 women were too ill, for 2 women it brought back too many painful memories (both non-carriers), and in 8 cases no reasons were given.

Table I. Patient characteristics and reproductive outcome of the 220 women who had been pregnant, according to factor V Leiden carrier status

	Factor V Leiden carrier ^a n = 80	Factor V Leiden non carrier n = 140
Age at thrombosis, years	45.1 (20-68)	44.4(20-70)
Age at questionnaire, years	47.6 (23-71)	46.8 (23-72)
Menstrual cycle, regular (%)	66 (84)	118 (84)
Education, college/ university (%)	32 (45)	52 (43)
Smoked, ever (%)	50 (63)	77 (55)
BMI, kg/m ²	27 (18-58)	27 (17-57)
Age at 1 st pregnancy attempt, years	24.1 (15-36)	24.1 (15-38)
Age at birth first child, years	25.2 (16-36)	25.3 (15-39)
Number of pregnancies	2.2	2.7
Number of liveborn children	1.7	1.9
Total number of miscarriages (%)	25 (14)	50 (14)
Stillbirths (%)	2 (1)	9 (2)
Ectopic pregnancies (%)	0 (-)	4 (1)
Planned Abortion (%)	10 (6)	17 (4)
Other, (twins eg.) (%)	4 (2)	9 (2)

Values are mean (range) or *n* (%). ^a One homozygous factor V Leiden mutation carrier.

Of the 102 factor V Leiden carriers who returned the completed questionnaire, 80 (78%) had been pregnant at least once compared to 140 (72%) of the 195 non-carriers (RR 1.1, 95% CI 0.95-1.2). Patient characteristics of these 220 women are listed in Table I. The reasons for not having had a pregnancy were similar; infertility was reported by 5% (1/22) of factor V Leiden carriers and 11% (6/55) of non-carriers (RR 0.4, 95% CI 0.1-3.3).




Figure 1. Fecundity (time to pregnancy) for all pregnancies in 220 venous thrombosis patients according to factor V Leiden mutation (179 pregnancies in factor V Leiden carriers and 365 pregnancies in non-carriers, number of pregnancies stated in the bars).

Time to first pregnancy was similar for factor V Leiden carriers and non-carriers. In both groups over 90% of those who had been pregnant could recall the time to their first pregnancy. Forty-two (58%) factor V Leiden carriers achieved their first pregnancy within 3 months compared to 70 (54%) of the non-carriers (RR 1.1, 95% CI 0.8-1.4). Nine (13%) factor V Leiden carriers reported a time to first pregnancy of more than 12 months, which had occurred in 21 (16%) of the non-carriers (RR 0.8, 95% CI 0.4-1.6). When all consecutive pregnancies per woman were combined, similar results were found (Figure 1).

The miscarriage proportion for all pregnancies was similar in both groups, 14% (25/179) for factor V Leiden carriers and 14% (50/365) for non-carriers (RR 1.0, 95% CI 0.6-1.6). Considering only the first pregnancy, 16% (13/80) of the factor V Leiden carriers had experienced a miscarriage compared to 15% (21/140) of the non-carriers (RR 1.1, 95% CI 0.6-2.0). However, the trimester in which the miscarriages occurred was different according to factor V Leiden status; factor V Leiden carriers who experienced a miscarriage during their first pregnancy had fewer miscarriages in the first trimester compared to non-carriers, respectively 46% and 95% (RR 0.5, 95% CI: 0.3-0.9). Subsequently, factor V Leiden carriers had more miscarriages in the second trimester (RR 10.8, 95% CI 1.5-77.7) (Table IIA). When all consecutive pregnancies per woman were combined the difference persisted (Table IIB). Stillbirth (fetal loss in the third trimester) was rare, and was equal in both groups. For the 80 first pregnancies of factor V Leiden carriers no stillbirths occurred compared to four (3%) stillbirths in 140 first pregnancies in non-carriers. For all pregnancies per woman combined, two (1%) stillbirths occurred in 179 pregnancies in factor V Leiden carriers compared to nine (2%) in 365 pregnancies in non-carriers (RR 0.4, 95% CI 0.1-2.1).

Table II. Miscarriages, per trimester, out of 220 women who had been pregnant, according to factor V Leiden (FVL) mutation

	First trimester (≤12 weeks)	Second trimester (13-24 weeks)	Total
First pregnancy only (80 pregnancies in FVL carriers and 140 in non-carriers)			
FVL+ (%)	6 (46)	7 (54)	13
FVL- (%)	19 (95)	1 (5)	20 ^a
Relative risk, (95% CI)	0.5 (0.3-0.9)	10.8 (1.5-77.7)	
All pregnancies combined (179 in FVL carriers and 365 in non-carriers)			
FVL+ (%)	16 (64)	9 (36)	25
FVL- (%)	43 (90)	5 (10)	48 ^b
Relative risk (95% CI)	0.7 (0.5-1.0)	3.5 (1.3-9.2)	

Values are *n* (%). ^a Of one non-carrier, the trimester in which the miscarriage took place was unknown. ^b Of two non-carriers, the trimester in which the miscarriage took place was unknown. CI = confidence interval.

The proportion of live births was similar in both groups; 74% (59/80) among factor V Leiden carriers compared to 76% (106/140) among non-carriers (RR 1.0, 95% CI 0.8-1.1). Term birth (37-42 weeks gestation) in first pregnancies was comparable, 64% (38/59) and 69% (73/106), respectively (RR 0.9, 95% CI 0.7-1.2). Mean birth weight for the children born at term was similar for factor V Leiden carriers (3644 gram, range 2500-4000) and non-carriers (3481gram, range 1200-3975).

DISCUSSION

In this study of 297 women with venous thrombosis, we found no association between factor V Leiden mutation and fecundity or the frequency of miscarriage. However, when miscarriages had occurred, they took place less often in the first trimester in factor V Leiden mutation carriers than in non-carriers.

Miscarriages occurred as often in factor V Leiden carriers (14%) and non-carriers (14%), in percentages that are similar to general population figures (10 to 15%) (Zinaman *et al.*, 1996). Published data on factor V Leiden in relation to miscarriages are conflicting. Recent meta-analyses have shown a relation between factor V Leiden and recurrent fetal loss, occurring both in early and in late in gestation (respectively OR 2.01, 95% CI 1.13-3.58 and OR 7.83, 2.83-21.67) (Rey *et al.*, 2003; Kovalevsky *et al.*, 2004). For non-recurrent early loss (< 19 weeks gestation) no clear association was found with factor V Leiden (OR 1.40, 0.66-2.97); and for non-recurrent isolated second/third trimester loss (stillbirth > 19 weeks) a positive association was found (OR 3.26, 1.82-5.83) (Rey *et al.*, 2003; Dudding *et al.*, 2004).

The focus in literature has mainly been on factor V Leiden in relation to second and third trimester loss, as thrombosis of the placental vessels is assumed to be an important factor in fetal loss. In our study factor V Leiden carriers experienced more fetal loss in the second trimester, as would be expected by the placental vessel thrombosis theory. A clear increase in third trimester loss would be expected too, with an even higher rate of fetal loss in third than in second trimester for factor V Leiden carriers. However, third trimester loss (stillbirth) was equally distributed over carriers and non-carriers, in a comparable small rate, which contradicts this theory. Moreover, in the present study, mean birth weight was not influenced

by factor V Leiden status, which is in line with earlier published data (Lindqvist *et al.*, 1999). If we consider hypercoagulability to lead to obstruction of placental vessels as a major pathological factor among pregnant factor V Leiden carriers, one would not expect an effect on first trimester miscarriages, since the placental circulation in the first trimester has not yet been fully established (Hustin and Schaaps, 1987; Burton *et al.*, 1999). Therefore, impaired placental perfusion might not be critical for the embryonic development and implantation during (very) early gestation. We can not differentiate between the types of early pregnancy loss in this study (eg presence or absence of fetal heart prior to the loss), as all the data were provided to us by the patients through interview. Furthermore, most would have not had an early first trimester scan, as they are not routinely done in The Netherlands.

Our results indicate a clear reduction in first trimester loss in factor V Leiden carriers, which is subsequently compensated by an increased loss in the second trimester. Similar findings were reported in a recent study where a decreased risk of recurrent pregnancy losses at less than 10 weeks gestation was found in factor V Leiden carriers (OR 0.23, 95% CI 0.07-0.77) with a subsequent increase in losses after 10 weeks (Roqué *et al.*, 2004). This suggests a protective effect on the embryo during the first trimester in factor V Leiden carriers, including less viable embryos that will eventually abort in the second trimester. This may explain the similar overall frequency of miscarriages in factor V Leiden carriers and non-carriers. A lower frequency of miscarriages in the first trimester may thus reflect a successful implantation. Approximately 50-70% of all miscarriages are attributable to detectable chromosomal abnormalities, furthermore 15-20% is thought to be due to morphological defect(s) in the embryo. A recent study confirmed this with a transcervical embryoscopy at the time of the curette and cytogenetic analysis of the products of conception (Philipp *et al.*, 2003). We do not have any cytogenetic information in our study, as karyotyping of the products of conception is not a routine consideration in The Netherlands. The possible protective effect of factor V Leiden early in pregnancy will require further study with among other things, cytogenetic testing of the miscarriage products and possibly transcervical embryoscopy prior to evacuation.

Our study did not show a clearly increased fecundity in factor V Leiden carriers. Many factors affect fecundity, including physiological, behavioural and environmental factors. Known

factors, such as age at pregnancy attempt, regularity of the menstrual cycle, smoking habits and educational level were similar in factor V Leiden carriers and non-carriers. As fecundity is a manifestation of both conception and implantation it is important to further distinguish these. For conception, factors such as sperm quality, coital frequency and timing are of great importance. However, this study was not designed to examine these factors and it remains unclear whether factor V Leiden has any effect on these factors. Aspects influencing implantation have not been fully elucidated yet, however an improved implantation could be due to an increase in the hypercoagulable state, related to the factor V Leiden mutation. This was suggested by a study that omitted conception by reporting only on ICSI pregnancies, reflecting implantation success (Göpel *et al.*, 2001). A higher incidence of implantation success was found if either the mother or the fetus was a factor V Leiden carrier. It is possible that the beneficial effect of factor V Leiden on implantation alone has a less clear effect on fecundity compared to various factors concerning conception and therefore, in our study, fails to show an overall difference in fecundity.

We investigated the reproductive histories of women who had suffered venous thrombosis. The choice of this design was opportunistic, for it offered the opportunity to study a large cohort of factor V Leiden carriers. We have considered whether this choice, rather than the ideal study of factor V Leiden carriers without thrombosis, could have distorted our results. Firstly, the period about which questions were asked preceded the thrombotic event, in most cases by many years. Hence, the thrombotic event itself did not influence our results. As the patients developed thrombosis, they will have more risk factors for thrombosis than other women. It is known that this is not only true for women without factor V Leiden but also for women with factor V Leiden (the majority of people with factor V Leiden never develop thrombosis, and there must be causes why some do). For this reason, we chose thrombosis cases without factor V Leiden as controls rather than healthy women without factor V Leiden, and therefore differences between the group can be attributed to factor V Leiden. One could argue that the groups differed more: women with thrombosis without factor V Leiden probably had more additional risk factors than those with factor V Leiden, for instance another, possibly still unknown, gene defect. This could, if that other prothrombotic factor also affected implantation and fetal loss, explain the absence of a difference between the groups in the frequency of miscarriage. However, we did find a difference. As the

reproductive data in the present study were collected by interview, recall bias is possible, but seems unlikely. Validation studies of fecundity and miscarriages have shown a good match between long-term recall of personal reproductive history through interview and medical data (Joffe *et al.*, 1993). Moreover, the women were unaware of their factor V Leiden status at the time of the questionnaire.

In conclusion, factor V Leiden mutation may support embryo implantation, as factor V Leiden carriers reported significantly fewer miscarriages in the first trimester. This was not reflected in an increased fecundity. The overall miscarriage proportion was not influenced by factor V Leiden status. These results suggest that factor V Leiden offers a protective effect on early pregnancy and that the miscarriage of embryos with poor viability in factor V Leiden carriers is postponed until the second trimester.

Acknowledgments

The authors wish to thank the directors of the Anticoagulation Clinics of Amersfoort (M.H.H. Kramer), Amsterdam (M. Remkes), Leiden (F.J.M. van der Meer), Rotterdam (A.A.H. Kasbergen), The Hague (E. van Meegen) and Utrecht (J. de Vries-Goldschmeding) who made the recruitment of patients possible. The interviewers S. van der Leden, M. Roosen and L. Willems of Brilman also performed the blooddraws. I. de Jonge, R. Roelofsen, M. Streevelaar, L. Timmers and J.J. Ververs are thanked for their secretarial, administrative and data-management support. The fellows J.W. Blom, A. van Hylckama Vlieg and L.W. Tick took part in the data collection. R. van Eck, J. van der Meijden, P.J. Noordijk and Th. Visser performed the laboratory measurements. H.L. Vos supervised the technical aspects of DNA analysis. We express our gratitude to all individuals who participated in the MEGA study. This research was supported by the Netherlands Heart Foundation (NHS 98.113) and the Dutch Cancer Foundation (RUL 99/1992).

REFERENCES:

- Bertina R.M., Koeleman B.P., Koster T., Rosendaal F., Dirven R., de Ronde H., van der Velden P., Reitsma P. (1994) Mutation in blood coagulation factor V associated with resistance to activated protein C. *Nature*, 369, 64-67.
- Burton G.J., Jauniaux E., Watson A.L. (1999). Maternal arterial connections to the placental intervillous space during the first trimester of human pregnancy: the Boyd collection revisited. *Am. J. Obstet. Gynecol.*, 181(3): 718-724.
- De Groot C.J., Bloemenkamp K.W., Duvekot E.J., Helmerhorst F.M., Bertina R.M., Van Der Meer F., De Ronde H., Oei S.G., Kanhai H.H., Rosendaal F.R. (1999) Preeclampsia and genetic risk factors for thrombosis: a case-control study. *Am.J.Obstet.Gynecol.*, 181, 975-980
- Dilley A., Benito C., Hooper W.C., Austin H., Miller C., El-Jamil M., Cottrell S., Benson J., Evatt B.L., Patterson-Bamett A., *et al.* (2002) Mutations in the factor V, prothrombin and MTHFR genes are not risk factors for recurrent fetal loss. *J. Mat. Fet. Neonat. Med.*, 11: 176-182
- Dudding T.E., Attia J. (2004) The association between adverse pregnancy outcomes and maternal factor V Leiden genotype: a meta-analysis. *Thromb. Haemost.*, 91(4): 700-11.
- Göpel W., Ludwig M., Junge A.K., Kohlmann T., Diedrich K., Moller J. (2001) Selection pressure for the factor-V-Leiden mutation and embryo implantation. *Lancet*, 358, 1238-1239
- Hundsdoerfer P., Vetter B., Stover B., Bassir C., Scholz T., Grimmer I., Monch E., Ziemer S., Rossi R., Kulozik A.E. (2003) Homozygous and double heterozygous Factor V Leiden and Factor II G20210A genotypes predispose infants to thromboembolism but are not associated with an increase of foetal loss. *Thromb. Haemost.*, 90(4):628-635
- Hustin J., Schaaps J.P. (1987) Echographic [corrected] and anatomic studies of the maternotrophoblastic border during the first trimester of pregnancy. *Am. J. Obstet. Gynecol.*, 157(1):162-168.
- Joffe M., Villard L., Li Z., Plowman R., Vessey M. (1993) Long-term recall of time-to-pregnancy. *Fertil. Steril.*, 60:99-104.
- Kovalevsky G., Gracia C.R., Berlin J.A., Sammel M.D., Barnhart K.T. (2004) Evaluation of the association between hereditary thrombophilias and recurrent pregnancy loss: a meta-analysis. *Arch. Intern. Med.*, 164(5):558-63
- Lindqvist P.G., Zöller B., Dählback B. (2001) Improved Hemoglobin status and reduced menstrual blood loss among female carriers of factor V Leiden – an evolutionary advantage? *Thromb. Haemost.*, 86:1122-1123
- Lindqvist P.G., Svensson P.J., Marsaal K., Grennert L., Luterkort M., Dählback B. (1999) Activated protein C resistance (FV:Q506) and pregnancy. *Thromb. Haemost.*, 81(4):532-537.
- Majerus P.W. (1994) Human genetics. Bad blood by mutation. *Nature*, 369, 14-15.
- Morrison E.R., Miedzybrodzka Z.H., Campbell D.M., Haites N.E., Wilson B.J., Watson M.S., Greaves M., Vickers M.A. (2002) Prothrombotic genotypes are not associated with pre-eclampsia and gestational hypertension: results from a large population-based study and systematic review. *Thromb. Haemost.*, 87: 779-785
- Pauer H.U., Voigt-Tschirschwitz T., Hinney B., Burfeind P., Wolf C., Emons G., Neesen J. (2003) Analyses of three common thrombophilic gene mutations in German women with recurrent abortions. *Acta Obst. Gynecol. Scand.*, 82: 942-947.

- Philipp T., Philipp K., Reiner A., Beer F., Kalousek D.K. (2003) Embryoscopic and cytogenetic analysis of 233 missed abortions: factors involved in the pathogenesis of developmental defects of early failed pregnancies. *Hum. Reprod.*, 18(8):1724-32.
- Rai R., Shlebak A., Cohen H., Backos M., Holmes Z., Marriott K., Regan L. (2001) Factor V Leiden and acquired activated protein C resistance among 1000 women with recurrent miscarriage. *Hum.Reprod.*, 16, 961-965.
- Rees D.C. (1996) The population genetics of factor V Leiden (Arg506Gln). *Br.J.Haematol.*, 95, 579-586.
- Rey E., Kahn S.R., David M., Shrier I. (2003) Thrombophilic disorders and fetal loss: a meta-analysis. *Lancet*, 361(9361): 901-8.
- Roqué H., Paidas M.J., Funai E.F., Kuczynski E., Lockwood C.J. (2004) Maternal thrombophilias are not associated with early pregnancy loss. *Thromb. Haemost.*;91(2):290-5
- Rosendaal F.R., Koster T., Vandenbroucke J.P., Reitsma P.H. (1995) High risk of thrombosis in patients homozygous for factor V Leiden (activated protein C resistance). *Blood*, 85; 6: 1504-1508
- Zinaman M.J., Clegg E.D., Brown C.C., O'Connor J., Selevan S.G. (1996) Estimates of human fertility and pregnancy loss. *Fertil.Steril.*, 65, 503-509.