



**Universiteit  
Leiden**  
The Netherlands

## **Matrix metalloproteinases in gastric inflammation and cancer : clinical relevance and prognostic impact**

Kubben, F.J.G.M.

### **Citation**

Kubben, F. J. G. M. (2007, September 27). *Matrix metalloproteinases in gastric inflammation and cancer : clinical relevance and prognostic impact*. Retrieved from <https://hdl.handle.net/1887/12356>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12356>

**Note:** To cite this publication please use the final published version (if applicable).

## CHAPTER 6

# Clinical evidence for a protective role of lipocalin-2 against MMP-9 autodegradation and the impact for gastric cancer

F.J.G.M. Kubben, C.F.M. Sier, L.J.A.C. Hawinkels, H. Tschesche<sup>1</sup>, W. van Duijn, K. Zuidwijk, J.J. van der Reijden, R. Hanemaaijer<sup>2</sup>, G. Griffioen, C.B.H.W. Lamers, H. W. Verspaget

Department of Gastroenterology and Hepatology, Leiden University Medical Centre, Leiden, The Netherlands; <sup>1</sup>Department of Biochemistry, University Bielefeld, Bielefeld, Germany; <sup>2</sup>TNO Quality of Life, Biomedical Research, Leiden, The Netherlands

*European Journal of Cancer 2007, in press*

## Abstract

Recently, complexes of matrix metalloproteinase MMP-9 with lipocalin-2 (neutrophil gelatinase-associated lipocalin) were found in the urine obtained from breast cancer patients, while these were completely absent in that obtained from healthy controls. *In vitro* data suggested a possible role for lipocalin-2 in the protection of MMP-9 against autolysis.

To establish this effect *in vivo*, we determined the presence of MMP-9, lipocalin-2 and their complex in tumour tissue from 81 gastric cancer patients. The effect of the presence of the individual parameters, the complexes, and the inhibitors TIMP-1 and TIMP-2 on MMP-9 activity was evaluated with a bioactivity assay. Immunohistochemical (double) staining identified epithelial cells as the most likely cellular source. Finally, evaluation of all these parameters with clinicopathological scores revealed that tumour MMP-9/lipocalin-2 complexes were significantly related with the classifications of Laurén and WHO, and highly associated with worse survival in Cox's univariate (HR 2.087,  $P=0.006$ ) and multivariate analysis (HR 2.095,  $P=0.025$ ).

## Introduction

Lipocalin-2 (also known as neutrophil gelatinase-associated lipocalin) is a member of the highly heterogeneous family of lipocalins, sharing a common tertiary structure [1, 2]. Lipocalin-2 has initially been discovered in specific granules of human neutrophils [3] and was later shown to be expressed also by certain epithelial cells, in particular during inflammatory or cancerous circumstances [4-10]. There is little information about the physiological functions of lipocalins, but lipocalin-2 has been associated with cellular iron uptake, antibacterial activity, and epithelial cell differentiation [2, 9].

Enhanced tissue, blood and urine levels of matrix metalloproteinase-9 (MMP-9) have been associated with the malignancy of various tumour types [11-14]. Using quantitative zymography and immunoassays we have previously shown that MMP-9 as well as MMP-2 are enhanced in gastric cancer tissue and that high levels are associated with worse survival of the patients [15, 16]. Next to MMP-9 and MMP-2, the zymograms revealed extra bands, particularly between 125-135 kDa. These bands have been described before in the urine obtained from cancer patients, and are most likely complexes of MMP-9 with lipocalin [17, 18]. *In vitro* experiments suggested a role for lipocalin-2 in the protection of MMP-9 against autolysis [17].

To investigate the suggested relevance of MMP-9/lipocalin-2 complexes *in vivo*, we determined the levels of MMP-9, lipocalin-2 and their complex in tissue homogenates from 81 gastric carcinomas in comparison with adjacent normal mucosa from the same patients. We used immunohistochemical staining of paraffin-embedded tissue sections to establish the cellular origin of MMP-9 and lipocalin-2. To confirm the histological findings, the levels of MMP-9, lipocalin-2 and the MMP-9/lipocalin-2 complexes in the homogenates were compared with markers for neutrophils, a known source of MMP-9 and lipocalin-2. The effect of complex formation between MMP-9 and lipocalin-2 on the MMP-9 activity state was evaluated using a specific MMP-9 bioactivity assay. Finally, the possible clinical consequence of the presence of MMP-9/lipocalin-2 complexes in gastric tumours was evaluated by examining for correlations with established clinicopathological parameters of the carcinoma patients, including univariate and multivariate Cox proportional hazard survival analyses.

## Materials & methods

### *Patients and study design*

Fresh tissue specimens from 81 patients (21 females and 60 males, mean age 65.9 years, range 35.1-91.3) who underwent resection for primary gastric adenocarcinoma between 1984 and 1996 at the department of Oncologic Surgery, Leiden University Medical Centre were collected prospectively. Samples from the mid-central non-necrotic part of the carcinoma and from normal mucosa, taken approximately 10 cm from the tumour, were snap-frozen and stored at  $-70^{\circ}\text{C}$  until extraction. All carcinomas were classified according to the TNM classification (UICC 1992), and localization as well as diameter of the tumour was registered. Microscopical histological parameters, including differentiation-grade, WHO-, Borrmann-, and Laurén-classification, as well as the presence of intestinal metaplasia in the normal gastric mucosa, were revised by a gastroenterologist and a pathologist. All patients entered the study at operation date, and the patient's time experience ended in the event of death or, when still alive, at the common closing date. The minimal follow-up was 33 months with a decreasing overall survival according to TNM stage, i.e. from TNM I (52.2%, n=23), to TNM II (26.9%, n=26), to TNM III (28%, n=25), and to TNM IV (0%, n=7). The study was performed according to the instructions and guidelines of the LUMC medical ethics committee.

### *Tissue preparation and protein concentration*

Homogenisation of tissue specimens and determination of protein concentrations were performed as described previously [15].

### *MMP-9/lipocalin-2 complex zymography*

Quantitative gelatin zymography for MMP-9/lipocalin-2 complexes was performed as described before [15], using an Ultrosan XL Laser Densitometer (LKB) for quantification. The MMP-9/lipocalin-2 complex levels in tissue homogenates were expressed in arbitrary units (AU) per mg protein.

### *ELISAs for MMP-9, lipocalin-2, MMP-9/lipocalin-2-complexes, MMP-8 and TIMPs*

Total antigen levels of MMP-9, lipocalin-2, and MMP-8 were determined using previously described ELISAs [19-22]. The concentrations of MMP-9/lipocalin-2 complexes, TIMP-1 and TIMP-2 were measured using commercial ELISAs according to the manufacturer instructions (R&D Systems Europe, Abingdon, UK). The MMP-9/lipocalin-2 ELISA immobilizes complexes via anti-MMP-9 antibodies followed by detection using anti-lipocalin-2 antibodies and does not detect MMP-9 or lipocalin-2 in their free forms.

### *MMP-9 activity assay*

The bioactivity assay (BIAs) for MMP-9 was done as described previously [14, 19, 22]. This assay detects active MMP-9 and total MMP-9 levels in parallel in 96-wells plates coated with MMP-9 specific antibodies and using modified MMP-sensitive pro-urokinase as substrate. The fraction of the latent MMP-9 proform is calculated by subtraction of active from total MMP-9.

### *Myeloperoxidase (MPO) activity assay*

MPO activity was measured as described previously [23]. In short, tissue homogenates were incubated with 0.5% hexadecyl-trimethylammonium bromide in 50 mM potassium phosphate buffer (pH 5.5), plus 0.026% *ortho*-dianisidine dihydrochloride substrate and 0.018% H<sub>2</sub>O<sub>2</sub>. The reaction kinetics were followed for 30 min at 450 nm in 96-well plates. The specificity of the reaction was checked with sodium azide (0.1 mM). All samples were analyzed in duplicate and standardized using a homogenate of pooled human neutrophils, and MPO activity was expressed in arbitrary units.


### *Immunohistochemistry and immunofluorescence double staining*

Paraffin sections (5 µm) from the same tumours as used for the homogenates were deparaffinized and stained for the localisation of MMP-9 and lipocalin-2. Antigen retrieval was performed through boiling in a 0.01 M citrate solution (pH 6.0) for 12 minutes in a microwave oven. After being rinsed in PBS and incubated with 10 % of normal goat serum (Dako) for 30 minutes, the sections were incubated with the primary antibody polyclonal rabbit anti-lipocalin-2 (1:100, from Drs H. Tschesche and O. Hiller) or polyclonal rabbit anti-MMP-9 (1:400, TNO, Leiden, The Netherlands) overnight at 4 degrees. After washing, the sections were incubated with biotinylated goat anti-rabbit 1:400 (Dako) for 30 minutes, followed by washing and incubation with Streptavidin/ABCcomplex/HRP (DakoCytomation) for 30 minutes. The brown colour was developed by 0.004 % H<sub>2</sub>O<sub>2</sub> (Merck) and 0.05 % diaminobenzidine tetrahydrochloride (Sigma) in 0.01 M Tris-HCl pH 6.0 for 10 minutes. The slides were counterstained with Mayer's haematoxylin (Merck). For specific cell recognition, i.e. epithelial cells, (myo)fibroblasts, neutrophils and endothelial cells, sequential tissue sections were stained with mouse anti-pan-cytokeratin (1:1000, clone C11, Santa Cruz biotechnologies, Santa Cruz, USA), mouse anti-vimentin (1:400, clone V9 Santa Cruz), mouse anti-smooth muscle actin (1:1000, clone ASM-1, Progen Heidelberg, Germany), rabbit anti-myeloperoxidase (1:1000, Dako) and mouse anti-CD31 (1:400, clone JC70A, Dako) followed by appropriate second antibodies and staining procedures. Immunofluorescence double staining was performed as described before [24]. In short, sections were incubated for 1 hr with rabbit polyclonal anti-lipocalin-2 and mouse monoclonal anti-MMP-9 (clone GE-213, 1:400, NeoMarkers, Fremont, CA) antibodies, appropriately

diluted in PBS with 1% BSA, washed, and incubated with respectively Alexa Fluor 488- and 546-conjugated anti-rabbit and anti-mouse antibodies (Molecular Probes, Leiden, The Netherlands) diluted in PBS-BSA. After incubation and washing, the sections were mounted in Mowiol. A Zeiss LSM 510 confocal microscope equipped with argon and He/Ne lasers and a 20x objective were used to obtain the images.


### Statistical analysis

Differences between normal and tumour values for all parameters were calculated using the Wilcoxon signed ranks test and visualized by Box-Whisker graphs using lower and upper margins of 5%. Correlations between parameters were determined according to Spearman's Rho test. For the survival analyses the clinicopathological parameters were dichotomized as described previously [15], unless indicated. Cut off values for MMPs and related factors were optimised. Survival analyses were performed with the Cox proportional hazards model using the SPSS Windows Release 12.0.1. Statistical Package (2004, SPSS Inc., Chicago, Illinois, USA). Multivariate survival analyses were performed using the Cox proportional hazards method by separately adding



**Figure 1.** Zymogram and immunoblot showing MMP-9, MMP-9/lipocalin-2 complexes and lipocalin-2 in a representative gastric cancer tissue homogenate (lane 2-4). MMP-9 activity is located in the zymograms (lane 2) between 70-92 kDa, representing active MMP-9 and pro-MMP-9, and at 135 kDa corresponding with MMP-9/lipocalin-2 complex standard (lane 1). The immunoblots show corresponding complex bands for MMP-9 (lane 3) and lipocalin-2 (lane 4) with extra bands at approximately 25 and 50 kDa representing respectively the monomer and homodimer forms of lipocalin-2. Lane 1 contains 20  $\mu$ l standard from the MMP-9/lipocalin-2 ELISA ( $\approx$  0.8 ng).

the significant MMP variables to the dichotomized clinicopathological parameters. Survival curves were constructed using the method of Kaplan and Meier including the Log-rank test. Differences were considered significant when  $P \leq 0.05$ .



**Figure 2.** Levels of a) MMP-9/lipocalin-2 complex in AU/mg protein, b) MMP-9/lipocalin-2 complex in ng/mg protein, c) MMP-9 in ng/mg protein, and d) lipocalin-2 in ng/mg protein in carcinoma tissue and adjacent normal mucosa from 81 gastric cancer patients.  $n=81$  unless indicated.



## Results

### *Quantification of MMP-9/lipocalin-2 complexes in gastric cancer tissue homogenates*

The presence of MMP-9/lipocalin-2 complexes in tissue homogenates from gastric cancer patients was determined using zymography and ELISA. Figure 1 shows a typical gastric cancer homogenate with in the zymogram abundant MMP-9 mediated lysis and a smaller band at molecular weight 135 kDa, corresponding with standard MMP-9/lipocalin-2 complex. The nature of this band was further verified using immunoblots for respectively MMP-9 and lipocalin-2 under normal (Figure 1) and reduced conditions (not shown). The amount of the MMP-9/lipocalin-2 complexes was quantified from the zymograms, using laser densitometry (Figure 2a). MMP-9/lipocalin-2 complexes were significantly enhanced in cancer tissue compared with control mucosa ( $27.3 \pm 2.0$  versus  $14.5 \pm 1.4$  AU/mg protein,  $P < 0.001$ ,  $n = 81$ ). The data from this semi-quantitative assay were compared with the results obtained with a commercial ELISA (Figure 2b). The correlation between both assays was highly significant ( $\rho = 0.488$ ,  $P < 0.0001$ ,  $n = 75$ , i.e. 5 normal mucosa and 70 carcinoma homogenates).

### *Levels of MMP-9 and lipocalin-2 in gastric cancer tissue homogenates*

The tissue levels of MMP-9 and lipocalin-2 are shown in figure 2c and d. The gastric carcinomas contained significant higher concentrations of MMP-9 ( $P < 0.001$ ) and


**Table 1 - Correlation coefficients ( $\rho$  plus  $P$ -values) for MMP-9, lipocalin-2 and MMP-9/lipocalin-2 complexes in relation to myeloperoxidase (MPO), MMP-8 and TIMP-1 in 162 gastric cancer tissue homogenates (81 normal/81 cancer).**

	MMP-9	Lipoc-2	MMP-9/Lipocalin-2 complex	MMP-9 active	MMP-9 latent
MMP-9 ng/mg protein		0.438 (0.000)	0.641 (0.000)	0.240 (0.003)	0.817 (0.000)
Lipocalin-2 ng/mg protein			0.273 (0.001)	-0.121 ns	0.443 (0.000)
MMP-9/Lipoc-2 AU/mg protein				0.166 (0.038)	0.586 (0.000)
MMP-9 active U/mg protein					0.263 (0.001)
MPO AU/mg protein	0.486 (0.000)	0.280 (0.000)	0.332 (0.000)	0.073 (ns)	0.462 (0.000)
MMP-8 ng/mg protein	0.810 (0.000)	0.482 (0.000)	0.578 (0.000)	0.128 ns	0.734 (0.000)
TIMP-1 ng/mg protein	0.358 (0.000)	0.363 (0.000)	0.315 (0.000)	-0.097 (ns)	0.240 (0.004)

lipocalin-2 ( $P=0.002$ ) than adjacent normal tissues. In general, lipocalin-2 was more abundantly present than MMP-9, in specific cases even more than 100 times higher.

*Correlation between MMP-9 and MMP-9/lipocalin-2 with MMP-9 activity state*

The correlation of MMP-9, lipocalin-2, and MMP-9/lipocalin-2-complex with MMP-9 activity in tissue homogenates of gastric cancer patients is shown in Table 1. Active MMP-9 levels correlated significantly with the total antigen level of MMP-9, but more interestingly also with the MMP-9/lipocalin-2 concentration ( $P=0.038$ ), suggesting a protective role for lipocalin-2-complex formation in MMP-9 (auto)activation. The tis-



**Figure 3.** Typical immunohistochemical staining of a human gastric intestinal type carcinoma for: a) MMP-9 (200x) and b) lipocalin-2 (200x). Black, red, green and yellow arrows indicate, respectively, epithelial cells, neutrophil-like cells, (myo)fibroblast like cells and endothelial cells. Protein levels in corresponding homogenate for MMP-9, lipocalin-2 and complex are respectively 29 ng/mg, 4928 ng/mg and 17 AU/mg protein. c) Immunofluorescence double staining (400x) for MMP-9 (red) and Lipocalin-2 (green). Yellow colour suggests complex formation.

sue concentration of TIMP-1, the most relevant tissue inhibitor of MMP-9, was equally correlated with the levels of MMP-9 and lipocalin-2, but not with MMP-9 activity.

#### *Immunohistochemical staining for MMP-9 and lipocalin-2*


To establish the cellular source of the MMP-9/lipocalin-2 complexes, sequential paraffin sections adjacent to the tissue used for homogenates were stained for MMP-9 and lipocalin-2. Normal mucosa showed barely any staining for MMP-9 nor lipocalin-2 (not shown). In carcinoma tissues staining for MMP-9 was found in neutrophils and a substantial part of the epithelial cells, occasionally in endothelial cells, and incidentally in muscle cells, macrophages, and fibroblasts (Figure 3a). In neutrophils and epithelial cells lipocalin-2 was similarly distributed compared with MMP-9, but lipocalin-2 was additionally present in tumour epithelial subgroups which lacked MMP-9 staining (Figure 3b). Endothelial cells and fibroblasts showed little or no staining for lipocalin-2. Immunofluorescence double staining confirmed that particular epithelial cells stained for lipocalin-2 but not for MMP-9 (Figure 3c red versus green). Furthermore this staining revealed that only a fraction of MMP-9 and lipocalin-2 was actually in close proximity (Figure 3c, yellow versus green). Yellow staining was found in particular at the periphery of cells, suggesting that the majority of both proteins is uncomplexed and presumably still compartmentalized within the cells, as suggested by zymographic analysis.

#### *Correlations between MMP-9, lipocalin-2, MMP-9/lipocalin-2, MMP-8 and MPO*

To confirm the similarities and the apparent difference between MMP-9 and lipocalin-2 in cellular origin, as found by immunohistochemistry, the concentrations of MMP-9, lipocalin-2 and MMP-9/lipocalin-2-complex in the tissue homogenates were evaluated for correlations with the levels of MPO and MMP-8 (Table 1). MPO, a commonly used cell marker for neutrophils, correlated strongly with MMP-8, a collagenase abundantly present in neutrophils ( $0.445, P < 0.0005$ ) as well as with MMP-9, but the correlation with lipocalin-2 was considerably less, suggesting a possible other source of lipocalin-2 than neutrophils only.

#### *Relation between MMP-9/lipocalin-2 complexes and clinicopathological parameters*

The MMP-9/lipocalin-2 levels were significantly enhanced in differentiated tumours according to the WHO classification ( $30.9 \pm 2.5$  vs.  $19.6 \pm 2.8$  AU/mg protein,  $P \leq 0.006$ ) and in tumours of the intestinal type ( $30.5 \pm 2.6$  vs.  $21.9 \pm 2.7$  AU/mg protein,  $P \leq 0.04$ ). MMP-9/lipocalin-2 levels showed a trend to increase with higher TNM stages. Dichotomization of the patients, based on low (AU < 36) or high (AU > 36) MMP-9/lipocalin-2 complex values in their tumour, showed a significant correlation with overall survival



**Figure 4.** Kaplan-Meier survival curve for a cohort of gastric cancer patients subdivided by low ( $\leq 36$  AU/mg protein) or high ( $> 36$  AU/mg protein) levels of MMP-9/lipocalin-2 complex in their tumour tissue homogenate.

**Table 2 - Univariate and multivariate Cox proportional hazard overall survival analyses for low or high levels of MMP-9/lipocalin-2 in tissue homogenates of gastric cancer versus different clinicopathological parameters.**

		Univariate				Multivariate		
		n	HR	CI 95%	P	HR	CI 95%	P
Gender	F/M	21/60	1.247	0.730-2.131	NS	1.622	0.900-2.923	NS
Age	<median>	40/41	1.323	0.815-2.149	NS	1.504	0.860-2.629	NS
TNM	1	23/81	1	-	-	1	-	-
	2	26/81	1.984	1.033-3.813	0.040	2.133	1.009-4.639	0.047
	3	25/81	1.586	0.804-3.130	NS	1.623	0.737-3.704	NS
	4	7/81	3.248	1.261-8.366	0.015	6.027	1.876-20.46	0.003
Laurén	dif/mx vs inte	30/50	1.103	0.671-1.816	NS	1.125	0.402-3.137	NS
WHO	diff vs undiff	54/26	0.881	0.525-1.480	NS	0.874	0.289-2.609	NS
Borrmann	fung. vs infiltr.	55/24	1.025	0.591-1.778	NS	0.846	0.457-1.567	NS
Localization	cardia vs rest	36/45	0.603	0.368-0.989	0.045	0.419	0.223-0.764	0.005
Diameter	$\leq 5$ vs $> 5$ cm	47/34	1.062	0.652-1.729	NS	0.695	0.403-1.195	NS
Eosinophils	few vs many	56/24	1.220	0.725-2.053	NS	1.846	1.023-3.544	0.042
Intest. metaplasia	not vs present	39/42	0.551	0.334-0.909	0.020	0.651	0.365-1.151	NS
MMP-9 antigen	<median>	40/40	1.143	0.701-1.863	NS	1.336	0.756-2.363	NS
Lipocalin-2	<median>	40/39	1.029	0.632-1.674	NS	0.772	0.422-1.413	NS
MMP-9/lipocalin-2	$\leq 36$ vs $> 36$ AU	58/23	2.087	1.229-3.544	0.006	2.095	1.099-4.031	0.025

NS: non significant

(Log Rank 8.04,  $P < 0.005$ ,  $n = 81$ ), as shown in figure 4. Analysis of the MMP-9/lipocalin-2 complex ELISA data showed a similar trend but did not reach statistical significance (Log Rank, 3.04,  $P = 0.0815$ ,  $n = 70$ ).

### *Survival analyses*

The relation of MMP-9/lipocalin-2 complexes with survival was further characterized with Cox's uni- and multivariate analyses against the clinicopathological parameters (Table 2). The level of MMP-9/lipocalin-2 was significantly associated with worse survival and kept its significance in multivariate analyses, indicating its value as an independent prognostic factor.

## **Discussion**

High levels of lipocalin-2 have been reported in various types of cancer [6-10]. Our study shows that lipocalin-2 levels are indeed significantly enhanced in gastric carcinomas compared to adjacent control tissue. Moreover and more interestingly, our data show that the complexes of lipocalin-2 with MMP-9 are also significantly enhanced in human gastric tumours.

*In vitro* experiments showed that lipocalin-2 is able to induce the expression of E-cadherin, to promote the formation of polarized epithelia, and to diminish the invasiveness and metastasis of Ras-transformed cells [25], suggesting a protective role against cancer. Other studies reported a positive correlation between lipocalin-2 expression levels and the growth rate of lipocalin-2 transfected MCF-7 human breast carcinoma cells, which were subcutaneously implanted in immuno-deficient mice [18]. Immunohistochemical analyses of these xenografted tumours showed that the over-expression of lipocalin-2 was accompanied by enhanced levels of MMP-9, suggesting the formation of complexes between MMP-9 and lipocalin-2. The formation of MMP-9/lipocalin-2 complexes has previously been shown to protect MMP-9 from auto-degradation *in vitro* [17, 18]. MMP-9/lipocalin-2 complex formation could result in increased extracellular, tumour-associated MMP-9, and hence in enhanced tumour growth as recently suggested by Fernández et al. [18]. We found that in gastric cancer tissue lipocalin-2 levels are in general 30 times higher than corresponding MMP-9 levels, presumably leading to MMP-9/lipocalin-2 complex formation of a substantial part of the MMP-9 fraction after it has been released from the cells. These complexes were significantly correlated with the active, as well as the latent fraction of MMP-9. Therefore, our data support the hypothesis that enhanced production of lipocalin-2 in cancerous tissue stimulates the formation of a complex with MMP-9, playing a role in the maintenance of an extracellular pool of a latent form of this powerful proteinase,

by prevention from auto-degradation. This latent pool of secreted, lipocalin-2-bound MMP-9 has previously been shown to be important for the spatial control of VEGF release from the ECM and hence for enhanced angiogenesis [26]. Our study does not provide information about the presence and/or role of MMP-9/lipocalin-2/TIMP-1 complexes. These ternary complexes have previously been isolated from phorbol myristate acetate stimulated neutrophils and showed low gelatinase activity, as expected [27]. In our study, total TIMP-1 levels correlated significantly with all the forms of MMP-9, except for the active form of MMP-9, suggesting that other factors are involved in regulating the activity of MMP-9, besides the ratio between MMP-9 and TIMP-1. TIMP-2 levels were weakly inversely correlated with MMP-9 antigen levels, suggesting little or no mutual interaction (data not shown).

The quantitative determination of MMP-9 and lipocalin-2 in tissue homogenates, as performed in this study, has several advantages compared to semi-quantitative immunohistological detection methods but obviously does not provide information about the localization of the proteins. Our immunohistochemical data revealed that lipocalin-2 as well as MMP-9 in gastric cancers are mainly present in neutrophils and epithelial cells, but that epithelial expression of MMP-9 is depending on the individual cancer and on the location within the tumour. MMP-9 was furthermore found in (myo) fibroblast-like cells and endothelial cells. These data are in accordance with what has been found previously in colonic cancer [6, 28]. Our fluorescent double-staining data suggest that, although MMP-9 and lipocalin-2 seem present in close proximity especially within the cells, overlap of green and red colours, presumably representing extra-cellular complex formation, is limited and mainly restricted to peri-cellular areas. Whether the enhancement of MMP-9/lipocalin-2 complexes in gastric cancer compared with adjacent normal mucosa was caused by the influx of neutrophils or alternatively by upregulated expression in malignant epithelial cells, could not be established in this study. The finding that high numbers of intra-tumoural neutrophils are associated with better survival of patients with gastric cancer [29], would suggest the latter.

From this study, the clinical relevance of MMP-9/lipocalin-2 complex formation appears most obvious from the correlation with overall survival of the patients. Enhanced levels of these complexes were highly prognostic for worse survival, whereas the levels of single MMP-9 and lipocalin-2 were not. The finding that MMP-9/lipocalin-2 levels are increased in gastric cancer tissue and that enhancement might be associated with clinical outcome of the patients is supported by a recent study reporting that similar complexes were present in approximately 90% of the urines obtained from breast cancer patients, but not in those from healthy controls [18]. The prognostic value of MMP-9/lipocalin-2 complexes is in accordance with the presumed role of lipocalin-2 in the protection of secreted MMP-9 against auto-degradation, which contributes to

an enhanced pool of potentially active MMP-9, a proteolytic enzyme associated with angiogenesis and tumour growth. High total MMP-9 levels were not associated with survival in the present study. This is not in agreement with what we have published previously [15], but those earlier data were based on a smaller group of patients and on detection of MMP-9 activity instead of total antigen level. The different outcome between both studies indicates the delicacy of the use of proteinase levels as prognostic indicators, as discussed before [16, 30]. Apparently not just the enhanced presence, but more the (potential) activation state of the proteinase, i.e. the result of, respectively, production, release, activation, and the inactivation by inhibitors, seems to be crucial, similar to what has been described for other enzymes playing a role in gastric cancer like urokinase and MMP-2 [16, 31]. Additionally, our data indicate that prevention of auto-degradation of MMP-9 by lipocalin-2 might play an important role too.

In conclusion, we have shown for the first time that complexes between MMP-9 and lipocalin-2 are present in enhanced levels in gastric cancer tissue and that high levels are associated with worse survival of the patients. The potential clinical value of our findings should be confirmed in larger groups of cancer patients. Recently the enzymatic activity of MMP-9/lipocalin-2 complex has indeed been found to correlate significantly with the depth of tumour invasion in esophageal squamous cell carcinomas [32].

## **Acknowledgements**

We are grateful to Dr. Oliver Hiller (Department of Biochemistry, Bielefeld University, Germany), Dr. Arko Gorter, Enno Dreef and Frans Prins (Department of Pathology, Leiden University Medical Centre, The Netherlands) for their helpful assistance with immunohistochemistry.

## **Conflict of interest**

None declared

## References

1. Flower DR. The lipocalin protein family: a role in cell regulation. *FEBS Lett* 1994; 354: 7-11
2. Xu S, Venge P. Lipocalins as biochemical markers of disease. *Biochim Biophys Acta* 2000; 1482: 298-307
3. Triebel S, Bläser J, Reinke H, Tschesche H. A 25 kDa alpha 2-microglobulin-related protein is a component of the 125 kDa form of human gelatinase. *FEBS Lett* 1992; 314: 386-388
4. Cowland JB, Sørensen OE, Sehested M, Borregaard N. Neutrophil gelatinase-associated lipocalin is up-regulated in human epithelial cells by IL-1 beta, but not by TNF-alpha. *J Immunol* 2003; 171: 6630-6639
5. Bartsch S, Tschesche H. Cloning and expression of human neutrophil lipocalin cDNA derived from bone marrow and ovarian cancer cells. *FEBS Lett* 1995; 357: 255-259
6. Nielsen BS, Borregaard N, Bundgaard JR, Timshel S, Sehested M, Kjeldsen L. Induction of NGAL synthesis in epithelial cells of human colorectal neoplasia and inflammatory bowel diseases. *Gut* 1996; 38: 414-420
7. Furutani M, Arii S, Mizumoto M, Kato M, Imamura M. Identification of a neutrophil gelatinase-associated lipocalin mRNA in human pancreatic cancers using a modified signal sequence trap method. *Cancer Lett* 1998; 122: 209-214
8. Friedl A, Stoesz SP, Buckley P, Gould MN. Neutrophil gelatinase-associated lipocalin in normal and neoplastic human tissues. Cell type-specific pattern of expression. *Histochem J* 1999; 31: 433-441
9. Mallbris L, O'Brien KP, Hulthén A, Sandstedt B, Cowland JB, Borregaard N, Ståhle-Bäckdahl M. Neutrophil gelatinase-associated lipocalin is a marker for dysregulated keratinocyte differentiation in human skin. *Exp Dermatol* 2002; 11: 584-591
10. Stoesz SP, Friedl A, Haag JD, Lindstrom MJ, Clark GM, Gould MN. Heterogeneous expression of the lipocalin NGAL in primary breast cancers. *Int J Cancer* 1998; 79: 565-572
11. Margulies IM, Höyhty M, Evans C, Stracke ML, Liotta LA, Stetler-Stevenson WG. Urinary type IV collagenase: elevated levels are associated with bladder transitional cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 1992; 1: 467-474
12. Roeb E, Dietrich CG, Winograd R, Arndt M, Breuer B, Fass J, Schumpelick V, Matern S. Activity and cellular origin of gelatinases in patients with colon and rectal carcinoma differential activity of matrix metalloproteinase-9. *Cancer* 2001; 92: 2680-2691
13. Kuyvenhoven JP, Van Hoek B, Blom E, Van Duijn W, Hanemaaijer R, Verheijen JH, Lamers CB, Verspaget HW. Assessment of the clinical significance of serum matrix metalloproteinases MMP-2 and MMP-9 in patients with various chronic liver diseases and hepatocellular carcinoma. *Thromb Haemost* 2003; 89: 718-725
14. Sier CF, Casetta G, Verheijen JH, Tizzani A, Agape V, Kos J, Blasi F, Hanemaaijer R. Enhanced urinary gelatinase activities (matrix metalloproteinases 2 and 9) are associated with early-stage bladder carcinoma: a comparison with clinically used tumor markers. *Clin Cancer Res* 2000; 6: 2333-2340
15. Sier CF, Kubben FJ, Ganesh S, Heerding MM, Griffioen G, Hanemaaijer R, van Krieken JH, Lamers CB, Verspaget HW. Tissue levels of matrix metalloproteinases MMP-2 and MMP-9 are related to the overall survival of patients with gastric carcinoma. *Br J Cancer* 1996; 74: 413-417



16. Kubben FJ, Sier CF, Van Duijn W, Griffioen G, Hanemaaijer R, van de Velde CJ, van Krieken JH, Lamers CB, Verspaget HW. Matrix metalloproteinase-2 is a consistent prognostic factor in gastric cancer. *Br J Cancer* 2006; 94: 1035-1040
17. Yan L, Borregaard N, Kjeldsen L, Moses MA. The high molecular weight urinary matrix metalloproteinase (MMP) activity is a complex of gelatinase B/MMP-9 and neutrophil gelatinase-associated lipocalin (NGAL). Modulation of MMP-9 activity by NGAL. *J Biol Chem* 2001; 276: 37258-37265
18. Fernández CA, Yan L, Louis G, Yang J, Kutok JL, Moses MA. The matrix metalloproteinase-9/neutrophil gelatinase-associated lipocalin complex plays a role in breast tumor growth and is present in the urine of breast cancer patients. *Clin Cancer Res* 2005; 11: 5390-5395
19. Hanemaaijer R, Visser H, Konttinen YT, Koolwijk P, Verheijen JH. A novel and simple immunocapture assay for determination of gelatinase- B (MMP-9) activities in biological fluids: saliva from patients with Sjogren's syndrome contain increased latent and active gelatinase-B levels. *Matrix Biol* 1998; 17: 657-665
20. Bläser J, Triebel S, Tschesche H. A sandwich enzyme immunoassay for the determination of neutrophil lipocalin in body fluids. *Clin Chim Acta* 1995; 235: 137-145
21. Bergmann U, Michaelis J, Oberhoff R, Knauper V, Beckmann R, Tschesche H. Enzyme linked immunosorbent assays (ELISA) for the quantitative determination of human leukocyte collagenase and gelatinase. *J Clin Chem Clin Biochem* 1989; 27: 351-359
22. Kubben FJ, Sier CF, Van Duijn W, Griffioen G, Hanemaaijer R, van de Velde CJ, van Krieken JH, Lamers CB, Verspaget HW. Matrix metalloproteinase-2 (MMP-2) is a consistent prognostic factor in gastric cancer. *Br J Cancer* 2006; 94: 1035-1040
23. Kruidenier L, Kuiper I, Van Duijn W, Mieremet-Ooms MA, Van Hogezaand RA, Lamers CB, Verspaget HW. Imbalanced secondary mucosal antioxidant response in inflammatory bowel disease. *J Pathol* 2003; 201: 17-27
24. Sier CF, Zuidwijk K, Zijlmans HJ, Hanemaaijer R, Mulder-Stapel AA, Prins FA, Dreef EJ, Kenter GG, Fleuren GJ, Gorter A. EMMPRIN-induced MMP-2 activation cascade in human cervical squamous cell carcinoma. *Int J Cancer* 2006; 118: 2991-2998
25. Hanai J, Mammoto T, Seth P, Mori K, Karumanchi SA, Barasch J, Sukhatme VP. Lipocalin 2 diminishes invasiveness and metastasis of Ras-transformed cells. *J Biol Chem* 2005; 280: 13641-13647
26. Mira E, Lacalle RA, Buesa JM, de Buitrago GG, Jimenez-Baranda S, Gomez-Mouton C, Martinez A, Manes S. Secreted MMP9 promotes angiogenesis more efficiently than constitutive active MMP9 bound to the tumor cell surface. *J Cell Sci* 2004; 117: 1847-1857
27. Kolkenbrock H, Hecker-Kia A, Orgel D, Kinawi A, Ulbrich N. Progelatinase B forms from human neutrophils. complex formation of monomer/lipocalin with TIMP-1. *Biol Chem* 1996; 377: 529-533
28. Nielsen BS, Timshel S, Kjeldsen L, Sehested M, Pyke C, Borregaard N, Danø K. 92 kDa type IV collagenase (MMP-9) is expressed in neutrophils and macrophages but not in malignant epithelial cells in human colon cancer. *Int J Cancer* 1996; 65: 57-62
29. Caruso RA, Bellocco R, Pagano M, Bertoli G, Rigoli L, Inferrera C. Prognostic value of intratumoral neutrophils in advanced gastric carcinoma in a high-risk area in northern Italy. *Mod Pathol* 2002; 15: 831-837
30. Duffy MJ. The role of proteolytic enzymes in cancer invasion and metastasis. *Clin Exp Metastasis* 1992; 10: 145-155

31. Sier CF, Verspaget HW, Griffioen G, Ganesh S, Vloedgraven HJ, Lamers CB. Plasminogen activators in normal tissue and carcinomas of the human oesophagus and stomach. *Gut* 1993; 34: 80-85
32. Zhang H, Xu L, Xiao D, Xie J, Zeng H, Wang Z, Zhang X, Niu Y, Shen Z, Shen J, Wu X, Li E. Up-regulation of Neutrophil Gelatinase-Associated Lipocalin in Esophageal Squamous Cell Carcinoma: Significantly Correlated with Cell Differentiation and Tumor Invasion. *J Clin Pathol* 2007; 60: 555-561