



Universiteit
Leiden
The Netherlands

Polyketide synthases in *Cannabis sativa* L

Flores-Sanchez, I.J.

Citation

Flores-Sanchez, I. J. (2008, October 29). *Polyketide synthases in Cannabis sativa L*. Retrieved from <https://hdl.handle.net/1887/13206>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13206>

Note: To cite this publication please use the final published version (if applicable).

References

- Abe I., Abe T., Wanibuchi K. and Noguchi H. (2006a) Enzymatic formation of quinolone alkaloids by a plant type III polyketide synthase. *Org Lett* 8: 6063–6065.
- Abe I., Oguro S., Utsumi Y., Sano Y. and Noguchi H. (2005b) Engineered biosynthesis of plant polyketides: Chain length control in an octaketide-producing plant type III polyketide synthase. *J Am Chem Soc* 127: 12709–12716.
- Abe I., Sano Y., Takahashi Y. and Noguchi H. (2003a) Site-directed mutagenesis of benzalacetone synthase: The role of Phe²¹⁵ in plant type III polyketide synthases. *J Biol Chem* 278: 25218–25226.
- Abe I., Takahashi Y. and Noguchi H. (2002) Enzymatic formation of an unnatural C₆–C₅ aromatic polyketide by plant type III polyketide synthases. *Org Lett* 4: 3623–3626.
- Abe I., Takahashi Y., Lou W. and Noguchi H. (2003b) Enzymatic formation of unnatural novel polyketides from alternate starter and nonphysiological extension substrate by chalcone synthase. *Org Lett* 5: 1277–1280.
- Abe I., Takahashi Y., Morita H. and Noguchi H. (2001) Benzalacetone synthase: A novel polyketide synthase that plays a crucial role in the biosynthesis of phenylbutanones in *Rheum palmatum*. *Eur J Biochem* 268: 3354–3359.
- Abe I., Utsumi Y., Oguro S. and Noguchi H. (2004a) The first plant type III polyketide synthase that catalyzes formation of aromatic heptaketide. *FEBS Lett* 562: 171–176.
- Abe I., Utsumi Y., Oguro S., Morita H., Sano Y. and Noguchi H. (2005a) A plant type III polyketide synthase that produces pentaketide chromone. *J Am Chem Soc* 127: 1362–1363.
- Abe I., Watanabe T. and Noguchi H. (2004b) Enzymatic formation of long-chain polyketide pyrones by plant type III polyketide synthases. *Phytochemistry* 65: 2447–2453.
- Abe I., Watanabe T. and Noguchi H. (2005c) Chalcone synthase superfamily of type III polyketide synthases from rhubarb (*Rheum palmatum*) *Proc Jpn Acad* 81(B): 434–440.
- Abe I., Watanabe T., Lou W. and Noguchi H. (2006b) Active site residues governing substrate selectivity and polyketide chain length in aloesone synthase. *FEBS J* 273: 208–218.
- Abe T., Morita H., Noma H., Kohno T., Noguchi H. and Abe I. (2007) Structure function analyses of benzalacetone synthase from *Rheum palmatum*. *Bioorg Med Chem Lett* 17: 3161–3166.

- Abe T., Noma H., Noguchi H. and Abe I. (2006c) Enzymatic formation of an unnatural methylated triketide by plant type III polyketide synthases. *Tetrahedron Lett* 47: 8727–8730.
- Achkar J., Xian M., Zhao H. and Frost J.W. (2005) Biosynthesis of phloroglucinol. *J Am Chem Soc* 127: 5332–5333.
- Adams M., Pacher T., Greger H. and Bauer R. (2005) Inhibition of leukotriene biosynthesis by stilbenoids from *Stemona* species. *J Nat Prod* 68: 83–85.
- Adawadkar P.D. and ElSohly M.A. (1981) Isolation, purification and antimicrobial activity of anacardic acids from *Ginkgo biloba* fruits. *Fitoterapia* 53: 129–135.
- Aida R., Kishimoto S., Tanaka Y. and Shibata M. (2000) Modification of flower color in torenia (*Torenia fournieri* Lind.) by genetic transformation. *Plant Sci* 153: 33–42.
- Akiyama T., Shibuya M., Liu H.M. and Ebizuka Y. (1999) *p*-Coumaroyltriacytic acid synthase, a new homologue of chalcone synthase, from *Hydrangea macrophylla* var. *thunbergii*. *Eur J Biochem* 263: 834–839.
- Alemanno L., Ramos T., Gargadenec A., Andary C. and Ferriere N. (2003) Localization and identification of phenolic compounds in *Theobroma cacao* L. somatic embryogenesis. *Ann Bot* 92: 613–623.
- Allwood E.G., Davies D.R., Gerrish C., Ellis B.E. and Bolwell G.P. (1999) Phosphorylation of phenylalanine ammonia-lyase: Evidence for a novel protein kinase and identification of the phosphorylated residue. *FEBS Lett* 457: 47–52.
- Ameri A. (1999) The effects of cannabinoids on the brain. *Prog Neurobiol* 58: 315–348.
- André C.L. and Vercruysse A. (1976) Histochemical study of the stalked glandular hairs of the female cannabis plants, using fast blue salt. *Planta Med* 29: 361–366.
- Aronne L.J. (2007) Rimonabant improves body weight and cardiometabolic risk factors in older adults. *J Am Coll Cardiol* 49–S1: 325A.
- Asahina Y. and Asano J. (1930) Über die constitution von hydrangenol und phyllodulcin (II.Mitteil.). *Chem Ber* 63: 429–437.
- Asakawa Y., Takikawa K., Toyota M. and Takemoto T. (1982) Novel bibenzyl derivatives and ent-cuparene-type sesquiterpenoids from *Radula* species. *Phytochemistry* 21: 2481–2490.

- Askari A., Worthen L.R. and Schimiza Y. (1972) Gaylussacin, a new stilbene derivative from species of *Gaylussacia*. *Lloydia* 35: 49–54.
- Atanassov I., Russinova E., Antonov L. and Atanassov A. (1998) Expression of an anther-specific chalcone synthase-like gene is correlated with uninucleate microspore development in *Nicotiana sylvestris*. *Plant Mol Biol* 38: 1169–1178.
- Austin M.B. and Noel J.P. (2003) The chalcone synthase superfamily of type III polyketide synthases. *Nat Prod Rep* 20: 79–110.
- Austin M.B., Bowman M.E., Ferrer J.L., Schröder J. and Noel J.P. (2004a) An aldol switch discovered in stilbene synthases mediates cyclization specificity of type III polyketide synthases. *Chem Biol* 11: 1179–1194.
- Austin M.B., Izumikawa M., Bowman M.E., Udway D.W., Ferrer J.L., Moore B.S. and Noel J.P. (2004b) Crystal structure of a bacterial type III polyketide synthase and enzymatic control of reactive polyketide intermediates. *J Biol Chem* 279: 45162–45174.
- Ayres D.C. and Loike J.D. (1990) Chemistry and Pharmacology of natural products. Lignans: Chemical, biological and clinical properties. Phillipson J.D., Ayres D.C. and Baxter H., eds. Cambridge University Press, UK.
- Back K., Jang S.M., Lee B.C., Schmidt A., Strack D. and Kim K.M. (2001) Cloning and characterization of a hydroxycinnamoyl-CoA:tyramine *N*-(hydroxycinnamoyl)transferase induced in response to UV-C and wounding from *Capsicum annuum*. *Plant Cell Physiol* 42: 475–481.
- Bahceevli A.K., Kurucu S., Kolak U., Topcu G., Adou E. and Kingston D.G.I. (2005) Alkaloids and aromatics of *Cyathobasis fruticulosa* (Bunge) Aellen. *J Nat Prod* 68: 956–958.
- Bangera M.G. and Thomashow L.S. (1999) Identification and characterization of a gene cluster for synthesis of the polyketide antibiotic 2,4-diacetylphloroglucinol from *Pseudomonas fluorescens* Q2-87. *J Bacteriol* 181: 3155–3163.
- Barrett M.L., Scutt A.M. and Evans F.J. (1986) Cannflavin A and B, prenylated flavones from *Cannabis sativa* L. *Experientia* 42: 452–453.
- Barron D. and Ibrahim R.K. (1996) Isoprenylated flavonoids—a survey. *Phytochemistry* 43: 921–982.
- Battelle B.A. and Hart M.K. (2002) Histamine metabolism in the visual system of the horseshoe crab *Limulus polyphemus*. *Comp Biochem Physiol A Mol Integr Physiol* 133: 135–142.

- Battelle B.A., Edwards S.C., Kass L., Maresch H.M., Pierce S.K. and Wishart A.C. (1988) Identification and function of octopamine and tyramine conjugates in the *Limulus* visual system. *J Neurochem* 51: 1240–1251.
- Batz O., Logemann E., Reinold S. and Hahlbrock K. (1998) Extensive reprogramming of primary and secondary metabolism by fungal elicitor or infection in parsley cells. *Biol Chem* 379: 1127–1135.
- Beckert C., Horn C., Schnitzler J.P., Lehning A., Heller W. and Veit M. (1997) Styrylpyrone biosynthesis in *Equisetum arvense*. *Phytochemistry* 44: 275–283.
- Beekwilder J., Wolswinkel R., Jonker H., Hall R., de Vos C.H.R. and Bovy A. (2006) Production of resveratrol in recombinant microorganism. *Appl Environ Microbiol* 72: 5670–5672.
- Beerhues L. (1996) Benzophenone synthase from cultured cells of *Centaurea erythraea*. *FEBS Lett* 383: 264–266.
- Bercht C.A.L., Lousberg R.J.J.C., Küppers F.J.E.M. and Salemink C.A. (1973) L-(+)-Isoleucine betaine in *Cannabis* seeds. *Phytochemistry* 12: 2457–2459.
- Bercht C.A.L., Samrah H.M., Lousberg R.J.J.C., Theuns H. and Salemink C.A. (1976) Isolation of vomifoliol and dihydrovomifoliol from *Cannabis*. *Phytochemistry* 15: 830–831.
- Bernards M.A. (2002) Demystifying suberin. *Can J Bot* 80: 227–240.
- Bienz S., Detterbeck R., Ensich C., Guggisberg A., Häusermann U., Meisterhans C., Wendt B., Werner C. and Hesse M. (2002) Putrescine, spermidine, spermine and related polyamine alkaloids. In: The alkaloids, chemistry and pharmacology. Vol. 58. Cordell G.A., ed. Academic Press, USA. 83–338.
- Binder M. and Popp A. (1980) Microbial transformation of cannabinoids, part 3: Major metabolites of (3R, 4R)- Δ^1 -Tetrahydrocannabinol. *Helv Chim Acta* 63: 2515–2518.
- Biondi S., Scaramagli S., Oksman-Caldentey K.M. and Poli F. (2002) Secondary metabolism in root and callus cultures of *Hyoscyamus muticus* L.: The relationship between morphological organization and response to methyl jasmonate. *Plant Sci* 163: 563–569.
- Blokhina O., Virolainen E. and Fagerstedt K.V. (2003) Antioxidants, oxidative damage and oxygen deprivation stress: A review. *Ann Bot* 91: 179–194.

- Bohlmann F. and Hoffmann E. (1979) Cannabigerol-ähnliche verbindungen aus *Helichrysum umbraculigerum*. *Phytochemistry* 18: 1371–1374.
- Borejsza-Wysocki W. and Hrazdina G. (1996) Aromatic polyketide synthases: Purification, characterization and antibody development to Benzalacetone synthase from raspberry fruits. *Plant Physiol* 110: 791–799.
- Bosabalidis A., Gabrieli C. and Niopas I. (1998) Flavone aglycones in glandular hairs of *Origanum x intercedens*. *Phytochemistry* 49: 1549–1553.
- Bourgaud F., Gravot A., Milesi S. and Gontier E. (2001) Production of plant secondary metabolites: A historical perspective. *Plant Sci* 161: 839–851.
- Bouvier F., Rahier A. and Camara B. (2005) Biogenesis, molecular regulation and function of plant isoprenoids. *Prog Lipid Res* 44: 357–429.
- Brady L.R. and Tyler V.E. (1958) Biosynthesis of hordenine in tissue homogenates of *Panicum miliaceum* L. *Plant Physiol* 33: 334–338.
- Braemer R. and Paris M. (1987) Biotransformation of cannabinoids by cell suspension culture of *Cannabis sativa* L. *Plant Cell Rep* 6: 150–152.
- Braemer R., Braut-Boucher F., Cosson L. and Paris M. (1985) Exemple de variabilite induite par biotransformation du cannabidiol par des cals et des suspensions cellulaires de *Cannabis sativa* L. *Bull Soc Bot Fr Actual Bot* 132: 148.
- Braemer R., Tsoutsias Y., Hurabielle M. and Paris M. (1986) Biotransformations of quercetin and apigenin by a cell suspension culture of *Cannabis sativa*. *Planta Med* 53: 225–226.
- Brand S., Holscher D., Schierhorn A., Svatos A., Schröder J. and Schneider B. (2006) A type III polyketide synthase from *Wachendorfia thyrsiflora* and its role in diarylheptanoid and phenylphenalenone biosynthesis. *Planta* 224: 413–428.
- Brenneisen R. and ElSohly M.A. (1988) Chromatographic and spectroscopic profiles of *Cannabis* of different origins: Part I. *J Forensic Sci* 33: 1385–1404.
- Bruneton J. (1999b) Lignans, neolignans and related compounds. *In*: Pharmacognosy, Phytochemistry, Medicinal plants. Second edition. Lavoisier Publishing Inc–Intercept Ltd, Paris. 279–293.
- Burstein S., Varanelli C. and Slade L.T. (1975) Prostaglandins and cannabis-III: Inhibition of biosynthesis by essential oil components of marihuana. *Biochem Pharmacol* 24: 1053–1054.

- Carchman R.A., Harris L.S. and Munson A.E. (1976) The inhibition of DNA synthesis by cannabinoids. *Cancer Res* 36: 95–100.
- Charles R., Garg S.N. and Kumar S. (1998) An orsellinic acid glucoside from *Syzygium aromaticum*. *Phytochemistry* 49: 1375–1376.
- Chen J.J., Huang S.Y., Duh C.Y., Chen I.S., Wang T.C. and Fang H.Y. (2006) A new cytotoxic amide from the stem wood of *Hibiscus tiliaceus*. *Planta Med* 72: 935–938.
- Cheng S.H., Sheen J., Gerrish C. and Bolwell P. (2001) Molecular identification of phenylalanine ammonia-lyase as a substrate of a specific constitutively active *Arabidopsis* CDPK expressed in maize protoplasts. *FEBS Lett* 503: 185–188.
- Chiron H., Drouet A., Lieutier F., Payer H.D., Ernst D. and Sanderman H.J. (2000) Gene induction of stilbene biosynthesis in Scot pine in response to ozone treatment, wounding and fungal infection. *Plant Physiol* 124: 865–872.
- Choi Y.H., Choi H.K., Hazekamp A., Bermejo P., Schilder Y., Erkelens C. and Verpoorte R. (2003) Quantitative analyses of bilobalide and ginkgolides from *Ginkgo biloba* leaves and ginkgo products using ^1H -NMR. *Chem Pharm Bull* 51: 158–161.
- Choi Y.H., Kim H.K., Hazekamp A., Erkelens C., Lefeber A.W.M. and Verpoorte R. (2004a) Metabolomic differentiation of *Cannabis sativa* cultivars using ^1H NMR spectroscopy and principal component analyses. *J Nat Prod* 67: 953–957.
- Choi Y.H., Kim H.K., Wilson E.G., Erkelens C., Trijzelaar B. and Verpoorte R. (2004b) Quantitative analyses of retinol and retinol palmitate in vitamin tablets using ^1H -nuclear magnetic resonance spectroscopy. *Anal Chim Acta* 512: 141–147.
- Christensen A.B., Gregersen P.L., Schröder J. and Collinge D.B. (1998) A chalcone synthase with an unusual substrate preference is expressed in barley leaves in response to UV light and pathogen attack. *Plant Mol Biol* 37: 849–857.
- Clark M.N. and Bohm B.A. (1979) Flavonoid variation in *Cannabis* L. *Bot J Linn Soc* 79: 249–257.
- Clarke R.C. (1981) Marijuana Botany: An advanced study, the propagation and breeding of distinctive cannabis. Ronin Publishing, Oakland, CA.
- Combet C., Jambon M., Deleage G. and Geourjon C. (2002) Geno3D: Automatic comparative molecular modeling of protein. *Bioinformatics* 18: 213–214.

- Contessotto M.G.G., Monteiro-Vitorello C.B., Mariani P.D.S.C. and Coutinho L.L. (2001) A new member of the chalcone synthase (CHS) family in sugarcane. *Genet Mol Biol* 24: 257–261.
- Courtney-Gutterson N., Napoli C., Lemieux C., Morgan A., Firoozabady E. and Robinson K.E.P. (1994) Modification of flower color in flower color in florist's chrysanthemum: Production of a white-flowering variety through molecular genetics. *Biotechnology* 12: 268–271.
- Crombie L. (1986) Natural products of Cannabis and Khat. *Pure Appl Chem* 58: 693–700.
- Crombie L. and Crombie W.M.L. (1982) Natural products of Thailand high Δ^1 -THC-strain *Cannabis*: The bibenzyl-spiran-dihydrophenanthrene group, relations with cannabinoids and canniflavones. *J Chem Soc Perkin Trans I* 1455–1466.
- Crombie L., Crombie W.M.L. and Firth D.F. (1988) Synthesis of bibenzyl cannabinoids, hybrids of two biogenetic series found in *Cannabis sativa*. *J Chem Soc Perkin Trans I* 1263–1270.
- Crombie L., Tuchinda P. and Powell M.J. (1982) Total synthesis of the spirans of *Cannabis*: Cannabispiradienone, cannabispirenone-A and -B, cannabispirone, α - and β -cannabispiranols and the dihydrophenanthrene cannithrene-1. *J Chem Soc Perkin Trans I* 1477–1484.
- Cutillo F., D'Abrosca B., DellaGreca M., Marino C.D., Golino A., Previtera L. and Zarrelli A. (2003) Cinnamic acid amides from *Chenopodium album*: Effects on seed germination and plant growth. *Phytochemistry* 64: 1381–1387.
- Dana C.D., Bevan D.R. and Winkel B.S.J. (2006) Molecular modeling of the effects of mutant alleles on chalcone synthase protein structure. *J Mol Model* 12: 905–914.
- Davies K.M. and Schwinn K.E. (2003) Transcriptional regulation of secondary metabolism. *Funct Plant Biol* 30: 913–925.
- Davies, K.M. and Schwinn K.E. (2006) Molecular biology and biotechnology of flavonoid biosynthesis. In: Flavonoids: Chemistry, biochemistry and applications. Andersen Ø.M. and Markham K.R., eds. CRC Press-Taylor & Francis Group, Boca Raton, FL. 143–218.
- Deroles S.C., Bradley J.M., Davies K., Schwinn K.E., Markham K.R., Bloor S., Manson D.G. and Davies K.M. (1998) An antisense chalcone synthase cDNA leads to novel colour patterns in lisianthus (*Eustoma grandiflorum*) flowers. *Mol Breed* 4: 59–66.
- Dewick P.M. (2002) Alkaloids. In: Medicinal natural products, a biosynthetic approach. 2nd edition. John Wiley & Sons. England. 291–403.

- Dhar A., Lee K.S., Dhar K. and Rosazza J.P.N. (2007) *Nocardia* sp. Vanillic acid decarboxylase. ***Enzyme Microb Technol*** 41: 271–277.
- Di Marzo V., Bisogno T. and De Petrocellis L. (2007) Endocannabinoids and related compounds: Walking back and forth between plant natural products and animal physiology. ***Chem Biol*** 14: 741–756.
- Di Marzo V. and De Petrocellis L. (2006) Plant, synthetic and endogenous cannabinoids in medicine. ***Annu Rev Med*** 57: 553–574.
- Dias A.C.P. (2003) The potential of *in vitro* cultures of *Hypericum perforatum* and of *Hypericum androsaemum* to produce interesting pharmaceutical compounds. *In: Hypericum: The genus Hypericum. Medical aromatic plants–industrial profiles. Vol. 31. Ernst E., ed. Taylor & Francis Group, London. 137–154.*
- Djoko B., Chiou R.Y.Y., Shee J.J. and Liu Y.W. (2007) Characterization of immunological activities of peanut stilbenoids, arachidin-1, piceatannol and resveratrol on lipopolysaccharide-induced inflammation of RAW 264.7 macrophages. ***J Agric Food Chem*** 55: 2376–2383.
- Dornenburg H. and Knorr D. (1994) Effectiveness of plant-derived and microbial polysaccharides as elicitors for anthraquinone synthesis in *Morinda citrifolia* cultures. ***J Agric Food Chem*** 42: 1048–1052.
- Douglas C.J. (1996) Phenylpropanoid metabolism and lignin biosynthesis: from weeds to trees. ***Trends Plant Sci*** 1: 171–178.
- Durbin M.L., McCaig B. and Clegg M.T. (2000) Molecular evolution of the chalcone synthase multigene family in the morning glory genome. ***Plant Mol Biol*** 42: 79–92.
- Durbin M.L., McCaig B. and Clegg M.T. (2000) Molecular evolution of the chalcone synthase multigene family in the morning glory genome. ***Plant Mol Biol*** 42: 79–92.
- Eckermann C., Schröder G., Eckermann S., Strack D., Schmidt J., Schneider B. and Schröder J. (2003) Stilbenecarboxylate biosynthesis: a new function in the family of chalcone synthase-related proteins. ***Phytochemistry*** 62: 271–286.
- Eckermann C., Schröder G., Schmidt J., Strack D., Edrada R.A., Helariutta Y., Elomaa P., Kotilainen M., Kilpelainen I., Proksch P., Teeri T.H. and Schröder J. (1998) New pathway to polyketides in plants. ***Nature*** 396: 387–390.
- Eilert U. and Constabel F. (1986) Elicitation of sanguinarine accumulation in *Papaver somniferum* cells by fungal homogenates—an induction process. ***J Plant Physiol*** 125: 167–172.

- El-Feraly F.S. and Turner C.E. (1975) Alkaloids of *Cannabis sativa* leaves. *Phytochemistry* 14: 2304.
- El-Feraly F.S., El-Sherei M.M. and Al-Muhtadi F.J. (1986) Spiro-indans from *Cannabis sativa*. *Phytochemistry* 25: 1992–1994.
- Elomma P., Honkanen J., Puska R., Seppanen P., Helariutta Y., Mehto M., Kotilainen M., Nevalainen L. and Teeri T.H. (1993) *Agrobacterium*-mediated transfer of antisense chalcone synthase cDNA to *Gerbera hybrida* inhibits flower pigmentation. *Biotechnology* 11: 508–511.
- ElSohly H.N., Turner C.E., Clark A.M. and ElSohly M.A. (1982) Synthesis and antimicrobial activities of certain cannabichromene and cannabigerol related compounds. *J Pharm Sci* 71: 1319–1323.
- ElSohly M.A. (1985) *Cannabis* alkaloids. In: Alkaloids, chemical and biological perspectives. Vol. 3. Pelletier S.W., ed. John Wiley & Sons, NY. 169–184.
- ElSohly M.A. and Slade D. (2005) Chemical constituents of marijuana: The complex mixture of natural cannabinoids. *Life Sci* 78: 539–548.
- ElSohly M.A., Turner C.E., Phoebe C.H., Knapp J.E., Schiff P.L. and Slatkin D.J. (1978) Anhydrocannabisativine, a new alkaloid from *Cannabis sativa*. *J Pharm Sci* 67: 124.
- Erhard D., Pohnert G. and Gross E.M. (2007) Chemical defense in *Elodea nuttallii* reduces feeding and growth of aquatic herbivorous Lepidoptera. *J Chem Ecol* 33: 1646–1661.
- Eriksson L., Johansson E., Kettaneh-Wold N., Trygg J., Wikstrom C. and Wold S. (2006) Multi- and megavariate data analysis. Part 1: Basic principles and applications. Second edition. Umetrics Academy, Umea, Sweden.
- Estrada-Soto S., Lopez-Guerrero J.J., Villalobos-Molina R. and Mata R. (2006) Endothelium-independent relaxation of aorta rings by two stilbenoids from the orchids *Scaphyglottis livida*. *Fitoterapia* 77: 236–239.
- Facchini P.J. and De Luca V. (1995) Phloem-specific expression of tyrosine/dopa decarboxylase genes and the biosynthesis of isoquinoline alkaloids in opium poppy. *Plant Cell* 7: 1811–1821.
- Facchini P.J., Johnson A.G., Poupart J. and De Luca V. (1996) Uncoupled defense gene expression and antimicrobial alkaloid accumulation in elicited opium poppy cell cultures. *Plant Physiol* 111: 687–697.

- Fellermeier M. and Zenk M.H. (1998) Prenylation of olivetolate by a hemp transferase yields cannabigerolic acid, the precursor of tetrahydrocannabinol. *FEBS Lett* 427: 283–285.
- Fellermeier M., Eisenreich W., Bacher A. and Zenk M.H. (2001) Biosynthesis of cannabinoids: Incorporation experiments with ^{13}C -labeled glucoses. *Eur J Biochem* 268: 1596–1604.
- Ferrer J.L., Jez J.M., Bowman M.E., Dixon R.A. and Noel J.P. (1999) Structure of chalcone synthase and the molecular basis of plant polyketide biosynthesis. *Nat Struct Biol* 6: 775–784.
- Fields P.G., Arnason J.T. and Fulcher R.G. (1990) The spectral properties of *Hypericum perforatum* leaves: The implications for its photoactivated defenses. *Can J Bot* 68: 1166–1170.
- Fischbach M.A. and Walsh C.T. (2006) Assembly-line enzymology for polyketide and nonribosomal peptide antibiotics: Logic, machinery, and mechanisms. *Chem Rev* 106: 3468–3496.
- Fischer R., Budde I. and Hain R. (1997) Stilbene synthase gene expression causes changes in flower colour and male sterility in tobacco. *Plant J* 11: 489–498.
- Fliegmann J., Schröder G., Schanz S., Britsch L. and Schröder J. (1992) Molecular analysis of chalcone and dihydropinosylvin synthase from Scots pine (*Pinus sylvestris*), and differential regulation of these and related enzyme activities in stressed plants. *Plant Mol Biol* 18: 489–503.
- Flores-Sanchez I.J., Ortega-Lopez J., Montes-Horcasitas M.C. and Ramos-Valdivia A.C. (2002) Biosynthesis of sterols and triterpenes in cell suspension cultures of *Uncaria tomentosa*. *Plant Cell Physiol* 43: 1502–1509.
- Formukong E.A., Evans A.T. and Evans F.J. (1988) Analgesic and antiinflammatory activity of constituents of *Cannabis sativa* L. *Inflammation* 12: 361–371.
- Fournier G., Richez-Dumanois C., Duvezin J., Mathieu J.P. and Paris M. (1987) Identification of a new chemotype in *Cannabis sativa*: Cannabigerol-dominant plants, biogenetic and agronomic prospects. *Planta Med* 53: 277–280.
- Fritzemeier K.H. and Kindl H. (1983) 9,10-dihydrophenanthrenes as phytoalexins of Orchidaceae: Biosynthetic studies *in vitro* and *in vivo* proving the route from L-phenylalanine to dihydro-*m*-coumaric acid, dihydrostilbene and dihydrophenanthrenes. *Eur J Biochem* 133: 545–550.
- Funa N., Awakawa T. and Horinouchi S. (2007) Pentaketide resorcylic acid synthesis by type III polyketide synthase from *Neurospora crassa*. *J Biol Chem* 282: 14476–14481.

- Funa N., Ohnishi Y., Fujii I., Ebizuka Y. and Horinouchi S. (2002) Properties and substrate specificity of RppA, a chalcone synthase-related polyketide synthase in *Streptomyces griseus*. *J Biol Chem* 277: 4628–4635.
- Funa N., Ohnishi Y., Fujii I., Shibuya M., Ebizuka Y. and Horinouchi S. (1999) A new pathway for polyketide synthesis in microorganism. *Nature* 400: 897–899.
- Fung S.Y., Brussee J., Van der Hoeven R.A.M., Niessen W.M.A., Scheffer J.J.C. and Verpoorte R. (1994) Analysis of proposed aromatic precursors of hop bitter acids. *J Nat Prod* 57: 452–459.
- Gaisser S., Trefzer A., Stockert S., Kirschning A. and Bechthold A. (1997) Cloning of an avilamycin biosynthetic gene cluster from *Streptomyces viridochromogenes* Tü57. *J Bacteriol* 179: 6271–6278.
- Gamborg O.L., Miller R.A., Ojima K. (1968) Nutrient requirements of suspension cultures of soybean root cells. *Exp Cell Res* 50: 151.
- Garcez W.S., Martins D., Garcez F.R., Marquez M.R., Pereira A.A., Oliveira L.A., Rondon J.N. and Peruca A.D. (2000) Effect of spores of saprophytic fungi on phytoalexin accumulation in seeds of frog-eye leaf spot and stem canker-resistant and -susceptible soybean (*Glycine max* L.) cultivars. *J Agric Food Chem* 48: 3662–3665.
- Garcia E.S. and Azambuja P. (2004) Lignoids in insects: Chemical probes for the study of ecdysis, excretion and *Trypanosoma cruzi*-triatomine interactions. *Toxicon* 44: 431–440.
- Gaucher G.M. and Shepherd M.G. (1968) Isolation of orsellinic acid synthase. *Biochem Biophys Res Commun* 32: 664–671.
- Gehlert R. and Kindl H. (1991) Induced formation of dihydrophenanthrenes and bibenzyl synthase upon destruction of orchid mycorrhiza, *Phytochemistry* 30: 457–460.
- Giuffrida A., Parsons L.H., Kerr T.M., Rodriguez de Fonseca F., Navarro M. And Piomelli D. (1999) Dopamine activation of endogenous cannabinoid signaling in dorsal striatum. *Nat Neurosci* 2: 358–363.
- Goldstein J.L. and Brown M.S. (1990) Regulation of mevalonate pathway. *Nature* 434: 425–430.
- Gorham J. (1977) Lunularic acid and related compounds in liverworts, algae and *Hydrangea*. *Phytochemistry* 16: 249–253.
- Gorham J. (1980) The Stilbenoids. In: Progress in Phytochemistry. Vol. 6. Reinhold L., Harborne J.B. and Swain T., eds. Pergamon Press, Oxford. 203–252.

- Gorham J., Tori M. and Asakawa Y. (1995) The biochemistry of stilbenoids. Biochemistry of natural products series. Vol.1. Harborne J.B. and Baxter H., eds. Chapman & Hall, London.
- Goto-Yamamoto N., Wan G.H., Masaki K. and Kobayashi S. (2002) Structure and transcription of three chalcone synthase genes of grapevine (*Vitis vinifera*). *Plant Sci* 162: 867–872.
- Gould K.S. and Lister C. (2006) Flavonoid functions in plants. *In*: Flavonoids: Chemistry, biochemistry and applications. Andersen Ø.M. and Markham K.R., eds. CRC Press–Taylor & Francis Group, Boca Raton, FL. 397–441.
- Grotenhermen F. (2002) Review of therapeutic effects. *In*: Cannabis and cannabinoids: Pharmacology, toxicology and therapeutic potential. Grotenhermen F. and Russo E., eds. The Haworth Integrative Healing Press, New York. 123–142.
- Guex N. and Peitsch M.C. (1997) Swiss-Model and the Swiss-PdbViewer: An environment for comparative protein modeling. *Electrophoresis* 18: 2714–2723.
- Hagel J.M. and Facchini P.J. (2008) Plant metabolomics: Analytical platforms and integration with functional genomics. DOI 10.1007/s11101-007-9086-9.
- Hamada T. (2005) New development of photo-induced electron transfer reaction and total synthesis of natural product. *Yakugaku Zasshi* 125: 1–16.
- Hammond C.T. and Mahlberg P.G. (1994) Phloroglucinol glucoside as a natural constituent of *Cannabis sativa*. *Phytochemistry* 37: 755–756.
- Hampson A.J., Grimaldi M., Axelrod J. and Wink D. (1998) Cannabidiol and (–) Δ^9 -tetrahydrocannabinol are neuroprotective antioxidants. *Proc Natl Acad Sci USA* 95: 8268–8273.
- Han S.H., Lee H.H., Lee I.S., Moon Y.H. and Woo E.R. (2002) A new phenolic amide from *Lycium chinense* Miller. *Arch Pharm Res* 25: 433–437.
- Harker C.L., Ellis T.H.N. and Coen E.S. (1990) Identification and genetic regulation of the chalcone synthase multigene family in pea. *Plant Cell* 2: 185–194.
- Hartsel S.C., Loh W.H.T. and Robertson L.W. (1983) Biotransformation of cannabidiol to cannabielsoin by suspension cultures of *Cannabis sativa* and *Saccharum officinarum*. *Planta Med* 48: 17–19.

- Hazekamp A., Peltenburg-Looman A., Verpoorte R. and Giroud C. (2005) Chromatographic and spectroscopic data of cannabinoids from *Cannabis sativa* L. *J Liq Chromatogr Relat Technol* 28: 2361–2382.
- Hazekamp A., Simons R., Peltenburg-Looman A., Sengers M., van Zweden R. and Verpoorte R. (2004) Preparative Isolation of cannabinoids from *Cannabis sativa* by centrifugal partition chromatography. *J Liq Chromatogr Relat Technol* 27: 2421–2439.
- Heath R.J. and Rock C.O. (2002) The Claisen condensation in biology. *Nat Prod Rep* 19: 581–596.
- Heitrich A. and Binder M. (1982) Identification of (3R, 4R)- $\Delta^1(6)$ -tetrahydrocannabinol as an isolation artifact of cannabinoid acids formed by callus cultures of *Cannabis sativa* L. *Experientia* 38: 898–899.
- Helariutta Y., Elomaa P., Kotilainen M., Griesbach R.J., Schröder J. and Teeri T. (1995) Chalcone synthase-like genes active during corolla development are differentially expressed and encode enzymes with different catalytic properties in *Gerbera hybrida* (Asteraceae). *Plant Mol Biol* 28: 47–60.
- Helariutta Y., Kotilainen M., Elomaa P., Kalkkinen N., Bremer K., Teeri T. and Albert V.A. (1996) Duplication and functional divergence in the chalcone synthase gene family of *Asteraceae*: Evolution with substrate change and catalytic simplification. *Proc Natl Acad Sci USA* 93: 9033–9038.
- Hendriks H., Malingre T.M., Batterman S. and Bos R. (1978) The essential oil of *Cannabis sativa* L. *Pharm Weekbl* 113: 413–424.
- Henness S., Robinson D.M. and Lyseng-Williamson K.A. (2006) Rimonabant. *Drugs* 66: 2109–2119.
- Henstrand J.M., McCue K.F., Brink K., Handa A.K., Herrmann K.M. and Conn E.E. (1992) Light and fungal elicitor induce 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase mRNA in suspension cultured cells of parsley (*Petroselinum crispum* L.). *Plant Physiol* 98: 761–763.
- Herderich M., Beckert C. and Veit M. (1997) Establishing styrylpyrone synthase activity in cell free extracts obtained from gametophytes of *Equisetum arvense* L. by high performance liquid chromatography–tandem mass spectrometry. *Phytochem Anal* 8: 194–197.
- Hillig K.W. (2004) A chemotaxonomic analysis of terpenoid variation in *Cannabis*. *Biochem Syst Ecol* 32: 875–891.

- Hillig K.W. (2005) Genetic evidence for separation in *Cannabis* (Cannabaceae). *Genet Resour Crop Evol* 52: 161–180.
- Hillis W.E. and Inoue T. (1968) The formation of polyphenols in trees–IV: The polyphenols formed in *Pinus radiata* after *Sirex* attack. *Phytochemistry* 7: 13–22.
- Hipskind J.D. and Paiva N.L. (2000) Constitutive accumulation of a resveratrol–glucoside in transgenic alfalfa increases resistance to *Phoma medicaginis*. *Mol Plant Microbe Interact* 13: 551–562.
- Hirner A.A. and Seitz H.U. (2000) Isoforms of chalcone synthase in *Daucus carota* L. and their differential expression in organs from the European wild carrot and in ultraviolet–A–irradiated cell cultures. *Planta* 210: 993–998.
- Hoelzl J. and Petersen M. (2003) Chemical constituents of *Hypericum* spp. *Med Aromat Plant–Ind Profiles* 31: 77–93.
- Höfig K.P., Moller R., Donaldson L., Putterill J. and Walter C. (2006) Towards male sterility in *Pinus radiata*: A stilbene synthase approach to genetically engineer nuclear male sterility. *Plant Biotechnol J* 4: 333–343.
- Hohlfeld H., Scheel D. and Strack D. (1996) Purification of hydroxycinnamoyl–CoA:tyramine hydroxycinnamoyltransferase from cell–suspension cultures of *Solanum tuberosum* L. cv. Datura. *Planta* 199: 166–168.
- Holley J.H., Hadley K.W. and Turner C.E. (1975) Constituents of *Cannabis sativa* L. XI: Cannabidiol and cannabichromene in samples of known geographical origin. *J Pharm Sci* 64: 892–895.
- Hopwood D.A. and Sherman D.H. (1990) Molecular genetics of polyketides and its comparison to fatty acid biosynthesis. *Annu Rev Genet* 24: 37–66.
- Horper W. and Marner F.J. (1996) Biosynthesis of primin and miconidin and its derivatives. *Phytochemistry* 41: 451–456.
- Huang Z., Dostal L. and Rosazza J.P.N. (1994) Purification and characterization of a ferulic acid decarboxylase from *Pseudomonas fluorescens*. *J Bacteriol* 176: 5912–5918.
- Huber S.C. and Hardin S.C. (2004) Numerous posttranslational modifications provide opportunities for the intricate regulation of metabolic enzymes at multiple levels. *Curr Opin Plant Biol* 7: 318–322.

- Hui L., Jin Z., Xiayu D., Baoqin S., Shuang J., Daowen W., Junwen O., Jiayang L., Liangcai L., Wenzhong T., Hain R. and Xu J. (2000) A transgenic wheat with a stilbene synthase gene resistant to powdery mildew obtained by biolistic method. *Chin Sci Bull* 45: 634–638.
- Husken A., Baumert A., Milkowski C., Becker H.C., Strack D. and Mollers C. (2005) Resveratrol glucoside (piceid) synthesis in seeds of transgenic oilseed rape (*Brassica napus* L.). *Theor Appl Genet* 111: 1553–1562.
- Iliya I., Akao Y., Matsumoto K., Nakagawa Y., Zulfiqar A., Ito T., Oyama M., Murata H., Tanaka T., Nozawa Y. and Iinuma M. (2006) Growth inhibition of stilbenoids in Welwitschiaceae and Gnetaceae through induction of apoptosis in human leukemia HL60 cells. *Biol Pharm Bull* 29: 1490–1492.
- Itokawa H., Takeya K. and Mihashi S. (1977) Biotransformation of cannabinoid precursors and related alcohols by suspension cultures of callus induced from *Cannabis sativa* L. *Chem Pharm Bull* 25: 1941–1946.
- Jabs T. (1999) Reactive oxygen intermediates as mediators of programmed cell death in plants and animals. *Biochem Pharmacol* 57: 231–245.
- Jacobs M. and Rubery P.H. (1988) Naturally occurring auxin transport regulators. *Science* 241:346–349.
- Jang S.M., Ishihara A. and Back K. (2004) Production of coumaroylserotonin and feruloylserotonin in transgenic rice expressing pepper hydroxycinnamoyl-coenzyme A:serotonin *N*-(hydroxycinnamoyl)transferase. *Plant Physiol* 135: 346–356.
- Jaroszewski J.W. (2005) Hyphenated NMR methods in natural products research, part 2: HPLC–SEP–NMR and other new trends in NMR hyphenation. *Planta Med* 71: 795–802.
- Jeandet P., Douillet-Breuil A.C., Bessis R., Debord S., Sbaghi M. and Adrian M. (2002) Phytoalexins from the Vitaceae: Biosynthesis, phytoalexin gene expression in transgenic plants, antifungal activity and metabolism. *J Agric Food Chem* 50: 2731–2741.
- Jekkel Z.S., Heszky L.E. and Ali A.H. (1989) Effect of different cryoprotectants and transfer temperatures on the survival rate of hemp (*Cannabis sativa* L.) cell suspension in deep freezing. *Acta Biol Hung* 40: 127–136.
- Jenke-Kodama H., Muller R. and Dittmann E. (2008) Evolutionary mechanism underlying secondary metabolite diversity. *Prog Drug Res* 65: 119, 121–140.

- Jez J.M., Austin M.B., Ferrer J.L., Bowman M.E., Schröder J. and Noel J.P. (2000a) Structural control of polyketide formation in plant-specific polyketide synthases. *Chem Biol* 7: 919–930.
- Jez J.M., Bowman M.E. and Noel J.P. (2001b) Structure-guided programming of polyketide chain-length determination in chalcone synthase. *Biochemistry* 40: 14829–14838.
- Jez J.M., Ferrer J.L., Bowman M.E., Austin M.B., Schröder J., Dixon R.A. and Noel J.P. (2001a) Structure and mechanism of chalcone synthase-like polyketide synthases. *J Ind Microbiol Biotechnol* 27: 393–398.
- Jez J.M., Ferrer J.L., Bowman M.E., Dixon R.A. and Noel J.P. (2000b) Dissection of malonyl-coenzyme A decarboxylation from polyketide formation in the reaction mechanism of a plant polyketide synthase. *Biochemistry* 39: 890–902.
- Jiang H.E., Li X., Zhao Y.X., Ferguson D.K., Hueber F., Bera S., Wang Y.F., Zhao L.C., Liu C.J. and Li C.S. (2006) A new insight into *Cannabis sativa* (Cannabaceae) utilization from 2500-year-old Yanghai Tombs, Xinjiang, China. *J Ethnopharmacol* 108: 414–422.
- Jin W. and Zjawiony J.K. (2006) 5-alkylresorcinols from *Merulius incarnates*. *J Nat Prod* 69: 704–706.
- Johnson J.M., Lemberger L., Novotny M., Forney R.B., Dalton W.S. and Maskarinec M.P. (1984) Pharmacological activity of the basic fraction of marijuana whole smoke condensate alone and in combination with delta-9-tetrahydrocannabinol in mice. *Toxicol Appl Pharmacol* 72: 440–448.
- Jones T.H., Brunner S.R., Edwards A.A., Davidson D.W. and Snelling R.R. (2005) 6-alkylsalicylic acids and 6-alkylresorcylic acids from ants in the genus *Crematogaster* from Brunei. *J Chem Ecol* 31: 407–417.
- Jorgensen K., Rasmussen A.V., Morant M., Nielsen A.H., Bjarnholt N., Zagrobelny M., Bak S. and Moller B.L. (2005) Metabolon formation and metabolic channeling in the biosynthesis of plant natural products. *Curr Opin Plant Biol* 8: 280–291.
- Junghanns K.T., Kneusel R.E., Gröger D. and Matern U. (1998) Differential regulation and distribution of acridone synthase in *Ruta graveolens*. *Phytochemistry* 49: 403–411.
- Junghans H., Dalkin K. and Dixon R.A. (1993) Stress response in alfalfa (*Medicago sativa* L.) 15: Characterization and expression patterns of members of a subset of the chalcone synthase multigene family. *Plant Mol Biol* 22: 239–253.

- Justesen U., Knuthsen P. and Leth T. (1998) Quantitative analyses of flavonols, flavones and flavanones in fruits, vegetables and beverages by high-performance liquid chromatography with photo-diode array and mass spectrometric detection. *J Chromatogr A* 799: 101–110.
- Kabarity A., El-Bayoumi A. and Habib A. (1980) C-tumours and polyploidy induced by some alkaloids of Opium and Cannabis. *Cytologia* 45: 497–506.
- Kaeberlein M., McDonagh T., Heltweg B., Hixon J., Westman E.A., Caldwell S.D., Napper A., Curtis R., DiStefano P.S., Fields S., Bedalov A. and Kennedy B.K. (2005) Substrate-specific activation of sirtuins by resveratrol. *J Biol Chem* 280: 17038–17045.
- Kajima M. and Piraux M. (1982) The biogenesis of cannabinoids in *Cannabis sativa*. *Phytochemistry* 21: 67–69.
- Karhunen T., Airaksinen M.S., Tuomisto L. and Panula P. (1993) Neurotransmitters in the nervous system of *Macoma balthica* (Bivalvia). *J Comp Neurol* 334: 477–488.
- Karst M., Salim K., Burstein S., Conrad I., Hoy L. and Schneider U. (2003) Analgesic effect of the synthetic cannabinoid CT-3 on chronic neuropathic pain. *JAMA* 290: 1757–1762.
- Katsuyama Y., Funa N., Miyahisa I. and Horinouchi S. (2007) Synthesis of unnatural flavonoids and stilbenes by exploiting the plant biosynthetic pathway in *Escherichia coli*. *Chem Biol* 14: 613–621.
- Katsuyama Y., Matsuzawa M., Funa N. and Horinouchi S. (2007) *In vitro* synthesis of curcuminoids by type III polyketide synthase from *Oryza sativa*. *J Biol Chem* 282: 37702–37709.
- Keller A., Leupin M. Mediavilla V. and Wintermantel E. (2001) Influence of the growth stage of industrial hemp on chemical and physical properties of the fibres. *Ind Crop Prod* 13: 35–48.
- Kettenes-van den Bosch J.J. (1978) New constituents of *Cannabis sativa* L. and its smoke condensate. Ph.D. Thesis. State University Utrecht, The Netherlands.
- Kettenes-van den Bosch J.J. and Salemink C.A. (1978) Cannabis XIX: Oxygenated 1,2-diphenylethanes from marihuana. *J R Netherlands Chem Soc* 97: 221–222.
- Kim E.S. and Mahlberg P.G. (1997) Immunochemical localization of tetrahydrocannabinol (THC) in cryofixed glandular trichomes of *Cannabis* (Cannabaceae). *Am J Bot* 84: 336–342.
- Kim J.I., Park J.E., Zarate X. and Song P.S. (2005) Phytochrome phosphorylation in plant light signaling. *Photochem Photobiol Sci* 4: 681–687.

- Kim Y., Han M.S., Lee J.S., Kim J. and Kim Y.C. (2003) Inhibitory phenolic amides on lipopolysaccharide-induced nitric oxide production in RAW 264.7 cells from *Beta vulgaris* var. *cicla* seeds. *Phytother Res* 17: 983–985.
- Kimura M. and Okamoto K. (1970) Distribution of tetrahydrocannabinolic acid in fresh wild *Cannabis*. *Experientia* 26: 819–820.
- Kindl H. (1985) Biosynthesis of stilbenoids. *In*: Biosynthesis and biodegradation of wood components. Higuchi T., ed. Academic Press Inc., New York. 349–377.
- King R.R. and Calhoun L.A. (2005) Characterization of cross-linked hydroxycinnamic acid amides isolated from potato common scab lesions. *Phytochemistry* 66: 2468–2473.
- Klein F.K. and Rapoport H. (1971) Cannabis alkaloids. *Nature* 232: 258–259.
- Klingauf P., Beuerle T., Mellenthin A., El-Moghazy S.A., Boubakir Z. and Beerhues L. (2005) Biosynthesis of the hyperforin skeleton in *Hypericum calycinum* cell cultures. *Phytochemistry* 66: 139–145.
- Knoller N., Levi L., Shoshan I., Reichenthal E., Razon N., Rappaport Z.H. and Biegon A. (2002) Dexanabinol (HU-211) in the treatment of severe closed head injury: A randomized, placebo-controlled, phase II clinical trial. *Crit Care Med* 30: 548–554.
- Kobayashi S., Ding C.K., Nakamura Y., Nakajima I. and Matsumoto R. (2000) Kiwifruits (*Actinidia deliciosa*) transformed with a *Vitis* stilbene synthase gene produce piceid (resveratrol-glucoside). *Plant Cell Rep* 19: 904–910.
- Koes R.E., Spelt C.E., van den Elzen P.J.M. and Mol J.N.M. (1989) Cloning and molecular characterization of the chalcone synthase multigene family of *Petunia hybrida*. *Gene* 81: 245–257.
- Kostecki K., Engelmeier D., Pacher T., Hofer O., Vajrodaya S. and Greger H. (2004) Dihydrophenanthrenes and other antifungal stilbenoids from *Stemona* cf. *pierrei*. *Phytochemistry* 65: 99–106.
- Kozubek A. and Tyman J.H.P. (1999) Resorcinolic lipids, the natural non-isoprenoid phenolic amphiphiles and their biological activity. *Chem Rev* 99: 1–25.
- Kraemer K.H., Schenkel E.P. and Verpoorte R (1999) Glucosylation of ethanol in *Ilex paraguariensis* cell suspension cultures. *Plant Cell Rep* 18: 509–513.

- Kreuzaler F. and Hahlbrock K. (1972) Enzymatic synthesis of aromatic compounds in higher plants: formation of naringenin (5,7,4'-trihydroxyflavanone) from *p*-coumaroyl coenzyme A and malonyl coenzyme A. *FEBS Lett* 28: 69–72.
- Kuethe J.T. and Comins D.L. (2004) Asymmetric total synthesis of (+)-cannabisativine. *J Org Chem* 69: 5219–5231.
- Kumar A. and Ellis B.E. (2003) A family of polyketide synthase genes expressed in ripening *Rubus* fruits. *Phytochemistry* 62: 513–526.
- Kurosaki F., Tsurusawa Y. and Nishi A. (1987) The elicitation of phytoalexins by Ca²⁺ and cyclic AMP in carrot cells. *Phytochemistry* 26: 1919–1923.
- Kushima H., Shoyama Y. and Nishioka I. (1980) Cannabis XII: Variations of cannabinoid contents in several strains of *Cannabis sativa* L. with leaf-age, season and sex. *Chem Pharm Bull* 28: 594–598.
- Lajide L., Escoubas P. and Mizutani J. (1995) Termite antifeedant activity in *Xylopi aethiopica*. *Phytochemistry* 40: 1105–1112.
- Lanyon V.S., Turner J.C. and Mahlberg P.G. (1981) Quantitative analysis of cannabinoids in the secretory product from capitate-stalked glands of *Cannabis sativa* L. (Cannabaceae). *Bot Gaz* 142: 316–319.
- Lanz T., Tropf S., Marner F.J., Schröder J. and Schröder G. (1991) The role of cysteines in polyketide synthases: Site-directed mutagenesis of resveratrol and chalcone synthases, two key enzymes in different plant-specific pathways. *J Biol Chem* 266: 9971–9976.
- Laurain-Mattar D., Gillet-Manceau F., Buchon L., Nabha S., Fliniaux M.A. and Jacquin-Dubreuil A. (1999) Somatic embryogenesis and rhizogenesis of tissue cultures of two genotypes of *Papaver somniferum*: Relationships to alkaloid production. *Planta Med* 65: 167–170.
- LeClere S., Schmelz E.A. and Chourey P.S. (2007) Phenolic compounds accumulate specifically in maternally-derived tissues of developing maize kernels. *Cereal Chem* 84: 350–356.
- Lee D.G., Park Y., Kim M.R., Jung H.J., Seu Y.B., Hahm K.S. and Woo E.R. (2004) Anti-fungal effects of phenolic amides isolated from the root bark of *Lycium chinense*. *Biotechnol Lett* 26: 1125–1130.
- Lee K.Y., Sung S.H. and Kim Y.C. (2006) Neuroprotective bibenzyl glycosides of *Stemona tuberosa* roots. *J Nat Prod* 69: 679–681.

- Lee S.K., Lee H.J., Min H.Y., Park E.J., Lee K.M., Ahn Y.H., Cho Y.J. and Pyee J.H. (2005) Antibacterial and antifungal activity of pinosylvins, a constituent of pine. *Fitoterapia* 76: 258–260.
- Lehmann T. and Brenneisen R. (1995) High performance liquid chromatographic profiling of Cannabis products. *J Liq Chromatogr* 18: 689–700.
- Leiro J., Arranz J.A., Fraiz N., Sanmartin M.L., Quezada E. and Orallo F. (2005) Effects of *cis*-resveratrol on genes involved in nuclear factor kappa B signaling. *Int Immunopharmacol* 5: 393–406.
- Lewis G.S. and Turner C.E. (1978) Constituents of *Cannabis sativa* L. XIII: Stability of dosage form prepared by impregnating synthetic (–) Δ^9 -*trans*-tetrahydrocannabinol on placebo Cannabis plant material. *J Pharm Sci* 67: 876–878.
- Lewis N.G. and Davin L.B. (1999) Lignans: Biosynthesis and function. *In: Comprehensive natural products chemistry*. Barton D.H.R., Nakanishi K. and Meth-Cohn O., eds. Vol. 1. Polyketides and other secondary metabolites including fatty acids and their derivatives. Sankawa U., ed. Elsevier Science Ltd., Oxford, UK. 639–712.
- Li T.L., Choroba O.W., Hong H., Williams D.H. and Spencer J.B. (2001) Biosynthesis of the vancomycin group of antibiotics: Characterization of a type III polyketide synthase in the pathway to (S)-3,5-dihydroxyphenylglycine. *Chem Commun* 20: 2156–2157.
- Linnaeus C. (1753) Species plantarum. T. I–II.
- Liswidowati, Melchior F., Hohmann F., Schwer B. and Kindl H. (1991) Induction of stilbene synthase by *Botrytis cinerea* in cultured grapevine cells. *Planta* 183: 307–314.
- Liu B., Falkenstein-Paul H., Schmidt W. and Beerhues L. (2003) Benzophenone synthase and chalcone synthase from *Hypericum androsaemum* cell cultures: cDNA cloning, functional expression and site-directed mutagenesis of two polyketide synthases. *Plant J* 34: 847–855.
- Liu B., Raeth T., Beuerle T. and Beerhues L. (2007) Biphenyl synthase, a novel type III polyketide synthase. *Planta* 225: 1495–1505.
- Lo C., Coolbaugh R.C. and Nicholson R.L. (2002) Molecular characterization and *in silico* expression analysis of a chalcone synthase gene family in *Sorghum bicolor*. *Physiol Mol Plant Pathol* 61: 179–188.
- Loh W.H.T., Hartsel S.C. and Robertson L.W. (1983) Tissue culture of *Cannabis sativa* L. and *in vitro* biotransformation of phenolics. *Z Pflanzenphysiol* 111: S395–400.

- Lois R. and Buchanan B.B. (1994) Severe sensitivity to ultraviolet radiation in an *Arabidopsis* mutant deficient in flavonoid accumulation. II. Mechanisms of UV-resistance in *Arabidopsis*. *Planta* 194: 504–509.
- Luan S. (2003) Protein phosphatases in plants. *Annu Rev Plant Biol* 54: 63–92.
- Lukacin R., Schreiner S. and Matern U. (2001) Transformation of acridone synthase to chalcone synthase. *FEBS Lett* 508: 413–417.
- Ma C.Y., Liu W.K. and Che C.T. (2002) Lignanamides and nonalkaloidal components of *Hyoscyamus niger* seeds. *J Nat Prod* 65: 206–209.
- Ma J., Jones S.H. and Hecht S.M. (2004) Phenolic acid amides: A new type of DNA strand scission agent from *Piper caninum*. *Bioorg Med Chem* 12: 3885–3889.
- MacCaman M.W., Stetzler J. and Clark B. (1985) Synthesis of γ -glutamyl dopamine and other peptidoamines in the nervous system of *Aplysia californica*. *J Neurochem* 45: 1828–1835.
- Macfarlane R.G., Macleod S.C., Midgley J.M. and Watson D.G. (1989) Analysis of biogenic amines in bovine retina by gas chromatography–negative ion chemical ionization mass spectrometry. *J Neurochem* 53: 1731–1736.
- Mahlberg P.G., Hammond C.T., Turner J.C. and Hemphill J.K. (1984) Structure, development and composition of glandular trichomes of *Cannabis sativa* L. *In: Biology and Chemistry of plant trichomes*. Rodriguez E., Healey P.L. and Mehta I., eds. Plenum Press, New York. 23–51.
- Majak W., Bai Y. and Benn M.H. (2003) Phenolic amides and isoquinoline alkaloids from *Corydalis sempervirens*. *Biochem Syst Ecol* 31: 649–651.
- Malingre T.H., Hendriks H., Batterman S., Bos R. and Visser J. (1975) The essential oil of *Cannabis sativa*. *Planta Med* 28: 56–61.
- Mandolino G. and Ranalli P. (1999) Advances in biotechnological approaches for hemp breeding and industry. *In: Advances in hemp research*. Ranalli P., ed. Food Products Press, NY. 185–212.
- Manthey J.A. and Buslig B.S. (1998) Flavonoids in the living system. *Adv Exp Med Biol* 439: 1–7.
- Martin–Tanguy J. (1985) The occurrence and possible function of hydroxycinnamoyl acid amides in plants. *Plant Growth Regul* 3: 381–399.

- Massi P., Vaccani A., Ceruti S., Colombo A., Abbracchio M.P. and Parolaro D. (2004) Antitumor effects of cannabidiol, a nonpsychoactive cannabinoid, on human glioma cell lines. *J Pharmacol Exp Ther* 308: 838–845.
- Matousek J., Novak P., Briza J., Patzak J. and Niedermeierova H. (2002a) Cloning and characterization of *chs*-specific DNA and cDNA sequences from hop (*Humulus lupulus* L.). *Plant Sci* 162: 1007–1018.
- Matousek J., Novak P., Patzak J., Briza J. and Krofta K. (2002b) Analyses of true chalcone synthase from *Humulus lupulus* L. and biotechnology aspects of medicinal hops. *Rostl Vyroba* 48: 7–14.
- Matsuda L.A., Lolait S.J., Brownstein M., Young A. and Bonner T.I. (1990) Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 346: 561–564.
- Maxwell G.D., Moore M.M. and Hildebrand J.G. (1980) Metabolism of tyramine in the central nervous system of the moth *Manduca sexta*. *Insect Biochem* 10: 657–665.
- McClanahan R.H. and Robertson L.W. (1984) Biotransformation of olivetol by *Syncephalastrum racemosum*. *J Nat Prod* 47: 828–834.
- McGarvey D.J. and Croteau R. (1995) Terpenoid metabolism. *Plant Cell* 7: 1015–1026.
- McNeil S.D., Nuccio M.L., Rhodes D., Shachar-Hill Y. and Hanson A.D. (2000) Radiotracer and computer modeling evidence that phospho-base methylation is the main route of choline synthesis in tobacco. *Plant Physiol* 123: 371–380.
- McPartland J.M. and Mediavilla V. (2002) Noncannabinoid components. *In: Cannabis and cannabinoids: Pharmacology, toxicology and therapeutic potential*. Grothenhermen F. and Russo E., eds. The Haworth Integrative Healing Press, New York. 401–409.
- McPartland J.M., Clarke R.C. and Watson D.P. (2000) Hemp diseases and pests: Management and biological control. CABI Publishing, Wallingford, UK.
- Mechoulam R. (1970) Marijuana chemistry. *Science* 168: 1159–1166.
- Mechoulam R. (1988) Alkaloids in *Cannabis sativa* L. *In: The alkaloids, chemistry and pharmacology*. Vol. 34. Brussi A., ed. Academic Press Inc, USA. 77–93.
- Mechoulam R. and Ben-Shabat S. (1999) From *gan-zi-gun-nu* to anandamide and 2-arachidonoylglycerol: the ongoing story of cannabis. *Nat Prod Rep* 16: 131–143.

- Mechoulam R., Fride E. and Di Marzo V. (1998) Endocannabinoids. *Eur J Pharm* 359: 1–18.
- Mediavilla V. and Steinemann S. (1997) Essential oil of *Cannabis sativa* L. strains. *J Int Hemp Assoc* 4: 82–84.
- Meigs T.E. and Simoni R.D. (1997) Farnesol as a regulator of HMG–CoA reductase degradation: Characterization and role of farnesyl pyrophosphatase. *Arch biochem Biophys* 345: 1–9.
- Miller I.J., McCallum N.K., Kirk C.M. and Peake B.M. (1982) The free radical oxidation of tetrahydrocannabinols. *Experientia* 38: 230–231.
- Mitscher L.A., Park Y.H., Al-Shamma A., Hudson P.B. and Hass T. (1981) Amorfrutin A and B, bibenzyl antimicrobial agents from *Amorpha fruticosa*. *Phytochemistry* 20: 781–785.
- Molnar J., Csiszar K., Nishioka I. and Shoyama Y. (1986) The effects of cannabispino compounds and tetrahydrocannabinolic acid on the plasmid transfer and maintenance in *E. coli*. *Acta Microbiol Hung* 33: 221–231.
- Money T., Comer F.W., Webster G.R.B., Wright I.G. and Scott A.I. (1967) Pyrone studies–I: Biogenetic–type synthesis of phenolic compounds. *Tetrahedron* 23: 3435–2448.
- Moore B.S. and Hopke J.N. (2001) Discovery of a new bacterial polyketide biosynthetic pathway. *Chem Biochem* 2: 35–38.
- Morelli R., Das S., Bertelli A., Bollini R., Scalzo R.L., Das D.K. and Falchi M. (2006) The introduction of the stilbene synthase gene enhances the natural antiradical activity of *Lycopersicon esculentum* mill. *Mol Cell Biochem* 282: 65–73.
- Moriguchi T., Kita M., Tomono Y., Endo–Inagaki T. and Mitsuo O. (1999) One type of chalcone synthase gene expressed during embryogenesis regulates the flavonoid accumulation in *Citrus* cell cultures. *Plant Cell Physiol* 40: 651–655.
- Morimoto S., Komatsu K., Taura F. and Shoyama Y. (1998) Purification and characterization of cannabichromenic acid synthase from *Cannabis sativa*. *Phytochemistry* 49: 1525–1529.
- Morimoto S., Tanaka Y., Sasaki K., Tanaka H., Fukamizu T., Shoyama Y., Shoyama Y. and Taura F. (2007) Identification and characterization of cannabinoids that induce cell death through mitochondrial permeability transition in *Cannabis* leaf cells. *J Biol Chem* 282: 20739–20751.
- Morimoto S., Taura F. and Shoyama Y. (1999) Biosynthesis of cannabinoids in *Cannabis sativa* L. *Curr Top Phytochem* 2: 103–113.

- Morita H., Kondo S., Kato R., Wanibuchi K., Noguchi H., Sugio S., Abe I. and Kohno T. (2007) Crystallization and preliminary crystallographic analysis of an acridone-producing novel multifunctional type III polyketide synthase from *Huperzia serrata*. *Acta Crystallogr* F63: 576–578.
- Morita H., Noguchi H., Schröder J. and Abe I. (2001) Novel polyketides synthesized with a higher plant stilbene synthase. *Eur J Biochem* 268: 3759–3766.
- Morita H., Takahashi Y., Noguchi H. and Abe I. (2000) Enzymatic formation of unnatural aromatic polyketides by chalcone synthase. *Biochem Biophys Res Commun* 279: 190–195.
- Morita H., Tanio M., Kondo S., Kato R., Wanibuchi K., Noguchi H., Sugio S., Abe I. and Kohno T. (2008) Crystallization and preliminary crystallographic analyses of a plant type III polyketide synthase that produces benzalacetone. *Acta Crystallogr* F64: 304–306.
- Mosbach K. and Ehrensvar U. (1966) Studies on lichen enzymes. Part I. Preparation and properties of a depside hydrolyzing esterase and of orsellinic acid decarboxylase. *Biochem Biophys Res Commun* 22: 145–150.
- Moss S.J., Martin C.J. and Wilkinson B. (2004) Loss of co-linearity by modular polyketide synthases: A mechanism for the evolution of chemical diversity. *Nat Prod Rep* 21: 575–593.
- Munro S., Thomas K.L. and Abu-Shaar M. (1993) Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 365: 61–65.
- Murashige T. and Skoog F. (1962) A revised medium for rapid growth and bioassays with tobacco cultures. *Physiol Plant* 15: 473–497.
- Musty R.E. (2004) Natural cannabinoids: Interactions and effects. *In*: The medicinal uses of cannabis and cannabinoids. Guy G.W., Whittle B.A. and Robson P.J., eds. Pharmaceutical Press, London, UK. 165–204.
- Nakatsuka A., Izumi Y. and Yamagishi M. (2003) Spatial and temporal expression of chalcone synthase and dihydroflavonol 4-reductase genes in the Asiatic hybrid lily. *Plant Sci* 165: 759–767.
- Napoli C.A., Fahy D., Wang H.Y. and Taylor L.P. (1999) *White anther*: A petunia mutant that abolishes pollen flavonol accumulation, induces male sterility, and is complemented by a chalcone synthase transgene. *Plant Physiol* 120: 615–622.

NCBI: <http://www.ncbi.nlm.gov/>

- Novak P., Krofta K. and Matousek J. (2006) Chalcone synthase homologues from *Humulus lupulus*: Some enzymatic properties and expression. *Biol Plant* 50: 48–54.
- Nurnberger T. (1999) Signal perception in plant pathogen defense. *Cell Mol Life Sci* 55: 167–182.
- O'Neill S.D., Tong Y., Sporlein B., Forkmann G. and Yoder J.I. (1990) Molecular genetic analyses of chalcone synthase in *Lycopersicon esculentum* and an anthocyanin-deficient mutant. *Mol Gen Genet* 224: 279–288.
- Office of Medicinal Cannabis, The Netherlands. Available from <http://www.cannabisbureau.nl>.
- Okada Y. and Ito K. (2001) Cloning and analyses of valerophenone synthase gene expressed specifically in lupulin gland of Hop (*H. lupulus* L.) *Biosci Biotechnol Biochem* 65: 150–155.
- Okada Y., Yamazaki Y., Suh D.Y. and Sankawa U. (2001) Bifunctional activities of valerophenone synthase in Hop (*Humulus lupulus* L.) *J Am Soc Brew Chem* 59: 163–166.
- Oliver J.M., Burg D.L., Wilson B.S., McLaughlin J.L. and Geahlen R.L. (1994) Inhibition of mast cell FcεR1-mediated signaling and effector function by the Syk-selective inhibitor, piceatannol. *J Biol Chem* 269: 29697–29703.
- Paniego N.B., Zuurbier K.W.M., Fung S.Y., Van der Heijden R., Scheffer J.J.C. and Verpoorte R. (1999) Phlorisovalerophenone synthase, a novel polyketide synthase from hop (*Humulus lupulus* L.) cones. *Eur J Biochem* 262: 612–616.
- Paris M., Boucher F. and Cosson L. (1975) The constituents of *Cannabis sativa* pollen. *Econ Bot* 29: 245–253.
- Pasqua G., Avato P., Monacelli B., Santamaria A.R. and Argentieri M.P. (2003) Metabolites in cell suspension cultures, calli and *in vitro* regenerated organs of *Hypericum perforatum* cv. Topas. *Plant Sci* 165: 977–982.
- Pastori G.M. and Del Rio L.A. (1997) Natural senescence of pea leaves: An activated oxygen-mediated function for peroxisomes. *Plant Physiol* 114: 411–418.
- Pate D.W. (1999) The phytochemistry of Cannabis: Its ecological and evolutionary implications. *In: Advances in hemp research*. Ranalli P., ed. Haworth Press, NY. 21–42.
- Paton W.D.M. and Pertwee R.G. (1973) The actions of Cannabis in man. *In: Marijuana: Chemistry, pharmacology, metabolism and clinical effects*. Mechoulam R., ed. Academic Press, NY. 287–333.

- Pedapudi S., Chin C.K. and Pedersen H. (2000) Production and elicitation of benzalacetone and the raspberry ketone in cell suspension cultures of *Rubus idaeus*. *Biotechnol Prog* 16: 346–349.
- Pellati F. and Benvenuti S. (2007) Fast high-performance liquid chromatography analyses of phenethylamine alkaloids in *Citrus* natural products on a pentafluorophenylpropyl stationary phase. *J Chromatogr A* 1165: 58–66.
- Peterson G.L. (1977) A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal Biochem* 83: 346–356.
- Petri G., Oroszlan P. and Fridvalszky L. (1988) Histochemical detection of hemp trichomes and their correlation with the THC content. *Acta Biol Hung* 39: 59–74.
- Pettersson G. (1965) An orsellinic acid decarboxylase isolated from *Gliocladium roseum*. *Acta Chem Scand* 19: 2013–2021.
- Pfeifer V., Nicholson G.J., Ries J., Recktenwald J., Schefer A.B., Shawky R.M., Schröder J., Wohlleben W. and Pelzer S. (2001) A polyketide synthase in glycopeptide biosynthesis: The biosynthesis of the non-proteinogenic amino acid (S)-3,5-dihydroxyphenylglycine. *J Biol Chem* 276: 38370–38377.
- Ponchet M., Martin-Tanguy J., Marais A. and Martin C. (1982) Hydroxycinnamoyl acid amides and aromatic amines in the inflorescences of some *Araceae* species. *Phytochemistry* 21: 2865–2869.
- Potter D. (2004) Growth and morphology of medicinal cannabis. *In: The medicinal uses of cannabis and cannabinoids*. Guy G.W., Whittle B.A. and Robson P.J., eds. Pharmaceutical Press, London, UK. 17–54.
- Preisig-Müller R., Gnau P. and Kindl H. (1995) The inducible 9,10-dihydrophenanthrene pathway: Characterization and expression of bibenzyl synthase and S-adenosylhomocysteine hydrolase. *Arch Biochem Biophys* 317: 201–207.
- Preisig-Müller R., Schwekendiek A., Brehm I., Reif H.J. and Kindl H. (1999) Characterization of a pine multigene family containing elicitor-responsive stilbene synthase genes. *Plant Mol Biol* 39: 221–229.
- Pryce R.J. (1971) Biosynthesis of lunularic acid—a dihydro-stilbene endogenous growth inhibitor of liverworts. *Phytochemistry* 10: 2679–2685.

- Pryce R.J. (1972) Metabolism of lunularic acid to a new plant stilbene by *Lunularia cruciata*. *Phytochemistry* 11: 1355–1364.
- Pryce R.J. and Linton L. (1974) Lunularic acid decarboxylase from the liverwort *Conocephalum conicum*. *Phytochemistry* 13: 2497–2501.
- Raharjo T.J. (2004) Studies of cannabinoid biosynthesis in *Cannabis sativa* L.: The polyketide synthase. Ph.D. Thesis. Leiden University, The Netherlands.
- Raharjo T.J., Chang W.T., Choi Y.H., Peltenburg–Looman A.M.G. and Verpoorte R. (2004a) Olivetol as product of a polyketide synthase in *Cannabis sativa* L. *Plant Sci* 166: 381–385.
- Raharjo T.J., Chang W.T., Verberne M.C., Peltenburg–Looman A.M.G., Linthorst H.J.M. and Verpoorte R. (2004b) Cloning and over-expression of a cDNA encoding a polyketide synthase from *Cannabis sativa*. *Plant Physiol Biochem* 42: 291–297.
- Raiber S., Schröder G. and Schröder J. (1995) Molecular and enzymatic characterization of two stilbene synthases from Eastern white pine (*Pinus strobus*): A single Arg/His difference determines the activity and the pH dependence of the enzymes. *FEBS Lett* 361: 299–302.
- Raman A. (1998) The Cannabis plant: Botany, cultivation and processing for use. *In*: Cannabis: the genus *Cannabis*. Brown D.T., ed. Harwood Academic Publishers, Amsterdam. 29–54.
- Ramawat K.G. and Mathur M. (2007) Factors affecting the production of secondary metabolites. *In*: Biotechnology, secondary metabolites, plants and microbes. Ramawat K.G. and Merillo J.M., eds. Science Publishers, Enfield, NH, USA. 59–102.
- Ranganathan M. and D'Souza D.C. (2006) The acute effects of cannabinoids on memory in humans: A review. *Psychopharmacology* 188: 425–444.
- Rawlings B.J. (1999) Biosynthesis of polyketides (other than actinomycete macrolides). *Nat Prod Rep* 16: 425–484.
- Razdan R.K., Puttick A.J., Zitko B.A. and Handrick G.R. (1972) Hashish VI: Conversion of (–)- $\Delta^1(6)$ -tetrahydrocannabinol to (–)- $\Delta^1(7)$ -tetrahydrocannabinol, stability of (–)- Δ^1 - and (–)- $\Delta^1(6)$ -tetrahydrocannabinols. *Experientia* 28: 121–122.
- Reinecke T. and Kindl H. (1994a) Characterization of bibenzyl synthase catalyzing the biosynthesis of phytoalexins of orchids. *Phytochemistry* 35: 63–66.

- Reinecke T. and Kindl H. (1994b) Inducible enzymes of the 9,10-dihydro-phenanthrene pathway: Sterile orchid plants responding to fungal infection. *Mol Plant Microbe Interact* 7: 449–454.
- Rhodes D. and Hanson A.D. (1993) Quaternary ammonium and tertiary sulfonium compounds in higher plants. *Annu Rev Plant Physiol Plant Mol Biol* 44: 357–384.
- Riedl W. (1954) Synthese einiger Lupulon-analoga mit abgewandeltem Acyl-rest. *Liebigs Ann Chem* 585: 38–42.
- Rix U., Fischer C., Remsing L.L. and Rohr J. (2002) Modification of post-PKS tailoring steps through combinatorial biosynthesis. *Nat Prod Rep* 19: 542–580.
- Robert N., Ferran J., Breda C., Coutos-Thevenot P., Boulay M., Buffard D. and Esnault R. (2001) Molecular characterization of the incompatible interaction of *Vitis vinifera* leaves with *Pseudomonas syringae* pv. *pisi*: Expression of genes coding for stilbene synthase and class 10 PR protein. *Eur J Plant Pathol* 107: 249–261.
- Robertson L.W., Koh S.W., Huff S.R., Malhotra R.K. and Ghosh A. (1978) Microbiological oxidation of pentyl side-chain of cannabinoids. *Experientia* 34: 1020–1022.
- Rolfs C.H. and Kindl H. (1984) Stilbene synthase and chalcone synthase: Two different constitutive enzymes in cultured cells of *Picea excelsa*. *Plant Physiol* 75: 489–492.
- Rolfs C.H., Fritzemeier K.H. and Kindl H. (1981) Cultured cells of *Arachis hypogaea* susceptible to induction of stilbene synthase (resveratrol-forming). *Plant Cell Rep* 1: 83–85.
- Ross A.B., Shepherd M.J., Schüpphaus M., Sinclair V., Alfaro B., Kamal-Eldin A. and Aman P. (2003) Alkylresorcinols in cereals and cereal products. *J Agric Food Chem* 51: 4111–4118.
- Ross S.A. and ElSohly M.A. (1995) Constituents of *Cannabis sativa* L. XXVIII a review of the natural constituents: 1980–1994. *Zagazig J Pharm Sci* 4: 1–10.
- Ross S.A. and ElSohly M.A. (1996) The volatile oil composition of fresh and air-dried buds of *Cannabis sativa*. *J Nat Prod* 59: 49–51.
- Ross S.A., ElSohly H.N., Elkashoury E.A. and ElSohly M.A. (1996) Fatty acids of Cannabis seeds. *Phytochem Anal* 7: 279–283.
- Ross S.A., ElSohly M.A., Sultana G.N.N., Mehmedic Z., Hossain C.F. and Chandra S. (2005) Flavonoid glycosides and cannabinoids from the pollen of *Cannabis sativa* L. *Phytochem Anal* 16: 45–48.

- Ross S.A., Mehmedic Z., Murphy T.P. and ElSohly M.A. (2000) GC–MS analysis of the total Δ^9 -THC content of both drug-and fiber-type cannabis seeds. *J Anal Toxicol* 24: 715–717.
- Rothschild M., Rowan M.R. and Fairbairn J.W. (1977) Storage of cannabinoids by *Arctia caja* and *Zonocerus elegans* fed on chemically distinct strains of *Cannabis sativa*. *Nature* 266: 650–651.
- Roy B. and Dutta B.K. (2003) *In vitro* lethal efficacy of leaf extract of *Cannabis sativa* on the larvae of *Chironomous samoensis* Edward: An insect of public health concern. *Indian J Exp Biol* 41: 1338–1341.
- Rozema J., Bjorn L.O., Bornman J.F., Gaberscik A., Hader D.P., Trost T., Germ M., Klisch M., Groniger A., Sinha R.P., Lebert M., He Y.Y., Buffoni–Hall R., de Bakker N.V.J., van de Staaij J. and Meijkamp B.B. (2002) The role of UV–B radiation in aquatic and terrestrial ecosystems—an experimental and functional analyses of the evolution of UV-absorbing compounds. *J Photochem Photobiol B: Biol* 66: 2–12.
- Ruhmann S., Treutter D., Fritsche S., Briviba K. and Szankowski I. (2006) Piceid (resveratrol glucoside) synthesis in stilbene synthase transgenic apple fruit. *J Agric Food Chem* 54: 4633–4640.
- Rupprich N. and Kindl H. (1978) Stilbene synthases and stilbenecarboxylate synthases, I: Enzymatic synthesis of 3,5,4'-trihydroxystilbene from *p*-coumaroyl coenzyme A and malonyl coenzyme A. *Hoppe Seylers Z Physiol Chem* 359: 165–172.
- Russo E. (2004) History of cannabis as a medicine. *In*: The medicinal uses of cannabis and cannabinoids. Guy G.W., Whittle B.A. and Robson P.J., eds. Pharmaceutical Press, London, UK. 1–16.
- Ryder T.B., Hedrick S.A., Bell J.N., Liang X., Clouse S.D. and Lamb C.J. (1987) Organization and differential activation of a gene family encoding the plant defense enzyme chalcone synthase in *Phaseolus vulgaris*. *Mol Gen Genet* 210: 219–233.
- Saeed A.I., Sharov V., White J., Li J., Liang W., Bhagabati N., Braisted J., Klapa M., Currier T., Thiagarajan M., Sturn A., Snuffin M., Rezantsev A., Popov D., Ryltsov A., Kostukovich E., Borisovsky I., Liu Z., Vinsavich A., Trush V., Quackenbush J. (2003) TM4: A free, open-source system for microarray data management and analysis. *Biotechniques* 34: 374–378.
- Sakakibara I., Ikeya Y., Hayashi K., Okada M. and Maruno M. (1995) Three acyclic *bis*-phenylpropane lignanamides from fruits of *Cannabis sativa*. *Phytochemistry* 38: 1003–1007.

- Samappito S., Page J., Schmidt J., De-Eknamkul W. and Kutchan T.M. (2003) Aromatic and pyrone polyketides synthesized by a stilbene synthase from *Rheum tataricum*. *Phytochemistry* 62: 313–323.
- Samappito S., Page J., Schmidt J., De-Eknamkul W. and Kutchan T.M. (2002) Molecular characterization of root-specific chalcone synthases from *Cassia alata*. *Planta* 216: 64–71.
- Sanchez-Sampedro A., Kim H.K., Choi Y.H., Verpoorte R. and Corchete P. (2007) Metabolomic alterations in elicitor treated *Silybum marianum* suspension cultures monitored by nuclear magnetic resonance spectroscopy. *J Biotechnol* 130: 133–142.
- Sankaranarayanan R., Saxena P., Marathe U.B., Gokhale R.S., Shanmugam V.M. and Rukmini R. (2004) A novel tunnel in mycobacterial type III polyketide synthase reveals the structural basis for generating diverse metabolites. *Nat Struct Mol Biol* 11: 894–900.
- Sankawa U. (1999) Polyketides and other secondary metabolites including fatty acids and their derivatives. *In*: Comprehensive natural products chemistry. Vol. 1. Barton D.H.R., Nakanishi K. and Meth-Cohn O., eds. Elsevier Science Ltd., Oxford, UK.
- Saxena P., Yadav G., Mohanty D. and Gokhale R. (2003) A new family of type III polyketide synthase in *Mycobacterium tuberculosis*. *J Biol Chem* 278: 44780–44790.
- Schijlen E., de Vos C.H.R., Jonker H., van den Broeck H., Molthoff J., van Tunen A., Martens S. and Bovy A. (2006) Pathway engineering for healthy phytochemicals leading to the production of novel flavonoids in tomato fruit. *Plant Biotechnol J* 4: 433–444.
- Schöppner A. and Kindl H. (1984) Purification and properties of a stilbene synthase from induced cell suspension cultures of peanut. *J Biol Chem* 259: 6806–6811.
- Schröder G. and Schröder J. (1992) A single change of histidine to glutamine alters the substrate preference of a stilbene synthase. *J Biol Chem* 267: 20558–20560.
- Schröder G., Brown J.W.S. and Schröder J. (1988) Molecular analyses of resveratrol synthase: cDNA, genomic clones and relationship with chalcone synthase. *Eur J Biochem* 172: 161–169.
- Schröder J. (1997) A family of plant-specific polyketide synthases: facts and predictions. *Trends Plant Sci* 2: 373–378.
- Schröder J. (1999) The chalcone/stilbene synthase-type family of condensing enzymes. *In*: Comprehensive natural products chemistry. Barton D.H.R., Nakanishi K. and Meth-Cohn O., eds. Vol. 1. Polyketides and other secondary metabolites including fatty acids and their derivatives. Sankawa U., ed. Elsevier Science Ltd., Oxford, UK. 749–771.

Schröder J. (2000) The family of chalcone synthase-related proteins: Functional diversity and evolution. *In: Evolution of metabolic pathways*. Vol. 34. Romeo J.T., Ibrahim R.K., Varin L. and De Luca V., eds. Pergamon Press, Amsterdam. 55–89.

Schröder J. and Schröder G. (1990) Stilbene and chalcone synthases: Related enzymes with functions in plant-specific pathways. *Z Naturforsch* 45c: 1–8.

Schröder J. Group, Freiburg University, Germany. <http://www.biologie.uni-freiburg.de/data/bio2/schroeder/stilbenecarboxylates.html>.

Schröder, J., Heller, W. and Hahlbrock, K. (1979) Flavanone synthase: simple and rapid assay for the key enzyme of flavonoid biosynthesis. *Plant Sci. Lett.* 14: 281–286.

Schröder J., Raiber S., Berger T., Schmidt A., Schmidt J., Soares-Sello A.M., Bardshiri E., Strack D., Simpson T.J., Veit M. and Schröder G. (1998) Plant polyketide synthases: A chalcone synthase-type enzyme which performs a condensation reaction with methylmalonyl-CoA in the biosynthesis of C-methylated chalcones. *Biochemistry* 37: 8417–8425.

Schultz K., Kuehne P., Häusermann U.A. and Hesse M. (1997) Absolute configuration of macrocyclic spermidine alkaloids. *Chirality* 9: 523–528.

Schütz R., Heller W. and Hahlbrock K. (1983) Substrate specificity of chalcone synthase from *Petroselinum hortense*. *J Biol Chem* 258: 6730–6734.

Segelman A.B., Segelman F.P. and Varma S. (1976) *Cannabis sativa* (marijuana) IX: Lens aldose reductase inhibitory activity of certain marijuana flavonoids. *J Nat Prod* 39: 475.

Segelman A.B., Segelman F.P., Star A.E., Wagner H. and Seligmann O. (1978) Structure of two C-diglycosylflavones from *Cannabis sativa*. *Phytochemistry* 17: 824–826.

Serazetdinova L., Oldach K.H. and Lorz H. (2005) Expression of transgenic stilbene synthases in wheat causes the accumulation of unknown stilbene derivatives with antifungal activity. *J Plant Physiol* 162: 985–1002.

Seshime Y., Juvvadi P.R., Fujii I. and Kitamoto K. (2005) Discovery of a novel superfamily of type III polyketide synthases in *Aspergillus oryzae*. *Biochem Biophys Res Commun* 331: 253–260.

Shimizu T., Akada S., Senda M., Ishikawa R., Harada T., Niizeki M. and Dube S.K. (1999) Enhanced expression and differential inducibility of soybean chalcone synthase genes by supplemental UV-B in dark-grown seedlings. *Plant Mol Biol* 39: 785–795.

- Shine W.E. and Loomis W.D. (1974) Isomerization of geraniol and geranyl phosphate by enzymes from carrot and peppermint. *Phytochemistry* 13: 2095–2101.
- Shirley B.W. (1996) Flavonoid biosynthesis: “new” functions for and a “old” pathway. *Trends Plant Sci* 1: 377–382.
- Shomura Y., Torayama I., Suh D.Y., Xiang T., Kita A., Sankawa U. and Miki K. (2005) Crystal structure of stilbene synthase from *Arachis hypogaea*. *Proteins* 60: 803–806.
- Shoyama Y. and Nishioka I. (1978) Cannabis, XIII: Two new spiro-compounds, cannabisirol and acetyl cannabisirol. *Chem Pharm Bull* 26: 3641–3646.
- Shoyama Y., Hirano H. and Nishioka I. (1984) Biosynthesis of propyl cannabinoid acid and its biosynthetic relationship with pentyl and methyl cannabinoid acids. *Phytochemistry* 23: 1909–1912.
- Shoyama Y., Hirano H., Makino H., Umekita N. and Nishioka I. (1977) Cannabis X: The isolation and structures of four new propyl cannabinoids acids, tetrahydrocannabivarinic acid, cannabidivarinic acid, cannabichromevarinic acid and cannabigerovarinic acid, from Thai cannabis, ‘Meao variant’. *Chem Pharm Bull* 25: 2306–2311.
- Shoyama Y., Takeuchi A., Taura F., Tamada T., Adachi M., Kuroki R., Shoyama Y. and Morimoto S. (2005) Crystallization of Δ^1 -tetrahydrocannabinolic acid (THCA) synthase from *Cannabis sativa*. *Acta Crystallogr* 61: 799–801.
- Shoyama Y., Yagi M. and Nishioka I. (1975) Biosynthesis of cannabinoid acids. *Phytochemistry* 14: 2189–2192.
- Sirikantaramas S., Morimoto S., Shoyama Y., Ishikawa Y., Wada Y., Shoyama Y. and Taura F. (2004) The gene controlling marijuana psychoactivity; molecular cloning and heterologous expression of Δ^1 -tetrahydrocannabinolic acid synthase from *Cannabis sativa* L. *J Biol Chem* 279: 39767–39774.
- Sirikantaramas S., Taura F., Tanaka Y., Ishikawa Y., Morimoto S. and Shoyama Y. (2005) Tetrahydrocannabinolic acid synthase, the enzyme controlling marijuana psychoactivity, is secreted into the storage cavity of the glandular trichomes. *Plant Cell Physiol* 46: 1578–1582.
- Skaltsa H., Verykokidou E., Harvala C., Karabourniotis G. and Manetas Y. (1994) UV-B protective potential and flavonoid content of leaf hairs of *Quercus ilex*. *Phytochemistry* 37: 987–990.
- Slatkin D.J., Doorenbos N.J., Harris L.S., Masoud A.N., Quimby M.W. and Schiff P.L.J. (1971) Chemical constituents of *Cannabis sativa* L. root. *J Pharm Sci* 60: 1891–1892.

- Sloley B.D., Juorio A.V., Durden D.A. (1990) High-performance liquid chromatographic analyses of monoamines and some of their γ -glutamyl conjugates produced by the brain and other tissues of *Helix aspersa*. *Cell Mol Neurobiol* 10: 175–192.
- Smith R.M. (1997) Identification of butyl cannabinoids in marijuana. *J Forensic Sci* 42: 610–618.
- Southon I.W. and Buckingham J. (1989) Dictionary of alkaloids. Vol I–II. Chapman & Hill Ltd, London.
- Springob K., Lukacin R., Ernwein C., Groning I. and Matern U. (2000) Specificities of functionally expressed chalcone and acridone synthases from *Ruta graveolens*. *Eur J Biochem* 267: 6552–6559.
- Springob K., Samappito S., Jindaprasert A., Schmidt J., Page J.E., De-Eknamkul W. and Kutchan T.M. (2007) A polyketide synthase of *Plumbago indica* that catalyzes the formation of hexaketide pyrones. *FEBS J* 274: 406–417.
- Stahl E. and Kunde R. (1973) Die leitsubstanzen der Haschisch–Suchhunde. *Kriminalistik* 9: 385–388.
- Stark–Lorenzen P., Nelke B., Hanbler G., Muhlbach H.P. and Thomzik J.E. (1997) Transfer of a grape stilbene synthase gene to rice (*Oryza sativa* L.). *Plant Cell Rep* 16: 668–673.
- Staunton J. and Weissman K.J. (2001) Polyketide biosynthesis: A millennium review. *Nat Prod Rep* 18: 380–416.
- Stivala L.A., Savio M., Carafoli F., Perucca P., Bianchi L., Magas G., Forti L., Pagnoni U.M., Albini A., Prosperi E. and Vannini V. (2001) Specific structural determinants are responsible for the antioxidant activity and the cell cycle effects of resveratrol. *J Biol Chem* 276: 22586–22594.
- Stöckigt J. and Zenk M.H. (1975) Chemical syntheses and properties of hydroxycinnamoyl–Coenzyme A derivatives. *Z Naturforsch C* 30: 352–358.
- Stratford M., Plumridge A. and Archer D.B. (2007) Decarboxylation of sorbic acid by spoilage yeasts is associated with the *PAD1* gene. *Appl Environ Microbiol* 73: 6534–6542.
- Suh D.Y., Fukuma K., Kagami J., Yamazaki Y., Shibuya M., Ebizuka Y. and Sankawa U. (2000) Identification of amino acid residues important in the cyclization reactions of chalcone and stilbene synthases. *Biochem J* 350: 229–235.

- Suh D.Y., Kagami J., Fukuma K. and Sankawa U. (2000) Evidence for catalytic cysteine–histidine dyad in chalcone synthase. *Biochem Biophys Res Commun* 275: 725–730.
- Suzuki Y., Kurano M., Esumi Y., Yamaguchi I. and Doi Y. (2003) Biosynthesis of 5-alkylresorcinol in rice: Incorporation of a putative fatty acid unit in the 5-alkylresorcinol carbon chain. *Bioorg Chem* 31: 437–452.
- Sweetlove L.J. and Fernie A.R. (2005) Regulation of metabolic networks: Understanding metabolic complexity in the systems biology era. *New Phytol* 168: 9–24.
- Szankowski I., Briviba K., Fleschhut J., Schonherr J., Jacobsen H.J. and Kiesecker J. (2003) Transformation of apple (*Malus domestica* Borkh.) with the stilbene synthase gene from grapevine (*Vitis vinifera* L.) and a *pgip* gene from kiwi (*Actinidia deliciosa*). *Plant Cell Rep* 22: 141–149.
- Tabor H., Rosenthal S.M. and Tabor C.W. (1958) The biosynthesis of spermidine and spermine from putrescine and methionine. *J Biol Chem* 233: 907–914.
- Taguchi C., Taura F., Tamada T., Shoyama Y., Shoyama Y., Tanaka H., Kuroki R. and Morimoto (2008) Crystallization and preliminary X-ray diffraction studies of polyketide synthase–1 (PKS–1) from *Cannabis sativa*. *Acta Crystallogr F* 64: 217–220.
- Tanaka H. and Shoyama Y. (1999) Monoclonal antibody against tetrahydrocannabinolic acid distinguishes *Cannabis sativa* samples from different plant species. *Forensic Sci Int* 106: 135–146.
- Tanaka H., Takahashi R., Morimoto S. and Shoyama Y. (1997) A new cannabinoid, Δ^6 -tetrahydrocannabinol 2'-O- β -D-glucopyranoside, biotransformed by plant tissue. *J Nat Prod* 60: 168–170.
- Taura F., Morimoto S. and Shoyama Y. (1995b) Cannabinerolic acid, a cannabinoid from *Cannabis sativa*. *Phytochemistry* 39: 457–458.
- Taura F., Morimoto S. and Shoyama Y. (1996) Purification and characterization of cannabidiolic acid synthase from *Cannabis sativa* L. *J Biol Chem* 271: 17411–17416.
- Taura F., Morimoto S., Shoyama Y. and Mechoulam R. (1995a) First direct evidence for the mechanism of Δ^1 -tetrahydrocannabinolic acid biosynthesis. *J Am Chem Soc* 117: 9766–9767.
- Taura F., Sirikantaramas S., Shoyama Y., Shoyama Y. and Morimoto S. (2007a) Phytocannabinoids in *Cannabis sativa*: Recent studies on biosynthetic enzymes. *Chem Biodivers* 4: 1649–1663.

- Taura F., Sirikantaramas S., Shoyama Y., Yoshikai K., Shoyama Y. and Morimoto S. (2007b) Cannabidiolic-acid synthase, the chemotype-determining enzyme in the fiber-type *Cannabis sativa*. **FEBS Lett** 581: 2929–2934.
- Taylor L.P. and Jorgensen R. (1992) Conditional male fertility in chalcone synthase-deficient petunia. **J Hered** 83: 11–17.
- Thakur G.A., Duclos R.I.Jr. and Makriyannis A. (2005) Natural cannabinoids: Templates for drug discovery. **Life Sci** 78: 454–466.
- Tropf S., Kärcher B., Schröder G. and Schröder J. (1995) Reaction mechanisms of homodimeric plant polyketide synthases (stilbene and chalcone synthase): A single active site for the condensing reaction is sufficient for synthesis of stilbenes, chalcones and 6'-deoxychalcones. **J Biol Chem** 270: 7922–7928.
- Tropf S., Lanz T., Schröder J. and Schröder G. (1994) Evidence that stilbene synthases have developed from chalcone synthases several times in the course of evolution. **J Mol Evol** 38: 610–618.
- Turner C.E. and ElSohly M.A. (1979) Constituents of *Cannabis sativa* L. XVI: A possible decomposition pathway of Δ^9 -tetrahydrocannabinol to cannabinol. **J Heterocycl Chem** 16: 1667–1668.
- Turner C.E. and Mole M.L. (1973) Chemical components of *Cannabis sativa*. **JAMA** 225: 639.
- Turner C.E., ElSohly M.A. and Boeren E.G. (1980) Constituents of *Cannabis sativa* L. XVII: A review of the natural constituents. **J Nat Prod** 43: 169–243.
- Turner J., Hemphill J. and Mahlberg P.G. (1977) Gland distribution and cannabinoid content in clones of *Cannabis sativa* L. **Am J Bot** 64: 687–693.
- Turner J., Hemphill J. and Mahlberg P.G. (1978) Quantitative determination of cannabinoids in individual glandular trichomes of *Cannabis sativa* L. (Cannabaceae). **Am J Bot** 65: 1103–1106.
- Turner J., Hemphill J. and Mahlberg P.G. (1981) Interrelationships of glandular trichomes and cannabinoid content. I: Developing pistillate bracts of *Cannabis sativa* L. (Cannabaceae). **Bull Narc** 33: 59–69.
- Uy R. and Wold F. (1977) Posttranslational covalent modification of proteins. **Science** 198: 890–896.

- Valant-Vetschera K.M. and Wollenweber E. (2006) Flavones and flavonols. *In: Flavonoids: Chemistry, biochemistry and applications*. Andersen Ø.M. and Markham K.R., eds. CRC Press–Taylor & Francis Group, Boca Raton, FL. 617–748.
- Valenzano D.R., Terzibasi E., Genade T., Cattaneo A., Domenici L. and Cellerino A. (2006) Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr Biol* 16: 296–300.
- Valio I.F.M. and Schwabe W.W. (1970) Growth and dormancy in *Lunularia cruciata* (L.) Dum. VIII: The isolation and bioassay of lunularic acid. *J Exp Bot* 21: 138–150.
- Van der Krol A., Lenting P.E., Veenstra J., van der Meer I.M., Koes R.E., Gerats A.G.M., Mol J.N.M. and Stuitje A.R. (1988) An anti-sense chalcone synthase gene in transgenic plants inhibits flower pigmentation. *Nature* 333: 866–869.
- Van Gaal L.F., Rissanen A.M., Scheen A.J., Ziegler O. and Rössner S. (2005) Effects of the cannabinoid-1 receptor blocker rimonabant on weight reduction and cardiovascular risk factors in overweight patients: 1-year experience from the RIO-Europe study. *Lancet* 365: 1389–1397.
- Vanhoenacker G., Van Rompaey P., De Keukeleire D. and Sandra P. (2002) Chemotaxonomic features associated with flavonoids of cannabinoid-free Cannabis (*Cannabis sativa* subsp. *sativa* L.) in relation to hops (*Humulus lupulus* L.) *Nat Prod Lett* 16: 57–63.
- Vastano B.C., Chen Y., Zhu N., Ho C.T., Zhou Z. and Rosen R.T. (2000) Isolation and identification of stilbenes in two varieties of *Polygonum cuspidatum*. *J Agric Food Chem* 48: 253–256.
- Velasco G., Galve-Roperh I., Sanchez C., Blazquez C., Haro A. and Guzman M. (2005) Cannabinoids and ceramide: Two lipids acting hand-by-hand. *Life Sci* 77: 1723–1731.
- Veliky I.A. and Genest K. (1972) Growth and metabolites of *Cannabis sativa* cell suspension cultures. *Lloydia* 35: 450–456.
- Vogt T., Wollenweber E. and Taylor L.P. (1995) The structural requirements of flavonols that induce pollen germination of conditionally male fertile *Petunia*. *Phytochemistry* 38: 589–592.
- Voirin B., Bayet C. and Colson M. (1993) Demonstration that flavone aglycones accumulate in the peltate glands of *Mentha x piperita* leaves. *Phytochemistry* 34: 85–87.
- Vree T.B., Breimer D.D. van Ginneken C.A.M. and van Rossum J.M. (1972) Identification in hashish of tetrahydrocannabinol, cannabidiol and cannabinol analogues with a methyl side-chain. *J Pharm Pharmacol* 24: 7–12.

- Wahby I., Arraez-Roman D., Segura-Carretero A., Ligerio F., Caba J.M. and Fernandez-Gutierrez A. (2006) Analysis of choline and atropine in hairy root cultures of *Cannabis sativa* L. by capillary electrophoresis-electrospray mass spectrometry. *Electrophoresis* 27: 2208–2215.
- Wanibuchi K., Zhang P., Abe T., Morita H., Kohno T., Chen G., Noguchi H. and Abe I. (2007) An acridone-producing novel multifunctional type III polyketide synthase from *Huperzia serrata*. *FEBS J* 274: 1073–1082.
- Ward R.S. (1999) Lignans, neolignans and related compounds. *Nat Prod Rep* 16: 75–96.
- Watanabe K., Yamaori S., Funahashi T., Kimura T. and Yamamoto I. (2007) Cytochrome P450 enzymes involved in the metabolism of tetrahydrocannabinols and cannabinol by human hepatic microsomes. *Life Sci* 80: 1415–1419.
- Watts K.T., Lee P.C. and Schmidt-Dannert C. (2004) Exploring recombinant flavonoid biosynthesis in metabolically engineered *Escherichia coli*. *Chem Biochem* 5: 500–507.
- Watts K.T., Lee P.C. and Schmidt-Dannert C. (2006) Biosynthesis of plant-specific stilbene polyketides in metabolically engineered *Escherichia coli*. *BMC Biotechnol* 6: 22–33.
- Wells L. and Hart G.H. (2003) O-GlcNAc turns twenty: Functional implications for post-translational modification of nuclear and cytosolic proteins with a sugar. *FEBS Lett* 546: 154–158.
- Werker E. (2000) Trichome diversity and development. *Adv Bot Res* 31: 1–35.
- Whitaker B.D. and Stommel J.R. (2003) Distribution of hydroxycinnamic acid conjugates in fruit of commercial eggplant (*Solanum melongena* L.) cultivars. *J Agric Food Chem* 51: 3448–3454.
- Whitehead I.M. and Dixon R.A. (1983) Chalcone synthase from cell suspension cultures of *Phaseolus vulgaris* L. *Biochem Biophys Acta* 747: 298–303.
- Whiting D.A. (2001) Natural phenolics compounds 1900–2000: A bird's eye view of a century's chemistry. *Nat Prod Rep* 18: 583–606.
- Widholm, J.M. (1972) The use of fluorescein diacetate and phenosafranine for determining viability of cultured plant cells. *Stain Technol* 47: 189–194.
- Wiese W., Vornam B., Krause E. and Kindl H. (1994) Structural organization and differential expression of three stilbene synthase genes located on a 13 kb grapevine DNA fragment. *Plant Mol Biol* 26: 667–677.

- Wilkinson B. and Micklefield J. (2007) Mining and engineering natural-product biosynthetic pathways. *Nat Chem Biol* 3: 379–386.
- Williamson E.M. and Evans F.J. (2000) Cannabinoids in clinical practice. *Drugs* 60: 1303–1314.
- Wills S. (1998) Cannabis use and abuse by man: An historical perspective. *In: Cannabis: the genus Cannabis*. Brown D.T., ed. Harwood Academic Publishers, Amsterdam. 1–27.
- Wilson I.B.H. (2002) Glycosylation of proteins in plants and invertebrates. *Curr Opin Struct Biol* 12: 569–577.
- Winkel-Shirley B. (1999) Evidence for enzyme complexes in the phenylpropanoid and flavonoid pathways. *Physiol Plant* 107: 142–149.
- Wollenweber W. (1980) The systematic implication of flavonoids secreted by plants. *In: Biology and Chemistry of plant trichomes*. Rodriguez E., Healey P.L. and Mehta I., eds. Plenum Press, New York. 53–69.
- Wu S., O’Leary S.J.B., Gleddie S., Eudes F., Laroche A. and Robert L.S. (2008) A chalcone synthase-like gene is highly expressed in the tapetum of both wheat (*Triticum aestivum* L.) and triticale (x *Triticosecale* Wiimack). *Plant Cell Rep* doi: 10.1007/s00299-008-0572-3.
- Xie D., Shao Z., Achkar J., Zha W., Frost J.W. and Zhao H. (2006) Microbial synthesis of triacetic acid lactone. *Biotechnol Bioeng* 93: 727–736.
- Yamada M., Hayashi K., Hayashi H., Ikeda S., Hoshino T., Tsutsui K., Tsutsui K., Iinuma M. and Nozaki H. (2006) Stilbenoids of *Kobresia nepalensis* (Cyperaceae) exhibiting DNA topoisomerase II inhibition. *Phytochemistry* 67: 307–313.
- Yamazaki Y., Suh D.Y., Sitthithaworn W., Ishiguro K., Kobayashi Y., Shibuya M., Ebizuka Y. and Sankawa U. (2001) Diverse chalcone synthase superfamily enzymes from the most primitive vascular plant, *Psilotum nudum*. *Planta* 214: 75–84.
- Yerger E.H., Grazzini R.A., Hesk D., Cox-Foster D.L., Craig R. and Mumma R.O. (1992) A rapid method for isolation glandular trichomes. *Plant Physiol* 99: 1–7.
- Ylstra B., Busscher J., Franken J., Hollman P.C.H., Mol J.N.M. and van Tunen A.J. (1994) Flavonols and fertilization in *Petunia hybrida*: Localization and mode of action during pollen tube growth. *Plant J* 6: 201–212.

- Yu M. and Facchini P.J. (1999) Purification, characterization and immunolocalization of hydroxycinnamoyl-CoA:tyramine *N*-(hydroxycinnamoyl) transferase from opium poppy. *Planta* 209: 33–44.
- Yusuf I., Yamaoka K., Otsuka H., Yamasaki K. and Seyama I. (1992) Block of sodium channels by tyramine and its analogue (*N*-feruloyl tyramine) in frog ventricular myocytes. *Jpn J Physiol* 42: 179–191.
- Zacares L., Lopez-Gresca M.P., Fayos J., Primo J., Belles J.M. and Conejero V. (2007) Induction of *p*-coumaroyldopamine and feruloyldopamine, two novel metabolites, in tomato by the bacterial pathogen *Pseudomonas syringae*. *Mol Plant Microbe Interact* 20: 1439–1448.
- Zha W., Rubin-Pitel S.B. and Zhao H. (2006) Characterization of the substrate specificity of PhID, a type III polyketide synthase from *Pseudomonas fluorescens*. *J Biol Chem* 281: 32036–32047.
- Zhang X. and Oppenheimer D.G. (2004) A simple and efficient method for isolation trichomes for downstream analyses. *Plant Cell Physiol* 45: 221–224.
- Zhang Y., Li S.Z., Li J., Pan X., Cahoon R.E., Jaworski J.G., Wang X., Jez J.M., Chen F. and Yu O. (2006) Using unnatural protein fusions to engineer resveratrol biosynthesis in yeast and mammalian cells. *J Am Chem Soc* 128: 13030–13031.
- Zhao J., Davis C.D. and Verpoorte R. (2005) Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol Adv* 23: 283–333.
- Zeng D. and Hrazdina G. (2008) Molecular and biochemical characterization of benzalacetone synthase and chalcone synthase genes and their proteins from raspberry (*Rubus idaeus* L.). *Arch Biochem Biophys* 470: 139–145.
- Zheng D., Schröder G., Schröder J. and Hrazdina G. (2001) Molecular and biochemical characterization of three aromatic polyketide synthase genes from *Rubus idaeus*. *Plant Mol Biol* 46: 1–15.
- Zheng X.Q., Nagai C. and Ashihara H. (2004) Pyridine nucleotide cycle and trigonelline (*N*-methylnicotinic acid) synthesis in developing leaves and fruits of *Coffea arabica*. *Physiol Plant* 122: 404–411.
- Zhou P., Watson D.G. and Midgley J.M. (1993) Identification and quantification of γ -glutamyl conjugates of biogenic amines in the nervous system of the snail, *Helix aspersa*, by gas chromatography–negative–ion chemical ionization mass spectrometry. *J Chromatogr* 617: 11–18.

- Zobayed S.M.A., Afreen F., Goto E. and Kozai T. (2006) Plant–environmental interactions: Accumulation of hypericin in dark glands of *Hypericum perforatum*. *Ann Bot* 98: 793–804.
- Zulak K.G., Cornish A., Daskalchuk T.E., Deyholos M.K., Goodenowe D.B., Gordon P.M.K., Klassen D., Pelcher L.E., Sensen C.W. and Facchini P.J. (2007) Gene transcript and metabolite profiling of elicitor-induced opium poppy cell cultures reveals the coordinated regulation of primary and secondary metabolism. *Planta* 225: 1085–1106.
- Zulak K.G., Weljie A.M., Vogel H.J. and Facchini P.J. (2008) Quantitative ¹H-NMR metabolomics reveals extensive metabolic reprogramming of primary and secondary metabolism in elicitor-treated opium poppy cell cultures. *BMC Plant Biol* 8: doi 10.1186/1471-2229-8-5. PMID: 18211706.
- Zuurbier K.W.M., Leser J., Berger T., Hofte A.J.P., Schröder G., Verpoorte R. and Schröder J. (1998) 4-hydroxy-2-pyrone formation by chalcone and stilbene synthase with nonphysiological substrates. *Phytochemistry* 49: 1945–1951.