



Universiteit  
Leiden  
The Netherlands

## The many phases of massive galaxies : a near-infrared spectroscopic study of galaxies in the early universe

Kriek, M.T.

### Citation

Kriek, M. T. (2007, September 26). *The many phases of massive galaxies : a near-infrared spectroscopic study of galaxies in the early universe*. Retrieved from <https://hdl.handle.net/1887/12353>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12353>

**Note:** To cite this publication please use the final published version (if applicable).

# **The Many Phases of Massive Galaxies**

A Near-Infrared Spectroscopic Study of Galaxies in the Early Universe



# The Many Phases of Massive Galaxies

A Near-Infrared Spectroscopic Study of Galaxies in the Early Universe

Proefschrift

ter verkrijging van  
de graad van Doctor aan de Universiteit Leiden,  
op gezag van Rector Magnificus prof. mr. P. F. van der Heijden,  
volgens besluit van het College voor Promoties  
te verdedigen op woensdag 26 september 2007  
klokke 16.15 uur

door

Mariska Therese Kriek

geboren te Leiden  
in 1979

Promotiecommissie

Promotores: Prof. dr. M. Franx  
Prof. dr. P. G. van Dokkum (Yale University, USA)

Referent: Prof. dr. P. Barthel (Rijksuniversiteit Groningen)

Overige leden: Prof. dr. P. T. de Zeeuw  
Prof. dr. G. D. Illingworth (University of California, Santa Cruz, USA)  
Prof. dr. K. Kuijken  
Dr. P. P. van der Werf





# Table of contents

	Page
<b>1 Introduction</b>	<b>1</b>
1.1 Structure Formation . . . . .	1
1.2 Galaxy Properties . . . . .	1
1.3 Massive Galaxies over Cosmic Time . . . . .	2
1.4 This Thesis . . . . .	4
1.5 Conclusions and Outlook . . . . .	6
<b>2 Stellar Continua and Balmer/4000Å Breaks of Red <math>z &gt; 2</math> Galaxies</b>	<b>9</b>
2.1 Introduction . . . . .	10
2.2 Data . . . . .	11
2.2.1 Target Selection and Photometry . . . . .	11
2.2.2 Observations . . . . .	12
2.2.3 Reduction of GNIRS Spectra . . . . .	13
2.2.4 Extraction of One-dimensional Spectra . . . . .	14
2.3 Spectral Modeling . . . . .	16
2.3.1 Fitting Procedure and Results . . . . .	17
2.3.2 Continuum Redshifts . . . . .	19
2.3.3 Comparison to Broadband SEDs . . . . .	20
2.3.4 Discussion of Modeling Results . . . . .	20
2.4 Spectral Diagnostics . . . . .	21
2.4.1 Break Indices . . . . .	21
2.4.2 Comparison to $H\alpha$ Equivalent Width . . . . .	22
2.4.3 Active Galactic Nuclei? . . . . .	23
2.5 Comparison to Other Galaxies . . . . .	24
2.5.1 Comparison to Other High-Redshift Galaxies . . . . .	24
2.5.2 Comparison with Low-Redshift Galaxies . . . . .	25
2.6 Summary and Conclusions . . . . .	26
<b>3 Massive Galaxies at <math>z \sim 2.3</math> with Strongly Suppressed Star Formation</b>	<b>29</b>
3.1 Introduction . . . . .	30
3.2 Galaxy Sample and Data . . . . .	30
3.3 Suppressed Star Formation . . . . .	32
3.4 Discussion . . . . .	34
<b>4 The Origin of Line Emission in Massive <math>z \sim 2.3</math> Galaxies</b>	<b>37</b>
4.1 Introduction . . . . .	38



4.2	Data . . . . .	39
4.2.1	Sample . . . . .	39
4.2.2	GNIRS Spectra . . . . .	40
4.2.3	SINFONI Spectra . . . . .	41
4.2.4	Extraction of One-Dimensional Spectra . . . . .	42
4.2.5	Line Measurements . . . . .	44
4.2.6	Line Maps . . . . .	46
4.3	Origin of the Line Emission . . . . .	47
4.3.1	Emission Line Ratios . . . . .	47
4.3.2	Other Diagnostics of AGN Activity . . . . .	49
4.3.3	Summary and AGN Fraction . . . . .	52
4.4	Implications . . . . .	53
4.4.1	Stellar Populations of AGN Host Galaxies . . . . .	53
4.4.2	Stellar Masses of AGN Host Galaxies . . . . .	55
4.4.3	Downsizing of AGN Host Galaxies . . . . .	56
4.4.4	AGNs and the Suppression of Star Formation . . . . .	58
4.5	Summary and Conclusions . . . . .	59
<b>5</b>	<b>A Near-Infrared Spectroscopic Survey of <math>K</math>-selected Galaxies at <math>z \sim 2.3</math></b>	<b>65</b>
5.1	Introduction . . . . .	66
5.2	Data . . . . .	67
5.2.1	Sample Selection . . . . .	67
5.2.2	NIR spectra . . . . .	70
5.3	Spectroscopic versus Photometric Redshifts . . . . .	75
5.3.1	Spectroscopic Redshifts and Galaxy Properties . . . . .	75
5.3.2	Photometric Redshifts and Galaxy Properties . . . . .	79
5.3.3	Direct Comparison . . . . .	79
5.3.4	Systematics and Catastrophic Failures . . . . .	80
5.3.5	An Empirically-Motivated Template Set . . . . .	83
5.4	Implications . . . . .	84
5.4.1	Implications for Derived Properties . . . . .	84
5.4.2	Implications for Photometric Samples . . . . .	87
5.4.3	Implications for Previous Studies . . . . .	91
5.5	Summary . . . . .	92
<b>6</b>	<b>The Detection of a Red Sequence at <math>z \sim 2.3</math> and its Evolution to <math>z \sim 0</math></b>	<b>97</b>
6.1	Introduction . . . . .	98
6.2	Data . . . . .	99
6.3	A Red Sequence at $z \sim 2.3$ . . . . .	99
6.3.1	The Detection of the Red Sequence at $z \sim 2.3$ . . . . .	99
6.3.2	Properties of Red-Sequence Galaxies . . . . .	101
6.4	Measuring the Evolution of the Red Sequence . . . . .	104
6.4.1	Spectroscopic Samples at Lower Redshifts . . . . .	104
6.4.2	The Color Evolution of the Red Sequence . . . . .	104
6.4.3	Evolution of the Mass and Number Density . . . . .	108

---

6.5	Modeling the Evolution of the Red Sequence . . . . .	109
6.5.1	Aging of Stellar populations . . . . .	109
6.5.2	Growth of the Red Sequence . . . . .	109
6.5.3	Red Mergers . . . . .	111
6.5.4	Other Influences . . . . .	112
6.6	Summary and Conclusions . . . . .	112
	<b>Nederlandse Samenvatting</b>	<b>115</b>
	<b>Curriculum Vitae</b>	<b>121</b>
	<b>Nawoord</b>	<b>123</b>

