



Universiteit
Leiden
The Netherlands

Pyrrolizidine alkaloid composition in the plant and its interaction with the soil microbial community

Joosten, L.

Citation

Joosten, L. (2012, September 20). *Pyrrolizidine alkaloid composition in the plant and its interaction with the soil microbial community*. Retrieved from <https://hdl.handle.net/1887/19839>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/19839>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/19839> holds various files of this Leiden University dissertation.

Author: Joosten, Lotte

Title: Pyrrolizidine alkaloid composition of the plant and its interaction with the soil microbial community

Issue Date: 2012-09-20

The Analysis of Pyrrolizidine Alkaloids in *Jacobaea vulgaris*; a Comparison of Extraction and Detection Methods

Lotte Joosten, Patrick P. J. Mulder, Klaas Vrieling, Johannes A. van Veen and Peter G. L. Klinkhamer

Phytochemical Analysis (2010) 21: 197-204

Abstract

Pyrrolizidine alkaloids (PAs) serve an important function in plant defence. In this study different extraction methods and detection techniques were compared, namely gas chromatography with nitrogen phosphorus detection (GC-NPD) and liquid chromatography tandem mass spectrometry (LC-MS/MS) with quadrupole analysers, for analysing PAs in *Jacobaea vulgaris*.

Both formic acid and sulfuric acid were tested for PA extraction from dry plant material. For GC-NPD, reduction is required to transform PA *N*-oxides into tertiary amines. Zinc and sodium metabisulfite were compared as reducing agents.

The lowest PA concentration measured with GC-NPD was approximately 0.03 mg/g and with LC-MS/MS 0.002 mg/g. The detection of major PAs by both techniques was comparable but a number of minor PAs were not detected by GC-NPD. With the LC-MS/MS procedure higher concentrations were found in plant extracts, indicating that losses may have occurred during the sample preparation for the GC-NPD method. Zinc proved a more effective reducing agent than sodium metabisulfite. The sample preparation procedure for LC-MS/MS analysis using formic acid extraction without any reduction and purification steps is far less complex and less time consuming compared to GC-NPD analysis with sulfuric acid extraction and PA *N*-oxide reduction with zinc and purification.

In terms of sensitivity and discrimination, formic acid extraction in combination with LC-MS/MS detection is the method of choice for analysing PAs (both free and *N*-oxide forms) in plant material.



Introduction

Pyrrolizidine alkaloids (PAs) are regarded as chemical defence substances against herbivores (Bentley et al. 1984; Dreyer et al. 1985; Vrieling et al. 1991; Leiss et al. 2009) and pathogenic fungi (Hol and van Veen 2002). Plants containing PAs such as *Jacobaea vulgaris* (also known as *Senecio jacobaea*) are an important food source for numerous insect species (Harper and Wood 1957). *Jacobaea vulgaris* originates from Europe and Asia (Cameron 1935; Harper and Wood 1957) and was introduced into North America, New Zealand and Australia (Pemberton and Turner 1990). In the introduced areas it is considered a pest species as it causes extensive livestock losses and is a potential contaminant in milk and honey. When parts of these plants are ingested in substantial quantities, PAs can cause acute and chronic liver damage in mammals including humans. It is therefore of great importance to develop analytical methods including sample preparation, extraction and detection techniques to get a better insight in the physiology, ecology and toxicology of PA containing plants. The mode of action in invertebrate species is not well known, but Frei et al. (1992) showed that PAs caused DNA damage. In general PAs deter generalist herbivores (van Dam 1995; Macel et al. 2005); however several specialist herbivores have adapted to PAs in their diet. In fact some specialists even prefer plants containing PAs (Boppré 1986; Hartmann and Witte 1995; Macel and Klinkhamer 2010).

PAs occur in free (tertiary amine) or *N*-oxide forms. Until now it has been generally assumed that in plants PAs are mainly present as *N*-oxide except in some seeds. However, recently we found that some PAs of *J. vulgaris* also occur in substantial amounts as tertiary amine (Chapter 4). This is of importance for understanding the role of PAs as plant defence compounds, as the tertiary PA form is known for its strong negative influence on several plant attackers. The *N*-oxide form is considered to be non-toxic (Lindigkeit et al. 1997).

For many years, PAs were typically isolated by acid-base extraction in combination with zinc reduction (Hartmann and Zimmer 1986; Witte et al. 1993). Gas chromatography (GC) with flame ionisation detection (FID) or nitrogen phosphorus detection (NPD) have been typically used as analytical methods (Hartmann and Toppel 1987; Witte et al. 1992; Hartmann and Dierich 1998; Kowalchuk et al. 2006). Over the years several other PA extraction methods (Pieters et al. 1989; Betteridge and Colegate 2005; Joosten et al. 2009; Jiang et al. 2009) and techniques for analysing PAs have been applied such as spectrophotometry (Mattocks 1967; de Boer 1999), TLC, HPLC-UV, GC-MS (Hartmann and Toppel 1987; Witte et al. 1992; Hartmann and Dierich 1998) and NMR (Pieters et al. 1989; Leiss et al. 2009). Recently liquid chromatography–tandem mass spectrometry (LC-MS/MS) has been introduced for measuring PAs in plant material (Wuilloud et al. 2004; Betteridge and Colegate 2005; Zhang et al. 2008; Jiang et al. 2009; Joosten et al. 2009). Unlike GC-related methods, LC-MS/MS and NMR can detect both tertiary amines and *N*-oxides without the time-consuming and tedious reducing step.

The aim of this study was to compare two different methods for the analysis of PAs; the traditional method consisting of sulfuric acid extraction, reduction of PA *N*-oxide with zinc and then purification by liquid-liquid extraction followed by GC-NPD analysis. The other LC-MS/MS method involves a formic acid extraction without reduction and further purification steps. In addition we also compared some critical steps involved in the two methods (Figure 1). For example we investigated whether, irrespective of the sample preparation procedure, detection of (reduced) PAs with GC-NPD and LC-MS/MS (procedure 4, 6, 8 vs 5, 7, 9) give similar results with respect to the concentration and composition in plant material. Furthermore, we tested the efficiency of formic acid extraction as an alternative to sulfuric acid extraction (procedures 6, 7 vs 8, 9). We also evaluated another reducing agent, sodium metabisulfite ($\text{Na}_2\text{S}_2\text{O}_5$) as an alternative to the commonly used zinc dust (procedures 2 vs 3 and 4, 5 vs 6, 7) for GC-based analysis.

Finally we investigated whether the traditional and the novel analytical methods give similar results with regards to PA concentration and composition in plant material (procedure 1 vs 8).




Figure 1. Extraction and detection of PAs from dried plant material and the different procedures applied. Dry plant material was extracted with formic acid (1-7) or sulfuric acid (8-9). The formic acid extract was either directly measured on the LC-MS/MS (1) or reduced with sodium metabisulfite (2, 4, 5) or reduced with zinc (3, 6, 7). The sulfuric acid extract was only reduced with zinc (8-9). The reduced extracts were also analysed directly by LC-MS/MS before liquid-liquid purification (2-3). Procedure 8 is the traditional method and procedure 1 the relatively novel method.

Material and Methods

Plant material samples

Five dried *J. vulgaris* samples were used for PA analysis. One of the samples was used as a reference as it contained a mixture of several different *J. vulgaris* plants (aerial parts) collected in Meijndel (the Netherlands) in July, 1997. This sample had been used in the past for identification of PAs detected by GC-NPD. The other four samples (two roots and two shoots) were made from a genotype that originated from a dune population in Meijndel in our tissue culture laboratory. Two plants were propagated from the genotype by tissue culture and grown for six weeks in a climate room (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C). After 6 weeks the two plants were harvested and cut above the root crown by scissors. The

plants were divided into roots (R1 and R2) and shoots (S1 and S2) and freeze-dried for 72 h under vacuum with a collector temperature of -55°C (Labconco Free Zone® 12 I Freeze Dry System, Kansas City, MO, USA).

Pyrrolizidine alkaloid analysis

Formic acid extraction (procedures 1-7)

Ten milligrams of dried and ground plant material was extracted with 2% formic acid in a 1.2:100 w/v ratio. The plant extract solution was shaken for 1 h. Solid plant material was removed by centrifugation at 2600 rpm for 10 min. Heliotrine (100 µg/mL in methanol) was added as an internal standard to a concentration of 1 µg/mL. An aliquot of the extract (25 µL) was diluted with water (975 µL) and injected in the LC-MS/MS system (procedure 1 in Figure 1).

Sulfuric acid extraction with zinc reduction (procedures 8 and 9)

Ten milligrams of dried and ground plant materials was extracted with 0.25 M sulfuric acid (H₂SO₄) in a 1.5:100 w/v volume ratio. The plant extract solution was shaken for 1 h and subsequently the plant material was removed by centrifugation at 2600 rpm for 10 min. Zinc dust was added to the extraction solution and shaken for 2 h to reduce the PA *N*-oxides. Sulfuric acid extraction and zinc reduction (procedures 8 and 9 in Figure 1) were performed following the 'standard' extraction method of Hartmann and Zimmer (1986) and adapted for small quantities by de Boer (1999) for GC analysis.

Sodium metabisulfite reduction (procedures 2, 4 and 5)

One molar Sodium metabisulfite was added to the formic acid extract solution in a 1:80 v/v ratio and shaken for 2 h to reduce the PAs *N*-oxides. At this stage an aliquot was taken for LC-MS/MS analysis (25 µL), diluted with water (975 µL) and heliotrine (100 µg/mL in methanol) was added as an internal standard to a concentration of 1 µg/mL (procedure 2 in Figure 1).

Zinc reduction (procedures 3, 6 and 7)

A small "knife tip" of zinc dust was added to the formic acid extract solution and shaken for 2 h to reduce the PAs *N*-oxides. If during shaking no visible hydrogen bubbles were formed anymore, additional zinc was added. At this stage an aliquot (25 µL) was diluted with water (975 µL) and after addition of heliotrine (100 µg/mL in methanol) as an internal standard to a concentration of 1 µg/mL (procedure 3 in Figure 1), it was analysed by LC-MS/MS.

Purification by liquid-liquid extraction (procedures 4-9)

To the reduced extract, 25% ammonium hydroxide was added to reach a pH 9-10. The extract was purified by applying it over a 0.60 x 30 mm Extrelut® NT 1 column (Merck, Darmstadt, Germany). The column was eluted with dichloromethane. After elution the dichloromethane fraction was evaporated overnight and the residue was redissolved in 150 µL methanol (MeOH) containing 0.5 mg/mL heliotrine as internal standard.

Analysis by LC-MS/MS (procedures 1-3, 5, 7 and 9)

The aqueous extracts obtained from procedures 1-3 were analysed. From the methanolic extracts obtained after liquid-liquid purification (procedures 5, 7 and 9 in Figure 1) aliquots (10 µL) were diluted with water (990 µL) and injected into the LC-MS/MS system. The LC-MS/MS system consisted of an Agilent HP1100 HPLC equipped with a binary pump system, an autoinjector and a column oven, coupled to a Waters Micromass Micro tandem mass spectrometer.

Chromatographic separation was achieved on a Waters Xbridge 150 x 3.0 mm HPLC column, run with a water-acetonitrile linear gradient containing 0.05% ammonia (pH 11 ± 0.5) at a flow rate of 0.4 mL/min. The gradient started at 100% water (2 min) and during analysis the acetonitrile percentage was raised

to 70% in 16 min. The column was kept at 40°C and the injection volume was 5 µL.

An electrospray ion source of the quadrupole mass analyser was used in positive mode, with the following instrument settings: capillary voltage, 2.7 kV; source temperature, 100°C; cone gas (N₂) flow, 50 L/h; desolvation gas (N₂) flow, 600 L/h; desolvation temperature, 400°C. Argon was used as collision gas at a pressure of 3.0 x 10⁻³ mbar. Data were recorded in multiple monitoring mode (MRM) using one selected precursor to product ion transition per compound. Based on the retention times of the individual PAs the transitions could be combined into three MRM windows. One window contained the early eluting PAs, and a second window the late eluting PAs. The PAs with precursor mass *m/z* 352 were combined into a third window, because they overlapped with the other two windows. In this way the number of simultaneously monitored transitions (channels) could be limited to 10 (dwell time 120 ms/transition) (Figure 2). Cone and collision energy settings were optimised for the individual compounds (Table 1). Obtained peak areas were internally calibrated using the internal standard heliotrine and the individual compounds were quantified against a standard solution of the PAs in water. Fourteen individual PA standards were available for this study, representing over 90% of the total amount of PAs present in the plant extracts. The remaining PAs, being tertiary amines as well as *N*-oxides, were quantified by using the mean response of the tertiary amine standards and the *N*-oxide standards, respectively. Data processing was conducted with Masslynx 4.0 software (Waters, Milford, MA, USA).

Analysis by GC-NPD (procedures 4, 6, 8)

The methanol extracts of procedure 3, 6 and 8 (Figure 1) were determined with GC-NPD (Hewlett Packard 6890 and a 30 m x 0.25 µm, HP-1) under the following conditions: injector 250°C, temperature programmed from 220°C (3 min) to 250 at 5°C/min, in the split mode (1:20), carrier gas flow (N₂) 0.9 mL/min, pressure 560 mbar, detector NPD. The injection volume was 1 µL. Quantitative analysis were performed via the NPD signals using heliotrine as internal standard.

Data processing was conducted with Microsoft Office Excel 2003 for Windows (Microsoft Corporation, Redmond, WA, USA). For identification of the tertiary PAs, retention times were compared with a reference sample (Table 2). The reference sample ('M') contained a mix of dried shoot material of several different *J. vulgaris* plants collected in Meijendel.

Data analysis

Data were all normally distributed. The correlation between the PA concentration measured by GC-NPD and LC-MS/MS was tested by the Pearson correlation coefficient with SPSS 15.0 for Windows (SPSS Inc., Chicago, IL, USA). Regression was based on 15 data points (5 samples x 3 different extraction methods). The regression coefficient was estimated by linear regression model II (Sokal and Rohlf 1981; de Jong and Klinkhamer 2005). We used 'two-factor ANOVA without replication' to analyse if the different methods and techniques did have a significant influence on the estimates of the total PA concentration and the estimates of the concentration of different major PAs. Regression and 'two-factor ANOVA without replication' were conducted with Microsoft Office Excel 2003 for Windows (Microsoft Corporation, Redmond, WA, USA).

Table 1. MS/MS fragmentation conditions and LC retention times for PA identified in *Jacobaea vulgaris* extracts. The mean PA concentration determined with the LC-MS/MS method.

Pyrrolizidine alkaloid	Abbreviation	Precursor to product ion transition (m/z)	Cone voltage (V)	Collision Energy (V)	Retention time (min)	Mean PA concentration (µg/g)	Occurrence in samples
Heliotrine	Hel	314.2 > 138.0	30	25	14.51		
Seneciphylline	Sp	334.2 > 120.0	40	30	15.35	56.3	5/5
Seneciphylline <i>N</i> -oxide	Spox	350.2 > 120.0	40	30	11.96	398.2	5/5
Spartoidine*	St	334.2 > 120.0	40	30	15.10	1.1	2/5
Integerrimine	Ir	336.2 > 120.0	40	30	16.00	13.2	5/5
Integerrimine <i>N</i> -oxide	Irox	352.2 > 120.0	40	30	12.52	153.6	5/5
Senecionine	Sn	336.2 > 120.0	40	30	16.24	90.9	5/5
Senecionine <i>N</i> -oxide	Snox	352.2 > 120.0	40	30	12.69	923.7	5/5
Jacozine*	Jz	350.2 > 94.0	40	40	13.19	7.6	5/5
Erucifoline*	Er	350.2 > 94.0	40	40	13.39	12.8	5/5
Erucifoline <i>N</i> -oxide*	Erox	366.2 > 94.0	40	40	9.97	33.3	5/5
Riddelliine	Rd	350.2 > 94.0	40	40	13.78	1.9	5/5
Riddelliine <i>N</i> -oxide	Rdox	366.2 > 94.0	40	40	10.83	21.4	5/5
Jacobine	Jb	352.2 > 120.0	40	30	13.81	482.5	5/5
Jacobine <i>N</i> -oxide	Jbox	368.2 > 94.0	40	40	10.88	600.7	5/5
Retrorsine	Rt	352.2 > 120.0	40	30	14.44	1.4	3/5
Retrorsine <i>N</i> -oxide	Rtox	368.2 > 94.0	40	40	11.53	3.15	2/5
Jacoline*	Jl	370.2 > 120.0	40	30	11.50	21.1	5/5
Jacoline <i>N</i> -oxide*	Jlox	386 > 94.0	40	40	9.20	9.6	5/5
Acetyl-seneciphylline	Acsp	376.2 > 120.0	40	30	18.54	6.0	5/5
Acetyl-seneciphylline <i>N</i> -oxide	Acspox	392.2 > 120.0	40	30	14.43	6.7	5/5
Jaconine*	Jn	388.2 > 120.0	40	30	14.82	20.0	5/5
Jaconine <i>N</i> -oxide*	Jnox	404.2 > 94.0	40	40	11.14	5.3	3/5
Acetyl-erucifoline*	Acer	392.2 > 120.0	40	40	16.65	8.1	5/5
Acetyl-erucifoline <i>N</i> -oxide*	Acerox	408.2 > 94.0	40	40	13.05	38.9	5/5

Table 2. GC-NPD retention time of individual PAs and the mean PA concentration (tertiary amines + reduced *N*-oxides) in *Jacobaea vulgaris* extracts, determined with the traditional method.

Pyrrolizidine alkaloid	Abbreviation	Retention time (min)	Mean PA concentration (µg/g)	Occurrence in samples
Heliotrine internal standard	Hel	5.4		
Senecionine	Sn	7.3	485.9	5/5
Seneciphylline	Sp	7.6	282.1	5/5
Integerrimine	Int	8.1	92.1	5/5
Jacobine	Jb	9.5	433.4	5/5
Jacozine	Jz	9.9	23.9 ^a	2/5
Jacoline	Jl	10.4	55.2 ^a	1/5
Erucifoline	Er	11.0	8.7 ^a	2/5

^a Concentration may be an underestimate, because in some samples the amount could not be quantified.




Figure 2. LC-MS/MS chromatograms of pyrrolizidine alkaloids in a *Jacobaea vulgaris* shoot extract. The various precursor to product ion transitions (channels) have been combined in this figure into three MRM windows. Mass spectrometer settings and abbreviations of individual PAs are as explained in Table 1. Some of the minor PAs are not visible in the combined MRM chromatograms. These PAs can be seen when the individual transitions are plotted (data not shown)

Results and Discussion

Detection of PAs by GC-NPD and LC-MS/MS

The GC-NPD and LC-MS/MS detection techniques for the determination of PAs in plant material were compared by the analysis of a number of extracts of dried *J. vulgaris* (procedures 4, 6, 8 vs 5, 7, 9). In this study the lowest PA concentration in dry plant material that could be quantified with GC-NPD was approximately 30 µg/g and with LC-MS/MS around 2 µg/g. The introduction of a tandem mass spectrometer as a detector in combination with liquid chromatographic separation thus greatly improves the possibilities of determination of individual PAs by lowering the detection limits (Betteridge and Colegate 2005; Wuilloud et al. 2004). It should be noted that, for LC-MS/MS analysis, the plant extracts had to be diluted 40 times, to keep the peak responses of the most abundant PAs within the linear dynamic range of the mass spectrometer, while for GC-NPD analysis the final extracts had to be concentrated approximately four times. As a result LC-MS/MS detected up to 13 different tertiary PAs in *J. vulgaris* extracts while GC-NPD detected only 7 PAs (Tables 1 and 2). With GC-NPD only four major PAs (senecionine, seneciophylline, integerrimine and jacobine) were detected in all samples (Figure 3). The concentrations of the other three PAs were sometimes just below the quantitation limit of the GC-NPD (Table 2).




Figure 3. The chemical structures of tertiary amine and N-oxide forms of four major PAs present in the *Jacobaea vulgaris* extracts

Omission of the reduction step resulted in the additional detection of 11 different PA N-oxides by LC-MS/MS. Thus the number of PAs detected is almost doubled. It is a major advantage of LC-MS/MS that it can determine both N-oxides and tertiary amines directly, without the necessity of reduction of N-oxides to the corresponding tertiary amines, as is required for GC-based methods.

GC-NPD and LC-MS/MS measurements of the reduced extracts (procedures 4-9) were found to be highly correlated with respect to the total PA concentration ($r^2=0.98$) as well as the four major individual PA concentrations separated (Figure 4).




Figure 4. Relationship between the concentrations of jacobine ($y=0.995x+0.006$ $r^2=0.996$ $p<0.01$ $n=15$), senecionine ($y=1.146x-0.008$ $r^2=0.972$ $p<0.01$ $n=15$), seneciophylline ($y=1.057x-0.009$ $r^2=0.955$ $p<0.01$ $n=15$) and integerrimine ($y=0.854x+0.007$ $r^2=0.904$ $p<0.01$ $n=15$) concentration (mg/g dry plant material) in samples analysed by GC-NPD and LC-MS/MS. Fifteen different points represent five samples (■ = Root 1, ◆ = Shoot 1, ▲ = Root 2, ● = Shoot 2 and + = reference sample) and three extraction/reduction methods (black = 4 and 5, grey = 6 and 7 and white = 8 and 9; Figure 1)

Formic acid versus sulfuric acid as extraction solvent

Formic acid is often used in analytical methods with LC-MS detection. Formic acid forms volatile salts when combined with a mobile phase that contains ammonia as modifier. Sulfuric acid, on the other hand, will form non-volatile salts that can precipitate in the source of the mass detector, and in the long run will affect sensitivity and reproducibility. For this reason direct injection of (diluted) sulfuric acid extracts into the LC-MS/MS system is not recommended. The formic acid and sulfuric acid extractions were compared (procedure 6 vs 8 and 7 vs 9 in Figure 1 and Figure 5). Exchange of sulfuric acid as extraction solvent by formic acid did not lead to significant differences in concentration of the total PAs or the individual major PAs when measured with GC-NPD or LC-MS/MS.

Zinc versus sodium metabisulfite as reduction method

Reduction of PA *N*-oxides with zinc dust is commonly used when the analysis is performed by GC-based methods. Hartmann and Toppel (1987) described the use of sodium dithionite ($\text{Na}_2\text{S}_2\text{O}_4$) in sulfuric acid as alternative to zinc as reducing agent. We observed that reduction with sodium dithionite in combination with formic acid proceeded very slow. As an alternative we tested sodium metabisulfite ($\text{Na}_2\text{S}_2\text{O}_5$). A 10 mM solution of sodium metabisulfite in dilute formic acid efficiently reduced PA *N*-oxides. To compare sodium metabisulfite reduction with zinc, the total PA concentration of formic acid extraction before purification was measured with LC-MS/MS (procedure 2 vs 3 in Figure 1). Zinc reduction gave a higher total PA concentration compared with sodium metabisulfite, respectively 1.96 and 1.43 mg/g dry plant material ($F_{1,4} = 2.1, p = 0.224$), but this was not significant due to the high levels of jacobine in the two shoot samples (Figure 6a). For jacobine no significant difference was found between the two reducing agents ($F_{1,4} = 0.7, p = 0.451$). The concentrations of other major PAs nearly doubled when zinc was used instead of sodium metabisulfite, respectively for senecionine 1.9 and 1.2 mg/g ($F_{1,4} = 11.7, p < 0.05$), seneciophylline 0.4 and 0.2 mg/g ($F_{1,4} = 16.4, p < 0.05$) and integerrimine 0.15 and 0.08 mg/g ($F_{1,4} = 22.6, p < 0.01$) dry plant material. A similar result was obtained when the total PA concentration among the three different extraction/reduction combinations (procedures 4-5, 6-7 and 8-9) was compared (Figure 5). Formic acid extraction in combination with sodium metabisulfite reduction resulted in the lowest mean total PA concentration, respectively 0.78 and 0.81 mg/g dry plant material according to GC-NPD and LC-MS/MS. Sulfuric acid extraction with zinc reduction produced the highest mean total PA concentration, 1.29 and 1.42 mg/g dry plant material, respectively. It can be concluded that on the whole zinc dust is somewhat more efficient in transforming PA *N*-oxides to the corresponding tertiary amines.

Comparison between GC-NPD and LC-MS/MS method

Sulfuric acid extraction in combination with zinc reduction of the *N*-oxides followed by liquid-liquid purification and GC-NPD analysis can be regarded as the traditional method of PA analysis (procedure 8 in Figure 1). Formic acid extraction without reduction and purification steps, followed by LC-MS/MS detection is the newer method (procedure 1 in Figure 1). For comparison of the LC-MS/MS data with those of the GC-NPD, the individual PA concentrations of the *N*-oxides and tertiary amines obtained for the LC-MS/MS samples were summed.

The LC-MS/MS method gave a significantly higher total PA concentration compared with the traditional method (Figure 6b), 2.18 and 1.29 mg/g dry plant material ($F_{1,4} = 10.7, p < 0.05$), respectively. For jacobine no significant difference was found between the two methods ($F_{1,4} = 4.4, p = 0.105$), but for the other three major PAs the concentration nearly doubled using the LC-MS/MS method. It should be noted that this difference was only a trend close to significance, for senecionine 0.49 and 1.01 mg/g ($F_{1,4} = 6.0, p = 0.07$), seneciophylline 0.28 and 0.45 mg/g ($F_{1,4} = 7.7, p = 0.05$) and integerrimine 0.09 and 0.17 mg/g ($F_{1,4} = 6.3, p = 0.07$) were obtained in dry plant material, respectively.




Figure 5. The mean PA concentration ($\pm$ SE-bar of total PA concentration) of formic acid extracts reduced by zinc analysed with GC-NPD (4) or LC-MS/MS (5). Formic acid reduced by sodium metabisulfite with GC-NPD (6) or LC-MS/MS (7). Sulfuric acid extracts reduced with zinc analysed with GC-NPD (8) or LC-MS/MS (9)




Figure 6. (A) The mean PA concentration ($\pm$ SE-bar of total PA concentration) analysed with LC-MS/MS of formic acid extracts reduced by sodium metabisulfite (2) and by zinc (3). (B) The mean PA concentration obtained with the traditional method: sulfuric acid extraction, PA *N*-oxide reduction with zinc, purification and GC-NPD analysis (8) and the newer method: formic acid extraction with LC-MS/MS analysis (1)

The PA concentrations obtained by the two different extraction methods (formic acid vs sulfuric acid) and the two detection techniques (GC-NPD vs LC-MS/MS) were more or less comparable as described above. Nevertheless with the LC-MS/MS method a higher PA concentration was obtained than with the traditional method. This can only be explained by the fact that losses occur during zinc reduction and/or in the liquid-liquid purification step of the traditional method. Therefore it is preferable to use the novel LC-MS/MS method; without reduction and purification to minimize PA loss.

For a high sample throughput it is desirable to minimize the complexity of the extraction procedure and to minimize the number of analytical steps to obtain the final extract. The extraction method of the novel technique, namely with formic acid without any extra steps (except a simple dilution step), is far less complex and time consuming compared with the traditional method: extraction with sulfuric acid, reduction with zinc and purification for GC-NPD analysis.

The traditional method for PA analysis by GC-NPD and the newer method for PA analysis by LC-MS/MS were compared by measuring a number of dried *J. vulgaris* samples. The latter technique performed better also with respect to sensitivity, simplicity and selectivity as analysis by LC-MS/MS tolerates a much simpler sample treatment procedure than the classical method with GC analysis. Reduction of analytical steps will in the end result a higher sample throughput and this will allow more comprehensive studies on PA analysis of plant material. The simultaneous detection of PA *N*-oxides and tertiary amines in extracts widens the possibility of investigation of these two forms in biological matrices (plants as well as invertebrates and mammals). LC-MS/MS intrinsically offers a much higher sensitivity than analysis by GC. It is not necessary to use this higher sensitivity to the full in case of plant extracts where there is a plentiful supply, but it will be very valuable when minute amounts of material have to be analysed or when trace levels in biological samples need to be determined. This study shows that the concentrations measured in plant material by LC-MS/MS are higher than those measured by GC-NPD. This strongly indicates that losses occur during the reduction and purification steps required for GC-NPD analysis. These losses are minimised in the LC-MS/MS method.

Based on the simple and rapid sample preparation, sensitivity and discrimination between the two PA forms, formic acid extraction in combination with LC-MS/MS is the method of choice for determining PAs in plant material.

Acknowledgement

We are grateful to Ms. Helene de Vos (Analyst IBL, Leiden University, Leiden, the Netherlands) for her technical assistance.

References

- Bentley MD, Leonard DE, Stoddard WF, Zalkow LH (1984) Pyrrolizidine alkaloids as larval feeding deterrents for spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Ann. Entomol. Soc. Am.* 77:393-397
- Betteridge KCY, Colegate SM (2005) Improved method for extraction and LC-MS analysis of pyrrolizidine alkaloids and their *N*-oxides in honey: application to *Echium vulgare* honeys. *J. Agric. Food Chem.* 53:1894-1902
- Boppré M (1986) Insect pharmacophagously utilizing defensive plant chemicals (pyrrolizidine alkaloids). *Naturwissenschaften* 73:17-26
- Cameron E (1935) A study of natural control of ragwort (*Senecio jacobaea* L.). *J. Ecol.* 23:265-322
- de Boer NJ (1999) Pyrrolizidine alkaloid distribution in *Senecio jacobaea* minimises losses to generalist feeding. *Ent. Exp. Appl.* 91:169-173
- de Jong TJ, Klinkhamer PGL (2005) Evolutionary ecology of plant reproductive strategies. University Press: Cambridge; pp 29-32
- Dreyer DL, Jones KC, Molyneux RJ (1985) Feeding deterrence of some pyrrolizidine, indolizidine, and quinolizidine alkaloids towards pea aphid (*Acyrtosiphon pisum*) and evidence for phloem transport of indolizidine alkaloid swainsonine. *J. Chem. Ecol.* 11:1045-1051
- Frei H, Lüthy J, Brauchli J, Zweifel U, Würigler FE, Schlatter C (1992) Structure/activity relationships of the genotoxic potencies of sixteen pyrrolizidine alkaloids assayed for the induction of somatic mutation and recombination in wing cells of *Drosophila melanogaster*. *Chem. Biol. Interactions* 83:1-22
- Harper JL, Wood WA (1957) Biological flora of the British Isles, *Senecio jacobaea* L. *J. Ecol.* 45:617-637
- Hartmann T, Toppel G (1987) Senecionine *N*-oxide, the primary product of pyrrolizidine alkaloid biosynthesis in root cultures of *Senecio vulgaris*. *Phytochemistry* 26:1639-1643
- Hartmann T, Zimmer M (1986) Organ-specific distribution and accumulation of pyrrolizidine alkaloids during the life-history of 2 annual *Senecio* species. *J. Plant Physiol.* 122:67-80
- Hartmann T, Witte L (1995) Pyrrolizidine alkaloids: chemical, biological and chemoeological aspects. In: Alkaloids: Chemical and biological Perspectives. Vol 9. Pelletier SW (Ed) Pergamon Press: Oxford; pp 155-233
- Hartmann T, Dierich B (1998) Chemical diversity and variation of pyrrolizidine alkaloids of the senecionine type: biological need or coincidence? *Planta* 206:443-451
- Hol WHG, van Veen JA (2002) Pyrrolizidine alkaloids from *Senecio jacobaea* affect fungal growth. *J. Chem. Ecol.* 28:1763-1772
- Jiang Z, Liu F, Goh JYL, Yu L, Li SFY, Ong ES, Ong CN (2009) Determination of senkirkine and senecionine in *Tussilago farfara* using microwave assisted extraction and pressurized hot water extraction with liquid chromatography tandem mass spectrometry. *Talanta* 79:539-546
- Joosten L, Mulder PPJ, Klinkhamer PGL, van Veen JA (2009) Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in *Jacobaea vulgaris*. *Plant Soil* 325:133-143
- Kowalchuk GA, Hol WHG, van Veen JA (2006) Rhizosphere fungal communities are influenced by *Senecio jacobaea* pyrrolizidine alkaloid content and composition. *Soil Biol. Biochem.* 38:2852-2859
- Leiss KA, Yuong CH, Abdel-Farid IFB, Verpoorte R, Klinkhamer PGL (2009) NMR metabolomics of thrips (*Frankliniella occidentalis*) resistance in *Senecio* hybrids. *J. Chem. Ecol.* 35:219-229
- Lindigkeit R, Biller A, Buch M, Schiebel, HM, Boppré M, Hartmann T (1997) The two faces of pyrrolizidine alkaloids: the role of tertiary amine and its *N*-oxide in chemical defense of insects with acquired plant alkaloids. *Eur. J. Biochem.* 245:626-636
- Macel M, Bruinsma M, Dijkstra SM, Ooijendijk T, Niemeyer HM, Klinkhamer PGL (2005) Differences in effects of pyrrolizidine alkaloids on five generalist insect herbivore species. *J. Chem. Ecol.* 31:1493-1508
- Macel M, Klinkhamer PGL (2010) Chemotype of *Senecio jacobaea* affects damage by pathogens and insect herbivores in the field. *Evol. Ecol.* 24:237-250
- Mattocks AR (1967) Spectrophotometric determination of unsaturated pyrrolizidine alkaloids. *Anal. Chem.* 34:443-447
- Pemberton RW, Turner CE (1990) Biological control of *Senecio jacobaea* in northern California, an enduring success. *Entomophaga* 35:71-77
- Pieters LAC, van Zoelen AM, Vrieling K and Vlietinck AJ (1989) Determination of the pyrrolizidine alkaloids from *Senecio jacobaea* by ¹H and ¹³C NMR spectroscopy. *Mag. Reson. Chem.* 27:754-759
- Sokal RR, Rohlf JF (1981) Biometry. W H. Freeman and company: New York; pp 547-555
- van Dam NM (1995) The 'raison d'être' of pyrrolizidine alkaloids in *Cynoglossum officinale*: deterrent effects against generalist herbivores. *J. Chem. Ecol.* 21:507-523
- Vrieling K, Soldaat LL, Smit W (1991) The influence of pyrrolizidine alkaloids of *Senecio jacobaea* on *Tyria jacobaeae*, *Brachycaudus cardii* and *Haplothrips senecionis*. *Neth. J. Zoology* 41:228-239
- Witte L, Ernst L, Adam H, Hartmann T (1992) Chemotypes of 2 pyrrolizidine alkaloid-containing *Senecio* species. *Phytochemistry* 31:559-565
- Witte L, Rubiolo P, Bicchi C, Hartmann T (1993) Comparative analysis of pyrrolizidine alkaloids from natural sources by gas chromatography-mass spectrometry. *Phytochemistry* 32:187-196
- Wuilloud JCA, Gratz GS, Gamble BM, Wolnik KA (2004) Simultaneous analysis of hepatotoxic pyrrolizidine alkaloids and *N*-oxides in comfrey root by LC-ion trap mass spectrometry. *Analyst* 129:150-156
- Zhang F, Wang CH, Wang W, Chen LX, Ma HY, Zhang CF, Zhang M, Bligh ASW, Wang ZT (2008) Quantitative analysis by HPLC-MS2 of the pyrrolizidine alkaloid adonifoline in *Senecio scandens*. *Phytochem. Anal.* 19:25-31