



Universiteit  
Leiden  
The Netherlands

## Quantitative modelling of the response of earthworms to metals

Qiu, H.

### Citation

Qiu, H. (2014, June 10). *Quantitative modelling of the response of earthworms to metals*. Retrieved from <https://hdl.handle.net/1887/25885>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/25885>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/25885> holds various files of this Leiden University dissertation

**Author:** Qiu, Hao

**Title:** Quantitative modelling of the response of earthworms to metals

**Issue Date:** 2014-06-10

---

**Bibliography**

- Akaike H. 1981. Likelihood of a model and information criteria. *J Econometrics* 16: 3-14.
- Ahlf W, Drost W, Heise S. 2009. Incorporation of metal bioavailability into regulatory frameworks - metal exposure in water and sediment. *J Soil Sediment* 9: 411-419.
- Allen HE, Hansen DJ. 1996. The importance of trace metal speciation to water quality criteria. *Water Environ Res* 68: 42-54.
- Alloway BJ. 1995. Soil processes and the behavior of heavy metals. In: Alloway BJ. (ed.) *Heavy Metals In Soils*. Blackie Academic and Professional Publisher, New York.
- Almas AR, Lofts S, Mulder J, Tipping E. 2007. Solubility of major cations and Cu, Zn and Cd in soil extracts of some contaminated agricultural soils near a zinc smelter in Norway: modelling with a multisurface extension of WHAM. *Eur J Soil Sci* 58: 1074-1086.
- Alsop DH, Wood CM. 2000. Kinetic analysis of zinc accumulation in the gills of juvenile rainbow trout: effects of zinc accumulation and implications for biotic ligand modelling. *Environ Toxicol Chem* 19: 1911-1981.
- Altenburger R, Walter H, Grote M. 2004. What contributes to the combined effect of a complex mixture? *Environ Sci Technol* 38: 6353-6362.
- Antunes PMC, Kreager NJ. 2009. Development of the terrestrial biotic ligand model for predicting nickel toxicity to barley (*Hordeum Vulgare*): Ion effects at low pH. *Environ Toxicol Chem* 28: 1704-1710.
- Antunes PMC, Berkelaar EJ, Boyle D, Hale BA, Hendershot W, Voigt A. 2006. The biotic ligand model for plants and metals: Technical challenges for field application. *Environ Toxicol Chem* 25: 875-882.
- Ardestani MM, Ortiz MD, Van Gestel CAM. 2013. Influence of Ca and pH on the uptake and effects of Cd in *Folsomia candida* exposed to simplified soil solutions. *Environ Toxicol Chem* 32: 1759-1767.
- Ardestani MM, Van Gestel CAM. 2013. Using a toxicokinetics approach to explain the effect of soil pH on cadmium bioavailability to *Folsomia candida*. *Environ Pollut* 180: 122-130.
- Baas J, Van Houte BPP, Van Gestel CAM, Kooijman SALM. 2007. Modelling the effects of binary mixtures on survival in time. *Environ Toxicol Chem* 26: 1320-1327.
- Backhaus T, Faust M. 2012. Predictive environmental risk assessment of chemical mixtures: a conceptual framework. *Environ Sci Technol* 46: 2564-2573.
- Baird DJ, Rubach MN, Van den Brinkt PJ. 2008. Trait-based ecological risk assessment (TERA): The new frontier? *Integr Environ Assess Manag* 4: 2-3.
- Barata C, Baird DJ, Nogueira AJA, Soares AMVM, Riva MC. 2006. Toxicity of binary mixtures of metals and pyrethroid insecticides to *Daphnia magna Straus*. Implications for multi-substance risks assessment. *Aquat Toxicol* 78: 1-14.
- Barnett AJ, Finlay K, Beisner BE. 2007. Functional diversity of crustacean zooplankton communities: towards a trait-based classification. *Freshw Biol* 52: 796-813.
- Baskin Y. 2005 *Under ground: how creatures of mud and dirt shape our world*. Shearwater Books, Island Press, Washington D.C. UAS.
- Batley GE, Apte SC, Stauber JL. 2004. Speciation and bioavailability of trace metals in water: progress since 1982. *Aust J Chem* 57: 903-919.

- Bengtsson G, Tranvik L. 1989. Critical metal concentrations for forest soil invertebrates. *Water Air Soil Pollut* 47: 381-417.
- Bliss CI. 1939. The toxicity of poisons applied jointly. *Ann Appl Biol* 26: 585-615.
- Blouin M, Hodson ME, Delgado EA, Baker G, Brussaard L, Butt KR, Dai J, Dendooven L, Peres G, Tondoh JE, Cluzeau D, Brun JJ. 2013. A review of earthworm impact on soil function and ecosystem services. *Eur J Soil Sci* 64: 161-182.
- Bongers M. 2007. Mixture toxicity of metals to *Folsomia candida* related to (bio)availability in soil. PhD thesis. Free University, Amsterdam.
- Borgmann U, Norwood WP, Dixon DG. 2004. Re-evaluation of metal bioaccumulation and chronic toxicity in *Hyalella azteca* using saturation curves and the biotic ligand model. *Environ Pollut* 131: 469-484.
- Bouché MB. 1977. Strategies lombriciennes. In: Lohm U, Persson T. (eds) Soil organisms as components of ecosystems. *Ecol Bull* 25: 122-132.
- Brock TC. 2013. Priorities to improve the ecological risk assessment and management for pesticides in surface water. *Integr Environ Assess Manag* 9: e64-74.
- Brown PL, Markich SJ. 2000. Evaluation of the free ion activity model of metal-organism interaction: extension of the conceptual model. *Aquat Toxicol* 51: 177-194.
- Brzóska MM, Moniuszko-Jakoniuk J. 2001. Interactions between cadmium and zinc in the organism. *Food Chem Toxicol* 39: 967-980.
- Butt KR. 1993. Reproduction and growth of three deep-burrowing earthworms (Lumbricidae) in laboratory culture in order to assess production for soil restoration. *Biol Fertil Soils* 16: 135-138.
- Calamari D, Alabaster JS. 1980. An approach to theoretical models in evaluating the effects of mixtures of toxicants in the aquatic environment. *Chemosphere* 9: 533-538.
- Campbell PGC. 1995. Interactions between trace metals and aquatic organisms: A critique of the free-ion activity model. In Tessier A, Turner DR. (eds) *Metal Speciation and Bioavailability in Aquatic Systems*. John Wiley, New York, USA, pp 46-102.
- Caren N, Panofsky A. 2005. TQCA: a technique for adding temporality to qualitative comparative analysis. *Socio Meth Res* 34: 147-172.
- Cataldo DA, Garland TR, Wildung RE. 1983. Cadmium uptake kinetics in intact soybean plants. *Plant Physiol* 73:844-848.
- Chapman PM, Allen HE, Godtfredsen K, Z'Graggen MN. 1996. Evaluation of bioaccumulation factors in regulating metals. *Environ Sci Technol* 30: 448-452.
- Conder JM, Lanno RP. 2000. Evaluation of surrogate measures of cadmium, lead, and zinc bioavailability to *Eisenia fetida*. *Chemosphere* 41: 1659-1668.
- Criel P, Lock K, Van Eeckhout H, Oorts K, Smolders E, Janssen CR. 2008. Influence of soil properties on copper toxicity for two soil invertebrates. *Environ Toxicol Chem* 27: 748-1755.
- Curry JP, Schmidt O. 2007. The feeding ecology of earthworms – A review. *Pedobiologia* 50: 463-477.
- Dallinger R. 1993. Strategies of metal detoxification in terrestrial invertebrates. In: Dallinger R, Rainbow PS, eds., *Ecotoxicology of Metals in Invertebrates*. Lewis, Boca Raton, FA, pp 245-289.
- D'Amore JJ, Al-Abed SR, Scheckel KG, Ryan JA. 2005. Methods for speciation of metals in

- soils: a review. *J Environ Qual* 34: 1707-1745.
- Das P, Samantaray S, Rout GR. 1997. Studies on cadmium toxicity in plants: a review. *Environ Pollut* 98:29-36.
- Davis ROE, Bennett HH. 1927. Grouping of soils on the basis of mechanical analysis. United States Department of Agriculture Departmental Circulation No. 419.
- De Boer TE, Taş N, Braster M, Temminghoff EJM, Röling WFM, Roelofs D. 2011. The influence of long-term copper contaminated agricultural soil at different pH levels on microbial communities and springtail transcriptional regulation. *Environ Sci Technol* 46: 60-68.
- De Lange HJ, Kramer K, Faber JH. 2013. Two approaches using traits to assess ecological resilience: A case study on earthworm communities. *Basic Appl Ecol* 14: 64-73.
- De Schamphelaere KAC, Janssen CR. 2002. A biotic ligand model predicting acute copper toxicity for *Daphnia magna*: The effects of calcium, magnesium, sodium, potassium, and pH. *Environ Sci Technol* 36: 48-54.
- De Schamphelaere KAC, Janssen CR. 2004 Development and field validation of a biotic ligand model predicting chronic copper toxicity to *Daphnia magna* *Environ Toxicol Chem* 23: 1365-1375.
- De Vries W, Curlik J, Muranyi A, Alloway B, Groenenberg BJ. 2005. Assessment of relationships between total and reactive concentrations of cadmium, copper, lead and zinc in Hungarian and Slovakian soils. *Ekologia-Bratislava* 24: 152-169.
- De Vries W, Loftis S, Tipping E, Meili M, Groenenberg J, Schütze G. 2007. Impact of soil properties on critical concentrations of cadmium, lead, copper, zinc, and mercury in soil and soil solution in view of ecotoxicological effects. In *Reviews of Environmental Contamination and Toxicology*, Springer, New York, pp 47-89.
- Deleebeeck NME, De Schamphelaere KAC, Janssen CR. 2009. Effects of  $Mg^{2+}$  and  $H^+$  on the toxicity of  $Ni^{2+}$  to the unicellular green alga *Pseudokirchneriella subcapitata*: Model development and validation with surface waters. *Sci Total Environ* 407: 1901-1914.
- Denmark. 2008, European Union risk assessment report. Nickel and nickel compounds. Prepared by The Danish Environmental Protection Agency, on behalf of the European Union.
- Di Toro DM, Allen HE, Bergman HL, Meyer JS, Paquin PR, Santore RC. 2001. Biotic ligand model of the acute toxicity of metals. 1. Technical basis. *Environ Toxicol Chem* 20: 2383-2396.
- Dickson KL, Giesy JP, Wolfe L. 1994. *Bioavailability: Physical, Chemical and Biological Interactions*. CRC, Boca Raton, FL, USA.
- Dittbrenner N, Schmitt H, Capowicz Y, Triebkorn R. 2011. Sensitivity of *Eisenia fetida* in comparison to *Aporrectodea caliginosa* and *Lumbricus terrestris* after imidacloprid exposure. Body mass change and histopathology. *J Soil Sediment* 11: 1000-1010.
- Domínguez J. 2004. State of the art and new perspectives on vermicomposting research. In: Edwards CA. (eds) *Earthworm Ecology*, 2nd ed. CRC Press LLC, Florida, USA, pp 401-424.
- Doube BM, Schmidt O, Killham K, Correll R. 1997. Influence of mineral soil on the palatability of organic matter for Lumbricid earthworms: A simple food preference study. *Soil Biol Biochem* 29: 569-575.

- Dudal Y, Gérard F. 2004. Accounting for natural organic matter in aqueous chemical equilibrium models: a review of the theories and applications. *Earth-Sci Rev* 66: 199-216.
- ECB. 2009. European union risk assessment report on nickel and nickel compounds. Ispra, Italy: European Chemicals Bureau.
- ECI. 2008. Voluntary European Union risk assessment report. Copper, copper II sulphate pentahydrate, copper(I)oxide, copper(II)oxide, dicopperchloride trihydroxide. European Copper Institute.
- Eisenhauer N, Marhan S, Scheu S. 2008. Assessment of anecic behavior in selected earthworm species: Effects on wheat seed burial, seedling establishment, wheat growth and litter incorporation. *Appl Soil Ecol* 38: 79-82.
- Edwards CA, Bohlen PJ. 1996. *Biology and Ecology of Earthworms*. Chapman and Hall, London
- Edwards CA, Arancon NQ. 2004. The use of earthworms in the breakdown of organic wastes to produce vermicomposts and animal feed protein. In: Edwards CA. (ed) *Earthworm Ecology*, 2nd edn. CRC Press LLC, Florida, pp 345-379.
- Ehlers LJ, Luthy RG. 2003. Peer reviewed: contaminant bioavailability in soil and sediment. *Environ Sci Technol* 37: 295A-302A.
- Emmerling C, Strunk H. 2012. Active dispersal of the endo-anecic earthworm *Aporrectodea longa* (Ude) in an experimental box. *Soil Organisms* 84: 491-498.
- Erickson RJ. 2013. The biotic ligand model approach for addressing effects of exposure water chemistry on aquatic toxicity of metals: Genesis and challenges. *Environ Toxicol Chem* 32: 1212-1214.
- Fitzpatrick LC, Muratti-Ortiz JF, Venables BJ, Goven AJ. 1996. Comparative toxicity in earthworms *Eisenia fetida* and *Lumbricus terrestris* exposed to cadmium nitrate using artificial soil and filter paper protocols. *Bull Environ Contam Toxicol* 57:63-68.
- Fraser PM, Boag B. 1998. The distribution of lumbricid earthworm communities in relation to flatworms: a comparison between New Zealand and Europe. *Pedobiologia* 42:542-553.
- Gimeno-García, E.; Andreu, V.; Boluda, R. 1996. Heavy metals incidence in the application of inorganic fertilizers and pesticides to rice farming soils. *Environ Pollut* 92: 19-25.
- Goody DC, Shand P, Kinniburgh DG, Van Riemsdijk WH. 1995. Field based partition coefficients for trace elements in soil solutions. *Eur J Soil Sci* 46: 265-285.
- Groenberg JE, Römkens PFAM, Comans RNJ, Luster J, Pampura T, Shotbolt L, Tipping E, De Vries W. 2010. Transfer functions for solid-solution partitioning of cadmium, copper, nickel, lead and zinc in soils: derivation of relationships for free metal ion activities and validation with independent data. *Eur J Soil Sci* 61: 58-73.
- Haanstra L, Doelman P, Voshaar JHO. 1985. The use of sigmoidal dose response curves in soil ecotoxicological research. *Plant Soil* 84: 293-297.
- Hamilton MA, Russo RC, Thurston RV. 1977. Trimmed Spearman-Kärber method for estimating median lethal concentrations in toxicity bioassays. *Environ Sci Technol* 11: 714-719.
- Hassler CS, Slaveykova VI, Wilkinson KJ. 2004. Some fundamental (and often overlooked) considerations underlying the free ion activity and biotic ligand model. *Environ Toxicol*

- Chem* 23: 283-291.
- Hassler CS, Wilkinson KJ. 2003. Failure of the biotic ligand and free-ion activity models to explain zinc bioaccumulation by *Chlorella kesslerii*. *Environ Toxicol Chem* 22: 620-626.
- Hauschild MZ, Huijbregts MAJ, Jolliet O, MacLeod M, Margni M, Van de Meent D, Rosenbaum RK, McKone TE. 2008. Building a model based on scientific consensus for life cycle impact assessment of chemicals: the search for harmony and parsimony. *Environ Sci Technol* 42: 7032-7037.
- He E, Qiu H, Van Gestel CAM. 2014. Modelling uptake and toxicity of nickel in solution to *Enchytraeus crypticus* with biotic ligand model theory. *Environ Pollut* 188: 17-26.
- Heijerick DG, De Schamphelaere KAC, Janssen CR. 2002. Predicting acute zinc toxicity for *Daphnia magna* as a function of key water chemistry characteristics: Development and validation of a biotic ligand model. *Environ Toxicol Chem* 21: 1309-1315.
- Hendrix PF, Bohlen PJ. 2002. Exotic Earthworm Invasions in North. America: Ecological and Policy Implications. *Bioscience* 52: 801-811.
- Hobbelen PHF, Koolhaas JE, Van Gestel CAM. 2006. Bioaccumulation of heavy metals in the earthworms *Lumbricus rubellus* and *Aporrectodea caliginosa* in relation to total and available metal concentrations in field soils. *Environ Pollut* 144: 639-646.
- Höfer T, Gerner I, Gundert-Remy U, Liebsch M, Schulte A, Spielmann H, Vogel R, Wettig K. 2004. Animal testing and alternative approaches for the human health risk assessment under the proposed new European chemicals regulation. *Arch. Toxicol.* 78: 549-64.
- Holmstrup M, Sørensen JG, Overgaard J, Bayley M, Bindsbøl AM, Slotsbo S, Fisker KV, Maraldo K, Waagner D, Labouriau R, Asmund G. 2011. Body metal concentrations and glycogen reserves in earthworms (*Dendrobaena octaedra*) from contaminated and uncontaminated forest soil. *Environ Pollut* 159: 190-197.
- Hopkin SP. 1989. *Ecophysiology of Metals in Terrestrial Invertebrates*. Elsevier Applied Science, London
- Houba VJG, Lexmond TM, Novozamsky I, Van der LJJ. 1996. State of the art and future developments in soil analysis for bioavailability assessment. *Sci Total Environ* 178: 21-28.
- ISO. 2012. Soil quality -- Effects of Pollutants on Earthworms -- Part 1: Determination of acute toxicity to *Eisenia fetida*/*Eisenia andrei* ISO 11268-1: 2012; Part 2: Determination of Effects on Reproduction of *Eisenia fetida*/*Eisenia andrei*, ISO 11268-2: 2012. International Standardization Organization, Geneva.
- Jager T. 2013. All individuals are not created equal; accounting for interindividual variation in fitting life-history responses to toxicants. *Environ Sci Technol* 47: 1664-1669.
- Jager T, Fleuren RHLJ, Hogendoorn EA, De Korte G. 2003. Elucidating the routes of exposure for organic chemicals in the earthworm, *Eisenia andrei* (Oligochaeta). *Environ Sci Technol* 37: 3399-3404.
- Janssen CR, Heijerick DG, De Schamphelaere KAC, Allen HE. 2003. Environmental risk assessment of metals: tools for incorporating bioavailability. *Environ Int* 28: 793-800.
- Janssen MPM, Bruins A, De Vries TH, Van Straalen NM. 1991. Comparison of cadmium kinetics in four soil arthropod species. *Arch Environ Contam Toxicol* 20: 305-312.

- Janssen RPT, Posthuma L, Baerselman R, Den Holander HA, Van Veen RPM, Peijnenburg WJGM. 1997. Equilibrium partitioning of heavy metals in Dutch soils. II. Prediction of metal accumulation in earthworms. *Environ Toxicol Chem* 16: 2479-2488.
- Jho EH, An J, Nam K. 2011. Extended biotic ligand model for prediction of mixture toxicity of Cd and Pb using single metal toxicity data. *Environ Toxicol Chem* 30: 1697-1703.
- Jihen EH, Fatima H, Nouha A, Baati T, Imed M, Abdelhamid K. 2010. Cadmium retention increase: A probable key mechanism of the protective effect of zinc on cadmium-induced toxicity in the kidney. *Toxicol Lett* 196: 104-109.
- Jihen EH, Messaoudi I, Hammouda F, Kerkeni A. 2009. Protective effects of selenium (Se) and zinc (Zn) on cadmium (Cd) toxicity in the liver of the rat: effects on the oxidative stress. *Ecotoxicol Environ Saf* 72: 1559-1564.
- Jihen EH, Messaoudi I, Fatima H, Kerkeni A. 2008. Protective effects of selenium (Se) and zinc (Zn) on cadmium (Cd) toxicity in the liver and kidney of the rat: histology and Cd accumulation. *Food Chem Toxicol* 46: 3522-3527.
- Jonker MJ, Svendsen C, Bedaux JJM, Bongers M, Kammenga JE. 2005. Significance testing of synergistic/antagonistic, dose level-dependent, or dose ratio-dependent effects in mixture dose-response analysis. *Environ Toxicol Chem* 24: 2701-2713.
- Jonker MJ, Sweijen RAJC, Kammenga JE. 2004. Toxicity of simple mixtures to the nematode *Caenorhabditis elegans* in relation to soil sorption. *Environ Toxicol Chem* 23: 480-488.
- Junghans M, Backhaus T, Faust M, Scholze M, Grimme LH. 2006. Application and validation of approaches for the predictive hazard assessment of realistic pesticide mixtures. *Aquat Toxicol* 76: 93-110.
- Kalis EJJ, Temminghoff EJM, Town RM, Unsworth ER, Van Riemsdijk WH. 2008. Relationship between metal speciation in soil solution and metal adsorption at the root surface of ryegrass. *J Environ Qual* 37: 2221-2231.
- Khalil MA, Abdel-Lateif HM, Bayoumi BM, Van Straalen NM. 1996. Analysis of separate and combined effects of heavy metals on the growth of *Aporrectodea caliginosa* (Oligochaeta; Annelida), using the toxic unit approach. *Appl Soil Ecol* 4: 213-219.
- Kinraide TB. 1998. Three mechanisms for the calcium alleviation of mineral toxicities. *Plant Physiol* 118: 513-520.
- Kördel W, Bernhardt C, Derz K, Hund-Rinke K, Harmsen J, Peijnenburg WJGM, Comans R, Tertytze K. 2013. Incorporating availability/bioavailability in risk assessment and decision making of polluted sites, using Germany as an example. *J Hazard Mater* 261: 854-862.
- Kortenkamp A, Backhaus T, Faust M. 2009. State of the art review of mixture toxicity. Report to the Commission of the European Union (Directorate General for the Environment).
- Koster M, De Groot A, Vijver MG, Peijnenburg WJGM. 2006. Copper in the terrestrial environment: Verification of a laboratory-derived terrestrial biotic ligand model to predict earthworm mortality with toxicity observed in field soils. *Soil Biol Biochem* 38: 1788-1796.
- Lahl U, Hawxwell KA. 2006. REACH: the new European chemicals law. *Environ Sci Technol* 40: 7115-7121.

- Langdon CJ, Hodson ME, Arnold RE, Black S. 2005. Survival, Pb-uptake and behaviour of three species of earthworm in Pb treated soils determined using an OECD-style toxicity test and a soil avoidance test. *Environ Pollut* 138: 368-375.
- Landrum PL, Hayton LW, Lee H, McCarthy LS, Mackay D, McKim JM. 1994. Synopsis of discussion session on the kinetics behind environmental bioavailability. In: Hamerlink JL, Landrum PF, Bergman HL, Benson. (eds) *Bioavailability: Physical, Chemical and Biological Interactions*. SETAC Pellston Workshop Series, Lewis, Boca Raton, FL, pp 203-219.
- Lavoie M, Le Faucheur S, Boulemant A, Fortin C, Campbell PGC. 2012. The influence of pH on algal cell membrane permeability and its implications for the uptake of lipophilic metal complexes. *J Phycol* 48: 293-302.
- Le TTY, Peijnenburg WJGM, Hendriks AJ, Vijver MG. 2012. Predicting effects of cations on copper toxicity to lettuce (*Lactuca sativa*) by the biotic ligand model. *Environ Toxicol Chem* 31: 355-359.
- Le TTY, Vijver MG, Jan Hendriks A, Peijnenburg WJGM. 2013. Modelling toxicity of binary metal mixtures ( $\text{Cu}^{2+}$ - $\text{Ag}^+$ ,  $\text{Cu}^{2+}$ - $\text{Zn}^{2+}$ ) to lettuce, *Lactuca sativa*, with the biotic ligand model. *Environ Toxicol Chem* 32: 137-43.
- Levy DB, Barbarick KA, Siemer EG, Sommers LE. 1992. Distribution and partitioning of trace metals in contaminated soil near Leadville, Colorado. *J Environ Qual* 21: 185-195.
- Li B, Zhang X, Wang X, Ma Y. 2009a. Refining a biotic ligand model for nickel toxicity to barley root elongation in solution culture. *Ecotoxicol Environ Saf* 72: 1760-1766.
- Li LZ, Zhou DM, Luo XS, Wang P, Wang QY. 2008. Effect of major cations and pH on the acute toxicity of cadmium to the earthworm *Eisenia fetida*: Implications for the biotic ligand model approach. *Arch Environ Contam Toxicol* 55: 70-77.
- Li LZ, Zhou DM, Peijnenburg WJGM, Wang P, Van Gestel CAM, Jin SY, Wang QY. 2010. Uptake pathways and toxicity of Cd and Zn in the earthworm *Eisenia fetida*. *Soil Biol Biochem* 42: 1045-1050.
- Li LZ, Zhou DM, Wang P, Allen HE, Sauvé S. 2009b. Predicting Cd partitioning in spiked soils and bioaccumulation in the earthworm *Eisenia fetida*. *Appl Soil Ecol* 42: 118-123.
- Lock K, De Schamphelaere KAC, Becaus S, Criel P, Van Eeckhout H, Janssen CR. 2006. Development and validation of an acute biotic ligand model (BLM) predicting cobalt toxicity in soil to the potworm *Enchytraeus albidus*. *Soil Biol Biochem* 38:1924-1932.
- Lock K, Vaneekhout H, De Schamphelaere KAC, Criel P, Janssen C. 2007. Development of a biotic ligand model (BLM) predicting nickel toxicity to barley (*Hordeum vulgare*). *Chemosphere* 66: 1346-1352.
- Lock K, Janssen CR. 2001. Zinc and cadmium body burdens in terrestrial Oligochaetes: Use and significance in environmental risk assessment. *Environ Toxicol Chem* 20: 2067-2072.
- Lock K, Janssen CR. 2002. Mixture toxicity of zinc, cadmium, copper, and lead to the potworm *Enchytraeus albidus*. *Ecotoxicol Environ Saf* 52: 1-7.
- Loewe S, Muischnek H. 1926. Effect of combinations: mathematical basis of problem. N-S. *Arch Ex Path Pharm* 114: 313-326.
- Lofts S, Spurgeon DJ, Svendsen C, Tipping E. 2004. Deriving Soil Critical Limits for Cu, Zn, Cd, and Pb: A method based on free ion concentrations. *Environ Sci Technol* 38:3623-3631.

- Lofts S, Criel P, Janssen CR, Lock K, McGrath SP, Oorts K, Rooney CP, Smolders E, Spurgeon DJ, Svendsen C, Van Eeckhout H, Zhao FZ. 2013. Modelling the effects of copper on soil organisms and processes using the free ion approach: Towards a multi-species toxicity model. *Environ Pollut* 178:244-253.
- Løkke H, Van Gestel CAM. 1998. *Handbook of Soil Invertebrate Toxicity Tests*. John Wiley & Sons, Chichester.
- Lukkari T, Aatsinki M, Vaisanen A, Haimi J. 2005. Toxicity of copper and zinc assessed with three different earthworm tests. *Appl Soil Ecol* 30: 133-146
- Luo XS, Li LZ, Zhou D-M. 2008. Effect of cations on copper toxicity to wheat root: Implications for the biotic ligand model. *Chemosphere* 73: 401-406.
- Luo XS, Zhou DM, Liu XH, Wang YJ. 2006. Solid/solution partitioning and speciation of heavy metals in the contaminated agricultural soils around a copper mine in eastern Nanjing city, China. *J Hazard Mater* 131: 19-27.
- Maleri RA, Reinecke AJ, Reinecke SA. 2007. A comparison of nickel toxicity to pre-exposed earthworms (*Eisenia fetida*, *Oligochaeta*) in two different test substrates. *Soil Biol Biochem* 39: 2849-2853.
- Mariño F, Morgan AJ. 1999. The time-course of metal (Ca, Cd, Cu, Pb, Zn) accumulation from a contaminated soil by three populations of the earthworm, *Lumbricus rubellus*. *Appl Soil Ecol* 12: 169-177.
- Markich J, Brown PL, Jeffree RA, Lim RP. 2003. The effects of pH and dissolved organic carbon on the toxicity of cadmium and copper to a freshwater bivalve: further support for the extended free ion activity model. *Arch Environ Contam Toxicol* 45: 479-491.
- Medlin EA. 1997. An in vitro method for estimating the relative bioavailability of lead in humans. Master's thesis, Department of Geological Sciences, University of Colorado at Boulder.
- Mench M, Martin E. 1991. Mobilization of cadmium and other metals from two soils by root exudates of *Zea mays* L., *Nicotiana tabacum* L. and *Nicotiana rustica* L. *Plant Soil* 132: 187-196.
- Menzies NW, Donn MJ, Kopittke PM. 2007. Evaluation of extractants for estimation of the phytoavailable trace metals in soils. *Environ Pollut* 145: 121-130.
- MEP. 2011. Measures for the Environmental Management of New Chemical Substances (Order 7). The Ministry of Environmental Protection of China.
- Mertens J, Degryse F, Springael D, Smolders E. 2007. Zinc toxicity to nitrification in soil and soilless culture can be predicted with the same biotic ligand model. *Environ Sci Technol* 41: 2992-2997.
- Meyer JS. 2002. The utility of the terms "bioavailability" and "bioavailable fraction" for metals. *Mar Environ Res* 53: 417-423.
- Meyer JS, Santore RC, Bobbitt JP, DeBrey LD, Boese CJ, Paquin PR, Allen HE, Bergman HL, Di Toro DM. 1999. Binding of nickel and copper to fish gills predicts toxicity when water hardness varies, but free-ion activity does not *Environ Sci Technol* 33: 913-916.
- Migula PA, Kedzioriski KA, Nakonieczny M. 1989. Combined and separate effects of cadmium, lead and zinc on growth and feeding in the house cricket (*Acheta domesticus*). *Biologia* 44: 911-921.
- Morel FMM. 1983. *Principles of Aquatic Chemistry*. John Wiley, New York, USA.

- Morgan JE, Richards SPG, Morgan A J. 2002. Contrasting accumulative patterns of two cationic analogues, Ca and Sr, in ecophysiologicaly contrasting earthworm species (*Aporrectodea longa* and *Allolobophora chlorotica*) from the field. *Appl. Soil Ecol.* 21: 11-22.
- Morgan JE, Morgan AJ. 1988. Earthworms as biological monitors of Cd, Cu, Pb, and Zn in metalliferous soils. *Environ Pollut* 54: 123-138.
- Morgan JE, Morgan AJ. 1991. Differences in the accumulated metal concentrations in two epigeic earthworm species (*Lumbricus rubellus* and *Dendrodrilus rubidus*) living in contaminated soils. *Bull Environ Contam Toxicol* 47: 296-301.
- Morgan JE, Morgan AJ. 1999. The accumulation of metals (Cd, Cu, Pb, Zn and Ca) by two ecologically contrasting earthworm species (*Lumbricus rubellus* and *Aporrectodea caliginosa*): implications for ecotoxicological testing. *Appl Soil Ecol* 13: 9-20.
- Muñoz A, Costa M. 2012. Elucidating the mechanisms of nickel compound uptake: A review of particulate and nano-nickel endocytosis and toxicity. *Toxicol Appl Pharmacol* 260: 1-16.
- Nahmani J, Hodson ME, Black S. 2007. A review of studies performed to assess metal uptake by earthworms. *Environ Pollut* 145: 402-424.
- Nannoni F, Protano G, Riccobono F. 2011. Uptake and bioaccumulation of heavy elements by two earthworm species from a smelter contaminated area in northern Kosovo. *Soil Biol Biochem* 43: 2359-2367.
- Niyogi S, Wood CM. 2004. Biotic ligand model, a flexible tool for developing site-specific water quality guidelines for metals. *Environ Sci Technol* 38: 6177-6192.
- Norwood WP, Borgmann U, Dixon DG, Wallace A. 2003. Effects of metal mixtures on aquatic biota: A review of observations and methods. *Hum Ecol Risk Assess* 9: 795-811.
- OECD. 1984. OECD Guidelines for the Testing of Chemicals No. 207. Earthworm, Acute toxicity tests. Organisation for Economic Cooperation and Development, Paris.
- OECD. 2004. OECD Guideline for the Testing of Chemicals No. 222. Earthworm Reproduction Test (*Eisenia fetida*/*Eisenia andrei*). Organization for Economic Cooperation and Development, Paris.
- Onianwa PC, Jaiyeola OM, Egekenze RN. 2003. Heavy metals contamination of topsoil in the vicinities of auto-repair workshops, gas stations and motor-parks in a Nigerian city. *Toxicol Environ Chem* 84: 33-39.
- Oorts K, Ghesquiere U, Swinnen K, Smolders E. 2006. Soil properties affecting the toxicity of CuCl<sub>2</sub> and NiCl<sub>2</sub> for soil microbial processes in freshly spiked soils. *Environ Toxicol Chem* 25: 836-844
- Ore, S.; Mertens, J.; Brandt, K. K.; Smolders, E. 2010 Copper toxicity to Bioluminescent *Nitrosomonas europaea* in soil is explained by the free metal ion activity in porewater. *Environ Sci Technol* 44: 9201-9206.
- Pagenkopf GK. 1983. Gill surface interaction model for trace-metal toxicity to fishes: role of complexation, pH, and water hardness. *Environ Sci Technol* 17: 342-347.
- Pansu M, Gautheyrou J. 2006. *Handbook of Soil Analysis: Mineralogical, Organic and Inorganic Methods*. Springer, Berlin, Germany.
- Paquin PR, Gorsuch JW, Apte S, Batley GE, Bowles KC, Campbell PGC, Delos CG, Di Toro DM, Dwyer RL, Galvez F, Gensemer RW, Goss GG, Hogstrand C, Janssen CR,

- McGeer JC, Naddy RB, Playle RC, Santore RC, Schneider U, Stubblefield WA, Wood CM, Wu KB. 2002. The biotic ligand model: a historical overview. *Comp Biochem Physiol, Part C: Toxicol Pharmacol* 133: 3-35.
- Patrick WH, Gambrell RP, Khalid RA. 1977. Physicochemical factors regulating solubility and bioavailability of toxic heavy metals in contaminated dredged sediment. *J Environ Sci Health, Part A* 12: 475-492.
- Peijnenburg WJGM, Baerselman R, De Groot AC, Jager T, Posthuma L, Van Veen RPM. 1999. Relating environmental availability to bioavailability: Soil-type-dependent metal accumulation in the Oligochaete *Eisenia andrei*. *Ecotoxicol Environ Saf* 44: 294-310.
- Peijnenburg WJGM, Capri E, Kula C, Liess M, Luttk R, Montforts M, Nienstedt K, Rombke J, Sousa JP, Jensen J. 2012. Evaluation of exposure metrics for effect assessment of soil invertebrates. *Crit Rev Environ Sci Technol* 42: 1862-1893.
- Peijnenburg WJGM, Jager T. 2003. Monitoring approaches to assess bioaccessibility and bioavailability of metals: Matrix issues. *Ecotoxicol Environ Saf* 56:63-77.
- Peijnenburg WJGM, Posthuma L, Eijsackers H J, Allen HE. 1997. A conceptual framework for implementation of bioavailability of metals for environmental management purposes. *Ecotoxicol Environ Saf* 37: 163-72.
- Peijnenburg WJGM, Vijver MG. 2007. Metal-specific interactions at the interface of chemistry and biology. *Pure Appl Chem* 79: 2351-2366.
- Peijnenburg WJGM, Vijver MG. 2009. Earthworm and their use in eco(toxico)logical modelling. In: Devillers J. (ed.) *Ecotoxicology Modelling*. Springer, New York, pp 177-204.
- Peijnenburg WJGM, Zablotzkaja M, Vijver MG. 2007. Monitoring metals in terrestrial environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicol Environ Saf* 67: 163-179.
- Pearce TP. 1972. The calcium relations of selected Lumbricidae. *J Anim Ecol* 41: 167-188.
- Playle RC, Dixon DG, Burnison K. 1993. Copper and cadmium binding to fish gills: estimates of metal-gill stability constants and modelling of metal accumulation. *Can J Fish Aquat Sci* 50: 2678-2687.
- Plette ACC, Benedetti MF, Van Riemsdijk WH. 1996. Competitive binding of protons, calcium, cadmium, and zinc to isolated cell walls of a gram-positive soil bacterium. *Environ Sci Technol* 30: 1902-1910.
- Plette ACC, Nederlof MM, Temminghoff EJM, Van Riemsdijk WH. 1999. Bioavailability of heavy metals in terrestrial and aquatic systems: a quantitative approach. *Environ Toxicol Chem* 18: 1882-1890.
- Plytycz B, Klimek M, Homa J, Kruk J, Morgan AJ. 2011a. Species-specific sensitivity of earthworm coelomocytes to dermal metal (Cd, Cu, Ni, Pb, Zn) exposures: Methodological approach. *Pedobiologia* 54S: S203-S210.
- Plytycz B, Cygal M, Lis-Molenda U, Klimek M, Mazur AI, Duchnowski M, Morgan AJ. 2011b. Characteristics of immune-competent amoebocytes non-invasively retrieved from populations of the sentinel earthworm *Lumbricus rubellus* (Annelida; Oligochaeta; Lumbricidae) inhabiting metal polluted field soils. *Ecotoxicol Environ Saf* 74: 719-726.

- Pop VV, Pop AA. 2004. A comprehensive study of the taxonomy and ecology of the Lumbricid earthworm genus *Octodrilus* from the Carpathians. In: Edwards CA. (ed.) *Earthworm Ecology*, 2nd edn. CRC Press LLC, Florida, pp 115-141.
- Posthuma L, Eijssacker HJP, Koelmans AA, Vijver M G. 2008. Ecological effects of diffuse mixed pollution are site-specific and require higher-tier risk assessment to improve site management decisions: A discussion paper. *Sci Total Environ* 406: 503-517.
- Prasad MNV. 1995. Cadmium toxicity and tolerance in vascular plants. *Environ Exp Bot* 35: 525-545.
- Qiu H, Vijver MG, He E, Peijnenburg WJGM. 2013. Predicting copper toxicity to different earthworm species using a multi-component Freundlich model. *Environ Sci Technol* 47:4796-4803.
- Qiu H, Vijver MG, Peijnenburg WJGM. 2011. Interactions of cadmium and zinc impact their toxicity to the earthworm *Aporrectodea caliginosa*. *Environ Toxicol Chem* 30:2084-2093.
- Reeder RJ, Schoonen MAA, Lanzirotti A. 2006. Metal speciation and its role in bioaccessibility and bioavailability. *Rev Mineral Geochem* 64: 59-113.
- Roberts D, Nachtgall M, Sparks DL. 2005. Speciation of metals in soil. In: Tabatabal MA, Sparks, DL. (eds) *Chemical Processes In Soil*. Book Series No.8 Madison WI: Soil Science Society of America, pp 619-654.
- Rooney CP, Zhao FJ, McGrath SP. 2006. Soil factors controlling the expression of copper toxicity to plants in a wide range of European soils. *Environ Toxicol Chem* 25: 726-732.
- Rooney CP, Zhao FJ, McGrath SP. 2007. Phytotoxicity of nickel in a range of European soils: Influence of soil properties, Ni solubility and speciation. *Environ Pollut* 145: 596-605.
- Rubach MN, Baird DJ, Boerwinkel MC, Maund S, Roessink I, Brink P. 2012. Species traits as predictors for intrinsic sensitivity of aquatic invertebrates to the insecticide chlorpyrifos. *Ecotoxicology* 21: 2088-2101.
- Rubach MN, Baird DJ, Van den Brink PJ. 2010. A new method for ranking mode-specific sensitivity of freshwater arthropods to insecticides and its relationship to biological traits. *Environ Toxicol Chem* 29: 476-487.
- Ruby MW, Davis A, Link TE, Schoof R, Chaney RL, Freeman GB, Bergstrom P. 1993. Development of an in vitro screening test to evaluate the in vivo bioaccessibility of ingested mine-waste lead. *Environ Sci Technol* 27: 2870-2877.
- Rüdiger H, Ralf-Rainer M. 2010. *Cell Biology of Metals and Nutrients*. Springer, Heidelberg, Germany.
- Santore RC, Mathew R, Paquin PR, DiToro D. 2002. Application of the biotic ligand model to predicting zinc toxicity to *Rainbow trout*, *Fathead minnow*, and *Daphnia magna*. *Comp Biochem Physiol, Part C: Toxicol Pharmacol* 133: 271-285.
- Santorufu L, Van Gestel CAM, Maisto G. 2012. Ecotoxicological assessment of metal-polluted urban soils using bioassays with three soil invertebrates. *Chemosphere* 88: 418-425.
- Sauvé S, Hendershot W, Allen HE. 2000. Solid-solution partitioning of metals in contaminated soils: dependence on pH, total metal burden, and organic matter. *Environ Sci Technol* 34: 1125-1131.

- Saxe JK, Impellitteri CA, Peijnenburg WJGM, Allen HE. 2001. Novel model describing trace metal concentrations in the earthworm, *Eisenia andrei*. *Environ Sci Technol* 35: 4522-4529.
- Sdepanian S. 2011. Modelling the effects of toxic metal mixtures on the reproduction of *Eisenia veneta* in different types of soil. PhD Thesis. Lancaster University, Lancaster Environment Centre.
- Sizmur T, Hodson ME. 2009. Do earthworms impact metal mobility and availability in soil? A review. *Environ Pollut* 157: 1981-1989.
- Slaveykova VI, Wilkinson KJ. 2005. Predicting the bioavailability of metals and metal complexes: critical review of the biotic ligand model. *Environ Chem* 2: 9-24.
- Smolders E, Oorts K, Van Sprang P, Schoeters I, Janssen CR, McGrath SP, McLaughlin MJ. 2009. Toxicity of trace metals in soil as affected by soil type and aging after contamination: using calibrated bioavailability models to set ecological soil standards. *Environ Toxicol Chem* 28: 1633-1642.
- Sprague JB. 1970. Measurement of pollutant toxicity to fish. II. Utilizing and applying bioassay results. *Water Res* 4: 3-32.
- Spurgeon DJ, Lofts S, Hankard PK, Toal M, McLellan D, Fishwick S, Svendsen C. 2006. Effect of pH on metal speciation and resulting metal uptake and toxicity for earthworms. *Environ Toxicol Chem* 25: 788-796.
- Spurgeon DJ, Hopkin SP. 1996. The effects of metal contamination on earthworm populations around a smelting works: Quantifying species effects. *Appl Soil Ecol* 4: 147-160
- Spurgeon DJ, Hopkin SP. 1999. Comparisons of metal accumulation and excretion kinetics in earthworms (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl Soil Ecol* 11: 227-243.
- Spurgeon DJ, Svendsen C, Rimmer VR, Hopkin SP, Weeks JM. 2000. Relative sensitivity of life-cycle and biomarker responses in four earthworm species exposed to zinc. *Environ Toxicol Chem* 19: 1800-1808.
- Steenbergen NTTM, Iaccino F, De Winkel M, Reijnders L, Peijnenburg WJGM. 2005. Development of a biotic ligand model and a regression model predicting acute copper toxicity to the earthworm *Aporrectodea caliginosa*. *Environ Sci Technol* 39: 5694-5702.
- Sunda WG, Huntsman SA. 1998. Processes regulating cellular metal accumulation and physiological effects: Phytoplankton as model systems. *Sci Total Environ* 219: 165-181.
- Tang W, Sadovic S, Shaikh ZA. 1998. Nephrotoxicity of cadmium-metallothionein: protection by zinc and role of glutathione. *Toxicol Appl Pharm* 151: 276-282.
- Taylor LN, McGeer JC, Wood CM, McDonald DG. 2000. Physiological effects of chronic copper exposure to rainbow trout (*Oncorhynchus mykiss*) in hard and soft water: evaluation of chronic indicators. *Environ Toxicol Chem* 19: 2298-2308.
- Temminghoff EJM, Plette ACC, Van Eck R, Van Riemsdijk WH. 2000. Determination of the chemical speciation of trace metals in aqueous systems by the Wageningen Donnan Membrane Technique. *Anal Chim Acta* 415: 149-157.
- Terhivuo J. 1989. The Lumbricidae (Ologochaeta) of southern Finland: species assemblages, numbers, biomass and respiration. *Ann Zool Fennici* 26: 1-23.

- Tessier A, Turner DR. 1995. *Metal speciation and bioavailability in aquatic systems*. John Wiley, Chichester.
- Thakali S, Allen HE, Di Toro DM, Ponizovsky AA, Rooney CP, Zhao F-J, McGrath SP. 2006a. A terrestrial biotic ligand model. 1. Development and application to Cu and Ni toxicities to barley root elongation in soils. *Environ Sci Technol* 40: 7085-7093.
- Thakali S, Allen HE, Di Toro DM, Ponizovsky AA, Rooney CP, Zhao F-J, McGrath SP, Criel P, Van Eeckhout H, Janssen CR, Oorts K, Smolders E. 2006b. Terrestrial biotic ligand model. 2. Application to Ni and Cu toxicities to plants, invertebrates, and microbes in soil. *Environ Sci Technol* 40: 7094-7100.
- The Netherlands. 2008. European Union risk assessment report on zinc and zinc compounds. Final report, Part 1 Environment; prepared by The Dutch National Institute for Public Health and the Environment (RIVM), on behalf of the European Union.
- Tipping E. 1998. Humic ion-binding model VI: an improved description of the interactions of protons and metal ions with humic substances. *Aquat Geochem* 4: 3-47.
- Tipping E, Rieuwerts J, Pan G, Ashmore MR, Lofts S, Hill MTR, Farago ME, Thornton I. 2003. The solid-solution partitioning of heavy metals (Cu, Zn, Cd, Pb) in upland soils of England and Wales. *Environ Pollut* 125: 213-225.
- Ure AM, Davidson CM. 1995. *Chemical speciation in the environment*. Blackie, Glasgow, UK.
- USEPA. 2007. Update of ambient water quality criteria for copper (EPA 822-R-03-026, February 2007). United States Environmental Protection Agency.
- Uvarov AV, Scheu S. 2004. Effects of density and temperature regime on respiratory activity of the epigeic earthworm species *Lumbricus rubellus* and *Dendrobaena octaedra* (Lumbricidae). *Eur J Soil Biol* 40: 163-167.
- Van Leeuwen CJ, Hermens JLM. 2007. *Risk Assessment of Chemicals - An Introduction*. Springer.
- Van Leeuwen HP, Pinheiro JP. 2001. Speciation dynamics and bioavailability of metals. Exploration of the case of two uptake routes. *Pure Appl Chem* 73: 39-44.
- Van Gestel CAM. 2012. Soil ecotoxicology: state of the art and future directions. *ZooKeys* 176: 275-296.
- Van Gestel CAM, Koolhaas JE. 2004. Water-extractability, free ion activity, and pH explain cadmium sorption and toxicity to *Folsomia candida* (Collembola) in seven soil-pH combinations. *Environ Toxicol Chem* 23: 1822-1833.
- Van Gestel CAM, Ortiz MD, Borgman E, Verweij RA. 2011. The bioaccumulation of molybdenum in the earthworm *Eisenia andrei*: Influence of soil properties and ageing. *Chemosphere* 82: 1614-1619.
- Van Gestel CAM, Van Dis WA. 1988. The influence of soil characteristics on the toxicity of four chemicals to the earthworm *Eisenia fetida andrei* (Oligochaeta), *Biol Fertil Soils* 6: 262-265.
- Van Gestel CAM, Hensbergen PJ. 1997. Interaction of Cd and Zn toxicity for *Folsomia candida* Willem (Collembola: Isotomidae) in relation to bioavailability in soil. *Environ Toxicol Chem* 16: 1177-1186.

- Van Straalen NM, Donker MH, Vijver MG, Van Gestel CAM. 2005. Bioavailability of contaminants estimated from uptake rates into soil invertebrates. *Environ Pollut* 136: 409-417.
- Van Zelm R, Huijbregts MAJ. 2013. Quantifying the trade-off between parameter and model structure uncertainty in life cycle impact assessment. *Environ Sci Technol* 47: 9274-9280.
- Verschoor AJ, Vink JPM, Vijver MG. 2012. Simplification of biotic ligand models of Cu, Ni, and Zn by 1-, 2-, and 3-parameter transfer functions. *Integr Environ Assess Manag* 8: 738-748.
- Vig K, Megharaj M, Sethunathan N, Naidu R. 2003. Bioavailability and toxicity of cadmium to microorganisms and their activities in soil: a review. *Adv Environ Res* 8: 121-135.
- Vijver MG, Elliott EG, Peijnenburg WJGM, De Snoo GR. 2011. Response predictions for organisms water-exposed to metal mixtures: A meta-analysis. *Environ Toxicol Chem* 30: 1482-1487.
- Vijver MG, Peijnenburg WJGM, De Snoo GR. 2010. Toxicological mixture models are based on inadequate assumptions. *Environ Sci Technol* 44: 4841-4842.
- Vijver MG, Van Gestel CAM, Lanno RP, Van Straalen NM, Peijnenburg WJGM. 2004. Internal metal sequestration and its ecotoxicological relevance: A review. *Environ Sci Technol* 38: 4705-4712
- Vijver MG, Vink JPM, Miermans CJH, Van Gestel CAM. 2003. Oral sealing using glue: a new method to distinguish between intestinal and dermal uptake of metals in earthworms. *Soil Biol Biochem* 35:125-132.
- Violle C, Navas ML, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E. 2007. Let the concept of trait be functional! *Oikos* 16: 882-892.
- Wang P, Kopittke PM, De Schamphelaere KAC, Zhao FJ, Zhou DM, Lock K, Ma YB, Peijnenburg WJGM, McGrath SP. 2011a. Evaluation of an electrostatic toxicity model for predicting Ni<sup>2+</sup> toxicity to barley root elongation in hydroponic cultures and in soils. *New Phytol* 192: 414-427.
- Wang P, Kinraide TB, Smolders E, Zhou D-M, Menzies NW, Thakali S, Xia W-W, Hao X-Z, Peijnenburg WJGM, Kopittke PM. 2013 An electrostatic model predicting Cu and Ni toxicity to microbial processes in soils. *Soil Biol Biochem* 57: 720-730.
- Wang P, Kinraide TB, Zhou D, Kopittke PM, Peijnenburg WJGM. 2011b. Plasma membrane surface potential: dual effects upon ion uptake and toxicity. *Plant Physiol* 155: 808-820.
- Wang P, Menzies NW, Wang YM, Zhou DM, Zhao FJ, Kopittke PM. 2012. Identifying the species of copper that are toxic to plant roots in alkaline nutrient solutions. *Plant Soil* 361: 317-327.
- Wang P, Zhou DM, Kinraide TB, Luo XS, Li LZ, Li D, Zhang H. 2008. Cell membrane surface potential plays a dominant role in the phytotoxicity of copper and arsenate. *Plant Physiol* 148: 2134-43.
- Wang P, Zhou DM, Luo XS, Li LZ. 2009. Effects of Zn-complexes on zinc uptake by wheat (*Triticum aestivum*) roots: a comprehensive consideration of physical, chemical and biological processes on biouptake. *Plant Soil* 316: 177-192.
- Wang X, Li B, Ma Y, Hua L. 2010. Development of a biotic ligand model for acute zinc toxicity to barley root elongation. *Ecotoxicol Environ Saf* 73: 1272-1278.

- Webb CT, Hoeting JA, Ames GM, Pyne MI, Poff NL. 2010. A structured and dynamic framework to advance traits-based theory and prediction in ecology. *Ecol Lett* 13: 267–283
- Weltje L. 1998. Mixture toxicity and tissue interactions of Cd, Cu, Pb and Zn in earthworms (Oligochaeta) in laboratory and field soils: a critical evaluation of data. *Chemosphere* 36: 2643-2660.
- Wood CM. 2001. Toxic responses of the gill. In: Schlenk DW, Benson WH. (eds) *Target Organ Toxicity in Marine and Freshwater Teleosts*. Vol 1-Organs. Taylor & Francis, Washington, DC, USA.
- Yan Z, Wang B, Xie D, Zhou Y, Guo G, Xu M, Bai L, Hou H, Li F. 2011. Uptake and toxicity of spiked nickel to earthworm *Eisenia fetida* in a range of Chinese soils. *Environ Toxicol Chem* 30: 2586-2593.
- Zhang H, Lombi E, Smolders E, McGrath S. 2004. Kinetics of Zn release in soils and prediction of Zn concentration in plants using diffusive gradients in thin films. *Environ Sci Technol* 38: 3608-3613.
- Zhang H, Zhao FJ, Sun B, Davison W, McGrath SP. 2001. A new method to measure effective soil solution concentration predicts copper availability to plants. *Environ Sci Technol* 35: 2602-2607.
- Zhang S, Hu F, Li H, Li X. 2009. Influence of earthworm mucus and amino acids on tomato seedling growth and cadmium accumulation. *Environ Pollut* 157: 2737-2742.
- Zidar P, Van Gestel CAM, Strus J. 2009. Single and joint effects of Zn and Cd on *Porcellio scaber* (Crustacea, Isopoda) exposed to artificially contaminated food. *Ecotoxicol Environ Saf* 72: 2075-2082.