



Universiteit  
Leiden  
The Netherlands

## GPCR and G protein mobility in *D. discoideum* : a single molecule study

Hemert, F. van

### Citation

Hemert, F. van. (2009, December 21). *GPCR and G protein mobility in D. discoideum : a single molecule study*. Casimir PhD Series. Retrieved from <https://hdl.handle.net/1887/14549>

Version: Corrected Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/14549>

**Note:** To cite this publication please use the final published version (if applicable).

# Bibliography

- [1] N. Andrew and R. H. Insall. Chemotaxis in shallow gradients is mediated independently of PtdIns 3-kinase by biased choices between random protrusions. *Nat Cell Biol*, 9(2):193–200, 2007.
- [2] N. L. Andrews, K. A. Lidke, J. R. Pfeiffer, A. R. Burns, B. S. Wilsona, J. M. Oliver, and D. S. Lidke. Actin restricts fceri diffusion and facilitates antigen-induced receptor immobilization. *Nat Cell Biol*, 10(8):955–963, 2008.
- [3] Y. Asano, A. Nagasaki, and T. Q. Uyeda. Correlated waves of actin filaments and PIP3 in Dictyostelium cells. *Cell Motil.Cytoskeleton*, 65(12):923–934, 2008.
- [4] H. C. Berg. *Random walks in biology*. Humana Press, 1993.
- [5] P. Bolourani, G. B. Spiegelman, and G. Weeks. Delineation of the roles played by RasG and RasC in camp-dependent signal transduction during the early development of Dictyostelium discoideum. *Mol Biol Cell*, 17(10):4543–4550, 2006.
- [6] L. Bosgraaf and P. J. V. Haastert. Navigation of chemotactic cells by parallel signaling to pseudopod persistence and orientation. *PLoS.One.*, 4(8):e6842, 2009.
- [7] H. R. Bourne, D. A. Sanders, and F. McCormick. The GTPase superfamily: conserved structure and molecular mechanism. *Nature*, 349(6305):117–127, 1991.

- [8] C. Bouzigues, M. Morel, A. Triller, and M. Dahan. Asymmetric redistribution of GABA receptors during GABA gradient sensing by nerve growth cones analyzed by single quantum dot imaging. *Proc.Natl.Acad.Sci U.S.A*, 104(27):11251–11256, 2007.
- [9] M. J. Caterina, J. L. Milne, and P. N. Devreotes. Mutation of the third intracellular loop of the cap receptor, cAR1, of Dictyostelium yields mutants impaired in multiple signaling pathways. *J Biol Chem.*, 269(2):1523–32, 1994.
- [10] S. Chandrasekhar. Stochastic problems in physics and astronomy. *Rev.Mod.Phys.*, 15:1–89, 1943.
- [11] L. Chen, M. Iijima, M. Tang, M. Landree, Y. Huang, Y. Xiong, P. Iglesias, and P. Devreotes. PLA2 and PI3K/PTEN pathways act in parallel to mediate chemotaxis. *Dev.Cell*, 12(4):603–614, 2007.
- [12] L. Chen, C. Janetopoulos, Y. E. Huang, M. Iijima, J. Borleis, , and P. N. Devreotes. Two phases of actin polymerization display different dependencies on PI(3,4,5)P3 accumulation and have unique roles during chemotaxis. *Mol Biol Cell*, 14(12):5028–5037, 2003.
- [13] M. Y. Chen, P. N. Devreotes, and R. E. Gundersen. Serine 113 is the site of receptor-mediated phosphorylation of the Dictyostelium G protein  $\alpha$ -subunit G $\alpha$ 2. *J Biol Chem.*, 269(19):20925–30, 1994.
- [14] J. Colicelli. Human RAS superfamily proteins and related GTPases. *Sci STKE.*, 2004(250):RE13, 2004.
- [15] F. I. Comer and C. A. Parent. PI 3-kinases and PTEN: how opposites chemoattract. *Cell*, 109(5):541–544, 2002.
- [16] E. L. de Hostos, B. Bradtke, F. Lottspeich, R. Guggenheim, and G. Gerisch. Coronin, an actin binding protein of Dictyostelium discoideum localized to cell surface projections, has sequence similarities to G protein  $\beta$  subunits. *EMBO J*, 10(13):4097–4104, 1991.

- [17] S. de Keijzer, A. Sergé, F. van Hemert, P. H. M. Lommerse, G. E. M. Lamers, H. P. Spaink, T. Schmidt, , and B. E. Snaar-Jagalska. A spatially restricted increase in receptor mobility is involved in directional sensing during *Dictyostelium discoideum* chemotaxis. *J Cell Sci*, 121(10):1750–1757, 2008.
- [18] T. Dertinger, V. Pacheco, I. von der Hocht, R. Hartmann, I. Gregor, and J. Endrelein. Two-focus fluorescence correlation spectroscopy: a new tool for accurate and absolute diffusion measurements. *Chemphyschem.*, 8(3):433–443, 2007.
- [19] P. N. Devreotes and T. L. Steck. Cyclic 3',5' AMP relay in *Dictyostelium discoideum*. II. Requirements for the initiation and termination of the response. *J Cell Biol*, 80(2):300–309, 1979.
- [20] L. Eichinger, J. A. Pachebat, G. Glöckner, M. A. Rajandream, R. Sucgang, M. Berriman, J. Song, R. Olsen, K. Szafranski, Q. Xu, B. Tunggal, S. Kummerfeld, M. Madera, B. A. Konfortov, F. Rivero, A. T. Bankier, R. Lehmann, N. Hamlin, R. Davies, P. Gaudet, P. Fey, K. Pilcher, G. Chen, D. Saunders, E. Sodergren, P. Davis, A. Kerhornou, X. Nie, N. Hall, C. Anjard, L. Hemphill, N. Bason, P. Farbrother, B. Desany, E. Just, T. Morio, R. Rost, C. Churcher, J. Cooper, S. Haydock, N. van Driessche, A. Cronin, I. Goodhead, D. Muzny, T. Mourier, A. Pain, M. Lu, D. Harper, R. Lindsay, H. Hauser, K. James, M. Quiles, M. M. Babu, T. Saito, C. Buchrieser, A. Wardrop, M. Felder, M. Thangavelu, D. Johnson, A. Knights, H. Loulseged, K. Mungall, K. Oliver, C. Price, M. A. Quail, H. Urushihara, J. Hernandez, E. Rabbinowitsch, D. Steffen, M. Sanders, J. Ma, Y. Kohara, S. Sharp, M. Simmonds, S. Spiegler, A. Tivey, S. Sugano, B. White, D. Walker, J. Woodward, T. Wincker, Y. Tanaka, G. Shaulsky, M. Schleicher, G. Weinstock, A. Rosenthal, E. C. Cox, R. L. Chisholm, R. Gibbs, W. F. Loomis, M. Platzer, R. R. Kay, J. Williams, P. H. Dear, A. A. Noegel, B. Barrell, and A. Kuspa. The genome of the social amoeba *Dictyostelium discoideum*. *Nature*, 435(7038):43–57, 2005.

- [21] L. Eichinger and F. Rivero. *Dictyostelium discoideum Protocols*. Humana Press, 2006.
- [22] C. A. Elzie, J. Colby, M. A. Sammons, and C. Janetopoulos. Dynamic localization of G proteins in Dictyostelium discoideum. *J Cell Sci*, 122(Pt 15):2597–2603, 2009.
- [23] C. J. L. et al. Loss of the Dictyostelium RasC protein alters vegetative cell size, motility and endocytosis. *Exp. Cell Res.*, 306(1):47–55, 2005.
- [24] P. Fey, P. Gaudet, T. Curk, B. Zupan, E. M. Just, S. Basu, S. N. Merchant, Y. A. Bushanova, G. Shaulsky, W. A. Kibbe, and R. L. Chisholm. dictyBase - a Dictyostelium bioinformatics resource update. *Nucl. Acids Res.*, 37(suppl1):515–519, 2009.
- [25] T. Fischer, L. Lu, H. T. Haigler, and R. Langen. Annexin B12 is a sensor of membrane curvature and undergoes major curvature-dependent structural changes. *Journal of Biological Chemistry*, 282(13):9996–10004, 2007.
- [26] J. Franca-Koh and P. N. Devreotes. Moving forward: mechanisms of chemoattractant gradient sensing. *Physiology.(Bethesda.)*, 19:300–308, 2004.
- [27] J. Franca-Koh, Y. Kamimura, and P. Devreotes. Navigating signaling networks: chemotaxis in Dictyostelium discoideum. *Current Opinion in Genetics & Development*, 16(4):333–338, 2006.
- [28] L. Frigeri and J. R. Apgar. The role of actin microfilaments in the down-regulation of the degranulation response in RBL-2H3 mast cells. *The Journal of Immunology*, 162(4):2243–2250, 1999.
- [29] T. Fujiwara, K. Ritchie, H. Murakoshi, K. Jacobson, and A. Kusumi. Phospholipids undergo hop diffusion in compartmentalized cell membrane. *J Cell Biol*, 157(6):1071–1081, 2002.
- [30] S. Funamoto, R. Meili, S. Lee, L. Parry, and R. A. Firtel. Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell*, 109(5):611–623, 2002.

- [31] N. J. Galvin, D. Stockhausen, B. L. Meyers-Hutchins, and W. A. Frazier. Association of the cyclic AMP chemotaxis receptor with the detergent-insoluble cytoskeleton of *Dictyostelium discoideum*. *J Cell Biol*, 98(2):584–595, 1984.
- [32] A. Gamba, A. de Candia, S. D. Talia, A. Coniglio, F. Bussolino, and G. Serini. Diffusion-limited phase separation in eukaryotic chemotaxis. *Proc.Natl.Acad.Sci U.S.A*, 102(47):16927–16932, 2005.
- [33] K. D. Girard, S. C. Kuo, and D. N. Robinson. Dictyostelium myosin II mechanochemistry promotes active behavior of the cortex on long time scales. *Proc.Natl.Acad.Sci U.S.A*, 103(7):2103–2108, 2006.
- [34] P. J. V. Haastert, J. D. Bishop, and R. H. Gomer. The cell density factor CMF regulates the chemoattractant receptor cAR1 in Dictyostelium. *J Cell Biol*, 134(6):1543–1549, 1996.
- [35] P. J. V. Haastert, I. Keizer-Gunnink, and A. Kortholt. Essential role of PI3-kinase and phospholipase A2 in *Dictyostelium discoideum* chemotaxis. *J Cell Biol*, 177(5):809–816, 2007.
- [36] G. S. Harms, L. Cognet, P. H. Lommerse, G. A. Blab, and T. Schmidt. Autofluorescent proteins in single-molecule research: applications to live cell imaging microscopy. *Biophys.J*, 80(5):2396–2408, 2001.
- [37] D. Hereld and P. N. Devreotes. The cAMP receptor family of Dictyostelium. *Int.Rev Cytol.*, 137B:35–47, 1992.
- [38] O. Hoeller and R. R. Kay. Chemotaxis in the absence of PIP3 gradients. *Curr.Biol*, 17(9):813–817, 2007.
- [39] L. Holtzer, T. Meckel, and T. Schmidt. Nanometric three-dimensional tracking of individual quantum dots in cells. *Applied Physics Letters*, 90(5):053902, Jan 2007.
- [40] M. Iijima and P. Devreotes. Tumor suppressor PTEN mediates sensing of chemoattractant gradients. *Cell*, 109(5):599–610, 2002.

- [41] M. Iijima, Y. E. Huang, H. R. Luo, F. Vazquez, and P. N. Devreotes. Novel mechanism of PTEN regulation by its phosphatidylinositol 4,5-bisphosphate binding motif is critical for chemotaxis. *J Biol Chem.*, 279(16):16606–16613, 2004.
- [42] T. Inoue and T. Meyer. Synthetic activation of endogenous PI3K and Rac identifies an AND-gate switch for cell polarization and migration. *PLoS One.*, 3(8):e3068, 2008.
- [43] V.-P. Jaakola, M. T. Griffith, M. A. Hanson, V. Cherezov, E. Y. T. Chien, J. R. Lane, A. P. IJzerman, and R. C. Stevens. The 2.6 angstrom crystal structure of a human A2A adenosine receptor bound to an antagonist. *Science*, 322(5905):1211–1217, 2008.
- [44] C. Janetopoulos, T. Jin, and P. Devreotes. Receptor-mediated activation of heterotrimeric G-proteins in living cells. *Science*, 291(5512):2408–2411, 2001.
- [45] P. M. Janssens and P. J. V. Haastert. Molecular basis of transmembrane signal transduction in *Dictyostelium discoideum*. *Microbiol.Rev*, 51(4):396–418, 1987.
- [46] T. Jin, N. Zhang, Y. Long, C. A. Parent, and P. N. Devreotes. Localization of the G protein complex in living cells during chemotaxis. *Science*, 287(5455):1034–1036, 2000.
- [47] H. Kae, C. J. Lim, G. B. Spiegelman, and G. Weeks. Chemoattractant-induced Ras activation during *Dictyostelium* aggregation. *EMBO Rep.*, 5(6):602–606, 2004.
- [48] Y. Kamimura, Yoichiro; Xiong, P. A. Iglesias, O. Hoeller, P. Bolourani, and P. N. Devreotes. PIP3-independent activation of TorC2 and PKB at the cell's leading edge mediates chemotaxis. *Curr.Biol*, 18(14):1034–1043, 2008.
- [49] A. R. Kimmel and C. A. Parent. The signal to move: *D. discoideum* go orienteering. *Science*, 300(5625):1525–1527, 2003.

- [50] P. S. Klein, T. J. Sun, C. L. S. 3rd, A. R. Kimmel, R. L. Johnson, and P. N. Devreotes. A chemoattractant receptor controls development in *Dictyostelium discoideum*. *Science*, 241(4872):1467–1472, 1988.
- [51] T. M. Konijn, J. G. van de Meene, J. T. Bonner, and D. S. Barkley. The acrasin activity of adenosine-3',5'-cyclic phosphate. *Proc.Natl.Acad.Sci U.S.A*, 58(3):1152–1154, 1967.
- [52] W. J. Koopmans, R. Buning, T. Schmidt, and N. J. van. spFRET using alternating excitation and FCS reveals progressive DNA unwrapping in nucleosomes. *Biophys.J*, 97(1):195–204, 2009.
- [53] A. Kortholt and P. J. V. Haastert. Highlighting the role of Ras and Rap during *Dictyostelium* chemotaxis. *Cell Signal.*, 20(8):1415–1422, 2008.
- [54] P. W. Kriebel, V. A. Barr, and C. A. Parent. Adenylyl cyclase localization regulates streaming during chemotaxis. *Cell*, 112(4):549–560, 2003.
- [55] A. Kusumi, C. Nakada, K. Ritchie, K. Murase, K. Suzuki, H. Murakoshi, R. S. Kasai, J. Kondo, and T. Fujiwara. Paradigm shift of the plasma membrane concept from the two-dimensional continuum fluid to the partitioned fluid: high-speed single-molecule tracking of membrane molecules. *Annu.Rev Biophys.Biomol.Struct.*, 34:351–378, 2005.
- [56] A. Kusumi and Y. Sako. Cell surface organization by the membrane skeleton. *Curr.Opin.Cell Biol*, 8(4):566–574, 1996.
- [57] A. Kusumi, Y. Sako, and M. Yamamoto. Confined lateral diffusion of membrane receptors as studied by single particle tracking (nanovid microscopy). effects of calcium-induced differentiation in cultured epithelial cells. *Biophys.J*, 65(5):2021–2040, 1993.
- [58] P. D. Langridge and R. R. Kay. Blebbing of *Dictyostelium* cells in response to chemoattractant. *Exp.Cell Res.*, 312(11):2009–2017, 2006.

- [59] H. Levine, D. A. Kessler, and W. J. Rappel. Directional sensing in eukaryotic chemotaxis: a balanced inactivation model. *Proc.Natl.Acad.Sci U.S.A*, 103(26):9761–9766, 2006.
- [60] P. Lilly, L. Wu, D. L. Welker, and P. N. Devreotes. A G-protein  $\beta$ -subunit is essential for Dictyostelium development. *Genes & Development*, 7(6):986–995, 1993.
- [61] C. J. Lim, G. B. Spiegelman, and G. Weeks. RasC is required for optimal activation of adenylyl cyclase and Akt/PKB during aggregation. *EMBO J*, 20(16):4490–4499, 2001.
- [62] C. J. Lim, G. B. Spiegelman, and G. Weeks. RasC is required for optimal activation of adenylyl cyclase and Akt/PKB during aggregation. *EMBO J*, 20(16):4490–4499, 2001.
- [63] L. Ma, C. Janetopoulos, L. Yang, P. N. Devreotes, and P. A. Iglesias. Two complementary, local excitation, global inhibition mechanisms acting in parallel can explain the chemoattractant-induced regulation of PI(3,4,5)P3 response in Dictyostelium cells. *Biophys.J*, 87(6):3764–3774, 2004.
- [64] R. Merkel, R. Simson, D. A. Simson, M. Hohenadl, A. Boulbitch, E. Wallraff, and E. Sackmann. A micromechanic study of cell polarity and plasma membrane cell body coupling in Dictyostelium. *Biophys.J*, 79(2):707–719, 2000.
- [65] Y. Miyanaga, S. Matsuoka, T. Yanagida, and M. Ueda. Stochastic signal inputs for chemotactic response in Dictyostelium cells revealed by single molecule imaging techniques. *Biosystems*, 88(3):251–260, 2007.
- [66] N. Morone, T. Fujiwara, K. Murase, R. S. Kasai, H. Ike, S. Yuasa, J. Usukura, and A. Kusumi. Three-dimensional reconstruction of the membrane skeleton at the plasma membrane interface by electron tomography. *J Cell Biol*, 174(6):851–862, 2006.

- [67] M. Nobles, A. Benians, and A. Tinker. Heterotrimeric G proteins precouple with G protein-coupled receptors in living cells. *Proc.Natl.Acad.Sci U.S.A.*, 102(51):18706–18711, 2005.
- [68] K. Okaichi, A. B. Cubitt, G. S. Pitt, and R. A. Firtel. Amino acid substitutions in the Dictyostelium G $\alpha$  subunit G $\alpha$ 2 produce dominant negative phenotypes and inhibit the activation of adenylyl cyclase, guanylyl cyclase, and phospholipase C. *Mol Biol Cell*, 3(7):735–747, 1992.
- [69] K. Palczewski, T. Kumashiro, T. Hori, C. A. Behnke, H. Motoshima, B. A. Fox, I. L. Trong, D. C. Teller, T. Okada, R. E. Stenkamp, M. Yamamoto, and M. Miyano. Crystal structure of Rhodopsin: A G protein-coupled receptor. *Science*, 289(5480):739–745, 2000.
- [70] C. A. Parent, B. J. Blacklock, W. M. Froehlich, D. B. Murphy, and P. N. Devreotes. G protein signaling events are activated at the leading edge of chemotactic cells. *Cell*, 95(1):81–91, 1998.
- [71] T. Pawson and J. D. Scott. Signaling through scaffold, anchoring, and adaptor proteins. *Science*, 278(5346):2075–2080, 1997.
- [72] T. D. Pollard. Regulation of actin filament assembly by Arp2/3 complex and formins. *Annu.Rev Biophys.Biomol.Struct.*, 36:451–477, 2007.
- [73] A. Ponti, M. Machacek, S. L. Gupton, C. M. Waterman-Storer, and G. Danuser. Two distinct actin networks drive the protrusion of migrating cells. *Science*, 305(5691):1782–1786, 2004.
- [74] M. Postma, J. Roelofs, J. Goedhart, T. W. Gadella, A. J. Visser, and P. J. Van Haastert. Uniform cAMP Stimulation of Dictyostelium Cells Induces Localized Patches of Signal Transduction and Pseudopodia. *Mol. Biol. Cell*, 14(12):5019–5027, 2003.
- [75] M. Postma and P. J. Van Haastert. A diffusion-translocation model for gradient sensing by chemotactic cells. *Biophysical Journal*, 2001.

- [76] E. O. Potma, W. P. de Boeij, L. Bosgraaf, J. Roelofs, P. J. van Haastert, and D. A. Wiersma. Reduced protein diffusion rate by cytoskeleton in vegetative and polarized Dictyostelium cells. *Biophys.J.*, 81(4):2010–2019, 2001.
- [77] S. G. F. Rasmussen, H.-J. Choi1, D. M. Rosenbaum, T. S. Kobilka, F. S. Thian, P. C. Edwards, M. Burghammer, V. R. P. Ratnala1, R. Sanishvili, R. F. Fischer, G. F. X. Schertler, W. I. Weis, and B. K. Kobilka. Crystal structure of the human [bgr]2 adrenergic g-protein-coupled receptor. *Nature*, 450(7168):383–387, 2007.
- [78] D. S. Rhoads, S. M. Nadkarni, L. Song, C. Voeltz, E. Bodenschatz, and J.-L. Guan. Using microfluidic channel networks to generate gradients for studying cell migration. *Methods Mol Biol*, 294:347–357, 2005.
- [79] A. T. Sasaki, C. Chun, K. Takeda, and R. A. Firtel. Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement. *J Cell Biol*, 167(3):505–518, 2004.
- [80] A. T. Sasaki, C. Janetopoulos, S. Lee, P. G. Charest, K. Takeda, L. W. Sundheimer, R. Meili, P. N. Devreotes, and R. A. Firtel. G protein-independent Ras/PI3K/F-actin circuit regulates basic cell motility. *J Cell Biol*, 178(2):185–191, 2007.
- [81] T. Schmidt, G. J. Schütz, W. Baumgartner, H. J. Gruber, and H. Schindler. Imaging of single molecule diffusion. *Proc.Natl.Acad.Sci U.S.A*, 93(7):2926–2929, 1996.
- [82] G. J. Schütz, H. Schindler, and T. Schmidt. Single-molecule microscopy on model membranes reveals anomalous diffusion. *Biophys.J.*, 73(2):1073–1080, 1997.
- [83] S. Semrau and T. Schmidt. Particle image correlation spectroscopy (PICS): retrieving nanometer-scale correlations from high-density single-molecule position data. *Biophys.J.*, 92(2):613–621, 2007.

- [84] L. Song, S. M. Nadkarni, H. U. Bödeker, C. Beta, A. Bae, C. Franck, W.-J. Rappel, W. F. Loomis, and E. Bodenschatz. *Dictyostelium discoideum* chemotaxis: threshold for directed motion. *Eur.J Cell Biol.*, 85(9-10):981–989, 2006.
- [85] B. L. Sprague, R. L. Pego, D. A. Stavreva, and J. G. McNally. Analysis of binding reactions by fluorescence recovery after photobleaching. *Biophys.J.*, 86(6):3473–3495, 2004.
- [86] K. Suzuki, K. Ritchie, E. Kajikawa, T. Fujiwara, and A. Kusumi. Rapid hop diffusion of a G-protein-coupled receptor in the plasma membrane as revealed by single-molecule techniques. *Biophys.J.*, 88(5):3659–3680, 2005.
- [87] M. Tsujioka, K. Yoshida, A. Nagasaki, S. Yonemura, A. Muller-Taubenberger, and T. Q. P. Uyeda. Overlapping functions of the two talin homologues in *Dictyostelium*. *Eukaryot.Cell*, 7(5):906–916, 2008.
- [88] R. I. Tuxworth, J. L. Cheetham, L. M. Machesky, G. B. Spiegelmann, G. Weeks, and R. H. Insall. *Dictyostelium RasG* is required for normal motility and cytokinesis, but not growth. *J Cell Biol.*, 138(3):605–614, 1997.
- [89] K. S. Uchida, T. Kitanishi-Yumura, and S. Yumura. Myosin II contributes to the posterior contraction and the anterior extension during the retraction phase in migrating *Dictyostelium* cells. *J Cell Sci.*, 116(Pt 1):51–60, 2003.
- [90] M. Ueda, Y. Sako, T. Tanaka, P. Devreotes, and T. Yanagida. Single-molecule analysis of chemotactic signaling in *Dictyostelium* cells. *Science*, 294(5543):864–867, 2001.
- [91] D. M. Veltman, I. Keizer-Gunnik, and P. J. V. Haastert. Four key signaling pathways mediating chemotaxis in *Dictyostelium discoideum*. *J Cell Biol.*, 180(4):747–753, 2008.
- [92] G. H. Wadhams and J. P. Armitage. Making sense of it all: bacterial chemotaxis. *Nat Rev Mol Cell Biol.*, 5(12):1024–1037, 2004.

- [93] D. J. Watts and J. M. Ashworth. Growth of myxameobae of the cellular slime mould *Dictyostelium discoideum* in axenic culture. *Biochem.J.*, 119(2):171–174, 1970.
- [94] I. Weisswange, T. Bretschneider, and K. I. Anderson. The leading edge is a lipid diffusion barrier. *J Cell Sci.*, 118(19):4375–4380, 2005.
- [95] D. Wessels, R. Brincks, S. Kuhl, V. Stepanovic, K. J. Daniels, G. Weeks, C. J. Lim, G. Spiegelman, D. Fuller, N. Iranfar, W. F. Loomis, and D. R. Soll. RasC plays a role in transduction of temporal gradient information in the cyclic-AMP wave of *Dictyostelium discoideum*. *Eukaryotic Cell*, 3(3):646–662, 2004.
- [96] D. Wessels, N. A. Schroeder, E. Voss, A. L. Hall, J. Condeelis, and D. R. Soll. cAMP-mediated inhibition of intracellular particle movement and actin reorganization in *Dictyostelium*. *J Cell Biol.*, 109(6 Pt 1):2841–2851, 1989.
- [97] N. Wettschureck and S. Offermanns. Mammalian G proteins and their cell type specific functions. *Physiological Reviews*, 85(4):1159–1204, 2005.
- [98] S. Wieser, M. Moertelmaier, E. Fuertbauer, H. Stockinger, and G. J. Schütz. (un)confined diffusion of CD59 in the plasma membrane determined by high-resolution single molecule microscopy. *Biophys.J.*, 92(10):3719–3728, 2007.
- [99] L. Wu, R. Valkema, P. J. V. Haastert, and P. N. Devreotes. The G protein  $\beta$  sub-unit is essential for multiple responses to chemoattractants in *Dictyostelium*. *J Cell Biol.*, 129(6):1667–1675, 1995.
- [100] X. Xu, M. Meier-Schellersheim, X. Jiao, L. E. Nelson, and T. Jin. Quantitative imaging of single live cells reveals spatiotemporal dynamics of multistep signaling events of chemoattractant gradient sensing in *Dictyostelium*. *Mol Biol Cell*, 16(2):676–688, 2005.
- [101] K. Yoshida and T. Soldati. Dissection of amoeboid movement into two mechanically distinct modes. *J Cell Sci.*, 119(Pt 18):3833–3844, 2006.

- [102] N. Zhang, Y. Long, and P. N. Devreotes. G $\gamma$  in Dictyostelium: Its role in localization of g $\beta$ g to the membrane is required for chemotaxis in shallow gradients. *Mol Biol Cell*, 12(10):3204–3213, 2001.
- [103] S. Zhang, P. G. Charest, and R. A. Firtel. Spatiotemporal regulation of Ras activity provides directional sensing. *Curr.Biol*, 18(20):1587–1593, 2008.

