



Universiteit
Leiden
The Netherlands

Breeding birds on organic and conventional arable farms

Kragten, S.

Citation

Kragten, S. (2009, December 2). *Breeding birds on organic and conventional arable farms*. Retrieved from <https://hdl.handle.net/1887/14458>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14458>

Note: To cite this publication please use the final published version (if applicable).

Chapter 7

Abundance of invertebrate prey for birds on organic and conventional arable farms in the Netherlands

Steven Kragten, Wil L.M. Tamis, Erik Gertenaar, Sharon M. Midcap Ramiro,

Rob J. van der Poll, Juanjuan Wang & Geert R. de Snoo

Accepted in Bird Conservation International



One of the nearly 1.000 pitfall traps used for invertebrate sampling

Abstract

As a result of agricultural intensification populations of farmland birds have been in steep decline for several decades now. Reduction of food abundance has been mentioned to be one factor behind these declines. Extensive farm management, such as organic, is expected to provide more food for birds. In this study, we compared invertebrate prey abundance for birds between organic and conventional arable farms during the breeding season. We made comparisons for three different groups of birds: (1) birds feeding on soil living invertebrates (earthworms), (2) birds feeding on ground-dwelling invertebrates and (3) birds feeding on aerial invertebrates. Invertebrate abundance was compared between organic and conventional farms, crops and non-crop habitats, and between crops and non-crop habitats under the same farm management. On organic sites earthworm abundance was 2-4 times higher compared to conventional sites, but no differences were found between crop types. Total abundance of ground-dwelling invertebrates did not differ between organic and conventional sites, but positive effects were found for several individual taxonomic groups, such as carabid beetles and spiders. On organic farms invertebrate abundance was higher in carrots, cereals and onions compared to other crops. On conventional farms this was true for onions. Compared with most crops, ground dwelling invertebrate abundance was low in uncropped field margins and on ditch banks. On organic farms aerial invertebrate abundance was approximately 70% higher compared to conventional farms. On cereal fields aerial invertebrates were especially abundant.

Keywords: Skylark; Lapwing; Barn Swallow; Food abundance; Earthworms

Introduction

The past few decades have seen a dramatic intensification of arable farming, characterised by increased pesticide usage, high inputs of artificial fertilizers, removal of non-crop habitats (such as hedgerows and ditches), larger fields and reduced crop diversity (Medley *et al.* 1995, Chamberlain *et al.* 2000, Stoate *et al.* 2001, Robinson and Sutherland 2002). At the same time, populations of bird species associated with arable landscapes have shown a marked decline (e.g. Krebs *et al.* 1999, Donald *et al.* 2001, 2006, Wretenberg *et al.* 2006).

One of the mechanisms believed to contribute to the decline of farmland bird populations is a reduction in the availability of invertebrate prey, one of the most important food sources for adults, and especially chicks, of most farmland bird species (Wilson *et al.* 1999, Benton *et al.* 2002, Holland *et al.* 2006). In agricultural areas invertebrate populations have declined as a result of several changes in farming practice (cf. Robinson and Sutherland 2002), viz.: (1) increased pesticide use, (2) a switch from organic manure to artificial fertilizers and (3) loss of semi-natural habitats.

An array of studies have shown that breeding bird densities are higher on organically managed farms (e.g. Christensen *et al.* 1996, Chamberlain *et al.* 1999, Beecher *et al.* 2002, Kragten and de Snoo 2008). One of the factors suggested as being responsible is the greater availability of food on organic farms (e.g. Christensen *et al.* 1996). Several studies have focussed on differences in invertebrate abundance between organically and conventionally managed farms e.g. (Hole *et al.* 2005). However, most studies focussed on only one taxonomic group of invertebrates and in most cases only one crop type was investigated. Farmland birds have different diets and diets often have a diverse prey composition (Holland *et al.* 2006). In order to assess the food abundance for farmland birds on farms it is therefore necessary to focus on a wide range of taxonomic groups of invertebrates. Besides this, crop type is likely to influence

food availability as a result of crop management and sward structure (e.g. Atkinson *et al.* 2005, Hole *et al.* 2005). However, so far it is unknown which crop types provide most food for farmland birds under organic or conventional management.

The present paper seeks to analyse whether organic arable farms provide insectivorous farmland birds with more prey items during the breeding season compared to conventionally managed farms. To this end three studies were designed, each focusing on a different group of invertebrates: soil-dwelling invertebrates (earthworms), ground-dwelling invertebrates (carabids, spiders, etc.) and aerial invertebrates. These groups represent the main prey items of different avian feeding guilds: (1) birds feeding mainly on earthworms (e.g. lapwing, *Vanellus vanellus*), (2) birds feeding mainly on invertebrates active on the ground surface (e.g. skylark, *Alauda arvensis*) and (3) birds feeding mainly on aerial invertebrates (e.g. barn swallow, *Hirundo rustica*). The abundance of each invertebrate group on organic and conventional farms was compared at both farm level and crop level. A comparison was also made of invertebrate abundance on different crop types under the same type of farm management. In this way not only was an overall picture obtained of the effects of organic farming on bird food abundance, but insights were also obtained into which types of crop potentially provide the greatest availability of food items. On the same farms included in this study, breeding bird surveys have been carried out as well (Kragten and de Snoo 2008, Kragten *et al.* 2009). Consequently, this study could give some more insight in the effects of differences in food abundance between organic and conventional arable farms on breeding bird densities.

Methods

Study site and data collection

The different studies were all carried out on organic and conventional arable farms in the province of Flevoland (52° 34' N, 5° 39' E), the Netherlands, in spring 2004 and 2005. Farms were selected in two sub-areas: Oostelijk Flevoland and Noordoostpolder, both polders, reclaimed during the 1950s and 1930s respectively. The soil type is therefore of marine clay origin. The main form of land use in the study area is agricultural (64% of total area). Of the farmland 75% is used for arable farming, 13% for grassland and 12% for other types of agriculture. Approximately 8% of the farmland is managed organically (Bakker 2007). The landscape is very homogeneous and open Natural habitats consist mainly of woodland patches, grass margins and artificial watercourses. Main differences between the two sub-areas are: (1) somewhat smaller parcels in Noordoostpolder and (2) trees are generally older in Noordoostpolder.

In both sub-areas 10 conventional and 10 organic farms were selected. A pairwise was used for farm selection with each pair consisting of one organic and one conventional farm. Both farms within a pair were surrounded by similar landscape elements, such as woodlots, tree lines, roads, power lines and wind turbines. Distance between farms was at least 600 m. Because of the homogeneous landscape between-pair variation of surrounding landscape was very limited. On-farm habitat factors, such as crops and non-crop habitats were not included in the pairing protocol as they constitute essential differences between the two farming systems and are a direct result of farm management. Consequently, there were large differences in crop composition between the two farm types (Table 12). On average, about 3–4% of the farm area consisted of non-crop habitats. Grassy, semi-natural elements were far more dominant than ditches, reed or woody elements. Grassy elements comprised grassy field

margins and ditch banks. Woody elements consisted mainly of solitary trees and scrub, though some farms had a small hedgerow. Organic farms had slightly more non-crop habitat than conventional farms (2004: 3.7% vs. 3.1%; 2005: 4.4% vs. 3.6%), although in both years differences were not significant (2004: Wilcoxon, $Z = 1.682$, NS, 2005: Wilcoxon, $Z = 1.717$, NS). When differences were analysed per habitat type only in 2005 more woody habitat elements were found on organic farms (Wilcoxon, $Z = 2.666$, $P < 0.01$), although the absolute difference was small. All the organic farms selected had been managed according to European Union Regulation 2092/91/EEC for at least 5 years. On these farms, farmyard manure was used and weeds were controlled mechanically. All the conventional farmers used artificial fertilizers (in some cases manure as well), herbicides, pesticides and fungicides. The farm layout (crop partitioning and abundance of non-crop habitats) has been described previously by Kragten and de Snoo (2008).

In spring 2005 the study on earthworms was carried out. On ten organic and ten conventional arable farms, which were all in the same sub-area, earthworms were sampled in two sampling rounds: the first from 30 March to 1 April and the second from 28 April to 3 May. These periods coincide with the breeding season of lapwings, which feed largely on earthworms (Sheldon, 2002). On each farm a maximum of 4 fields were sampled by taking 4 $30 \times 30 \times 30$ cm soil cores on each field. Only fields with potatoes, onions, sugar beet, organic spring cereals or bare ploughed fields were sampled, as these were the dominant crop types. During the first sampling round 61 fields were sampled (33 on organic and 28 on conventional farms) and during the second round 55 fields (29 on organic and 26 on conventional farms).

Table 12 Differences in crop type between organic and conventional arable farms, showing mean relative farm area (\pm SD) with each crop and percentage of farmers growing the crop. Crop diversity is expressed as the Shannon-Wiener index. N = number of farms. PO = potatoes, SC = spring cereals, ON = onions, SB = sugar beet, WC = winter cereals, CA = carrots, BE = Belgian endive, BEA = beans, PE = peas, OC = other crops, CD = crop diversity. *** = $P < 0.001$, ** = $P < 0.005$, * = $P < 0.05$, NS = $P > 0.05$.

Year	2004					2005				
	Organic (N=10)		Conventional (N=10)		Sig.	Organic (N=20)		Conventional (N=20)		Sig.
Area (%)	Farms (%)	Area (%)	Farms (%)	Area (%)		Farms (%)	Area (%)	Farms (%)	Area (%)	
PO	19 \pm 4	100	28 \pm 6	100	*	16 \pm 9	85	27 \pm 8	95	**
SC	28 \pm 8	100	4 \pm 6	30	**	27 \pm 11	100	5 \pm 9	30	***
ON	11 \pm 7	70	11 \pm 9	70	NS	11 \pm 7	75	11 \pm 10	65	NS
SB	5 \pm 11	20	16 \pm 9	80	*	2 \pm 5	15	15 \pm 10	80	***
WC	0 \pm 0	0	15 \pm 11	70	*	0	0	12 \pm 14	50	*
CA	7 \pm 8	50	4 \pm 5	40	NS	7 \pm 8	55	4 \pm 6	35	NS
BE	1 \pm 3	10	6 \pm 8	40	NS	3 \pm 6	25	8 \pm 11	45	NS
BEA	5 \pm 7	40	3 \pm 11	10	NS	5 \pm 6	50	3 \pm 8	15	NS
PE	3 \pm 8	20	0 \pm 0	0	NS	6 \pm 8	40	1 \pm 4	15	*
OC	21 \pm 17	90	12 \pm 16	60	*	23 \pm 15	85	14 \pm 17	45	NS
CD	2.5 \pm 0.3		2.3 \pm 0.3		NS	2.6 \pm 0.5		2.2 \pm 0.4		*

The study on ground-dwelling invertebrates took place on twenty organic and twenty conventional farms. These invertebrates were sampled for one week (June 1-8) in 2004. During this period many farmland passerines have chicks (e.g. Wilson *et al.* 1997, Kragten *et al.* 2008). Ground-dwelling invertebrates were sampled using pitfall traps (diameter 11.6 cm, depth 7 cm) filled with ethylene glycol diluted with water (1:1). Pitfalls were placed in the dominant crop types of each farm (maximum of 6 crop types per farm) and, if these were present, in one grassy field margin and on one ditch bank. In all, 25 different crop types were sampled. In each plot four pitfalls were placed, separated by a distance of 15 m. Within the crops the pitfalls were placed at 15, 30, 45 and 60 m from the field edge.

Aerial invertebrates were sampled during the same period and on the same farms as ground-dwelling invertebrates were sampled. The sampling

periods coincided with the peak of the barn swallow breeding season (Evans *et al.* 2007). On each farm aerial invertebrates were sampled in the dominant crop types (maximum 6 crop types per farm) and, if present, on one ditch bank and in one field margin using two Sticky Traps (*Pherobank*®). Sticky Traps are plastic plates (25 x 10 cm) covered on both sides with non-drying and non-drip ‘insect-glue’. These traps were attached to 1 m high bamboo poles, thus protruding above the vegetation, and placed at 30 and 60 m from the field edge, where they were left for a period of 7 days. The invertebrates trapped were then counted.

Data analysis

For all three invertebrate groups (soil-dwelling, ground-dwelling and aerial) a three-part analysis of the results was carried out. First, invertebrate abundance on organic and conventional farms was compared. Second, their abundance on organic and conventional crops was compared (for example, organically versus conventionally managed cereal fields). Similar analyses were performed for non-cropped field margins and ditch banks. Third, invertebrate abundance on different crop types was compared with that found in non-crop habitats under the same type of farm management.

Mean invertebrate abundance was calculated per sampled field after $\log_{10}(x+1)$ transformation and used for statistical analysis. For the farm-level comparisons the mean value per crop per farm was multiplied by the relative area occupied by that crop on the farm concerned. In this way a correction was made for inter-farm differences in crop composition. Comparisons were only made when the number of sampled fields was at least five per farm type (Table 13). As aerial invertebrates were sampled in field margins on only four conventional farms, the data on field margins and ditch banks were pooled. In this way aerial invertebrate abundance in non-crop habitats could be compared with abundance in crops. To analyse differences in invertebrate abundance

between organic and conventional farms or crop types, General Linear Mixed Models (GLMM) with Poisson error and logarithm link function were used with farm management (organic/conventional) set as the fixed effect and farm pair as random effect.

The farms on which ground-dwelling and aerial invertebrates were sampled were located in two different regions of the study area and therefore, ‘region’ was included as a random effect in the analyses for these two groups. Because on some farms earthworms were sampled on more than one bare ploughed field, ‘farm’ was included as a random effect as well. To assess whether the sampling position in the field was of influence on invertebrate abundance, differences in invertebrate number per position were analysed by means of a Kruskal-Wallis test. No systematic effects were found and therefore sampling position was not included in the analyses. Differences in invertebrate abundance between different crop types under the same farm management were analysed using a Kruskal-Wallis test followed by a Bonferroni-like procedure as described by Neter *et al.* (1996). Analyses were carried out using Genstat 10.0 (GLMM) and SPSS 12.0 (Kruskal-Wallis).

Table 13 Number of fields sampled per crop type and per invertebrate group. -- = no fields available for sampling. Other crops are mainly vegetable crops.

Crop type	Earthworms				Aerial invertebrates		Ground surface invertebrates	
	Period 1		Period 2		O	C	O	C
	O	C	O	C				
Bare ploughed	30	23	3	6	--	--	--	--
Potatoes	--	--	6	5	15	16	18	19
Onions	1	3	8	8	8	12	14	14
Spring cereals	4	--	10	--	19	4	20	3
Sugar beet	--	3	2	8	4	12	5	17
Carrots	--	--	--	--	7	4	7	5
Winter cereals	--	--	--	--	--	9	--	11
Other crops	--	--	--	--	18	11	21	20
Field margins	--	--	--	--	10	4	13	5
Ditch banks	--	--	--	--	12	16	20	19

Results

Earthworms

Earthworm abundance was generally 2 to 4 times higher on organic farms and fields. At the farm level this difference was significant during the first sampling period only (Table 14). At the crop level, too, earthworm abundance on organic farms was generally higher than on conventional farms. During the first sampling period the difference was significant for bare ploughed fields and during the second period for all cropped fields combined (Table 14). Earthworm abundance did not differ significantly among the various crop types.

Table 14 Mean numbers (\pm SE) of earthworms caught on organic and conventional farms and fields. Significant results are indicated by * ($P < 0.05$) and ** ($P < 0.01$). -- = no fields available for sampling. No significant differences between crop types were found.

		Organic	Conventional	F
			al	
Period 1 (March 30-April 1)	Farm level	4.3 \pm 1.9	1.1 \pm 0.3	5.54 *
	Bare ploughed	4.5 \pm 1.0	1.0 \pm 0.3	11.67 **
	All crops	5.4 \pm 5.0	1.4 \pm 0.4	2.30
Period 2 (April 28-May 3)	Farm level	4.4 \pm 1.0	2.3 \pm 0.8	2.82
	Bare ploughed	--	4.6 \pm 3.5	--
	All crops	4.5 \pm 0.7	1.6 \pm 0.4	10.98 **
	Onions	4.3 \pm 0.9	2.0 \pm 0.8	2.80
	Potatoes	5.2 \pm 2.4	1.0 \pm 0.7	3.97
	Spring cereals	4.8 \pm 1.1	--	--

Ground-dwelling invertebrates

No evidence was found for greater total abundance of ground-dwelling invertebrates on organic farms (Table 15). In total, 18 taxonomic groups of

invertebrates were distinguished, with Carabidae, Diptera and Collembola predominating. At the farm level the abundance of 13 groups was greater on organic farms and for Carabidae Araneae, Aphididae, Hymenoptera and Cicadellidae these differences were significant. The abundance of 'other Coleoptera' was greater on conventional farms. For all the other groups analysed no significant effects of farm type were found.

At the crop level there was no evidence that organic management led to a greater total invertebrate abundance. However, certain individual taxonomic groups were found to more abundant in organic crops, viz.: Carabidae (cereals and potatoes), Araneae (cereals), Staphylinidae (potatoes), Formicidae (carrots) and 'other invertebrates' (carrots). In cereals, Staphylinidae and Collembola showed the opposite trend. With respect to non-crop habitats, Isopoda were found in greater numbers in organically managed field margins. In contrast, Carabidae and 'other Coleoptera' were found to be more abundant on conventionally managed ditch banks compared to organically managed ditch banks.

Within a given farm type, total ground-dwelling invertebrate abundance differed significantly between crop types (organic: Kruskal-Wallis, df 6, $\chi^2 = 16.90$, $P = 0.010$; conventional: Kruskal-Wallis, df 6, $\chi^2 = 14.74$, $P = 0.022$). On organic farms total invertebrate abundance was greatest in carrots, onions and cereals and least on ditch banks. On conventional farms total invertebrate abundance in onions exceeded that on any other crop.

Aerial invertebrates

At the farm level aerial invertebrate abundance was significantly greater on organic farms (Table 16). At the crop level all comparisons showed greater aerial invertebrate abundance on organic fields as well, but only in the case of potatoes was the difference significant. In non-crop habitats aerial invertebrates

were more abundant on organic farms, but this difference was not significant (Table 16). On both organic and conventional farms the numbers of aerial invertebrates caught differed from crop to crop (organic: Kruskal-Wallis, df 4, $\chi^2 = 19.22$, $P = 0.001$; conventional: Kruskal-Wallis, df 4, $\chi^2 = 13.70$, $P = 0.008$). On both types of farm aerial invertebrate abundance was greatest over cereal fields. On organic farms aerial invertebrates were least abundant over carrots and onions and on conventional farms over potatoes and sugar beet.

Table 15 Mean number of ground-dwelling invertebrates on organic and conventional farms and crops. *P* indicates level of significance for difference in abundance between organic and conventional plots. *** < 0.001, ** < 0.01, * < 0.05, NS not significant. Ca = Carabidae, Di = Diptera, Co = Collembola, Ot = other Coleoptera, St = Staphylinidae, Ar = Arachnida, Ap = Aphididae, Ch = Chilopoda, Hy = Hymenoptera, Fo = Formicidae, Ac = Acari, Ga = Gastropoda, Ci = Cicadellidae, Is = Isopoda, Lu = Lumbricina, Ph = Phalangida, In = other invertebrates, He = Heteroptera.

	Farm-level			Carrots			Cereals			Onions		
	O	C	F	O	C	F	O	C	F	O	C	F
Total	140.9	116.5	NS	150.0	85.2	NS	158.7	100.1	NS	133.2	187.5	NS
Ca	61.9	34.2	*	44.9	14.0	NS	79.9	17.9	*	37.7	69.5	NS
Di	23.8	29.4	NS	19.5	26.9	NS	7.4	6.0	NS	33.8	37.1	NS
Co	15.9	25.1	NS	46.0	15.5	NS	4.4	23.1	**	33.2	50.2	NS
Ot	7.8	10.0	*	10.9	12.7	NS	3.9	4.4	NS	9.3	15.0	NS
St	9.8	6.7	NS	6.8	4.3	NS	10.4	22.2	*	8.6	6.0	NS
Ar	8.1	3.9	**	4.0	2.6	NS	12.8	5.7	*	3.1	3.1	NS
Ap	6.5	1.5	***	11.3	1.9	NS	13.2	4.3	NS	1.8	1.7	NS
Ch	2.3	2.2	NS	1.5	5.2	NS	2.8	4.6	NS	1.8	2.0	NS
Hy	1.7	0.9	*	2.3	1.0	NS	1.2	1.2	NS	1.8	0.9	NS
Fo	1.2	0.9	NS	1.5	0.4	*	0.9	1.1	NS	1.2	1.0	NS
Ac	0.6	0.8	NS	0.5	0.1	NS	0.4	1.7	NS	0.3	0.2	NS
Ga	0.3	0.3	NS	0.0	0.0	NS	0.0	0.1	NS	0.0	0.0	NS
Ci	0.3	0.1	***	0.2	0.4	NS	0.1	0.1	NS	0.1	0.1	NS
Is	0.2	0.2	NS	0.2	0.0	NS	0.0	0.1	NS	0.0	0.1	NS
Lu	0.2	0.2	NS	0.0	0.1	NS	0.2	0.1	NS	0.2	0.5	NS
Ph	0.1	0.1	NS	0.2	0.0	NS	0.1	0.4	NS	0.0	0.0	NS
In	0.1	0.1	NS	0.2	0.0	*	0.0	0.1	NS	0.3	0.1	NS
He	0.0	0.0	NS	0.0	0.0	NS	0.0	0.0	NS	0.0	0.0	NS
O>C		13			11			7			8	
C>O		5			5			10			10	

	Potatoes			Sugar beet			Field margins			Ditch banks			O>C	C>O
	O	C	F	O	C	F	O	C	F	O	C	F		
Total	122.6	104.6	NS	100.6	113.8	NS	110.8	82.9	NS	71.3	83.1	NS	4	3
Ca	55.5	21.7	*	36.7	46.1	NS	20.0	21.5	NS	4.4	9.4	*	3	4
Di	33.9	45.9	NS	22.1	22.3	NS	13.2	8.0	NS	6.3	6.5	NS	2	5
Co	14.2	20.6	NS	15.3	13.4	NS	3.2	6.5	NS	4.1	6.5	NS	2	5
Ot	5.6	6.5	NS	10.7	17.6	NS	7.4	8.8	NS	3.2	5.3	**	0	7
St	5.1	2.9	*	4.0	3.4	NS	8.6	4.6	NS	4.2	4.9	NS	5	2
Ar	2.3	1.5	NS	2.3	4.6	NS	21.3	13.5	NS	12.6	10.9	NS	6	1
Ap	1.4	1.0	NS	1.7	0.9	NS	3.2	3.1	NS	2.0	1.2	NS	7	0
Ch	1.8	2.0	NS	4.5	3.3	NS	1.2	1.9	NS	1.4	2.9	NS	1	6
Hy	1.4	0.8	NS	0.9	0.7	NS	4.0	2.4	NS	3.1	2.5	NS	7	0
Fo	0.6	0.5	NS	1.4	0.8	NS	7.7	3.9	NS	7.5	8.7	NS	5	2
Ac	0.4	0.8	NS	0.5	0.3	NS	3.4	2.2	NS	3.4	4.7	NS	4	3
Ga	0.0	0.0	NS	0.0	0.0	NS	4.6	3.6	NS	7.5	10.1	NS	2	3
Ci	0.0	0.0	NS	0.0	0.0	NS	3.9	1.1	NS	3.6	3.8	NS	2	5
Is	0.0	0.0	NS	0.2	0.0	NS	1.2	0.2	**	4.8	5.9	NS	4	3
Lu	0.1	0.1	NS	0.1	0.2	NS	0.4	0.5	NS	0.7	0.7	NS	2	5
Ph	0.1	0.0	NS	0.0	0.1	NS	0.5	0.2	NS	0.4	0.3	NS	5	2
In	0.0	0.1	NS	0.0	0.0	NS	0.8	0.9	NS	0.7	0.5	NS	3	4
He	0.1	0.0	NS	0.0	0.0	NS	0.2	0.1	NS	0.2	0.2	NS	2	2
O>C		12			8			12			5			
C>O		6			8			6			13			

Table 16 Mean numbers (\pm SE) of aerial invertebrates caught on organic and conventional farms and over crops and non-crop habitats. Significant results are indicated by * ($P < 0.05$), ** ($P < 0.01$) and *** ($P < 0.001$). Letters indicate significant differences between crop types within the same type of farm management. Same letters indicate no difference.

	Organic	Conventional	F
Farm level	169.3 \pm 19.2	100.1 \pm 7.9	15.14 ***
Non-crop habitat	149.3 \pm 21.4 ^b	121.3 \pm 8.9 ^{ab}	2.62
Onions	106.4 \pm 17.7 ^c	106.1 \pm 12.1 ^{bc}	0.32
Potatoes	153.8 \pm 28.4 ^b	92.9 \pm 10.1 ^{cd}	4.87 *
Cereals	228.9 \pm 26.3 ^a	140.7 \pm 18.0 ^a	12.27 **
Carrots	99.4 \pm 9.4 ^c	--	--
Sugar beet	--	82.5 \pm 15.7 ^d	--

Discussion

This study found positive effects of organic farming on the abundance of earthworms and aerial and ground-dwelling invertebrates. Earthworm abundance was 2-4 times higher on organic farms and fields. For a given type of farm management no differences in earthworm abundance between crop types were found. The total abundance of ground dwelling invertebrates did not differ significantly between farm types, although Carabidae, Araneae, Aphididae, Hymenoptera and Cicadellidae were all more abundant on organic farms. The opposite was true for the group of 'other invertebrates'. In carrot, cereal and potato fields certain groups were found to be more abundant on organically managed farms. On organic farms ground-dwelling invertebrates were most abundant in carrots, cereals and onions. On conventional farms this held true for onion fields. Compared with most crops, ground-dwelling invertebrate abundance was low in uncropped field margins and on ditch banks. Aerial invertebrates were more abundant on organic farms. At the crop level significantly higher abundances were found in organically managed cereal and potato fields. Compared with other crop types, aerial invertebrate abundance

was greatest in cereal fields. Some caution should be applied when interpreting the results of this study as ground-dwelling and aerial invertebrates were sampled during a period of one week only.

The greater abundance of earthworms on organic farms may be due to the use of farmyard manure rather than artificial fertilizers, manure constituting an important food resource for earthworms (Pfiffner and Mäder 1997). The absence of pesticide use may also be beneficial, especially for earthworms close to the soil surface (Pfiffner and Mäder, 1997). Many taxonomic groups of ground-dwelling invertebrates were found in greater abundance on organically managed compared with conventionally managed sites. Possible causes of these differences are absence of pesticide use, richer understory vegetation and increased food supply (Hole *et al.* 2005), but these parameters were not measured during the present study. The effects of organic farming on ground-dwelling invertebrates were somewhat inconsistent among crop types. This may be due to differences in farming practice among the various crops, including differences in tillage and pesticide application (Hole *et al.* 2005).

The greater abundance of aerial invertebrates on organic farms was probably caused by the absence of pesticide use and by higher inputs of organic fertilizers. Pesticides inputs are known to have damaging effects on invertebrate populations (e.g. Aebischer 1990, Anderson and Lydy 2002). Higher inputs of organic material are known to have positive effects on the numbers of decomposers like many Diptera species (Smeding and de Snoo 2003).

Landscape composition is known to affect differences in invertebrate abundance and diversity between organic and conventional farming systems (Purtauf *et al.* 2005, Schmidt *et al.* 2005, Holzschuh *et al.* 2007). In this study, the surrounding landscape composition was part of the farm pairing protocol. Besides this, landscape composition of the entire study area was relatively homogeneous. Therefore, it is unlikely that this has been of influence on the results. On-farm landscape composition differed largely as a result of different

crop rotation systems (Table 12), but this is a direct effect of different farm management strategies.

Several studies have shown the importance of grassy or herbaceous field margins as foraging sites for birds (e.g. Marshall and Moonen 2002). In our study the abundance of ground-dwelling invertebrates in field margins was much lower than in crops. This could be a bias resulting from the sampling method adopted. With pitfalls, insect activity density is measured (e.g. Winder *et al.* 2005). Invertebrate activity depends on food availability, vegetation structure and micro-climatic conditions, and therefore comparing invertebrate abundance between different habitats using pitfalls could be biased. Also comparisons between crops could suffer to some extent from this bias, so some caution in interpreting these results should be taken.

Although overall abundance of ground-dwelling invertebrates was generally lower in uncropped habitats, certain taxonomic groups were more abundant here than in crops (Table 15). Groups characteristic of stable habitats, in particular, were more abundant in field margins and on ditch banks. In addition, most of these groups are detritivorous, hydrophilous or associated with dense vegetation structures.

Implications for birds

In 2004 and 2005 breeding bird surveys for field-breeding species were carried out on the same farms as where the invertebrates surveys took place. Besides that, barn swallow nests were counted on all these farms during spring 2005. Skylark and lapwing were both breeding in higher densities on organic farms (Kragten and de Snoo 2008), but there was no difference in the number of breeding barn swallows between the two farming types (Kragten *et al.* 2009). In contrast to skylarks, breeding densities of other species feeding mainly on ground-dwelling invertebrates, such as yellow wagtail *Motacilla flava* and

meadow pipit *Anthus pratensis*, did not differ between the two farm systems. For lapwing indications were found that differences in food abundance could play a role in this. Lapwing densities were higher on organic onion fields compared to conventional onion fields, probably an effect of differences in crop management and consequently food abundance. However, for skylarks no such indications were found. Differences in skylark densities were mainly caused by differences in crop rotation schemes between the two farming systems.

The present study shows that lapwing food (earthworms) is indeed more abundant on organically managed farms. In addition to this, Baines (1990) found correlations between lapwing densities and food abundance. This reinforces the hypothesis that higher lapwing densities on organic farms are due to greater earthworm abundance. Besides lapwing, other species feeding on earthworms (e.g. blackbird, *Turdus merula*, song thrush, *Turdus philomelos*) are also likely to benefit from organic farm management.

Carabidae, Araneae and Aphididae are all relatively abundant in the diets of farmland birds (Wilson *et al.* 1999, Holland *et al.* 2006). These groups were found in greater abundance on organic farms. However, no indications were found that these differences have caused differences in densities of birds feeding on these prey items (Kragten and de Snoo 2008). It is likely, though, that skylarks and other insectivorous passerines will benefit from the greater food abundance in terms of improved breeding success (e.g. Boatman *et al.* 2004, Hart *et al.* 2006).

Christensen *et al.* (1996) found greater numbers of barn swallows flying over organically managed fields than conventionally managed fields. In the Netherlands however, no difference was found in the number of breeding barn swallows (Lubbe and de Snoo 2007, Kragten *et al.* 2009). The number of breeding barn swallows is probably correlated with the availability of suitable breeding sites. This is likely to be the same on organic and conventional farms, as the types of building on both are more or less equivalent (Lubbe and de Snoo

2007). However, the number of foraging swallows shows a positive relationship with prey abundance (Evans *et al.* 2007). Greater food abundance may therefore result in improved barn swallow breeding success, but is not likely to result in higher breeding densities.

Especially for birds feeding on ground-dwelling invertebrates, greater food abundance does not necessarily mean greater food availability as well, for differences in sward structure can lead to differences in availability even if food is equally abundant. Dense, high swards generally limit accessibility (Atkinson *et al.* 2005) and many birds prefer to forage in short swards (Devereux *et al.* 2004). As organic farms grow more spring-sown crops compared with conventional farms (e.g. Hole *et al.* 2005, Kragten and de Snoo 2008), swards are generally lower and food accessibility therefore probably higher.

The present study shows that food abundance for insectivorous breeding farmland birds is higher on organically managed arable farms. It is likely that this will result in higher adult survival rates, breeding success and better fledgling body condition of breeding birds on organic farms. Therefore, organic farming systems could potentially be beneficial for farmland bird populations. However, data on these topics are scarce and therefore future studies should focus on this.

Acknowledgements

We are grateful to all farmers who gave us permission to work on their land. Bart Goossen and Chantal Kersten assisted during the field work. Nigel Harle edited the English. The comments of two anonymous referees seriously improved the manuscript.

References

- Aebischer, N. J., 1990. Assessing pesticide effects on non-target invertebrates using long-term monitoring and time-series modelling. *Functional Ecology* 4: 369-373.
- Anderson, T. D., Lydy, M. J., 2002. Increased toxicity to invertebrates associated with a mixture of atrazine and organophosphate insecticides. *Environmental Toxicity and Chemistry* 21: 1507-1514.
- Atkinson, P. W., Fuller, R. J., Vickery, J. A., Conway, G. J., Tallowin, J. R. B., Smith, R. E. N., Haysom, K. A., Ings, T. C., Asteraki, E. J., Brown, V.K., 2005. Influence of agricultural management, sward structure and food resources on grassland field use by birds in lowland England. *Journal of Applied Ecology* 42: 932-942.
- Baines, D., 1990. The roles of predation, food and agricultural practice in determining the breeding success of the lapwing (*Vanellus vanellus*) on upland grasslands. *Journal of Animal Ecology* 59: 915-929.
- Bakker, J., 2007. Bio-Monitor Jaarrapport 2006. Biologica, Utrecht.
- Beecher, N. A., Johnson, R. J., Brandle, J. R., Case, R. M., Young, L.J., 2002. Agroecology of birds in organic and nonorganic farmland. *Conservation Biology* 16: 1620-1631.
- Benton, T. G., Bryant, D. M., Cole, L., Crick, H.Q.P., 2002. Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology* 39: 673-687.
- Boatman, N. D., Brickle, N. W., Hart, J. D., Milsom, T. P., Morris, A. J., Murray, A. W. A., Murray, K. A., Robertson, P. A., 2004. Evidence for the indirect effects of pesticides on farmland birds. *Ibis* 146 (suppl. 2): 131-143.
- Chamberlain, D. E., Wilson, J.D., Fuller, R.J., 1999. A comparison of bird populations on organic and conventional farm systems in southern Britain. *Biological Conservation* 88: 307-320.
- Chamberlain, D. E., Fuller, R. J., Bunce, R. G. H., Duckworth, J. C., Shrubbs, M., 2000. Changes in the abundance of farmland birds in relation to the timing of agricultural intensification in England and Wales. *Journal of Applied Ecology* 37: 771-788.

- Christensen, K. D, Jacobsen, E. M., Nøhr, H., 1996. A comparative study of bird faunas in conventionally and organically farmed areas. *Dansk Ornitologisk Forenings Tidsskrift* 90: 21-28.
- Devereux, C. L., McKeever, C. U., Benton, T. G., Whittingham, M. J., 2004. The effect of sward height and drainage on Common Starlings *Sturnus vulgaris* and Northern Lapwings *Vanellus vanellus* foraging in grassland habitats. *Ibis* 146: 115-122.
- Donald, P. F, Green, R. E., Heath, M. F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proceedings of the Royal Society B* 268: 25-29.
- Donald, P. F., Sanderson, F. J., Burfield, I. J., van Bommel, F.P.J., 2006. Further evidence of continent-wide impacts of agricultural intensification on European farmland birds, 1990-2000. *Agriculture, Ecosystems and Environment* 116: 189-196.
- Evans, K. L., Wilson, J. D., Bradbury, R.B., 2007. Effects of crop type and aerial invertebrate abundance on foraging barn swallows *Hirundo rustica*. *Agriculture, Ecosystems and Environment* 122: 267-273.
- Hart, J. D., Milsom, T. P., Fisher, G., Wilkins, V., Moreby, S. J., Murray, A. W. A., Robertson, P. A., 2006. The relationship between yellowhammer breeding performance, arthropod abundance and insecticide applications on arable farmland. *Journal of Applied Ecology* 43: 81-91.
- Hole, D. G., Perkins, A. J., Wilson, J. D., Alexander, I. H., Grice, P. V., Evans, A.D., 2005. Does organic farming benefit biodiversity? *Biological Conservation* 122: 113-130.
- Holland, J. M., Hutchison, M. A. S., Smith, B., Aebischer, N. J., 2006. A review of invertebrates and seed-bearing plants as food for farmland birds in Europe. *Annals of Applied Biology* 148: 49-71.
- Holzschuh, A., Steffan-Dewenter, I., Kleijn, D., Tschardtke, T., 2007. Diversity of flower-visiting bees in cereal fields: effects of farming system, landscape composition, and regional context. *Journal of Applied Ecology* 44: 41-49.
- Kragten, S., de Snoo, G. R., 2008. Field-breeding birds on organic and conventional arable farms in the Netherlands. *Agriculture, Ecosystems and Environment* 126: 270-274.

- Kragten, S., Reinstra, E., Gertenaar, E., 2009. Breeding barn swallows *Hirundo rustica* on organic and conventional arable farms in the Netherlands. *Journal of Ornithology* 150: 515-518.
- Kragten, S., Trimbos, K. B., de Snoo, G. R., 2008. Breeding activity of skylarks (*Alauda arvensis*) on organic and conventional arable farms in The Netherlands. *Agriculture, Ecosystems and Environment* 126: 163-167.
- Krebs, J. R., Wilson, J. D., Bradbury, R. B., Siriwardena, G. M., 1999. The second silent spring? *Nature* 400: 611-612.
- Lubbe, S. K., de Snoo, G. R., 2007. Effect of dairy farm management on Swallow *Hirundo rustica* abundance in The Netherlands. *Bird Study* 54: 176-181.
- Marshall, E. J. P., Moonen, A. C., 2002. Field margins in northern Europe: their functions and interactions with agriculture. *Agriculture, Ecosystems and Environment* 89: 5-21.
- Medley, K. E., Okey, B. W., Barrett, G. W., Lucas, M. F., Renwick, W. H., 1995. Landscape change with agricultural intensification in a rural watershed, southwestern Ohio, U.S.A. *Landscape Ecology* 10: 161-176.
- Neter, J., Kutner, M. H., Nachtsheim, C. J., 1996. *Applied Linear Statistical Models*. Times Mirror Higher Education Group, Chicago.
- Pfiffner, L., Mäder, P., 1997. Effects of biodynamic, organic and conventional production systems on earthworm populations. *Biological Agriculture and Horticulture* 15: 3-10.
- Purtauf, T., Roschewitz, I., Dauber, J., Thies, C., Tschardtke, T., Wolters, V., 2005. Landscape context of organic and conventional farms: Influences on carabid beetle diversity. *Agriculture, Ecosystems and Environment* 108: 165-174.
- Robinson, R. A., Sutherland, W. J., 2002. Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology* 39: 157-176.
- Schmidt, M. H., Roschewitz, I., Carsten, T., Tschardtke, T., 2005. Differential effects of landscape and management on diversity and density of ground-dwelling farmland spiders. *Journal of Applied Ecology* 42: 281-287.

Sheldon, R. D., 2002, Factors affecting the Distribution, Abundance and Chick Survival of the Lapwing (*Vanellus vanellus*). Harper Adams University College, Newport.

Smeding, F. W., de Snoo, G. R., 2003. A concept of food-web structure in organic arable farming systems. *Landscape and Urban Planning* 65: 219-236.

Stoate, C., Boatman, N. D., Borralho, R. J., Rio Carvalho, C., de Snoo, G. R., Eden, P., 2001. Ecological impacts of arable intensification. *Journal of Environmental Management* 63: 337-365.

Wilson, J. D., Evans, J., Browne, S. J., King, J. R., 1997. Territory distribution and breeding success of skylarks *Alauda arvensis* on organic and intensive farmland in southern England. *Journal of Applied Ecology* 34: 1462-1478.

Wilson, J. D., Morris, A. J., Arroyo, B. E., Clark, S. C., Bradbury, R. B., 1999. A review of the abundance and diversity of invertebrate and plant food of granivorous birds in northern Europe in relation to agricultural change. *Agriculture, Ecosystems and Environment* 75: 13-30.

Winder, L., Alexander, C. J., Holland, J. M., Symondson, W. O. C., Perry, J. N., Woolley, C., 2005. Predatory activity and spatial pattern: the response of generalist carabids to their aphid prey. *Journal of Applied Ecology* 74: 443-454.

Wretenberg, J., Lindström, Å., Svensson, S., Thierfelder, T., Pärt, T., 2006. Population trends of farmland birds in Sweden and England: similar trends but different patterns of agricultural intensification. *Journal of Applied Ecology* 43: 1110-1120.

