



Universiteit
Leiden
The Netherlands

Regional LCA in a global perspective

Wegener Sleeswijk, A.

Citation

Wegener Sleeswijk, A. (2010, September 2). *Regional LCA in a global perspective*. Uitgeverij BOX Press, Oisterwijk. Retrieved from <https://hdl.handle.net/1887/15921>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/15921>

Note: To cite this publication please use the final published version (if applicable).

REGIONAL LCA IN A GLOBAL PERSPECTIVE

Colophon

Regional LCA in a global perspective
A basis for spatially differentiated environmental life cycle assessment

PhD thesis Leiden University, The Netherlands

annekesleeswijk@globright.nl

This work has been made possible by the financial support of Unilever.

ISBN 978-90-8891-187-3

Cover: world map 'political world' (Central Intelligence Agency, World Factbook, online version 2010); 'Earth in a Box' (iStockphoto); T-O map from the Etymologiae of Isidorus, 1472 (Wikimedia Commons)

Cover design: Anneke Wegener Sleeswijk/
Proefschriftmaken.nl | | Printyourthesis.com

Printed by: Proefschriftmaken.nl | | Printyourthesis.com

Published by: Uitgeverij BOX Press, Oosterwijk, The Netherlands

© 2010 by Anneke Wegener Sleeswijk, except for the chapters 2 to 6. Copyrights of these chapters belong to the publishers as noted in the beginning of each chapter.

Regional LCA

in a global perspective

**A basis for spatially differentiated
environmental life cycle assessment**

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden
op gezag van de Rector Magnificus prof. mr. P.F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op donderdag 2 september 2010
klokke 16.15 uur

door

Anneke Wegener Sleeswijk

geboren te Naarden in 1960

Promotiecommissie

PROMOTOR

Prof. dr. H.A. Udo de Haes (Universiteit Leiden)

CO-PROMOTOR

Dr. G. Huppes (Universiteit Leiden)

OVERIGE LEDEN

Prof. dr. W.J.G.M. Peijnenburg (Universiteit Leiden)

Prof. dr. G.R. de Snoo (Universiteit Leiden)

Prof. dr. K. Blok (Universiteit Utrecht)

Prof. dr. M.L. Diamond (University of Toronto, Canada)

Contents

Preface	9
Synopsis	11
1 General introduction	21
1.1 Introduction	21
1.2 Fate	23
1.3 Human intake.....	27
1.4 Effect.....	28
1.5 LCA characterisation methods.....	29
1.6 Normalisation	30
1.7 Environmental parameters	30
1.8 Goal of this thesis	31
References	31
2 General prevention and risk minimization	39
2.1 Introduction	40
2.2 Risk minimization and general prevention in the context of LCA.....	41
2.3 Combining general prevention with risk minimization.....	45
2.4 A methodological framework.....	48
2.5 Conclusion.....	53
References	54
3 HERA and LCA	57
3.1 Introduction	58
3.2 Level 1: basic equations.....	60
3.3 Level 2: overall model structure.....	61
3.4 Level 3: applications.....	67

3.5	Discussion	68
3.6	Conclusions	71
	References	72
	Appendix: Mathematical analysis of HERA and LCA	76
4	Metals in the ocean	81
4.1	Introduction	81
4.2	The GLOBOX model	83
4.3	Adaptations	84
4.4	Results and discussion	88
	References	90
5	GLOBOX	93
5.1	Introduction	94
5.2	The GLOBOX model	97
5.3	Fate	99
5.4	Human intake	102
5.5	Toxic impacts	103
5.6	Results for nitrobenzene	106
5.7	Discussion	118
	Appendix: Supplementary data	125
	References	125
6	Normalisation in LCA	131
6.1	Introduction	132
6.2	General methodological choices	134
6.3	Guidelines for data source prioritisation and data estimation	138
6.4	Results	142
6.5	Discussion	147
6.6	Conclusions	151
	References	152

7	General discussion and conclusions.....	159
7.1	Introduction	159
7.2	Actual versus potential impacts in connection to LCA and RA	159
7.3	LCA characterisation factors for metals.....	162
7.4	Regional differentiation.....	164
7.5	Normalisation	166
7.6	Main achievements and conclusions	168
	References	170
	Samenvatting.....	175
	Bibliography.....	189
	Curriculum Vitae.....	191

Preface

When I started writing this PhD thesis I had already been working on LCA for a number of years. I contributed to the development of LCA methodology as well as to a number of LCA case studies. During these years, I got well acquainted with this tool, both with respect to its strengths and with respect to what I considered its limitations.

The strengths of LCA, as I saw them, were in its ‘looking behind the obvious’ with respect to environmental impacts of products over their entire life cycles, in accounting for the a large spectrum of environmental impacts, in describing the connection between environmental interventions on the one hand and products on the other in an exact manner, in estimating the quantitative relationships between environmental interventions and their impacts, and in preventing dilution from being considered as a solution for pollution. The quantitative aspects of LCA intrigued me: how could we get it right? And what is ‘right’?

The limitations that struck me were the points at which I felt we were not yet right in our quantification. These points mainly concerned LCA toxicity assessment: our lack of a measure for ‘actual’ toxic impacts (beside the potential ones), the fact that metal emissions heavily dominated the toxicity impact scores in LCA, while experts stated this corresponded in no way to their relative environmental harmfulness, and the fact that the assessment of environmental impacts did not account for regional differences, even though the range of processes of a single product life cycle might span the world. These were the three aspects that I felt I should work on to get them right, or at least more right. I was lucky to get the freedom to address all these issues – and one more – in a PhD-project on environmental fate modelling in the context of LCA toxicity assessment.

In 2006, I was involved in a project on LCA normalisation, a subject which I had only had superficial attention for, despite its quantitative character. While working on this project, I discovered an interesting methodological issue, and felt we should adapt our methodology and introduce a new principle, concerning the definition of the reference emissions. Again, I was lucky that there was support for my ideas to include this principle in our normalisation study, and to include the normalisation study in my PhD thesis. With this, I broadened the scope of my thesis from mere LCA toxicity characterisation to life cycle impact assessment as such, be it that the overall focus is still on toxicity assessment. The global character links the different aspects together.

LCA toxicity assessment cannot replace human and environmental risk assessment, not even in a spatially differentiated form. Risk assessment tools are designed for the assessment of ambient concentration dependent effects, which are not part of LCA. With this, risk assessment tools can help assess whether processes in the product life cycle meet environmental standards, and whether they can be considered as environmentally responsible. What LCA toxicity assessment can add is an answer to the question which product alternative is optimal with respect to overall environmental burdening. Spatial differentiation can help to model this as well as possible.