



Universiteit
Leiden
The Netherlands

Maternal reflective functioning: influence on parenting practices and the early development of externalizing behavior

Smaling, H.J.A.

Citation

Smaling, H. J. A. (2017, March 14). *Maternal reflective functioning: influence on parenting practices and the early development of externalizing behavior*. Retrieved from <https://hdl.handle.net/1887/46723>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/46723>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/46723> holds various files of this Leiden University dissertation.

Author: Smaling, H.J.A.

Title: Maternal reflective functioning: influence on parenting practices and the early development of externalizing behavior

Issue Date: 2017-03-14

Chapter 2

The Mother-Infant Neurodevelopment Study (*MINDS*) – Leiden: Background, design and study population.

Smaling, H. J. A., Suurland, J., Huijbregts, S. C. J., Van der Heijden, K. B., Van Goozen, S. H. M., & Swaab, H. The Mother-Infant Neurodevelopment Study (*MINDS*) – Leiden: Background, design and study population. *Under review*.

Chapter 2

Abstract

This paper describes the background, design, and sample characteristics of the Mother-Infant Neurodevelopment Study – Leiden, a longitudinal study investigating 1) mechanisms through which neurobiological, neurocognitive and social-environmental factors increase risk for emotional and behavioral problems in early childhood, 2) effects of an intensive home-visiting program for first-time mothers at high-risk on child neurobiological, neurocognitive and emotional and behavioral development, and 3) which factors (neurobiological, neurocognitive and social-environmental) predict variation in effects of the home-visiting program on child emotional and behavioral outcomes. A total of 275 families (128 low-risk and 147 high-risk) were included in the study. High-risk women were randomly assigned to the intervention ($n = 65$) or high-risk control group ($n = 82$). Six assessment waves were conducted within a four-year period. Demographic and mental health characteristics of the low-risk and high-risk group, collected during the first assessment at 27 gestational weeks, are presented. This study will help identifying specific biomarkers, precursors of neurocognitive functions and temperamental factors in infancy, facilitating the detection of children at risk for later emotional and behavior problems. Furthermore, this study may yield insights into effective, targeted, and tailor-made components of prevention programs, ultimately reducing the psychological and economic costs of mental health problems to society.

Introduction

Children growing up in families struggling with multiple complex issues, including maternal psychiatric problems, substance (ab)use, single parenthood, and poverty, are at high risk for developing emotional and behavioral problems (Cabaj, McDonald, & Tough, 2014; Côté, Vaillancourt, LeBlanc, Nagin, & Tremblay, 2006; Hay, Mundy, et al., 2011; Huijbregts, Seguin, Zoccolillo, Boivin, & Tremblay, 2008). As these problems raise major public health concerns and increase costs to society, it is important to gain insight into developmental mechanisms and effectiveness of prevention approaches. Recent theoretical models have emphasized the complex interactions between neurobiological vulnerabilities and environmental risk factors (Beauchaine, Neuhaus, Brenner, & Gatzke-Kopp, 2008; Belsky & Pluess, 2009; Boyce & Ellis, 2005), and the mediating role of neurocognitive and neurobiological factors in the link between early adversity and emotional and behavioral outcomes (Van Goozen, Fairchild, Snoek, & Harold, 2007).

However, there are several important gaps in the current literature. First, most studies so far have involved toddlers, school-aged children and adolescents, and little is known about earlier manifestations of neurobiological and neurocognitive vulnerabilities, and how these might increase risk for emotional and behavioral problems later in life. This is particularly important given that the neurobiological systems underlying later emotional and behavioral regulation rapidly develop during the prenatal period and first years of life, resulting in increased sensitivity to environmental influences (Beauchaine et al., 2008; Laurent, Harold, Leve, Shelton, & Van Goozen, 2016). Prevention efforts initiated early in life are therefore presumed to be more effective (Beauchaine et al., 2008). Second, most longitudinal studies investigating the neurobiological and neurocognitive mechanisms underlying emotional and behavioral problems in early childhood involved community samples. Because evidence is accumulating that shows that different neurodevelopmental processes underlie emotional and behavioral development in normative versus high-risk or clinical samples (Beauchaine, 2001; Beauchaine et al., 2008), it is of critical importance to extend the current body of literature with studies focusing on high-risk samples in order to fully understand the mechanisms that are related to the development of emotional and behavioral problems.

The overarching aim of the Mother-Infant Neurodevelopment Study (*MINDS*) – Leiden study is to examine which neurobiological, neurocognitive and social-environmental factors

Chapter 2

increase risk or confer protection for developing emotional and behavioral problems in the first years of life in a heterogeneous sample of low- and high-risk families. Based on integrative models of the neurobiological bases of early-onset antisocial behavior (Van Goozen et al., 2007), and theories of differential susceptibility (Belsky & Pluess, 2009) and biological sensitivity to context (Boyce & Ellis, 2005), we considered neurobiological and neurocognitive factors as potential mediating and moderating mechanisms that lead to emotional and behavioral problems. We adopted a rigorous and systematic approach, assessing a wide range of social-environmental (i.e., maternal psychiatric problems, substance (ab)use, poverty, social support, parenting, maternal reflective functioning), neurocognitive (i.e., precursors of executive functioning, Theory of Mind, language, empathy) and neurobiological factors (i.e., autonomic nervous system and hypothalamic pituitary adrenal axis) that may either directly, indirectly, or in interaction increase risk for emotional and behavioral problems in young children.

Families at risk need substantial support to promote good enough care for their child. However, it is often difficult to engage these families in intervention programs. Home-visiting programs for first-time mothers at high-risk have the advantage of serving families at their home, thereby increasing the likelihood that they will (continue to) participate. Home-visiting programs have been found to be effective in improving maternal prenatal health behaviors (e.g., reductions in tobacco use), maternal life course (e.g., fewer rapid second pregnancies, returning to school/ seeking education), sensitive parenting behavior and parenting attitudes, and child physical abuse (e.g., number of emergency room visits, injuries or ingestions treated, and accidents requiring medical attention) (Mejdoubi et al., 2014; Mejdoubi et al., 2015; Olds, Sadler, & Kitzman, 2007; Ordway et al., 2014; Sweet & Appelbaum, 2004). Further, positive effects of home-visiting programs have been reported for a wide range of child outcomes (Avellar & Supplee, 2013; Filene, Kaminski, Valle, & Cachat, 2013; Mejdoubi et al., 2015; Olds et al., 2007; Ordway et al., 2014; Peacock, Konrad, Watson, Nickel, & Muhajarine, 2013; Sweet & Appelbaum, 2004). For example, home-visiting programs have been found to positively affect cognitive development (e.g., academic and arithmetic achievement, intellectual functioning, executive functioning, receptive language), socio-emotional development (e.g., attachment security, social development, emotional vulnerability), and to reduce behavioral problems (e.g., externalizing and internalizing behavior problems). Also, favorable results have been reported for birth outcomes (e.g., birth weight, gestational age) and physical health, although these effects have been less consistent among different home-visiting programs

(Avellar & Supplee, 2013; Filene et al., 2013; Robling et al., 2016). Moreover, at 15 year follow-up, adolescents who had received home visitation early in life had fewer arrests and convictions, and committed fewer violations of probation (Olds et al., 1998).

Despite these promising results, the overall effect sizes of studies evaluating home-visiting programs are relatively small (Filene et al., 2013), and many studies report non-significant findings (Avellar & Supplee, 2013). Currently, an important question is which factors predict this variability in the effects of home-visiting programs on child outcomes? While it is generally acknowledged that effectiveness of prevention programs may vary as a function of neurobiological and neurocognitive vulnerabilities (Beauchaine et al., 2008), there are no studies that have examined child neurobiological and neurocognitive factors as moderators of the effects of home-visiting programs. Moreover, surprisingly little is known about the effects of home-visiting programs on child neurobiological and neurocognitive development. Therefore, a second overarching aim of *MINDS* – Leiden was to evaluate the effects of an intensive home-visiting program for first-time mothers at high-risk on child neurobiological, neurocognitive and emotional and behavioral development, and to evaluate the predictive value of these child neurobiological and neurocognitive factors on the effectiveness of such a program.

Study aims

MINDS – Leiden is part of a large research program funded by the Dutch government called ‘Brain & Cognition – Social innovation in health, education, and safety’ (<http://www.nwo.nl/en/research-and-results/programmes/nihc/hcmi/index.html>). The purpose of this Brain & Cognition program is to promote neurobiological and neurocognitive research to develop and improve intervention efforts aimed at reducing aggression and violence in society.

In the present study three important questions are addressed. First, which neurobiological and neurocognitive factors predict (directly, indirectly or in interaction with social-environmental factors) emotional and behavioral problems – specifically aggressive behavior – in the first three-and-a-half years of life? Second, what are the effects of an intensive home-visiting program for first-time mothers at high-risk on neurobiological, neurocognitive and emotional and behavioral development in children in the first years of life? Third, which factors (neurobiological, neurocognitive and social-environmental) predict variation in effects of the home-visiting program on child

Chapter 2

emotional and behavioral outcomes? This paper describes the design of the study, the measures used, the intervention program and the sample characteristics.

Methods

Study design

The *MINDS* – Leiden study was designed as a longitudinal randomized control trial (see Figure 1 for an overview of the design of the study). Based on elaborate screening for the presence of risk factors during the first prenatal home visit (27 weeks gestation), pregnant women were assigned to either the high-risk (HR) or low-risk (LR) group (see *screening for risk factors* for a description of allocation criteria). Women in the HR group were randomly assigned to the high-risk intervention (HR-I) group (see *The intervention* for more details) or the high-risk control (HR-C) group. All participating families were followed over a period of approximately four years, consisting of six assessment waves, (the first assessment took place in the third trimester of pregnancy and the last assessment took place when the children were 3.5-4 years of age). A total of 65 families were included in the HR-I group. Families in the HR-I group took part in a home-visiting program starting within 2 weeks after the prenatal assessment until 2.5 years post-partum.

Recruitment

Recruitment of pregnant women took place between February 2011 and April 2015, via hospitals, midwifery clinics, prenatal classes, pregnancy fairs, and social workers. Dutch-speaking primiparous women between 17 and 25 years old with uncomplicated pregnancies were eligible to participate. Exclusion criteria were heavy drug addiction or severe psychiatric or psychotic disorder, an intelligence quotient (IQ) below 70, major acute or significant chronic illness in the mother or a disorder or syndrome in the child, which would affect normal development. We oversampled families from a high-risk background in order to obtain sufficient variance in risk factors that might influence children's early socio-emotional and cognitive development. This was done by collaborating with midwifery/obstetric clinics in areas with a low average social-economic status and/or by recruitment through social workers. All participating women provided written informed

The Mother-Infant Neurodevelopment Study – Leiden

consent. The study was approved by the Medical Research Ethics Committee at the Leiden University Medical Centre (NL39303.058.12), and by the ethics committee of the Department of Education and Child Studies at the Faculty of Social and Behavioral Sciences, Leiden University (ECPW-2011/025).

Screening for risk factors

Classification to the HR-group was based on the following risk factors (Smaling et al., 2015; World Health Organization, 2005, 2016): 1) positive screening on current psychiatric disorder(s) or substance use (alcohol, tobacco and/or drugs) during pregnancy; or 2) presence of two or more of the following risk factors: single status (biological father not involved), unemployment, financial problems, no secondary education, limited social support network (<4 individuals listed in network), and young maternal age (<20 years). In case only one risk factor was present - other than an indication for current psychiatric disorder(s) or substance use - women were discussed in a clinical expert meeting to determine whether placement in the HR-group was appropriate ($n = 6$).

Positive screening on current psychiatric disorder(s) was established by the Mini-International Neuropsychiatric Interview – plus (M.I.N.I. - plus; Sheehan et al., 1997; Van Vliet, Leroy, & Van Megen, 2000) by screening for the following disorders: depressive disorder (current), dysthymic disorder (past 2 years), suicidality, mania (current), panic disorder (current and lifetime), agoraphobia (current), social phobia (current), other type of phobia (current), obsessive-compulsive disorder (current), generalized anxiety disorder (GAD; current), posttraumatic stress disorder (current), alcohol dependence and abuse (current and lifetime), drug dependence and abuse (non-alcohol, current and lifetime), attention-deficit/hyperactivity disorder (lifetime), and antisocial personality disorder (lifetime). Current is defined as 'in the past month' for all diagnoses except GAD, which has a 6-month time frame, and alcohol abuse/dependence and drug abuse/dependence for which a 12-month time frame is used.

The size of the social support network was established by using the Norbeck Social Support Questionnaire (NSSQ; Norbeck, Lindsey, & Carrieri, 1981, 1983). The presence of other risk factors (i.e., substance use during pregnancy, no secondary education, unemployment, financial problems, single status, and maternal age) were assessed by means of the Dutch translation of the 'Becoming a mother' questionnaire (Hay et al., 2011; Smaling et al., 2015).

Chapter 2

Procedures

The study comprised of six assessment waves (see Figure 1). During the third trimester of pregnancy (T1), at 6 months (T2) and at 20 months (T4) post-partum 2-to-2.5 hour home visits were carried out by two female researchers. One leading researcher conducted all the tasks with the infant and guided the mother-infant interaction tasks, while a second researcher digitally recorded the whole session and administered the questionnaires to the mother. At 12 months (T3), 30 months (T5) and 40 months (T6), the mother-infant dyads visited the Babylab at the Faculty of Social and Behavioral Sciences, Leiden University. The lab test sessions were carried out by one researcher, while a second researcher was seated behind a one-way screen, recorded the session and provided observational records.

Table 1 gives an overview of the main research areas of the study. In addition, Table 2 gives a detailed overview of instruments and its content used in the study. T1 included an interview regarding the emotional impact of the pregnancy to assess prenatal reflective functioning (RF), a structured interview to assess current psychiatric disorders, and various questionnaires to assess demographic information, mental and physical health, life style, social support, self-efficacy in the nurturing role, executive functioning, emotion regulation, life events, and antenatal attachment. T2 to T5 started with a free play session to measure maternal interactive behavior. For T2, T3, and T5, this was followed by attachment of the cardiac monitoring equipment to the child to measure their autonomic nervous system (ANS) reactivity and regulation during mildly stressful events (see Table 2 for the paradigms that were used to measure stress reactivity and regulation). In addition, during the lab visits saliva samples were taken several times (before and after stress) from mother and child to measure hypothalamic pituitary adrenal (HPA)-axis functioning. Moreover, each wave consisted of various age-appropriate tasks to measure precursors of executive functioning, temperament and emotion, empathy, Theory of Mind, and language skills, and ended with the mother completing various questionnaires to assess demographic information and multiple mother (i.e., mental and physical health, obstetric characteristics, life style, social support, self-efficacy in the nurturing role, executive functioning, emotion regulation, life events, parenting cognitions and parenting stress) and child domains (language, temperament, aggression, behavioral problems, executive functioning) (see Table 1 and 2 for more details). Mother-child interaction during normal, playful interaction, teaching tasks and following mildly stressful events was observed at T2 to T5. T4 also included an interview regarding maternal representations of the relationship with their child and the emotional impact of parenting to measure postnatal RF. At T6, children

were assessed for approximately 1 hour in individual testing rooms, and then brought together with their mother and one or two other families for a simulated birthday party and a 20 minute free play session with peers, designed to provide an acceptable yet emotionally arousing setting in which to observe children's social behavior (and specifically aggressive behavior) with their peers.

The intervention

Families in the HR-I group ($n = 65$) participated in an intensive home-visiting program based on 'Minding the Baby' (MTB) (Sadler et al., 2013; Slade et al., 2005; Slade, Sadler, & Mayes, 2005). MTB is an interdisciplinary home-visiting program developed at Yale University Child Study Center and Yale University School of Nursing (Slade et al., 2010). MTB focuses on young vulnerable first-time parents, primarily mothers, who are at high risk due to multiple complex issues, including psychiatric problems, poverty, young maternal age, single motherhood, or limited social support. MTB combines two well-researched early-intervention models; home visiting and infant-parent-psychotherapy, in order to meet the holistic, complex, multilayered care needs of vulnerable families (Sadler et al., 2013). The program specifically aims to enhance maternal reflective functioning (RF) and the development of secure attachment relationships, as well as to address maternal (mental) health issues. RF refers to the mother's capacity to 'keep the baby in mind', to make sense of his/her internal states, emotions, thoughts, and intentions, as well as her own (Slade, 2002). Particularly in high-risk mothers, RF is often compromised, leading to disrupted interactions, insecure attachment relationships, and long term emotional difficulties. In the present study (see Table 3), RF skills were significantly lower among women in the HR-group compared to women in LR-group. In MTB, parents are encouraged to be curious, to try and figure out what the child needs or is thinking or feeling even in early infancy. RF is seen as a key to maternal sensitivity and plays an important role in the development of the child's capacity for Theory of Mind and adaptive socio-emotional development in young children (Ordway et al., 2014; Laranjo, Bernier, Meins, & Carlson, 2010; Sadler et al., 2013; Sadler, Slade, & Mayes, 2006; Slade, Sadler, & Mayes, 2005). Programs that are specifically aimed at improving parental RF in 'at-risk' parents indeed appear to improve RF-skills and parenting behavior (Katznelson, 2014; Suchman et al., 2010). For a more detailed description of the conceptual framework underlying the MTB-model, see Sadler and colleagues (2013).

Chapter 2

The MTB-program offers a treatment manual with a set of well-developed protocols and guidelines (Slade et al., 2010). At the same time, the program can be adapted to the individual needs of the family and the circumstances of each home visit. For implementation of the MBT-program in The Netherlands, one of the PI's (HS) and social workers of the *MINDS*-Leiden team were thoroughly trained in the basic constructs and techniques of the reflective parenting model used in MTB. This MTB 'Introductory training institute' provides the basis for implementation of the MTB-program in other settings. A difference between the original MTB and the intervention used in this study is that we chose to work only with clinical social workers ("coaches"), instead of alternating the home visits between a nurse practitioner and clinical social worker. This decision has been based on the fact that in The Netherlands, mother and child pay regular visits to a pediatric nurse in the first four years of life. In the MTB-program, the visits by the nurse practitioner are mostly centered around health-related issues, which in The Netherlands are monitored by pediatric nurses at child health and welfare centers.

Home visits generally lasted about one hour, although at times of crisis home visits could be extended or increased in frequency. The home visits were conducted by a trained coach, starting during the last trimester of pregnancy until the child was 2.5 years old, and were scheduled weekly during the first year and continued two-weekly after that. Apart from the planned home visits, the coaches were available for their families when needed (by phone or 'Whatsapp'). The main objective of the coaches was to promote parental RF, support the mother-infant attachment relationship, and stimulate adequate parenting skills. Further, the coaches aimed to reinforce prenatal health care and health education, supported both mother's and child's health and development, helped mothers to extend or build a stronger social support network, educated mothers about the safety of their child, referred to a range of treatments as appropriate when psychiatric complaints were detected, and helped mothers negotiate issues involving legal, financial and housing problems. However, the coaches conferred regularly about their families during monthly supervision meetings, and maintained close contact with each other and their supervisor in case of crisis, or other family problems.

Table 1. The main research areas of the MINDS - Leiden Study.

Maternal variables	Wave 1: pregnancy	Wave 2: 6 months	Wave 3: 12 months	Wave 4: 20 months	Wave 5: 30 months	Wave 6: 3.5 years
Demographics	X	X	X	X	X	X
Reflective functioning	X			X		X
Mental and physical health	X	X	X	X	X	X
Life style	X	X			X	X
Social support	X				X	X
Self-efficacy in the nurturing role	X	X		X		
Executive functioning	X				X	
Emotion regulation	X				X	
Life events	X					X
Antenatal attachment	X					
Obstetric characteristics		X				
Parenting		X	X	X	X	X
Parenting stress					X	
IQ estimate		X				
Stress reactivity and regulation			X		X	

Chapter 2

Child variables	Wave 1: pregnancy	Wave 2: 6 months	Wave 3: 12 months	Wave 4: 20 months	Wave 5: 30 months	Wave 6: 3.5 years
Physical health		X	X	X	X	X
Cognitive development		X				
Stress reactivity and regulation		X	X		X	X
Language development			X	X	X	X
(Precursors of) executive functioning			X	X	X	X
(Precursors of) Theory of Mind			X	X	X	X
(Precursors of) empathy				X	X	X
Aggression		X	X	X	X	X
Behavioral problems (other)				X	X	X
Temperament		X	X	X		
Peer social interaction						X

Note. IQ = intelligence quotient.

Table 2. Research domains, instruments, and contents.

Domain (mother)	Instrument / data source	Content
Demographics, lifestyle, physical health & obstetric characteristics	'Becoming a mother'-questionnaire and 'Being a mother'-questionnaire (Hay, Mundy, et al., 2011; Smaling et al., 2015)	Key information regarding demographics, obstetric characteristics, life style, and physical health
Life events	List of Threatening Experiences – questionnaire (LTE-Q; Brugha & Cragg, 1990)	Stressful life events over the past year to self and close others, and whether they currently have impact
Social support	Norbeck Social Support Questionnaire (NSSQ; Norbeck, Lindsey, & Carrieri, 1981, 1983)	Functional variables (affect, aid, affirmation) and total network variables of social support (number of persons in the network, duration of relationships, frequency of contact), and loss of support within past year
Self-efficacy in the nurturing role	Multidimensional Scale of Perceived Social Support (MSPSS; Zimet, Dahlem, Zimet, & Farley, 1988)	Subjective assessment of social support adequacy
Mental health	Self-efficacy in the nurturing role scale (SENR; Pedersen, Bryan, Huffman, & Del Carmen, 1989)	Mothers' perceptions of their competence on basic skills required in caring for an infant
	Mini-International Neuropsychiatric Interview (M.I.N.I.) – plus (Sheehan et al., 1997)	Screening for psychiatric disorders
	Beck depression inventory (BDI) – II (Beck, Steer, & Brown, 1996)	Intensity of depression

	State-Trait Anxiety Inventory (STAI; Spielberger, 2010; Spielberger, Gorsuch, & Lushene, 1970)	Indication of transient anxiety and tendency to experience general anxiety
	Aggression Questionnaire (AQ; Buss & Perry, 1992)	Verbal aggression, physical aggression, anger, and hostility
	Borderline personality checklist (Arntz et al., 2003; Giesen-Bloo, Arntz, & Schouten, 2005)	Severity of borderline-related symptoms during past month
Emotion regulation	Difficulties in Emotion Regulation Scale (DERS; Gratz & Roemer, 2004)	Clinically relevant difficulties in emotion regulation.
Antenatal attachment	Maternal antenatal attachment scale (MAAS; Condon, 1993; Van Bussel, Spitz, & Demyttenaere, 2010)	Quality of mother's affective experiences towards the fetus and intensity of preoccupation with the fetus
Reflective functioning	Pregnancy Interview (PI) – Revised (Slade, 2007; Slade, Patterson, & Miller, 2007) and Parent Development Interview (PDI) – Revised (Slade et al., 2003; Slade et al., 2005)	Parental reflective functioning
Parenting	Adapted version of the Mother Infant Coding System (Miller, McDonough, Rosenblum, & Sameroff, 2002) to code maternal behavior during Free play, Teaching tasks, and Still-Face Paradigm (SFP; Tronick et al., 1978)	Maternal interactive behavior (e.g. sensitivity, intrusiveness) with her child during various contexts.
	Parental Cognitions and Conduct Toward the Infant Scale (PACOTIS; Boivin et al., 2005)	Parental perceptions and behavioral tendencies towards infant

Parenting stress	Nijmeegse Ouderlijke Stress Index (NOSI; Abidin, 1990; De Brock, Vermulst, Gerris, & Abidin, 1992)	Parenting stress
IQ	Wechsler Adult Intelligence Scale (WAIS-III-NL; Wechsler, 2005) – subtests Vocabulary, Matrix Reasoning, and Digit Span	Global indicators of intellectual functioning (IQ estimate) - mother
Executive functioning	Behavior Rating Inventory of Executive Function – adult version (BRIEF - A; Roth, Isquith, & Gioia, 2005)	Difficulties in executive functioning in daily life
Stress reactivity and regulation – baseline, stress and recovery	Maternal response to mildly stressful situation (i.e., short mother-child separation and child exposure to fear paradigm (see child variables))	<i>Hypothalamic pituitary adrenal (HPA)-axis:</i> Cortisol and Alpha-amylase (saliva) <i>Subjective experience of stress (mother):</i> Visual Analogue Scale (VAS; Lesage, Berjot, & Deschamps, 2012)
Domain (child)	Instrument / data source	Content
Cognitive development	Infant Mental Development Index of the Bayley Scales of Infant Development (BSID), 2nd version (Bayley, 1993)	Global measure of infant cognitive development
Stress reactivity and regulation – baseline, stress and recovery	Stressors: - <i>Social:</i> Still-Face Paradigm (Mesman, Van IJzendoorn, & Bakermans-Kranenburg, 2009; Tronick et al., 1978)	<i>Autonomic nervous system (child):</i> Heart rate, pre-ejection period, skin conductance, respiratory sinus arrhythmia with the Vrije

	-	<i>Frustration:</i> Laboratory Temperament Assessment Battery (Lab-TAB; Goldsmith & Rothbart, 1999a, 1999b), Car seat, Gentle arm restraint	Universiteit Ambulatory Monitoring System (VU-AMS; De Geus, Willemsen, Klaver, & Van Doornen, 1995; Willemsen, De Geus, Klaver, Van Doornen, & Carrofl, 1996)
	-	<i>Fear:</i> Adapted version of unpredictable-mechanical-toy from the Lab-TAB (Baker, Shelton, Baibazarova, Hay, & van Goozen, 2013; Goldsmith & Rothbart, 1999a)	<i>Hypothalamic pituitary adrenal (HPA)-axis:</i> Cortisol and Alpha-amylase (saliva) <i>Behavioral stress reactivity and regulation (child):</i> observation (coding based on Lab-TAB)
Aggression		Cardiff Infant Contentiousness Scale (CICS; Hay et al., 2010) Physical Aggression Scale for Early Childhood (PASEC; Alink et al., 2006) Peer Interaction Coding System (PICS; Caplan, Vespo, Pedersen, & Hay, 1991; Hay, Nash, et al., 2011)	Early manifestations of aggression Physical aggression Peer aggression
Behavioral problems		Child Behavior Check List (CBCL) 1 ½ -5 year (Achenbach & Rescorla, 2000; Achenbach & Ruffle, 2000)	Internalizing and externalizing behavioral problems
Temperament		Short versions of the Infant Behavior Questionnaire (IBQ; Rothbart, 1981) and Early Childhood Behavior Questionnaire (ECBQ; Putnam & Rothbart, 2006)	Dimensions of temperament (i.e., negative emotionality, effortful control)
(Precursors of) executive functioning		BRIEF – preschool version (Gioia, Espy, & Isquith, 2003)	Difficulties in executive functioning in daily life

	Music box (Goldsmith & Rothbart, 1999a)	Sustained attention
	Don't Paradigm (adapted from Kochanska, Tjebkes, & Fortnan, 1998)	Infants' spontaneous restraint to maternal prohibition
	Self-restraint (Friedman, Miyake, Robinson, & Hewitt, 2011), and Inhibitory control tasks (Carlson & Moses, 2001)	Inhibitory control
	Snack task and Gift task (Kochanska, Coy, & Murray, 2001; Spinrad et al., 2007)	Delay of gratification
	Hide the pots (Bernier, Carlson, & Whipple, 2010), and Spin the pots (Hughes & Ensor, 2005)	Working memory
	Dimensional Change Card Sort (Zelazo, 2006)	Mental flexibility
(Precursors of Theory of Mind)	Pleasure-disgust task (Slaughter & McConnell, 2003)	Social referencing
	Discrepant desires tasks (Atance, Bélanger, & Meltzoff, 2010; Laranjo, Bernier, Meins, & Carlson, 2010; Repacholi & Gopnik, 1997)	Understanding of discrepant desires
	Early Social Communication Scales (ESCS) – subtask joint attention (Mundy et al., 2003)	Joint attention
	Visual Perspectives (Bigelow & Dugas, 2009; Carlson, Mandell, & Williams, 2004; Laranjo et al., 2010)	Visual perspective taking
	Subtest functional and symbolic imitation - Autism Diagnostic Observation Schedule (ADOS; Luyster et al., 2009)	Imitation

(Precursors of empathy)	<p>False belief tasks (Bigelow & Dugas, 2009)</p> <p>Pain task (Young, Fox, & Zahn-Waxler, 1999; Zahn-Waxler, Robinson, & Emde, 1992)</p> <p>Mishaps (Kochanska, Gross, Lin, & Nichols, 2002)</p> <p>Emotion recognition task (Pollak, Cicchetti, Hornung, & Reed, 2000)</p> <p>Emotion eliciting film clips of ecologically valid situations (e.g., fear, sadness)</p>	<p>False belief</p> <p>Empathy</p> <p>Guilt</p> <p>Emotion recognition</p> <p>Social attention with the Tobii T120 eye-tracker</p>
Language development	<p>MacArthur communicative development inventory (Fenson et al., 2000)</p> <p>Reynell Developmental Language Scales(Reynell, 1985), Peabody Picture Vocabulary Test (PPVT; Dunn & Dunn, 1997)</p>	<p>Communicative skills in infants and toddlers</p> <p>Receptive language skills</p>
	<p>Schlichting Expressive Language Test (Schlichting, Van Eldik, Lutje Spelberg, Van der Meulen, & Van der Meulen, 1995)</p>	<p>Expressive vocabulary skills</p>

Group characteristics

Table 3 provides an overview of demographic and obstetric characteristics of the LR-and HR-groups. Compared to women in the LR-group, women in the HR-group were significantly younger, lower educated, had a lower income, were more often non-Caucasian and single, and had a smaller social support network ($ps < .01$). Further, pregnancies in the HR-group were more often unplanned, and women in the HR-group more often experienced miscarriages or had undergone abortion ($ps < .05$).

Among women in the LR-group ($n = 128$), 9% ($n = 11$) had one risk factor present, which was mostly limited social support ($n = 5$), followed by single parenthood ($n = 2$), young maternal age ($n = 2$), unemployment ($n = 1$) or no secondary education ($n = 1$). In the HR-group, 64% of the women ($n = 94$) had an indication for current psychiatric disorder(s), with 26% ($n = 38$) having two or more diagnoses on the M.I.N.I.-plus. See Table 4 for an overview of the diagnoses on the M.I.N.I.-plus in the HR-group. Substance use during pregnancy was the second most frequent observed risk factor in the HR-group. Of the women who used substances during pregnancy, 33% ($n = 48$) continued to smoke, 5% ($n = 8$) drank alcohol, 1% ($n = 1$) continued to use (other) drugs, 2% ($n = 3$) were smoking and drinking alcohol, 3% ($n = 5$) were smoking and using drugs, and 1% ($n = 1$) used all these substances (smoking, alcohol and drugs). Drugs used during pregnancy were cannabis ($n = 4$), cocaine ($n = 1$), methadone ($n = 1$), and cocaine and cannabis ($n = 1$).

Table 3. Demographic characteristics of the low-risk and high-risk groups.

Variables	LR (<i>n</i> = 128)		HR (<i>n</i> = 147)		Group comparisons ^d
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Maternal age (years)	23.42	1.74	21.56	2.32	$F(1,273)=55.59^{***}$, $\eta^2=.17$
Maternal education (% high ^b)	41%		7%		$\chi^2(1)=44.10^{***}$
Monthly family income (Euros)	2,944.02	964.50	1,609.00	1,145.00	$F(1,273)=107.47^{***}$, $\eta^2=.28$
Ethnicity (% Caucasian)	91%		76%		$\chi^2(1)=10.06^{**}$
Married or living with partner (%)	98%		61%		$\chi^2(1)=54.59^{***}$
First-time pregnant (%)	87%		76%		$\chi^2(1)=5.53^*$
Unplanned pregnancy (%)	23%		72%		$\chi^2(1)=65.15^{***}$
Expecting twins (%)	4%		5%		<i>ns</i>
Gestational weeks at assessment	30.03	3.77	29.91	3.68	<i>ns</i>
Size social support network	9.19	3.84	6.94	3.54	$F(1,273)=25.40^{***}$, $\eta^2=.09$
Prenatal reflective functioning	4.27	1.03	3.53	0.86	$F(1,273)=42.58^{***}$, $\eta^2=.14$
Number of risk factors	0.09	0.29	2.24	1.35	$F(1,273)=311.72^{***}$, $\eta^2=.53$
Current psychiatric disorder(s) (%)	0%		64%		$\chi^2(1)=124.36^{***}$
Substance use pregnancy (%)	0%		44%		$\chi^2(1)=70.78^{***}$
Alcohol	0%		7%		$\chi^2(1)=7.36^{**}$

Tobacco	0%	39%	$\chi^2(1)=62.61^{***}$
(Other) drugs	0%	5%	$\chi^2(1)=6.25^*$
Single parenthood (%)	2%	21%	$\chi^2(1)=24.70^{***}$
Unemployed (%)	1%	24%	$\chi^2(1)=31.89^{***}$
No secondary education (%)	1%	9%	$\chi^2(1)=9.21^{**}$
Financial problems (%)	0%	25%	$\chi^2(1)=37.23^{***}$
Limited social support (<4 persons)	4%	12%	$\chi^2(1)=6.21^*$
Young maternal age (<20 years)	2%	20%	$\chi^2(1)=23.63^{***}$

Note: * $p<.05$, ** $p<.01$, *** $p<.001$ ^aANOVA or Chi-square test, ^bMaternal education (% high) represents percentage with a bachelor's or master's degree, LR = low-risk group, HR = high-risk group.

Chapter 2

Table 4. Overview of the current psychiatric problems in the HR-group.

Current psychiatric problems*	N
Depressive disorder	17
Dysthymic disorder	8
Suicidality	11
Mania	1
Panic disorder	15
Agoraphobia	17
Social phobia	7
Other type of phobia	8
Obsessive-compulsive disorder	5
Generalized anxiety disorder	4
Posttraumatic stress disorder	4
Alcohol dependence / abuse	9
Drug dependence / abuse (non-alcohol)	18
Attention-deficit/hyperactivity disorder	7
Antisocial personality disorder	14

Note: N high-risk group = 147; * = positive screening based on the M.I.N.I.-plus (Sheehan et al., 1997; Van Vliet & De Beurs, 2007).

Discussion

This paper presented an overview of the background, methods and study population of the *MINDS* - Leiden study, a longitudinal study investigating social-environmental, neurobiological, and neurocognitive mechanisms predicting emotional and behavioral problems in young children and variation in effects of an intensive RF-based home-visiting program for first-time mothers with a high-risk background. The scope for changing behavior is greatest in the early years because neurobiological systems involved in emotional and behavioral regulation are presumed to be most sensitive for environmental influences early in development (Beauchaine et al., 2008). The outcomes of this study may result in the identification of specific biomarkers, precursors of neurocognitive functions and temperamental factors in infancy, which directly, indirectly or interactively with social-environmental risk factors, may help to detect children who are at risk for later emotional and behavior problems.

Home-visiting programs hold considerable promise for improving child outcomes (Avellar & Supplee, 2013; Filene et al., 2013; Olds et al., 2007; Peacock et al., 2013; Sweet & Appelbaum, 2004). From a socio-economic and ethical perspective, intervention programs should only be offered to those families who are expected to benefit and an alternative should be offered to those who are not expected to benefit. The outcomes of our study may result in a better understanding of the individual (neurocognitive and neurobiological) factors that explain variation in effects of home-visiting programs. Ultimately, this may contribute to more efficient matching of families to intervention programs while at the same time provide relevant information to further enhance the effectiveness of the home-visiting program.

A considerable strength of the study includes the random controlled trial and the use of multi-method approach, including a combination of (semi-)structured interviews, questionnaires, behavioral observations, and physiological measures. An important aspect of the current study, compared to previous studies evaluating the effects of home-visiting programs (Olds et al., 2007; Ordway et al., 2014; Robling et al., 2016), is the use of both a high-risk and a low-risk control group when examining the effects of home-visitation, which allows us to more thoroughly determine the extent to which neurobiological and neurocognitive development in children in the HR-I group normalizes (as observed deviations in HR-children compared to LR-children throughout development may, in part, depend on their ‘starting levels’).

Chapter 2

A limitation of this study is that we lack information about the number of subjects that refused to participate before the research team tried to contact them for scheduling the first assessment. This is due to our sampling strategy in which we relied on external parties for the recruitment of potential participants. It cannot be ruled out that non-participants may differ from participants in several respects, especially as participation in a longitudinal study with 6 assessments requires a strong commitment of the mothers and might be perceived as a burden. This may have resulted in the loss of more severe cases. For example, in our high-risk sample, only a small group of participants had three or more risk factors present (17%). We do have information about the women who were recruited by their healthcare provider and gave permission to contact them for partaking in the study, but declined participation when they were contacted (16%). The most common reasons for not participating were: too busy/too time-consuming, partner does not want to participate, personal problems, medical problems, or miscarriage.

A special point of interest is the potential loss of participants to follow-up over time. Although some loss to follow-up is inevitable, we used various strategies for maximizing retention in order to retain as many participants as possible. For example, mothers received gift certificates (respectively 15, 20, 25, 30, 35, and 25 EUR) and a present for their child after each assessment. After completion of the fifth assessment, they received a personalized DVD with video material from all five assessments. Also, over the course of the study regular contact was maintained by telephone calls after birth, birth- and birthday cards, and a newsletter (every 3 months). Travel expenses were covered for lab visits (wave three, five, and six). Moreover, we offered to pick mother and child up from their homes and drive them to our baby lab when they had difficulties coming to the lab, and mothers in the HR-intervention group were often accompanied to the lab by their coach. With data collection for the first assessment completed, attrition rates between the first and second assessment were 5%. Therefore, we are optimistic that our strategies are working and our attrition rates in the long run will be limited.

At this moment, funding has been obtained for the first five waves of follow up, and we plan to seek funding for additional follow-up cycles. Now that the study is well underway, efforts are being made to disseminate the results of our study to obstetric care providers, health care workers, pediatricians and policymakers on a regular basis. Further, once we have gained insight into factors that predict the effectiveness of the home-visiting program, we will look for ways to implement the program in clinical practice. For this, we have

The Mother-Infant Neurodevelopment Study – Leiden

already sought cooperation with health care organizations in the area of Leiden, The Hague, and Amsterdam.

Taken the potential restraints into account, we believe that this study program may provide detailed insight in the factors associated with (very) early child development and treatment success for interventions aiming to reduce and/or prevent emotional and behavioral problems. We hope that by effectively addressing behavioral problems from infancy onwards, improvements in (mental) health status (of children and their mothers), and public health policy and decision making may be achieved.

Chapter 2

References

- Abidin, R. R. (1990). *Parenting Stress Index (PSI)*. Charlottesville, VA: Pediatric Psychology Press.
- Achenbach, T. M., & Rescorla, L. A. (2000). *Manual for the ASEBA preschool forms and profiles*. Burlington: University of Vermont, Research Center for Children, Youth, and Families.
- Achenbach, T. M., & Ruffle, T. M. (2000). The Child Behavior Checklist and related forms for assessing behavioral/emotional problems and competencies. *Pediatrics in Review, 21*(8), 265-271.
- Alink, L. R. A., Mesman, J., Van Zeijl, J., Stolk, M. N., Juffer, F., Koot, H. M., . . . Van IJzendoorn, M. H. (2006). The early childhood aggression curve: Development of physical aggression in 10- to 50-month-old children. *Child Development, 77*(4), 954-966. doi: 10.1111/j.1467-8624.2006.00912.x
- Arntz, A., Van den Hoorn, M., Cornelis, J., Verheul, R., Van den Bosch, W., & De Bie, A. (2003). Reliability and validity of the borderline personality disorder severity index. *Journal of Personality Disorders, 17*, 45-59.
- Atance, C. M., Bélanger, M., & Meltzoff, A. N. (2010). Preschoolers' understanding of others' desires: Fulfilling mine enhances my understanding of yours. *Developmental Psychology, 46*(6), 1505-1514.
- Avellar, S. A., & Supplee, L. H. (2013). Effectiveness of home visiting in improving child health and reducing child maltreatment. *Pediatrics, 132*(Supplement 2), S90-S99. doi: 10.1542/peds.2013-1021G
- Baker, E., Shelton, K. H., Baibazarova, E., Hay, D. F., & Van Goozen, S. H. M. (2013). Low skin conductance activity in infancy predicts aggression in toddlers 2 years later. *Psychological Science, 24*(6), 1051-1056. doi: 10.1177/0956797612465198
- Bayley, N. (1993). *Bayley Scales of Infant Development-Second Edition*. San Antonio, TX: The Psychological Corporation.
- Beauchaine, T. P. (2001). Vagal tone, development, and Gray's motivational theory: Toward an integrated model of autonomic nervous system functioning in psychopathology. *Development and Psychopathology, 13*(02), 183-214.
- Beauchaine, T. P., Neuhaus, E., Brenner, S. L., & Gatzke-Kopp, L. (2008). Ten good reasons to consider biological processes in prevention and intervention research. *Development and Psychopathology, 20*(3), 745-774. doi: 10.1017/s0954579408000369
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck depression inventory-II*. San Antonio, TX: Psychological Corporation.
- Belsky, J., & Pluess, M. (2009). Beyond diathesis stress: Differential susceptibility to environmental influences. *Psychological Bulletin, 135*(6), 885-908.
- Bernier, A., Carlson, S. M., & Whipple, N. (2010). From external regulation to self-regulation: Early parenting precursors of young children's executive functioning. *Child Development, 81*(1), 326-339. doi: 10.1111/j.1467-8624.2009.01397.x
- Bigelow, A. E., & Dugas, K. (2009). Relations among preschool children's understanding of visual perspective taking, false belief, and lying. *Journal of Cognition and Development, 9*(4), 411-433. doi: 10.1080/15248370802678299

The Mother-Infant Neurodevelopment Study – Leiden

- Boivin, M., Pérusse, D., Dionne, G., Saisset, V., Zoccolillo, M., Tarabulsy, G. M., . . . Tremblay, R. E. (2005). The genetic-environmental etiology of parents' perceptions and self-assessed behaviours toward their 5-month-old infants in a large twin and singleton sample. *Journal of Child Psychology and Psychiatry*, 46(6), 612-630. doi: 10.1111/j.1469-7610.2004.00375.x
- Boyce, W. T., & Ellis, B. J. (2005). Biological sensitivity to context: I. An evolutionary-developmental theory of the origins and functions of stress reactivity. *Development and Psychopathology*, 17, 271-301.
- Brugha, T. S., & Cragg, D. (1990). The List of Threatening Experiences: The reliability and validity of a brief life events questionnaire. *Acta Psychiatrica Scandinavica*, 82(1), 77-81. doi: 10.1111/j.1600-0447.1990.tb01360.x
- Buss, A. H., & Perry, M. (1992). The Aggression Questionnaire. *Journal of Personality and Social Psychology*, 63(3), 452-459. doi: 10.1037/0022-3514.63.3.452
- Cabaj, J. L., McDonald, S. W., & Tough, S. C. (2014). Early childhood risk and resilience factors for behavioural and emotional problems in middle childhood. *BMC pediatrics*, 14(1), 166-177.
- Caplan, M., Vespo, J., Pedersen, J., & Hay, D. F. (1991). Conflict and its resolution in small groups of one- and two-year-olds. *Child Development*, 62(6), 1513-1524. doi: 10.1111/1467-8624.ep9202105177
- Carlson, S. M., Mandell, D. J., & Williams, L. (2004). Executive function and theory of mind: Stability and prediction from ages 2 to 3. *Developmental Psychology*, 40(6), 1105-1122. doi: 10.1037/0012-1649.40.6.1105
- Carlson, S. M., & Moses, L. J. (2001). Individual differences in inhibitory control and children's theory of mind. *Child Development*, 72(4), 1032-1053. doi: 10.1111/1467-8624.00333
- Condon, J. T. (1993). The assessment of antenatal emotional attachment: Development of a questionnaire instrument. *British Journal of Medical Psychology*, 66(2), 167-183. doi: 10.1111/j.2044-8341.1993.tb01739.x
- Côté, S., Vaillancourt, T., LeBlanc, J. C., Nagin, D. S., & Tremblay, R. E. (2006). The development of physical aggression from toddlerhood to pre-adolescence: A nation wide longitudinal study of Canadian children. *Journal of Abnormal Child Psychology*, 34(1), 68-82.
- De Brock, A. J. L. L., Vermulst, A. A., Gerris, J. R. M., & Abidin, R. R. (1992). *Nijmeegse Ouderlijke Stress Index (NOSI)-manual*. Lisse: Swets en Zeitlinger.
- De Geus, E. J. C., Willemsen, G. H. M., Klaver, C. H. A. M., & Van Doornen, L. J. P. (1995). Ambulatory measurement of respiratory sinus arrhythmia and respiration rate. *Biological Psychology*, 41, 205-227. doi: 10.1016/0301-0511(95)05137-6
- Dunn, L. M., & Dunn, L. M. (1997). *PPVT-III: Peabody picture vocabulary test* (4th ed.). Circle Pines, MN: American Guidance Service.
- Fenson, L., Pethick, S., Renda, C., Cox, J. L., Dale, P. S., & Reznick, J. S. (2000). Short-form versions of the MacArthur communicative development inventories. *Applied Psycholinguistics*, 21(1), 95-116.
- Filene, J. H., Kaminski, J. W., Valle, L. A., & Cachat, P. (2013). Components associated with home

Chapter 2

visiting program outcomes: A meta-analysis. *Pediatrics*, 132(Supplement 2), S100-S109. doi: 10.1542/peds.2013-1021H

Friedman, N. P., Miyake, A., Robinson, J. L., & Hewitt, J. K. (2011). Developmental trajectories in toddlers' self-restraint predict individual differences in executive functions 14 years later: A behavioral genetic analysis. *Developmental Psychology*, 47(5), 1410-1430. doi: 10.1037/a0023750

Giesen-Bloo, J. H., Arntz, A., & Schouten, E. (2005). *The borderline personality checklist: Psychometric evaluation and factorial structure in clinical and nonclinical samples*. Internal Document. Maastricht University.

Gioia, G. A., Espy, K. A., & Isquith, P. K. (2003). *Behavior rating inventory of executive function – Preschool version*. Odessa, Florida: Psychological Assessment Resources.

Goldsmith, H. H., & Rothbart, M. K. (1999a). *The Laboratory Temperament Assessment Battery: Description of procedures (Locomotor Version 3.1)*. Manual. Madison, Wisconsin.

Goldsmith, H. H., & Rothbart, M. K. (1999b). *The Laboratory Temperament Assessment Battery: Description of procedures (Prelocomotor Version 3.1)*. Manual. Madison, Wisconsin.

Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment*, 26(1), 41-54. doi: 10.1023/b:joba.0000007455.08539.94

Hay, D. F., Mundy, L., Roberts, S., Carta, R., Waters, C. S., Perra, O., . . . Van Goozen, S. (2011).

Known risk factors for violence predict 12-month-old Infants' aggressiveness with peers. *Psychological Science*, 22(9), 1205-1211. doi: 10.1177/0956797611419303

Hay, D. F., Nash, A., Caplan, M., Swartzentruber, J., Ishikawa, F., & Vespo, J. E. (2011). The emergence of gender differences in physical aggression in the context of conflict between young peers. *British Journal of Developmental Psychology*, 29(2), 158-175. doi: 10.1111/j.2044-835X.2011.02028.x

Hay, D. F., Perra, O., Hudson, K., Waters, C. S., Mundy, L., Phillips, R., . . . Van Goozen, S. (2010). Identifying early signs of aggression: psychometric properties of the Cardiff infant contentiousness scale. *Aggressive Behavior*, 36(6), 351-357. doi: 10.1002/ab.20363

Hughes, C., & Ensor, R. (2005). Executive function and theory of mind in 2 year olds: A family affair? *Developmental Neuropsychology*, 28(2), 645-668. doi: 10.1207/s15326942dn2802_5

Huijbregts, S. C. J., Seguin, J. R., Zoccolillo, M., Boivin, M., & Tremblay, R. E. (2008). Maternal prenatal smoking, parental antisocial behavior, and early childhood physical aggression. *Development and Psychopathology*, 20(2), 437-453. doi: 10.1017/s0954579408000217

Katznelson, H. (2014). Reflective functioning: A review. *Clinical Psychology Review*, 34(2), 107-117. doi: 10.1016/j.cpr.2013.12.003

Kochanska, G., Coy, K. C., & Murray, K. T. (2001). The development of self-regulation in the first four years of life. *Child Development*, 72(4), 1091-1111. doi: 10.1111/1467-8624.00336

Kochanska, G., Gross, J. N., Lin, M.-H., & Nichols, K. E. (2002). Guilt in young children: Development,

The Mother-Infant Neurodevelopment Study – Leiden

- determinants, and relations with a broader system of standards. *Child Development*, 73(2), 461-482. doi: 10.1111/1467-8624.00418
- Kochanska, G., Tjebkes, J. L., & Fortnan, D. R. (1998). Children's emerging regulation of conduct: Restraint, compliance, and internalization from infancy to the second year. *Child Development*, 69(5), 1378-1389. doi: 10.1111/j.1467-8624.1998.tb06218.x
- Laranjo, J., Bernier, A., Meins, E., & Carlson, S. M. (2010). Early manifestations of children's theory of mind: The roles of maternal mind-mindedness and infant security of attachment. *Infancy*, 15(3), 300-323. doi: 10.1111/j.1532-7078.2009.00014.x
- Laurent, H. K., Harold, G. T., Leve, L., Shelton, K. H., & Van Goozen, S. H. M. (2016). Understanding the unfolding of stress regulation in infants. *Development and Psychopathology, FirstView*, 1-10. doi: 10.1017/S0954579416000171
- Lesage, F. X., Berjot, S., & Deschamps, F. (2012). Clinical stress assessment using a visual analogue scale. *Occupational medicine*, 62(8), 600-605. doi: 10.1093/occmed/kqs140
- Luyster, R., Gotham, K., Guthrie, W., Coffing, M., Petrak, R., Pierce, K., . . . Lord, C. (2009). The Autism Diagnostic Observation Schedule—toddler module: A new module of a standardized diagnostic measure for autism spectrum disorders. *Journal of Autism & Developmental Disorders*, 39(9), 1305-1320. doi: 10.1007/s10803-009-0746-z
- Mejdoubi, J., Van den Heijkant, S. C. C. M., Van Leerdam, F. J. M., Crone, M., Crijnen, A., & HiraSing, R. A. (2014). Effects of nurse home visitation on cigarette smoking, pregnancy outcomes and breastfeeding: A randomized controlled trial. *Midwifery*, 30(6), 688-695. doi: 10.1016/j.midw.2013.08.006
- Mejdoubi, J., Van den Heijkant, S. C. C. M., Van Leerdam, F. J. M., Heymans, M. W., Crijnen, A., & HiraSing, R. A. (2015). The effect of Voorzorg, the Dutch Nurse-Family Partnership, on child maltreatment and development: A randomized controlled trial. *PLoS ONE*, 10(4), 1-14. doi: 10.1371/journal.pone.0120182
- Mesman, J., Van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2009). The many faces of the Still-Face Paradigm: A review and meta-analysis. *Developmental Review*, 29(2), 120-162. doi: 10.1016/j.dr.2009.02.001
- Miller, A. L., McDonough, S. C., Rosenblum, K. L., & Sameroff, A. J. (2002). Emotion regulation in context: Situational effects on infant and caregiver behavior. *Infancy*, 3(4), 403-433. doi: 10.1207/S15327078IN0304_01
- Mundy, P., Delgado, C., Block, J., Venezia, M., Hogan, A., & Seibert, J. (2003). *A manual for the abridged Early social communication scales (ESCS)*. Manual. Coral Gables, FL.
- Norbeck, J. S., Lindsey, A. M., & Carrieri, V. L. (1981). The development of an instrument to measure social support. *Nursing research*, 30(5), 264-269.
- Norbeck, J. S., Lindsey, A. M., & Carrieri, V. L. (1983). Further development of the Norbeck social support questionnaire: Normative data and validity testing. *Nursing research*, 32(1), 4-9.
- Olds, D. L., Henderson, C. R. J., Cole, R., Eckenrode, J., Kitzman, H., Luckey, D., . . . Powers, J. (1998). Long-term effects of nurse home visitation on behavior - 15-year follow-up of a

Chapter 2

- randomized controlled trial. *JAMA*, 280(14), 1238-1244. doi: 10.1001/jama.280.14.1238
- Olds, D. L., Sadler, L., & Kitzman, H. (2007). Programs for parents of infants and toddlers: Recent evidence from randomized trials. *Journal of Child Psychology and Psychiatry*, 48(3-4), 355-391. doi: 10.1111/j.1469-7610.2006.01702.x
- Ordway, M. R., Sadler, L. S., Dixon, J., Close, N., Mayes, L. C., & Slade, A. (2014). Lasting effects of an interdisciplinary home visiting program on child behavior: Preliminary follow-up results of a randomized trial. *Journal of Pediatric Nursing*, 29(1), 3-13. doi: 10.1016/j.pedn.2013.04.006
- Peacock, S., Konrad, S., Watson, E., Nickel, D., & Muhajarine, N. (2013). Effectiveness of home visiting programs on child outcomes: a systematic review. *BMC Public Health*, 13(1), 1-14. doi: 10.1186/1471-2458-13-17
- Pedersen, F. A., Bryan, Y. E., Huffman, L., & Del Carmen, R. (1989). *Construction of self and offspring in the pregnancy and early infancy periods*. Paper presented at the Society for Research in Child Development, Kansas City, MO.
- Putnam, S. P., & Rothbart, M. K. (2006). Development of short and very short forms of the Children's Behavior Questionnaire. *Journal of Personality Assessment*, 87(1), 102-112. doi: 10.1207/s15327752jpa8701_09
- Repacholi, B. M., & Gopnik, A. (1997). Early reasoning about desires: Evidence from 14- and 18-month-olds. *Developmental Psychology*, 33(1), 12-21. doi: 10.1037/0012-1649.33.1.12
- Reynell, J. K. (1985). *Reynell Developmental Language Scales* (2nd ed.). Windsor: NFER Nelson.
- Robling, M., Bekkers, M.-J., Bell, K., Butler, C. C., Cannings-John, R., Channon, S., . . . Torgerson, D. (2016). Effectiveness of a nurse-led intensive home-visitation programme for first-time teenage mothers (Building Blocks): A pragmatic randomised controlled trial. *The Lancet*, 387(10014), 146-155. doi: 10.1016/S0140-6736(15)00392-X
- Roth, R. M., Isquith, P. K., & Gioia, G. A. (2005). *BRIEF-A. Behavior rating inventory of executive function-adult version professional manual*. Lutz: Psychological Assessment Resources.
- Rothbart, M. K. (1981). Measurement of temperament in infancy. *Child Development*, 52, 569-578.
- Sadler, L. S., Slade, A., Close, N., Webb, D. L., Simpson, T., Fennie, K., & Mayes, L. C. (2013). Minding the Baby: Enhancing reflectiveness to improve early health and relationship outcomes in an interdisciplinary home-visiting program. *Infant Mental Health Journal*, 34(5), 391-405. doi: 10.1002/imhj.21406
- Schlichting, J. E. P. T., Van Eldik, M. C. M., Lutje Spelberg, H. C., Van der Meulen, S., & Van der Meulen, B. F. (1995). *Schlichting Test voor Taalproductie*. Nijmegen: Berkhout Nijmegen BV.
- Sheehan, D. V., Lecrubier, Y., Harnett Sheehan, K., Janavs, J., Weiller, E., Keskiner, A., . . . Dunbar, G. C. (1997). The validity of the Mini International Neuropsychiatric Interview (MINI) according to the SCID-P and its reliability. *European Psychiatry*, 12(5), 232-241. doi: 10.1016/S0924-9338(97)83297-X
- Slade, A. (2002). Keeping the baby mind: A critical factor in perinatal mental health. *Zero to three*, 22(6), 10-16.

The Mother-Infant Neurodevelopment Study – Leiden

- Slade, A., Aber, J. L., Berger, B., Bresgi, I., & Kaplan, M. (2003). *The parent development interview – revised*. New York: The City University of New York.
- Slade, A., Bernbach, E., Grienenberger, J., Levy, D., & Locker, A. (2005). *Manual for scoring reflective functioning on the Parent Development Interview*. New York: The City University of New York.
- Slade, A., Patterson, M., & Miller, M. R. (2007). *Addendum to reflective functioning scoring manual for use with the pregnancy interview, version 2.0*. New York: The City University of New York.
- Slade, A., Sadler, L., De Dios-kenn, C., Webb, D., Currier-Ezepchick, J., & Mayes, L. C. (2005). Minding the Baby: A reflective parenting program. *The Psychoanalytic Study of the Child*, 60, 74-100.
- Slade, A., Sadler, L. S., Close, N., Webb, D., de Dios-Kenn, C., Ezepchick, J., et al. (2010). *Minding the Baby treatment manual*. New Haven, CT: Yale University.
- Slade, A., Sadler, L. S., & Mayes, L. C. (2005). Minding the Baby: Enhancing parental reflective functioning in a nursing/mental health home visiting program. In L. J. Berlin, Y. Ziv, L. Amaya-Jackson, & M. T. Greenberg (Eds.), *Enhancing early attachments: Theory, research, intervention, and policy* (pp. 152–177). New York: Guilford Press.
- Slaughter, V., & McConnell, D. (2003). Emergence of joint attention: Relationships between gaze following, social referencing, imitation, and naming in infancy. *Journal of Genetic Psychology*, 164(1), 54.
- Smaling, H. J. A., Huijbregts, S. C. J., Suurland, J., Van der Heijden, K. B., Van Goozen, S. H. M., & Swaab, H. (2015). Prenatal reflective functioning in primiparous women with a high-risk profile. *Infant Mental Health Journal*, 36(3), 251-261. doi: 10.1002/imhj.21506
- Spielberger, C. D. (2010). *State-Trait Anxiety Inventory*. The Corsini Encyclopedia of Psychology: John Wiley & Sons, Inc.
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *The state-trait anxiety inventory*. Palo Alto, Calif: Consulting Psychologists Press Inc.
- Spinrad, T. L., Eisenberg, N., Gaertner, B., Popp, T., Smith, C. L., Kupfer, A., . . . Hofer, C. (2007). Relations of maternal socialization and toddlers' effortful control to children's adjustment and social competence. *Developmental Psychology*, 43(5), 1170-1186. doi: 10.1037/0012-1649.43.5.1170
- Suchman, N. E., DeCoste, C., Castiglioni, N., McMahan, T. J., Rounsaville, B., & Mayes, L. C. (2010). The Mothers and Toddlers Program, an attachment-based parenting intervention for substance using women: Post-treatment results from a randomized clinical pilot. *Attachment & Human Development*, 12(5), 483-504. doi: 10.1080/14616734.2010.501983
- Sweet, M. A., & Appelbaum, M. I. (2004). Is home visiting an effective strategy? A meta-analytic review of home visiting programs for families with young children. *Child Development*, 75(5), 1435-1456. doi: 10.1111/j.1467-8624.2004.00750.x
- Tronick, E., Als, H., Adamson, L., Wise, S., & Brazelton, T. B. (1978). The infant's response to entrapment between contradictory messages in face-to-face interaction. *Journal of the American Academy of Infant and Adolescent Psychiatry*, 17(1), 1-13.

Chapter 2

- Van Bussel, J. C. H., Spitz, B., & Demyttenaere, K. (2010). Reliability and validity of the Dutch version of the maternal antenatal attachment scale. *Archives of Women's Mental Health, 13*(3), 267-277. doi: 10.1007/s00737-009-0127-9
- Van Goozen, S. H. M., Fairchild, G., Snoek, H., & Harold, G. T. (2007). The evidence for a neurobiological model of childhood antisocial behavior. *Psychological Bulletin, 133*(1), 149-182. doi: 10.1037/0033-2909.133.1.149
- Van Vliet, I. M., & De Beurs, E. (2007). The MINI-International Neuropsychiatric Interview. A brief structured diagnostic psychiatric interview for DSM-IV and ICD-10 psychiatric disorders. *Tijdschrift voor Psychiatrie, 49*(6), 393-397.
- Van Vliet, I. M., Leroy, H., & Van Megen, H. J. G. M. (2000). *MINI plus. International neuropsychological interview. Nederlandse Versie 5.0. 0.[Dutch Version 5.0. 0.]*.
- Wechsler, D. (2005). *WAIS-III NL. Wechsler Adult Intelligence Scale WAIS-III. Dutch version. (3rd ed.)*. Amsterdam: Harcourt Test Publishers.
- Willemsen, G. H. M., De Geus, E. J. C., Klaver, C. H. A. M., Van Doornen, L. J. P., & Carrofl, D. (1996). Ambulatory monitoring of the impedance cardiogram. *Psychophysiology, 33*(2), 184-193. doi: 10.1111/j.1469-8986.1996.tb02122.x
- World Health Organization. (2005). *Child abuse and neglect*. Retrieved November 21, 2013, from http://who.int/violence_injury_prevention/violence/neglect/en/print.html
- World Health Organization. (2016). *Adolescent pregnancy*. Retrieved May 4, 2016, from http://www.who.int/maternal_child_adolescent/topics/maternal/adolescent_pregnancy/en/
- Young, S. K., Fox, N. A., & Zahn-Waxler, C. (1999). The relations between temperament and empathy in 2-year-olds. *Developmental Psychology, 35*(5), 1189-1197. doi: 10.1037/0012-1649.35.5.1189
- Zahn-Waxler, C., Robinson, J. L., & Emde, R. N. (1992). The development of empathy in twins. *Developmental Psychology, 28*(6), 1038-1047. doi: 10.1037//0012-1649.28.6.1038
- Zelazo, P. D. (2006). The Dimensional Change Card Sort (DCCS): A method of assessing executive function in children. *Nature Protocols Electronic Edition, 1*(1), 297-301. doi: 10.1038/nprot.2006.46
- Zimet, G. D., Dahlem, N. W., Zimet, S. G., & Farley, G. K. (1988). The Multidimensional Scale of Perceived Social Support. *Journal of Personality Assessment, 52*(1), 30-41.