



Universiteit
Leiden
The Netherlands

The role of BDNF in depression : will the neurotrophin hypothesis sparkle on, long after the glitter of the firework is gone?

Molendijk, M.L.

Citation

Molendijk, M. L. (2014, June 3). *The role of BDNF in depression : will the neurotrophin hypothesis sparkle on, long after the glitter of the firework is gone?*. Retrieved from <https://hdl.handle.net/1887/25851>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/25851>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/25851> holds various files of this Leiden University dissertation

Author: Molendijk, M.L.

Title: The role of BDNF in depression : will the neurotrophin hypothesis sparkle on, long after the glitter of the firework is gone?

Issue Date: 2014-06-03

REFERENCES

- A**
- aan het Rot M, Zarate CA, Charney DS, Mathew SJ. Ketamine for depression: where do we go from here? *Biol Psychiatry* 2012; **72**: 537-547.
- Agartz I, Sedvall GC, Terenius L, Kulle B, Frigessi A, Hall H *et al*. BDNF gene variants and brain morphology in schizophrenia. *Am J Med Gen* 2006; **114**: 513-523.
- Aguilera M, Arias B, Wichers M, Barrantes-Vidal N, Moja J, Villa H *et al*. Early adversity and 5-HTT/BDNF genes: new evidence of gene-environment interactions on depressive symptoms in a general population. *Psychol Med* 2009; **39**: 1425-1432.
- Aid T, Kazantseva A, Piirsoo M, Palm K, Timmusk T. Mouse and rat BDNF gene structure and expression revisited. *J Neurosci Res* 2007; **85**: 525-535.
- Almeida OP, Lautenschlager N, Vasikaram S, Leedman P, Flicker L. Association between physiological serum concentration of estrogen and the mental health of community dwelling postmenopausal women age 70 years and over. *Am J Ger Psychiatry* 2005; **13**: 142-149.
- Altar CA, Laeng P, Jurata LW, Brockman JA, Lemire A, Bullard J *et al*. Electroconvulsive seizures regulate gene expression of distinct neurotrophic signaling pathways. *J Neurosci* 2004; **24**: 2667-2677.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th edn. *American Psychiatric Association*: Washington, DC, 1994.
- American Psychiatric Association taskforce on laboratory tests in psychiatry. The dexamethasone suppression test: an overview of its current status in psychiatry. *Am J Psychiatry* 1987; **144**: 1353-1262.
- Anacker C, Zunszain PA, Carvalho PA, Pariante CM. The glucocorticoid receptor: pivot of depression and of antidepressant treatment? *Psychoneuroendocrinol* 2011; **36**: 415-425.
- Andrews G, Brugh T, Thase ME, Duffy FF, Rucci P, Slade T. Dimensionality and the category of major depressive episode. *Int J Methods Psychiatr Res* 2007; **16**: S41-S51.
- Angelucci F, Brene S, Mathe AA. BDNF in schizophrenia, depression and corresponding animal models. *Mol Psychiatry* 2005; **10**: 345-352.
- Anttila SAK, Leinonen EVJ. A review of the pharmacological and clinical profile of mirtazapine. *CNS Drug Rev* 2001; **7**: 249-264.
- Ashburner J. A fast diffeomorphic image registration algorithm. *Neuroimage* 2007; **38**: 95-113.
- Autry AE, Adachi M, Cheng P, Monteggia LM. Genders specific impact of brain-derived neurotrophic factor signaling on stress-induced depression-like behavior. *Biol Psychiatry* 2009; **66**: 84-90.
- Autry AE, Monteggia LM. Brain-derived neurotrophic factor and neuropsychiatric disorders. *Pharmacol Rev* 2012; **64**: 238-258.
- Aydemir C, Yalcin ES, Aksaray S, Kisa C, Yildirim SG, Uzbay T *et al*. Brain-derived neurotrophic factor (BDNF) changes in the serum of depressed women. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; **30**: 1256-1260.
- Aydemir O, Deveci A, Taskin OE, Taneli F, Esen-Danaci A. Serum brain-derived neurotrophic factor level in dysthymia: a comparative study with major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2007; **31**: 1023-1026.
- Azoulay D, Vachapova V, Shihman B, Miller A, Karni A. Lower brain-derived neurotrophic factor in serum of relapsing remitting MS: reversal by glatiramer acetate. *J Neuroimmunol* 2005; **167**: 215-218.
- B**
- Babor TF, Epstein N, Brown G, Steer RA. Early detection of harmful alcohol consumption: comparison of clinical, laboratory, and self-report screening procedures. *Addict Behav* 1989; **14**: 139-157.
- Ball S, Marangell LB, Lipsius S, Russell JM. Brain-derived neurotrophic factor in generalized anxiety disorder: results from a duloxetine trial. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2013; **43**: 217-221.
- Barde YA, Edgar D, Thoenen H. Purification of a new neurotrophic factor from mammalian brain. *J EMBO* 1982; **1**: 549-553.
- Barnham KJ, Masters CL, Bush AI. Neurodegenerative diseases and oxidative stress. *Nat Rev Neurosci* 2004; **3**: 205-214.
- Basterzi AD, Yazici K, Aslan E, Delialioglu N, Tasdelen B, Acar ST *et al*. Effects of fluoxetine and venlafaxine on serum brain-derived neurotrophic factor concentrations in depressed patients. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2009; **33**: 281-285.
- Bath KG, Chuang J, Spencer-Segal JL, Amso D, Altemus M, McEwen BS *et al*. Variant brain-derived neurotrophic factor (valine⁶⁶methionine) polymorphism contributes to developmental and estrous stage-specific expression of anxiety-like behavior in female mice. *Biol Psychiatry* 2012; **72**: 499-504.
- Beck AT, Kovacs M, Weissman A. Assessment of suicidal intention: the Scale for Suicide Ideation. *J Consult Clin Psychol* 1979; **47**: 343-352.
- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 1988; **56**: 893-897.
- Beevers CG, Wells TT, McGeary JE. The BDNF val⁶⁶met polymorphism is associated with rumination in healthy adults. *Emotion* 2009; **9**: 579-584.
- Begliuomini S, Casarosa E, Pluchino N, Lenzi E, Centofanti M, Freschi L *et al*. Influence of endogenous and exogenous sex hormones on plasma brain-derived neurotrophic factor. *Hum Reprod* 2007; **22**: 995 - 1002.
- Begliuomini S, Lenzi E, Ninni F, Casarosa E, Merlini S, Pluchino N *et al*. Plasma brain-derived neurotrophic factor daily variations in men: correlation with cortisol circadian rhythm. *J Endocrinol* 2008; **197**: 429-435.
- Bennett MR, Lagopoulos J. Stress and trauma: BDNF control of dendritic-spine formation and regression. *Prog Neurobiol* 2013; AOP doi: <http://dx.doi.org/10.1016/j.pneurobio.2013.10.005>.
- Berton O, Nestler EJ. New approaches to antidepressant drug discovery. *Nat Rev Neurosci* 2006; **7**: 137-151.
- Bejot Y, Mossiat C, Giroud M, Prigent-Tessier A, Marie C. Circulating and Brain BDNF levels in stroke rats. Relevance for clinical studies. *PLoS One* 2011; **6**: e29405.
- Benjamin S, McQuoid DR, Potter G, Payne ME, MacFall JR, Steffens DC *et al*. The brain-derived neurotrophic factor val⁶⁶met polymorphism, hippocampal volume, and cognitive function in geriatric depression. *Am J Ger Psychiatry* 2010; **18**: 323-331.
- Berkemeier LR, Winslow JW, Kaplan DR, Nikolics K, Goeddel DV *et al*. Neurotrophin-5 a novel neurotrophic factor that activates trk and trkB. *Neuron* 1991; **7**: 857-66.
- Bhang SY, Choi SW, Ahn JH. Changes in plasma brain-derived neurotrophic factor levels in smokers after smoking cessation. *Neurosci Lett* 2010; **468**: 7-11.
- Binder DK, Scharfman HE. Brain-derived neurotrophic factor. *Growth Factors* 2004; **22**: 123-131.
- Birkenhäger TK, Geldermans S, Van den Broek WW, van Beveren N, Fekkes D. Serum brain-derived neurotrophic factor level in relation to illness severity and episode duration in patients with major depression. *J Psychiatr Res* 2012; **46**: 285-289.
- Bocchi-Chiavetto L, Zanardini R, Bortolomasi M, Abate M, Segala M, Giacomuzzi M *et al*. Electroconvulsive therapy increases serum BDNF in drug resistant depressed patients. *Eur Neuropsychopharmacol* 2006; **16**: 56-59.

Bocchio-Chiavetto L, Bagnardi V, Zanardini R, Molteni R, Nielsen MG, Placentino A et al. Serum and plasma BDNF levels in major depression: a replication study and meta-analyses. *W J Biol Psychiatry* 2010; **11**: 763-773.

Bogdan R, Hariri AR. Neural embedding of stress reactivity. *Nature Rev Neurosci* 2012; **15**: 1605-1607.

Bogdan R, Hyde LW, Hariri AR. A neurogenetics approach to understanding individual differences in brain, behavior, and risk for psychopathology. *Mol Psychiatry* 2013; **18**: 288-299.

Boomsma DI, Willemsen G, Sullivan PF, Heutink P, Meijer P, Sondervan D et al. Genomewide association of major depression: description of samples for the GAIN major depressive disorder study: NTR and NESDA biobank projects. *Eur J Hum Genet* 2008; **16**: 335-342.

Borenstein M, Hedges LV, Higgins JPT. 2009. *Introduction to meta-analyses*. Chichester; Wiley, NH, USA.

Boulle F, van den Hove DLA, Jakob SB, Rutten BP, Hamon M, van Os J et al. Epigenetic regulation of the BDNF gene: implications for psychiatric disorders. *Mol Psychiatry* 2012; **17**: 584-596.

Braff L. The neuropsychiatric translational revolution: still very early and very challenging. *JAMA Psychiatry* 2013; **70**: 777-779.

Bramham CR, Messaoudi E. BDNF function in adult synaptic plasticity: the synaptic consolidation hypothesis. *Prog Neurobiol* 2005; **76**: 99-125.

Bremner JD, Elzinga BM, Schmahl C, Vermetten E. Structural and functional plasticity of the human brain in post-traumatic stress disorder. *Prog Brain Res* 2007; **167**: 171-186.

Brett M, Anton J-L, Valabregue R, Poline JB. Region of interest analysis using an SPM toolbox. Presented at the 8th International Conference on Functional Mapping of the Human Brain, June 2-6, 2002, Sendai, Japan. Available on CD-ROM in *Neuroimage* 2002; **16**.

Brink CB, Harvey BH, Brand L. A novel atypical antidepressant that may provide new insights into the biomolecular basis of depression. *CNS Drug Discov* 2006; **1**: 29-41.

Bruce SE, Yonkers KA, Otto MW, Eisen JL, Weisberg JB, Pagano M et al. Influence of psychiatric comorbidity on recovery and recurrence in generalized anxiety disorder, social phobia, and panic disorder: a 12-year prospective study. *Am J Psychiatry* 2005; **162**: 1179 – 1187.

Brugha TS, Cragg D. The list of threatening experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand* 1990; **82**: 77-81.

Brugha T, Bebbington P, Tennant C, Hurry J. The list of threatening experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med* 1985; **15**: 189-194.

Brunoni AR, Lopes M, Fregni F. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. *Int J Neuropsychopharmacol* 2008; **11**: 1169-1180.

Bueller JA, Aftab M, Sen S, Gomez-Hassan D, Burmeister M, Zubieta JK. BDNF val⁶⁶met allele is associated with reduced hippocampal volume in healthy subjects. *Biol Psychiatry* 2006; **59**: 812-815.

Bunny WE Jr, Davis JM. Norepinephrine in depressive reactions: a review. *Arch Gen Psychiatry* 1965; **13**: 483-494.

Bus BAA, Molendijk ML, Penninx BJWH, Buitelaar JK, Kenis G, Prickaerts J et al. 2011. Determinants of serum brain derived neurotrophic factor. *Psychoneuroendocrinol* 2011; **36**: 228 – 239.

Bus BAA, Tendolkar I, Franke B, de Graaf J, Den Heijer M, Buitelaar JK et al. Serum brain-derived neurotrophic factor: determinants and relationship with depressive symptoms in a community population of middle-aged and elderly people. *W J Biol Psychiatry* 2012; **13**: 39-47.

Bus BAA, Molendijk ML, Penninx BJWH, Buitelaar JK, Prickaerts J, Elzinga BM et al. Low serum BDNF concentrations in depressed patients cannot be attributed to individual depressive symptoms or symptom cluster. *W J Biol Psychiatry*, In Press

Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**: 365-376.

C

Cai D, Holm, JM, Duignan JJ, Zheng J, Xaymardan M, Chin A et al. BDNF-mediated enhancement of inflammation and injury in the aging heart. *Physiol Genomics* 2006; **24**: 191-197.

Cappiello A, Malison RT, McDougle CJ, Vegso SJ, Charney DS, Heninger GR et al. Seasonal variation in neuroendocrine and mood responses to IV L-tryptophan in depressed patients and healthy subjects. *Neuropsychopharmacol* 1996; **15**: 475-483.

Carlsson A, Svennerholm L, Winblad B. Seasonal and circadian monoamine variations in human brains examined post mortem. *Acta Psychiatr Scand* 1980; **280**: S75-S85.

Casey BJ, Craddock N, Cutbert BN, Hyman SE, Lee FS, Ressler KJ. DSM-5 and RDoC: progress in psychiatry research. *Nat Rev Neurosci* 2013; **14**: 810-814.

Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H et al. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 2003; **301**: 386-389.

Caspi A, Moffitt TE. Gene-environment interactions in psychiatry: joining forces with neuroscience. *Nat Rev Neurosci* 2006; **7**: 583-590.

Castren E, Zafr F, Thoenen H, Lindholm D. Light regulates expression of brain-derived neurotrophic factor mRNA in rat visual cortex. *Proc Natl Acad Sci USA* 1992; **89**: 9444-9448.

Castren E, Vöikar V, Rantamäki T. Role of neurotrophic factors in depression. *Curr Opin Pharmacol* 2007; **7**: 18-21.

CBS 04/21/2010 URL: <http://www.cbs.nl/nl-NL/menu/methoden/>.

Chalakov GN, Fiore M, Stankulov IS, Manni L, Hristova MG, Antonelli A et al. Neurotrophin presence in human coronary atherosclerosis and metabolic syndrome: a role for NGF and BDNF in cardiovascular disease? *Prog Brain Res* 2004; **146**: 279-289.

Chalakov GN, Tonchev AB, Manni L, Hristova MG, Nikolova V, Fiore M et al. 2007. Comment on: Krabbe KS, Nielsen AR, Krogh-Madsen R. BDNF and type 2 diabetes. *Diabetologia* 2007; **50**: 431-438.

Chen B, Dowlathshahi D, McQueen GM, Wang JF, Young LT. Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry* 2001; **50**: 260-265.

Cardon LR, Palmer LJ. Population stratification and spurious allelic association. *Lancet* 2003; **361**: 598-604.

Campbell S, Marriott M, Nahmias C, Macqueen GM. Lower hippocampal volumes in patients suffering from depression: a meta-analysis. *Am J Psychiatry* 2004; **161**: 598-607.

Cassiman D, Denef C, Desmet VJ, Roskams T. Human and rat hepatic stellate cells express neurotrophins and neurotrophin receptors. *Hepatology* 2001; **33**: 148-158.

Chao MV. Neurotrophins and their receptors: a convergence point for many signaling pathways. *Nat Rev Neurosci* 2003; **4**: 299-309.

Chan KL, Tong KY, Yip SP. Relationship of serum brain-derived neurotrophic factor (BDNF) and health-related lifestyle in healthy human subjects. *Neurosci Lett* 2008; **447**: 124-128.

Charney DS. Psychobiological mechanisms of resilience and vulnerability. *Focus* 2004; **2**: 368-391.

Chen ZY, Patel PD, Sant G, Meng CX, Teng KK, Hempstead BL et al. Variant BDNF (met66) alters the intracellular trafficking and activity-dependent secretion of wild-type BDNF in neurosecretory cells and cortical neurons. *J Neurosci* 2004; **24**: 4401-4411.

Chepenik LG, Fredericks C, Papademetris X, Spencer L, Lacadie C, Wang F et al. Effects of brain-derived neurotrophic growth factor val⁶⁶met variation on hippocampus morphology in bipolar disorder. *Neuropsychopharmacol* 2009; **34**: 944-951.

Chen Z-Y, Jing D, Bath KG, Ieraci A, Khan T, Siao C-J et al. Genetic variant BDNF (val⁶⁶met) polymorphism alters anxiety-related behavior. *Science* 2006; **314**: 140-143.

Choi DC, Maguschak KA, Ye K, Jang S-W, Myers KM, Ressler KJ. Prelimbic cortical BDNF is required for memory of learned fear but not extinction or innate fear. *Proc Natl Acad Sci USA* 2010; **107**: 2675-2680.

Ciammola A, Sassone J, Cannella M, Calza S, Poletti B, Frati L et al. Low BDNF levels in serum of Huntington's disease patients. *Am J Med Genet B: Neuropsychiatr Genet* 2007; **144B**: 574-577.

Coe R. It's the effect size stupid: what effect size is and why it is important. Paper presented at the Annual conference of the British Educational Research Association 2002. www.leeds.ac.uk/educol/documents

Cohen S, Levi-Montalcini R, Hamburger V. A nerve growth-stimulating factor isolated from sarcomas 37 and 180. *Proc Natl Acad Sci USA* 1954; **40**: 1014-1018.

Cohen J. Statistical Power Analysis for the Behavioral Sciences. Lawrence Erlbaum Associates: Hillsdale, NJ, 1988.

Cole J, Weinberger DR, Mattay VS, Cheng X, Toga AW, Thompson PM et al. No effect of 5HTTLPR or BDNF val⁶⁶met polymorphism on hippocampal morphology in major depression. *Genes Brain Beh* 2011; **10**: 756-764.

Conti AC, Cryan JF, Dalvi A, Lucki I, Blendy JA. CAMP response element-binding protein is essential for the upregulation of brain-derived neurotrophic factor, but not the behavioral or endocrine responses to antidepressant drugs. *J Neurosci* 2002; **22**: 3262-3268.

Corominas-Roso M, Ramos-Quiroga JA, Ribases M, Sanchez-Mora C, Palomar G, Valero S et al. Decreased serum levels of brain-derived neurotrophic factor in adults with attention-deficit hyperactivity disorder. *Int J Neuropsychopharmacol* 2013; AOP March 2012: doi: 10.1017/S1461145712001629.

Couzin-Frankel J. When mice mislead. *Science* 2013; **342**: 922-925.

Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003; **35**: 1381-1395.

Christley RM. Power and error: increased risk of false positive results in underpowered studies. *Open Epidemiol* 2010; **3**: 16-19.

Cronbach LJ, Meehl PE. Construct validity in psychological tests. *Psychol Bull* 1955; **52**: 281-302.

Cryan JF, Holmes A. The ascent of mouse: advances in modeling human depression and anxiety. *Nat Rev Drug Discov* 2005 **4**: 775-790.

Currie J, Ramsbottom R, Ludlow H, Nevill A, Gilder M. Cardio-respiratory fitness, habitual physical activity and serum brain derived neurotrophic factor (BDNF) in men and women. *Neurosci Lett* 2009; **451**: 152-155.

D

Daselaar SM, Veltman DJ, Rombouts SA, Raaijmakers JG, Jonker C. Neuroanatomical correlates of episodic encoding and retrieval in young and elderly subjects. *Brain* 2003; **126**: 43-56.

David DJ, Samuels BA, Rainer Q, Wang J-W, Marsteller D, Mendez I et al. Neurogenesis-dependent and independent effects of fluoxetine in an animal model of anxiety/depression. *Neuron* 2009; **62**: 479-493.

Davidson J, Turnbull CD, Strickland R, Miller R, Graves K. The Montgomery-Asberg depression scale: reliability and validity. *Acta Psychiatr Scand* 1986; **73**: 544-548.

Davis MI. Ethanol-BDNF interactions: still more questions than answers. *Pharmacol Ther* 2008; **118**: 36-57.

Dawood T, Anderson J, Barton D, Lambert E, Esler M, Hotchkin E et al. Reduced overflow of BDNF from the brain is linked with suicide risk in depressive illness. *Mol Psychiatry* 2007; **12**: 981-983.

de Graaf R, Bijl RV, ten Have M, Beekman ATF, Vollebergh WAM. Pathways to comorbidity: the transition of pure mood, anxiety and substance abuse disorders into comorbid conditions in a longitudinal population based study. *J Affect Disord* 2004a; **82**: 461-467.

de Graaf R, Bijl RV, ten Have M, Beekman ATF, Vollebergh WAM. Rapid onset of comorbidity of common mental disorders: findings from the Netherlands Mental Health Survey and Incidence Study. *Acta Psychiatr Scand* 2004b; **109**: 55-63.

De Kloet ER, Vreugdenhil E, Oitzl MS, Joëls M. Brain corticosteroid receptor balance in health and disease. *Endocr Rev* 1998; **19**: 269-301.

Delgado PL. Depression: the case for a monoamine deficiency. *J Clin Psychiatry* 2000; **61**: S7-S11.

Dennis NA, Cabeza R, Need AC, Waters-Metenier S, Goldstein DB, LaBar KS. BDNF val⁶⁶met Polymorphism and hippocampal activation during episodic memory encoding and retrieval tasks. *Hippocampus* 2011; **21**: 980-989.

Deuschle M, Gilles M, Scharnholtz B, Lederbogen F, Lang U, Hellweg R. Changes of serum concentrations of BDNF during treatment with venlafaxine and mirtazapine: role of medication and response to treatment. *Pharmacopsychiatry* 2013; **46**: 54-58.

Deveci A, Aydemir O, Taskin O, Taneli F, Esen-Danaci A. Serum BDNF concentrations in suicide attempters related to psychosocial stressors: a comparative study with depression. *Neuropsychobiol* 2007; **56**: 93-97.

Dickersin K. The existence of publication bias and risk factors for its occurrence. *JAMA* 1990; **263**: 1385-1359.

Diniz BS, Teixeira AL, Talib LL, Mendoca VA, Gattaz WF, Forlenza OV. Serum brain-derived neurotrophic factor level is reduced in antidepressant-free patients with late-life depression. *W J Biol Psychiatry* 2010; **11**: 550-555.

Dodds CM, Henson RN, Suckling J, Miskowiak KW, Ooi C, Tait R et al. Effect of the BDNF val⁶⁶met polymorphism and met allele load on declarative memory related neural networks. *PLoS ONE* 2013; **8**: e74133.

Donovan MJ, Lin MI, Wiegand P. Brain-derived neurotrophic factor is an endothelial cell survival factor required for intramyocardial vessel stabilization. *Development* 2000; **127**: 4531-4540.

Dranovsky A, Hen R. Hippocampal neurogenesis: regulation by stress and antidepressants. *Biol Psychiatry* 2006; **59**: 1136-1143.

Driscoll I, Martin B, An Y, Maudsley S, Ferruci L, Mattson M et al. Plasma BDNF is associated with age-related white matter atrophy but not with cognitive function in older, non-demented adults. *PLoS One* 2012; **7**: e35217.

Duman RS, Heninger GR, Nestler EJ. A molecular and cellular theory of depression. *Arch Gen Psychiatry* 1997; **54**: 597-606.

Duman RS, Malberg J, Nakagawa S, D'Sa C. Neuronal plasticity and survival in mood disorders. *Biol Psychiatry* 2000; **48**: 732-739.

Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 2006; **59**: 1116-1127.

Duncan LE, Hutchison KE, Carey G, Craighead WE. Variation in brain-derived neurotrophic factor (BDNF) gene is associated with symptoms of depression. *J Affect Disord* 2009; **115**: 215-219.

Dunham JS, Deakin JFW, Miyajima F, Payton A, Toro CT. Expression of hippocampal BDNF and its receptors in Stanley consortium brains. *J Psychiatric Res* 2009; **43**: 1175-1184.

Dutt A, McDonald C, Dempster E, Prata D, Shaikh M, Schulze K et al. The effects of COMT, BDNF, 5-HTT, NRG1 and DTNBP1 genes on hippocampal and lateral ventricular volume in psychosis. *Psychological Med* 2009; **39**: 1783-1797.

Duval S, Tweedie RL. A nonparametric trim and fill method of accounting for publication bias in meta-analysis. *J Am Stat Soc* 2000; **95**: 89-98.

Dwivedi Y, Rizavi HS, Conley RR, Roberts RC, Tamminga CA, Pandey GN. Altered gene expression of BDNF and receptor tyrosine kinase B in postmortem brains of suicide victims. *Arch Gen Psychiatry* 2003; **60**: 804–815.

E

- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A *et al*. The BDNF val⁶⁶met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 2003; **112**: 257–269.
- Egger, M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634.
- Eisch AJ, Bolanos CA, de Wit J, Simonak RD, Pudiak CM, Barrot M *et al*. Brain-derived neurotrophic factor in the ventral midbrain-nucleus accumbens pathway: a role in depression. *Biol Psychiatry* 2003; **54**: 994-1005.
- Ejiri J, Inoue N, Kobayashi S, Shiraki R, Otsui K, Honjo T *et al*. Possible role of brain-derived neurotrophic factor in the pathogenesis of coronary artery disease. *Circulation* 2005; **112**: 2114-2120.
- Eker C, Kitis O, Taneli F, Eker DE, Ozan E, Yucel K *et al*. Correlation of serum BDNF levels with hippocampal volumes in first episode, medication-free depressed patients. *Eur Arch Psychiatry Clin Neurosci* 2010; **260**: 527-533.
- Elfving B, Plougmann PH, Muller HK, Mathe AA, Rosenberg R, Wegener G. Inverse correlation of brain and blood BDNF levels in a genetic rat model of depression. *Int J Neuropsychopharmacol* 2010; **13**: 563-572.
- Elfving B, Butterschön HN, Foldager L, Poulsen PHP, Andersen JH, Grynderup MB *et al*. Depression, the val⁶⁶met polymorphism, age, and gender influence the serum BDNF level. *J Psychiatric Res* 2012; **46**: 1118-1125.
- Elzinga BM, Molendijk ML, Oude Voshaar RC, Bus AA, Prickaerts J, Spinhoven P *et al*. The impact of childhood abuse and recent stress on serum BDNF and the moderating role of BDNF val⁶⁶met. *Psychopharmacol* 2011; **214**: 319-328.
- Erickson KI, Kim JS, Suever BL, Voss MW, Francis BM, Kramer AF. Genetic contributions to age-related decline in executive function: a 10-year longitudinal study of COMT and BDNF polymorphisms. *Front in Hum Neurosci* 2008; **2**: 1-9.
- Erickson KI, Prakash RS, Voss MW, Chaddock L, Heo S, McLaren M. Brain-derived neurotrophic factor is associated with age-related decline in hippocampal volume. *J Neurosci* 2010; **30**: 5368-5375.
- Esteban I, Hannestad J, Levanti B, Del Valle ME, Naves FJ, Vega JA. Neurotrophin receptor proteins immuno-reactivity in human gastrointestinal endocrine cells. *Brain Res Bull* 1995; **38**: 539-543.
- Eyler LT, Sherzai A, Kaup AR, Jeste DV. A review of functional brain imaging correlates of successful cognitive aging. *Biol Psychiatry* 2011; **70**: 115-122.

F

- Farrag A-K, Khedr EM, Abdel-Aleem T, Rageh A. Effects of surgical menopause on cognitive functions. *Dement Geriatr Cogn Disord* 2002; **13**: 193-198.
- Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Beh Res Methods* 2009; **41**: 1149-1160.
- Fernandes B, Gama CS, Massuda R, Torres M, Camargo D, Kunz M *et al*. Serum brain-derived neurotrophic factor is not associated with response to electroconvulsive therapy: a pilot in drug resistant depressed patients. *Neurosci Lett* 2009; **453**: 195-198.
- Fernandes BS, Gama CS, Cereser KM, Yatham LN, Fries GR, Colpo G *et al*. Brain-derived neurotrophic factor as a state marker of mood episodes in bipolar disorders: a systematic review and meta-regression analysis. *J Psychiatric Res* 2011; **45**: 995-1004.
- Friston K. Ten ironic rules for non-statistical reviewers. *Neuroimage* 2012; **61**: 1300-1310.
- Frodl T, Schule C, Schmitt G, Born C, Baghai T, Zill P *et al*. Association of brain-derived neurotrophic factor val⁶⁶met polymorphism with reduced hippocampal volumes in major depression. *Arch Gen Psychiatry* 2007; **64**: 410-416.
- Fujimura H, Altar CA, Chen R, Nakamura T, Nakahashi T, Kambayashi J *et al*. Brain-derived neurotrophic factor is stored in human platelets and released by agonist stimulation. *Thromb Haemost* 2002; **87**: 728-734.
- Fujinami A, Ohta K, Obayashi H, Fukui M, Hasegawa G, Nakamura N *et al*. Serum BDNF in patients with type 2 diabetes mellitus: relationship to glucose metabolism and biomarkers of insulin resistance. *Clin Biochem* 2008; **41**: 812-817.

G

- Gass P, Hellweg R. Peripheral BDNF levels as a biomarker for affective disorders. *Int J Neuropsychopharmacol* 2010; **13**: 1–4.
- Gaster B, Holroyd J. St John's wort for depression. *Arch Intern Med* 2000; **160**: 152–156.
- Gatt JM, Nemeroff CB, Dobson-Stone C, Paul RH, Bryant RH, Schofield R *et al*. Interactions between BDNF val⁶⁶met polymorphism predicts brain and arousal pathways to syndromal depression and anxiety. *Mol Psychiatry* 2009; **14**: 681–695.
- Gazal M, Motta LS, Wiener CD, Fernandes JC, Quevedo LA, Jansen K *et al*. Brain-derived neurotrophic factor in post-partum depressive mothers. *Neurochem Res* 2012; **37**: 583-587.
- Gedge L, Beaudoin A, Lazowski L, Toit R, Jokic R, Milev R. Effects of electro-convulsive therapy and repetitive transcranial magnetic stimulation on serum BDNF levels in patients with depression. *Front Psychiatry* 2012; **3**: 1-8.
- Genazzani RA, Spinetti A, Gallo R, Bernardi F. Menopause and the central nervous system: intervention options. *Maturitas* 1999; **31**: 103-110.
- Geroldi D, Minoretto P, Emanuele E. Brain-derived neurotrophic factor and the metabolic syndrome: more than just a hypothesis. *Med Hypotheses* 2006; **67**: 195-196.
- Gerritsen L, Tendolkar I, Franke B, Vasquez AA, Buitelaar J, Fernandez G *et al*. BDNF val⁶⁶met genotype modulates the effect of childhood adversity on subgenual anterior cingulate cortex volume in healthy subjects. *Mol Psychiatry* 2012; **17**: 597-603.
- Gervasoni N, Aubry JM, Bondolfi G, Osiek C, Schwald M, Bertschy G, Karege F. Partial normalization of serum brain-derived neurotrophic factor in remitted patients after a major depressive episode. *Neuropsychobiol* 2005; **51**: 234-238.
- Giese M, Unternaher E, Huttig H, Beck J, Brand S, Calabrese *et al*. BDNF: an indicator of insomnia? *Mol Psychiatry*; AOP doi: 10.1038/mp.2013.10.
- Godfrey R, Julien M. Urbanisation and health. *Clin Med* 2005; **5**: 137-141.
- Golden E, Emiliano A, Maudsley S, Windham GB, Carlson OD, Egan JM *et al*. Circulating BDNF and indices of metabolic and cardiovascular health: data from the Baltimore longitudinal study of aging. *PLoS ONE* 2010; **5**: e10099.
- Gonul AS, Akdeniz F, Taneli F, Donat O, Eker Ç, Vahip S. Effect of treatment on serum BDNF levels in depressed patients. *Eur Arch Psychiatry Clin Neurosci* 2005; **255**: 381-386.
- Gonul AS, Kitis O, Eker MC, Eker OD, Ozan E, Coburn K. Association of the BDNF val⁶⁶met polymorphism with hippocampus volumes in drug-free depressed patients. *W J Biol Psychiatry* 2011; **12**: 110-118.
- Gonzalez MM, Aston-Jones G. Light deprivation damages monoamine neurons and produces a depressive behavioural phenotype in rats. *Proc Natl Acad Sci USA* 2008; **105**: 4898-4903.

Goodman LA, Thompson KM, Weinfurt K, Corl S, Acker P, Mueser KY, Rosenberg SD. Reliability of reports of violent victimization and posttraumatic stress disorder among men and women with serious mental illness. *J Trauma Stress* 1999; **12**: 587–599

Gordon DJ, Trost DC, Hyde J, Whaley FS, Hannan PJ, Jacobs DR *et al.* Seasonal cholesterol cycles: the lipid research clinics coronary primary prevention trial placebo group. *Circulation* 1987; **76**: 1224–1231.

Gorgulu Y, Caliyurt O. Rapid antidepressant effects of sleep deprivation therapy correlates with serum BDNF changes in major depression. *Brain Res Bull* 2009; **80**: 158–162.

Govindarajan A, Rao BS, Nair D, Trinh M, Mawjee N, Tonegawa S *et al.* Transgenic brain-derived neurotrophic expression causes both anxiogenic and antidepressant effects. *Proc Natl Acad Sci USA* 2006; **103**: 13208–13213.

Grassi-Oliveira R, Stein LM, Lopes RP, Teixeira AL, Bauer ME. Low plasma BDNF and childhood physical neglect are associated with verbal memory impairment in major depression - a preliminary report. *Biol Psychiatry* 2008; **64**: 281–285

Gratacos M, Gonzalez JR, Mercader JM, Urretavizcaya M, Estivill X. BDNF val⁶⁶met and psychiatric disorders: meta-analysis of case-control studies confirm association to substance-related disorders, eating disorders, and schizophrenia. *Biol Psychiatry* 2007; **61**: 911–922.

Green EK, Draddock N. Brain-derived neurotrophic factor as a potential risk factor for bipolar disorder: evidence, limitations, and implications. *Current Psychosis and therapeutics Reports* 2005; **2**: 1530159.

Green MJ, Matheson SL, Sheperd A, Weickert CS, Carr VJ. Brain-derived neurotrophic factor levels in schizophrenia: a systematic review with meta-analysis. *Mol Psychiatry* 2011; **16**: 960–972.

Greenberg P, Siglin L, Finkelstein L, Berndt E. The economic burden of depression in 1990. *J Clin Psychiatry* 1993; **54**: 405–418.

Grønli O, Stensland GØ, Wynn R, Olstad R. Neurotrophic factors in serum following ECT: a pilot study. *World J Biol Psychiatry* 2009; **10**: 295–301.

Groves JO. Is it time to reassess the BDNF hypothesis of depression? *Mol Psychiatry* 2007; **12**: 1079–1088.

Gruber O, Hasan A, Scherk H, Wobrock T, Schneider-Axmann T, Ekwardhani S *et al.* Association of the BDNF val⁶⁶met polymorphism with magnetic resonance spectroscopic markers in the human hippocampus: in vivo evidence for effects on the glutamate system. *Eur Arch Psychiatry Clin Neurosci* 2011; **262**: 23–31.

Grundey SM, Cleeman JJ, Daniels SR, Donato KA, Eckel RH, Franklin BA *et al.* Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005; **112**: 2735–2752.

Guilloux J-P, Douillard-Guilloux G, Kota R, Wang X, Gardier AM, Martinovich K *et al.* Molecular evidence for BDNF-GABA related dysfunctions in the amygdala of female subjects with major depression. *Mol Psychiatry* 2012; **17**: 1130–1142.

Gunstad J, Benitez A, Smith J, Glickman E, Spitznagel MB, Alexander T *et al.* Serum BDNF is associated with cognitive function in healthy older adults. *J Geriatr Psychiatry Neurol* 2008; **21**: 166–170.

Gustafsson G, Lira CM, Johansson J, Wisen A, Wohlfart B, Ekman N *et al.* The acute response of plasma brain-derived neurotrophic factor as a result of exercise in major depressive disorder. *Psychiatry Res* 2009; **169**: 244–248.

Gyekis JP, Yu W, Dong S, Wang H, Qian J, Kota P *et al.* No association of genetic variants in BDNF with major depression: a meta- and gene-based analysis. *Am J Med Genet* 2013; **162**: 61–70.

H

Hajek T, Kopecek M, Hoschl C. Reduced hippocampal volumes in healthy carriers of brain-derived neurotrophic factor val⁶⁶met polymorphism: meta-analysis. *W J Biol Psychiatry* 2012; **13**: 178–187.

Hallböök F. Evolution of the vertebrate neurotrophin and Trk receptor gene families. *Curr Opin Neurobiol* 1999; **6**: 616–621.

Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960; **23**: 56–62.

Hanna GL, Himle JA, Curtis GC, Koram DQ, Van der Weele J, Leventhal BL *et al.* Serotonin transporter and seasonal variation in blood serotonin in families with obsessive-compulsive disorder. *Neuropsychopharmacol* 1998; **18**: 102–111.

Harraguchi S, Sasahara K, Shikmi H, Honda S, Harada N, Tsutsui K. Estradiol promotes purkinje dendritic growth, spinogenesis, and synaptogenesis during neonatal life by inducing the expression of BDNF. *Cerebellum* 2012; **11**: 416–417.

Hariri AR, Goldberg TE, Mattay VS, Kolchana BS, Callicot JH, Egan MF *et al.* Brain-derived neurotrophic factor val⁶⁶met polymorphism affects human-memory related hippocampal activity and predicts memory performance. *J Neurosci* 2003; **23**: 6690–6694.

Harris AHS, Cronkite R, Moos R. Physical activity, exercise coping, and depression in a 10-year cohort study of depressed patients. *J Affect Disord* 2006; **93**: 79–85.

Harris EC, Barraclough B. Excess mortality of mental disorder. *Br J Psychiatry* 1998; **173**: 11–53.

Harvey BH, Hamer M, Louw R, van der Westhuizen FH, Malan L. Metabolic and glutathione redox markers associated with brain-derived neurotrophic factor in depressed African men and women: evidence for counter-regulation. *Neuropsychobiol* 2013; **67**: 33–40.

Hashimoto K, Iwata Y, Nakamura K, Tsujii M, Tsuchiya KJ, Sekine Y *et al.* Reduced serum levels of brain-derived neurotrophic factor in adult male patients with autism. *Prog Neuropsychopharmacol Biol Psychiatry* 2006; **30**: 1529–1531.

Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom Test for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict* 1991; **86**: 1119–1127.

Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biol Psychiatry* 2001; **49**: 1023–1039

Heim C, Shugart M, Craighead WE, Nemeroff CB. Neurobiological and psychiatric consequences of childhood abuse and neglect. *Dev Psychobiology* 2010; **52**: 671–690.

Hellweg R, Ziegenhorn A, Heuser I, Deuschle M. Serum concentrations of nerve growth factor and brain-derived neurotrophic factor in depressed patients before and after antidepressant treatment. *Pharmacopsychiatry* 2008; **41**: 66–71.

Henninger G, Delgado P, Charney D. The revised monoamine theory of depression: a modulatory role for monoamines, based on new findings from monoamine depletion experiments in humans. *Pharmacopsychiatry* 1996; **29**: 2–11.

Higgins JPT, Thompson SG. Quantifying heterogeneity in meta-analysis. *Statist Med* 2002; **21**: 1539–

Hiltunen JO, Arumae U, Moshnyakov M, Saarna M. Expression of mRNAs for neurotrophins and their receptors in developing rat heart. *Circ Res* 1996; **79**: 930–939.

Hirschfeld RM. History and evolution of the monoamine hypothesis of depression. *J Clin Psychiatry* 2000; **61**: S4–S6.

Hohn A, Leibrock J, Bailey K, Barde Y-L. Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family. *Nature* 1990; **344**: 339–341.

Hopkins ME, Davis FC, Vantighem MR, Whalen PJ, Bucci DJ. Differential effects of acute and regular exercise on cognition and affect. *Neurosci* 2012; **215**: 59–68.

Hoshaw BA, Malberg JE, Lucki I. Central administration of IGF-I and BDNF leads to long-lasting antidepressant-like effects. *Brain Res* 2005; **1037**: 204–208.

Hovens JGFM, Wiersma JE, Giltay EJ, van Oppen P, Spinhoven P, Penninx BWJH *et al.* Childhood life events and childhood trauma in adult patients with depressive, anxiety and comorbid disorders versus controls. *Acta Psychiatr Scand* 2009; **122**: 66–74.

Hristova M, Aloe L. Metabolic syndrome - neurotrophic hypothesis. *Med Hypotheses* 2006; **66**: 545-549.

Hu Y, Yu X, Yang F, Si T, Wang W, Tan Y *et al.* The brain-derived neurotrophic factor is associated with the therapeutic efficacy of modified electroconvulsive therapy in Chinese patient with depression. *J ECT* 2010; **26**: 121-125.

Huang TL, Lee CT, Liu YL. Serum brain-derived neurotrophic factor levels in patients with major depression: effects of antidepressants. *J Psychiatr Res* 2008; **42**: 521–525.

Huang M-C, Chen C-H, Liu C-H, Chen C-C, Ho C-C, Leu S-J. Differential patterns of serum brain-derived neurotrophic factor levels in alcoholic patients with and without delirium tremens during acute withdrawal. *Alcoholism: Clin Exp Res* 2011; **35**: 126-131.

I

Ikeda Y, Yahata N, Ito I, Nagano M, Toyota T, Yoshikawa T *et al.* Low serum levels of brain-derived neurotrophic factor and epidermal growth factor in patients with chronic schizophrenia. *Schizophr Res* 2008; **101**: 58-66.

Impey S, McCorkie SR, Cha-Molstad H, Dwyer JM, Yochum GS, Boss J *et al.* Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell* 2004; **119**: 1041-1054.

Ingre M. Why small low-powered studies are worse than large high-powered studies and how to protect against ‘trivial’ findings in research: Comment on Friston. *Neuroimage* 2013; **81**: 496-498.

Ioannidis JPA. Why most published research findings are false. *PLoS MED* 2005; **2**: e124.

Ioannidis JPA. Excess significance bias in the literature on brain volume abnormalities. *Arch Gen Psychiatry* 2011; **68**: 773-780.

J

Jessen F, Schuhmacher A, von Widdern O, Guttenthaler V, Hofels S, Suliman H *et al.* No association of the val⁶⁶ met polymorphism of the BDNF gene with hippocampal volume in major depression. *Psychiatric Genet* 2009; **19**: 99-101.

Jeon HJ, Kang E-S, Lee EH, Jeong E-H, Jeon JR, Mischoulon D *et al.* Childhood trauma and platelet brain-derived neurotrophic factor (BDNF) after a three month follow-up in patients with major depressive disorder. *J Psychiatric Res* 2012; **46**: 966-972.

Joffe RT, Gatt JM, Kemp AH, Grieve S, Dobson-Stone C, Kuan SA, Schofield PR *et al.* BDNF val⁶⁶ met polymorphism, the five-factor model of personality and hippocampal volume: implications for depressive illness. *Hum Brain Mapping* 2009; **30**: 1246-1256.

Johnson VE. Revised standards for statistical evidence. *Proc Natl Acad Sci USA*; AOP November 2013; doi: 10.1073/pnas.1313476110.

Judd FK, Hickey M, Bryant C. Depression and midlife: are we overpathologising the menopause? *J Affect Disord* 2012; **136**: 199-211.

Juhasz G, Dunham JS, McKie S, Thomas E, Downey D, Chase D *et al.* The CREB1-BDNF-NTRK2 pathway in depression: Multiple gene-cognition-environment interactions. *Biol Psychiatry* 2010; **69**: 762-771.

K

Kabacoff RI, Segal DL, Hersen M, van Hasselt VB. Psychometric properties and diagnostic utility of the Beck anxiety inventory and the state-trait anxiety inventory with older adult psychiatric outpatients. *J Anxiety Disord* 1997; **11**: 33–47.

Kachur SG, Hannan CL, Ward KE. Antidepressant-induced weight gain. *Med Health Res* 2005; **88**: 359–361.

Kambeitz JP, Bhattacharya S, Kambeitz-Llankovic LM, Valli I, Collier DA, McGuire P. Effect of BDNF val⁶⁶ met polymorphism on declarative memory and its neural substrate: a meta-analysis. *Neurosci Biobehavioral Rev* 2012; **36**: 2165-2177.

Kanellopoulos D, Gunning FM, Morimoto SS, Hoptman MJ, Murphy GF, Kelly RE *et al.* Hippocampal volumes and the brain-derived neurotrophic factor val⁶⁶ met polymorphism in geriatric major depression. *Am J Geriatr Psychiatry* 2011; **19**: 13-22.

Kang HJ, Voleti B, Hajszan T, Rajkowska G, Stockmeier CA, Licnerski P *et al.* Decreased expression of synapse related genes and loss of synapses in major depressive disorder. *Nature Med* 2011; **18**: 1413-1417.

Kapczinski F, frey BN, Andreazza AC, Kauer-Sant’Anna M, Cunha ABM, Post RM. Increased oxidative stress as a mechanism for decreased BDNF levels in acute manic episodes. *Rev Bras Psiquiatr* 2008; **30**: 243-245.

Kapczinski F, Dal-Pizzol F, Teixeira AL, Magalhaes PVS, Kauer-Sant’Anna M, Klamt F *et al.* A systematic toxicity index developed to assess peripheral changes in mood episodes. *Mol Psychiatry* 2010; **15**: 784-786.

Kapur S, Phillips AG, Insel TR. Why has it taken so long for biological psychiatry to develop clinical tests and what to do about it? *Mol Psychiatry* 2012; **17**: 1174-1179.

Karege F, Perret G, Bondolfi G, Schwald M, Bertschy G, Aubry JM. Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Res* 2002a; **109**: 143-148.

Karege F, Schwald M, Cisse M. Postnatal developmental profile of brain-derived neurotrophic factor in rat brain and platelets. *Neurosci Lett* 2002b; **328**: 261-264.

Karege F, Bondolfi G, Gervasoni N, Schwald M, Aubry J-M, Bertschy G. Low brain-derived neurotrophic factor levels in serum of depressed patients probably result from lower platelet BDNF release unrelated to platelet reactivity. *Biol Psychiatry* 2005; **57**: 1068-1072.

Karg K, Burtmeister M, Shedden K, Sen S. The serotonin transporter promoter variant (5-HTTLPR), stress, and depression. Meta-analysis revisited. *Arch Gen Psychiatry* 2011; **68**: 444-454.

Karlović D, Serretti A, Jevtović S, Vrkić N, Šerić V, Peleš AM. Diagnostic accuracy of serum brain derived neurotrophic factor concentration in antidepressant naïve patients with first major depression episode. *J Psychiatric Res* 2012; **47**: 162-167.

Karnik MS, Wang L, Barch DM, Morris JC, Csernansky JG. BDNF polymorphism rs6265 and hippocampal structure and memory performance in healthy control subjects. *Psychiatry Res* 2010; **178**: 425-429.

Karpova NN, Rantamaki T, Di Lieto A, Lindenmann L, Hoener MC, Castren E. Darkness reduces BDNF expression in the visual cortex and induces repressive chromatin remodelling at the BDNF gene in both hippocampus and visual cortex. *Cell Mol Neurobiol* 2010; **30**: 1117-1123.

Kasper S, Wehr TA, Gartko JJ, Gaist PA, Rosenthal NE. Epidemiological findings of seasonal changes in mood and behavior: a telephone survey of Montgomery County, Maryland. *Arch Gen Psychiatry* 1989; **46**: 823-833.

Kato M, Serretti A. Review and meta-analysis of antidepressant pharmacogenetic findings in major depressive disorder. *Mol Psychiatry* 2008; **15**: 473-500.

Kato-Sembah R, Takeuchi IK, Semba R, Kato K. Distribution of brain-derived neurotrophic factor in rats and its changes with development in the brain. *J Neurochem* 1997; **69**: 34-42.

Katz LC, Schatz CJ. Synaptic activity and the construction of cortical circuits. *Science* 1996; **274**: 1133-1138.

Kauer-Sant’Anna M, Tramontina J, Andreazza AC, Cereser K, daCosta S, Santin A *et al.* Traumatic life events in bipolar disorder: impact on BDNF concentrations and psychopathology. *Bipolar Disord* 2007; **9**: S128–S135.

Kaufman J, Yang B-Z, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshvar S *et al.* Krystal JH, Gelertner J. BDNF–5-HTTLPR gene interactions and environmental modifiers of depression in children. *Biol Psychiatry* 2006; **59**: 673–680.

Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ. Major depressive disorder and generalized anxiety disorder. Same genes, (partly) different environments? *Arch Gen Psychiatry* 1992; **49**: 716–722.

Kendler KS, Neale MC, MacLean CJ, Heath AC, Eaves LJ, Kessler RC. Smoking and major depression. *Arch Gen Psychiatry* 1993; **50**: 36–43.

Kendler KS, Neale MC. Endophenotype: a conceptual analysis. *Mol Psychiatry* 2010; **15**: 789–797.

Kendler KS, Walters EE, Neale MC, Kessler RC, Heath AC, Eaves LJ. The structure of the genetic and environmental risk factors for six major psychiatric disorders in women: phobia, generalized anxiety disorder, panic disorder, bulimia, major depression, and alcoholism. *Arch Gen Psychiatry* 1995; **52**: 374–383.

Kendler KS, Karowski L, Prescott CA. Causal relationships between stressful life events and the onset of major depression. *Am J Psychiatry* 1999; **156**: 837–841.

Kendler KS, Thornton LM, Gardner CO. Stressful life events and previous episodes in the etiology of major depressive disorder in women: and evaluation of the ‘kindling’ hypothesis. *Am J Psychiatry* 2000; **157**: 1243–1252.

Kendler KS. “A gene for ...”: the nature of gene action in psychiatric disorders. *Am J Psychiatry* 2005; **162**: 1243–1252.

Kendler KS. Levels of explanation in psychiatric and substance use disorders: implications for the development of an etiologically based nosology. *Mol Psychiatry* 2012; **17**: 11–21.

Kent JM. SNaRIs, NaSSAs, and NaRIs: new agents for the treatment of depression. *Lancet* 2000; **355**: 911–918.

Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR *et al.* The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA* 2003; **289**: 3095–3105.

Kessler RC, Gruber M, Hettema JM, Hwang I, Sampson N, Yonkers KA. Comorbid major depression and generalized anxiety disorder in the National Comorbidity Survey follow-up. *Psychol Med* 2008; **38**: 365–374.

Khan AK, Jacobson KC, Gardner CO, Prescott CA, Kendler KS. Personality and comorbidity of common psychiatric disorders. *B J Psychiatry* 2005; **186**: 190–196.

Kanzode SD, Dakhale GN, Khanzide SS, Anan PR. Oxidative damage and major depression: the potential antioxidant action of selective serotonin re-uptake inhibitors. *Redox Rep* 2003; **8**: 365–370.

Kim TS, Kim DJ, Lee H, Kim YK. Increased plasma brain-derived neurotrophic factor levels in chronic smokers following unaided smoking cessation. *Neurosci Lett* 2007; **423**: 53–57.

Kim Y-K, Lee H-P, Won S-D, Park E-U, Lee H-Y, Lee B-H *et al.* Low plasma BDNF is associated with suicidal behavior in major depression. *Progress Neuro Psychopharmacol Biol Psychiatry* 2007; **31**: 78–85.

Kirschner PB, Jenkins BG, Schulz JB, Finkelstein SP, Matthews RT, Rosen BR *et al.* NGF, BDNF and NT-5, but not NT-3 protect against MPP⁺ toxicity and oxidative stress in neonatal animals. *Brain Res* 1996; **713**: 178–185.

Klaassen T, Klumperbeek J, Deutz NE, van Praag HM, Griez E. Effects of tryptophan depletion on anxiety and on panic provoked by carbon dioxide challenge. *Psychiatr Res* 1998; **77**: 167–174.

Klein AB, Williamson R, Santini MA, Clemmensen C, Ettrup A, Rios M *et al.* Blood BDNF concentrations reflect brain-tissue BDNF concentrations across species. *Int J Neuropsychopharmacol* 2011; **14**: 347–353.

Kobayakawa M, Inagaki M, Fujimori M, Hamazaki K, Hamazaki T, Akechi T *et al.* Serum brain-derived neurotrophic factor and antidepressant-naïve major depression after lung cancer diagnosis. *Jpn J Clin Oncol* 2011; **4**: 1233–1237.

Kobayashi K, Shimizu E, Hashimoto K, Mitsumori M, Koike K, Okamura N *et al.* Serum Brain-Derived Neurotrophic Factor (BDNF) levels in patients with panic disorder: As a biological predictor of response to group cognitive behavioral therapy. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2005; **29**: 658–663.

Komulainen P, Pedersen M, Hanninen T, Bruunsgaard H, Lakka TA, Kivipelto M *et al.* BDNF is a novel marker of cognitive function in ageing women: the DR’s EXTRA study. *Neurobiol Learning Mem* 2008; **4**: 596–603.

Koolschijn CPMP, van Haren NEM, Bakker SC, Hoogendoorn MLC, Hulshoff Pol HE, Kahn RS. Effects of Brain-derived neurotrophic factor polymorphism on hippocampal volume change in schizophrenia. *Hippocampus* 2010; **20**: 1010–1017.

Krabbe KS, Nielsen AR, Krogh-Madsen R, Plomgaard P, Rasmussen, Erikstrup C *et al.* Brain-derived neurotrophic factor (BDNF) and type 2 diabetes. *Diabetologia* 2007; **50**: 431–438.

Krishnan V, Nestler EJ. Linking molecules to mood: new insight into the biology of depression. *Am J Psychiatry* 2010; **167**: 1305–1320.

Krishnan V, Konakondla S, Nicholas J, Varma A, Kindy M, Wen X. Biomaterial-based interventions for neuronal regeneration and functional recovery in rodent model of spinal cord injury: a systematic review. *J Spin Cord Med* 2013; **36**: 174–190.

Kroenke K, Spitzer RL, Williams JBW, Monahan PO, Lowe B. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Ann Int Med* 2007; **146**: 317–325.

Kubo T, Nonomura T, Enokido Y, Hatanaka H. Brain-derived neurotrophic factor (BDNF) can prevent apoptosis of rat cerebellar granule neurons in culture. *Dev Brain Res* 1995; **85**: 249–258.

Kulkarni J, de Castella A, Fitzgerald PB, Gurvich CT, Bailey M, Burger H. Estrogen in severe mental illness: a potential new treatment approach. *Arch Gen Psychiatry* 2008; **65**: 955–960.

Kuzumaki N, Ikegami D, Tamura R, Hareyama N, Imai S, Narita M *et al.* 2010. Hippocampal epigenetic modification at the brain-derived neurotrophic factor gene induced by an enriched environment. *Hippocampus* 2010; **21**: 127–132.

L

Lacasse JR, Leo J. Serotonin and depression: a disconnect between the advertisements and the scientific literature. *PLoS MED* 2005; **2**: e392. doi:10.1371/journal.pmed.0020392.

Lambert GW, Reid C, Kaye DM, Jennings GL, Esler MD. Effect of sunlight and season on serotonin turnover in the brain. *The Lancet* 2002; **360**: 1840–1842.

Lang UE, Hellweg R, Gallinat J. BDNF serum concentrations in healthy volunteers are associated with depression-related personality traits. *Neuropsychopharmacol* 2004; **29**: 795–798.

Lang UE, Sander T, Lohoff FW, Hellweg R, Bajbouj M, Winterer G *et al.* Association of the met66 allele of brain-derived neurotrophic factor (BDNF) with smoking. *Psychopharmacol (Berl)* 2007a; **190**: 433–439.

Lang UE, Hellweg R, Seifert F, Schubert F, Gallinat J. Correlation between serum brain-derived neurotrophic factor level and an in vivo marker of cortical integrity. *Biol Psychiatry* 2007b; **62**: 530–535.

Lang UE, Hellweg R, Gallinat J, Bajbouj M. Acute prefrontal cortex transcranial magnetic stimulation in healthy volunteers: no effects on brain-derived neurotrophic factor concentrations in serum. *J Affect Disord* 2008; **107**: 255–258.

Lang UE, Bajbouj M, Gallinat J, Hellweg R. Brain-derived neurotrophic factor serum concentrations in depressive patients during vagus nerve stimulation and repetitive transcranial magnetic stimulation. *Psychopharmacol* 2006; **187**: 56-59.

Lang UE, Hellweg R, Sander T, Gallinat J. The Met allele of the BDNF val⁶⁶ met polymorphism is associated with increased serum BDNF concentrations. *Mol Psychiatry* 2009; **14**: 120-122.

Lanz TA, Bove SE, Pilsmaier CD, Mariga A, Drummond EM, Cadelina GW *et al*. Robust changes in expression of brain-derived neurotrophic factor mRNA and protein across the brain do not translate to detectable changes in BDNF levels in CSF or plasma. *Biomarkers* 2012; **17**: 524-531.

Laske C, Stransky E, Eschweiler GW. Increased BDNF serum concentration in fibromyalgia with or without depression or antidepressants. *J Psychiatr Res* 2007; **41**: 600-605.

Lau JYF, Goldman D, Buzas B, Hodgkinson C, Leibenluft E, Nelson E *et al*. BDNF gene polymorphism (val⁶⁶ met) predicts amygdala and anterior hippocampus responses to emotional faces in anxious and depressed adolescents. *Neuroimage* 2010; **53**: 952-961.

Lavebratt C, Aberg E, Sjoholm LK, Forsell Y. Variations in FKBP5 and BDNF genes are suggestively associated with depression in a Swedish population-based cohort. *J Affect Disord* 2010; **125**: 249-255.

Lee R, Kermani P, Teng KK, Hempstead BL. Regulation of the survival secreted proneurotrophins. *Science* 2001; **294**: 1945-1948.

Lee B-H, Kim Y, Park S-H, Kim Y-K. Decreased plasma BDNF level in depressive patients. *J Affect Disord* 2007; **101**: 239-244.

Lee H-Y, Kim Y-K. Plasma brain-derived neurotrophic factor as a peripheral marker for the action mechanism of antidepressants. *Neuropsychobiology* 2008; **57**: 194-199.

Lee JG, Shin BS, You YS, Kim JE, Yoon SW, Jeon DW *et al*. Decreased serum brain-derived neurotrophic factor levels in elderly Korean with dementia. *Psychiatry Invest* 2009; **6**: 299-305.

Lenth RV. Some practical guidelines for effective sample size determination. *Am Statistician* 55: 187-193.

Levi-Montalcini R. The nerve growth factor: its mode of action on sensory and sympathetic nerve cells. *Harvey Lect* 1966; **60**: 217-259.

Levi-Montalcini R. The nerve growth factor: thirty-five years later. *Biosci Rep* 1987; **9**: 681-699.

Lewy AJ, Lefler BJ, Emens JS, Bauer VK. The circadian rhythm of winter depression. *Proc Natl Acad Sci USA* 2006; **103**: 7414-7419.

Li N, Liu G-T. The novel squamosome derivative FLZ enhances BDNF/TrkB/CREB signaling and inhibits neuronal apoptosis in APP/PS1 mice. *Acta Pharmacologica Sin* 2010; **31**: 265-272.

Liao G-Y, An JJ, Gharami K, Waterhouse EG, Vanevski F, Jones KR *et al*. Dendritically targeted BDNF mRNA is essential for energy balance and response to leptin. *Nat Med* 2012; **18**: 564-572.

Licinio J, Dong C, Wong M-L. Novel sequence variations in the brain-derived neurotrophic gene and association with major depression and antidepressant treatment response. *Arch Gen Psychiatry* 2009; **66**: 488-497.

Lindsay RM, Wiegand SJ, Altar CA, Di Stefano PS. Neurotrophic factors: from molecule to man. *Trends Neurosci* 1994; **17**: 182-190.

Little J, Higgins JP, Ioannidis JPA, Moher D, Gagnon F, von Elm E *et al*. Strengthening the Reporting of Genetic Association Studies (STREGA): an extension of the STROBE statement. *Hum Gen* 2009; **125**: 131-151.

Liu QR, Walther D, Drgon T, Polesskaya O, Lesnick TG, Strain KJ *et al*. Human brain-derived neurotrophic genes, splicing patterns, and assessments of associations with substance abuse and Parkinson's disease. *Am J Med Gen* 2005; **134**: 93-103.

Lieverse R, van Someren EJW, Nielen MMA, Uitdehaag BMJ, Smit JH, Hoogendijk WJG. Bright light treatment in Elderly patients with non-seasonal major depressive disorder: a randomized placebo-controlled trial. *Arch Gen Psychiatry* 2011; **68**: 61-70.

Lohmueller KE, Pearce CL, Pike M, Lander ES, Hirschhorn JN. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to disease. *Nat Genet* 2003; **33**: 177-182.

Lommatzsch M, Schloetcke K, Klotz J, Schuhbaeck K, Zingler D, Zingler C *et al*. Brain-derived neurotrophic factor in platelets and airflow limitation in asthma. *Am J Respir Crit Care Med* 2005; **171**: 115-120.

Lommatzsch M, Hornych K, Zingler C, Schuff-Werner P, Hoppner J, Virchow JC *et al*. Maternal serum concentrations of BDNF and depression in the perinatal period. *Psychoendocrinol* 2006; **31**: 388-394.

Longo FM, Massa SM. Small-molecule modulation of neurotrophin receptors: a strategy for the treatment of neurological disease. *Nat Rev Drug Discov* 2013; **12**: 507-525.

Lu B, Gottschalk W. Modulation of hippocampal synaptic transmission and plasticity by neurotrophins. *Prog Brain Res* 2000; **128**: 231-241.

Lu B. BDNF and activity-dependent synaptic modulation. *Learn Mem* 2003; **10**: 86-98.

Lu B, Pang PT, Woo NH. The Yin and Yang of neurotrophin action. *Nat Rev Neurosci* 2005; **6**: 603-614.

Lu B, Nagappan G, Guan X, Nathan PJ, Wren P. BDNF based synaptic repair as a disease-modifying strategy for neurodegenerative diseases. *Nat Rev Neurosci* 2013; **14**: 401-416.

Luykx J, Bakker SC, Lentjes E, Moks MP, van Geloven N, Eijkemans MJ *et al*. Season of sampling and season of birth influence serotonin metabolite levels in human cerebrospinal fluid. *PLoS ONE* 2012; **e7**: e30497.

Lyketsos CG, Nestadt G, Cwi J, Heithoff K, Eaton WW. The life chart interview: a standardized method to describe the course of psychopathology. *Int J Methods Psychiatr Res* 1994; **4**: 143-155.

M

Ma Y, Olendzki BC, Li W, Hafner AR, Chiriboga D, Hebert JR *et al*. Seasonal variation in food intake, physical activity, and body weight in a predominantly overweight population. *Eur J Clin Nutr* 2006; **60**: 519-528.

Machado-Vieira R, Dietrich MO, Leke R, Cereser VH, Zanatto V, Kapczinski F *et al*. Decreased plasma brain derived neurotrophic factor levels in unmedicated bipolar patients during manic episode. *Biol Psychiatry* 2007; **61**: 142-144.

Machado-Vieira R, Yuan P, Brutsche N, Diaz-Granados N, Luckenbaugh D, Manji HK *et al*. BDNF and initial antidepressant response to an N-methyl-D-aspartate antagonist. *J Clin Psychiatry* 2009; **70**: 1662-1666.

Maes M, Gelecki P, Chang Y-S, Berk M. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; **35**: 676-692.

Magarinos AM, Li CJ, Gal Toth J, Bath KG, Jing D, Lee FS *et al*. Effect of brain-derived neurotrophic factor haploinsufficiency on stress-induced remodeling of hippocampal neurons. *Hippocampus* 2011; **21**: 253-264.

MacQueen G, Frodl T. The hippocampus in major depression: evidence for the convergence of the bench and bedside in psychiatric research? *Mol Psychiatry* 2011; **16**: 252-264.

Malberg JE, Eisch AJ, Nestler EJ, Duman RS. Chronic antidepressant treatment increases neurogenesis in adult neurogenesis in adult rat hippocampus. *J Neurosci* 2000; **20**: 9104-9110.

Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 2003; **19**: 1233-1239.

Maller JJ, Daskalakis ZJ, Thomson RHS, Daigle M, Barr MS, Fitzgerald PB. Hippocampal volumetrics in Treatment-Resistant Depression and Schizophrenia: The Devil's in De-Tail. *Hippocampus* 2012; **22**: 9-16.

Maisonpierre PC, Le Beau MM, Espinosa III R, Ip NY, Belluscio L, de la Monte SM et al. Human and rat brain-derived neurotrophic factor and neurotrophin-3: gene structures, distributions, and chromosomal localizations. *Genomics* 1991; **10**: 558-568.

Maisonpierre PC, Belluscio L, Squinto S, Furth M, Lindsay R, Yancopoulos G. Neurotrophin-3: a neurotrophic factor related to NGF and BDNF. *Science* 1990; **247**: 1446-1451.

Mandelman SD, Grigorenko EL. BDNF val⁶⁶met and cognition: all, none, or some? A meta-analysis of the genetic association. *Genes, Brain, Beh* 2012; **11**: 127-136.

Mann JJ. The medical management of depression. *N Engl J Med* 2005; **353**: 1819-1834.

Manni L, Nikolova V, Vyagova D, Chaldakov GN, Aloe L. Reduced plasma levels of NGF and BDNF in patients with acute coronary syndromes. *Int J Cardiol* 2005; **102**: 169-171.

Marano CM, Phatak P, Vemulapalli UR, Sasan A, Nalbandyan MR, Ramanujam S et al. 2007. Increased plasma concentration of BDNF with electroconvulsive therapy: a pilot study in patients with major depression. *J Clin Psychiatry* 2007; **68**: 512-517.

Maron E, Toru I, Vasar V, Shlik J. The effect of 5-hydroxy tryptophan on cholecystokinin-4-induced panic attacks in healthy volunteers. *J Psychopharmacol* 2004; **18**: 194-199.

Martin DW. Doing psychological experiments (1996; 4th ed.). Pacific Grove, CA: Brooks Colorado.

Martinowich K, Hattori D, Wu H, Fouse S, He F, Fan G et al. DNA methylation-related chromatin remodeling in activity-dependent BDNF gene regulation. *Science* 2003; **302**: 890-893.

Martinowich K, Manji H, Lu B. New insights into BDNF function in depression and anxiety. *Nat Neurosci* 2007; **10**: 1089-1093.

Martinowich K, Lu B. Interaction between BDNF and serotonin: role in mood disorders. *Neuropsychopharmacol* 2008; **33**: 73-83.

Matrisciano F, Bonaccorso S, Ricciardi A, Scaccianoce S, Panaccione I, Wang L et al. Changes in BDNF serum concentrations in patients with major depression disorder after 6 months treatment with sertraline, citalopram, or venlafaxine. *J Psych Res* 2009; **43**: 247-254.

Mattson MP, Maudsley S, Martin B. BDNF and 5-HT: a dynamic duo in age related neuronal plasticity and neurodegenerative disorders. *Trends Neurosci* 2004; **27**: 589-594.

Mattson MP. Evolutionary aspects of human exercise – born to run purposefully. *Ageing Res Rev* 2012; **11**: 347-352.

McLaughlin KA, Green J, Gruber MJ, Sampson MA, Zaslavsky AM, Kessler RC. Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication II. *Arch Gen Psychiatry* 2010; **67**: 124-132.

McConnel MJ, Lindberg MR, Brennand KJ, Piper JC, Voet T, Cowing-Zitron C et al. Mosaic copy number variation in human neurons. *Science* 2013; **342**: 632-637.

Mendels J, Stinnett J, Burns D, Frazer A. Amine precursors and depression. *Arch Gen Psychiatry* 1975; **32**: 22-30.

Millan Sanchez M, Das D, Taylor JL, Noda A, Yesavage JA, Salehi A. BDNF polymorphism predicts the rate of decline in skilled task performance and hippocampal volume in healthy individuals. *Trans Psychiatry* 2011; **1**: e51; doi:10.1038/tp. 2011.47.

Miyajima F, Ollier W, Mayes A, Jackson A, Thacker N, Rabbitt R et al. BDNF val⁶⁶met influences cognitive abilities in the elderly. *Genes Brain Beh* 2008; **7**: 411-417.

Moffet JR, Ross B, Arun P, Madhavarao CN, Namboodiri AM. N-Acetylaspate in the CNS: from neurodiagnostics to neurobiology. *Progress in Neurobiology* 2007; **81**: 89-131.

Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the prisma statement. *Ann Intern Med* 2009; **151**: 264-269.

Molendijk ML, Bus BAA, Spinhoven P, Penninx, BJWH, Kenis G, Prickaerts J et al. Serum BDNF concentrations in major depressive disorder: state and trait issues, clinical features, and pharmacological treatment. *Mol Psychiatry* 2011; **16**: 1088-1095.

Molendijk ML, Bus BAA, Spinhoven P, Penninx BWJH, Prickaerts J, Oude Voshaar RC et al. Gender specific associations of serum BDNF concentrations in anxiety. *W J Biol Psychiatry* 2012; **13**: 535-543.

Molendijk ML, van Tol M-J, Penninx BWJH Penninx, van der Wee NJA, Aleman A et al. BDNF val⁶⁶met affects hippocampal volume and memory related hippocampal activity. *Trans Psychiatry* 2010; **2**: e74.

Molendijk ML, Haffmans JPM, Bus BAA, Spinhoven P, Penninx BWJH, Prickaerts J et al. BDNF concentrations show strong seasonal variation and correlations with the amount of ambient sunlight *PLoS ONE* 2012; **7**: e48046.

Molendijk ML, Bus BAA, Spinhoven P, Kaimatzoglou A, Oude Voshaar RC, Penninx BWJH et al. A systematic review and meta-analysis on the association between BDNF val⁶⁶met and hippocampal volume – a genuine effect or a winners curse? *Am J Med Genet* 2012; **159**: 731-740.

Montag C, Basten U, Stelzel C, Fiebach CJ, Reuter M. The BDNF val⁶⁶met polymorphism and smoking. *Neurosci Lett* 2008; **442**: 30-33.

Montag C, Reuter M, Newport B, Elger C, Weber B. The BDNF val⁶⁶met polymorphism affects amygdala activity in response to emotional stimuli: evidence from a genetic imaging study. *Neuroimage* 2008; **42**: 1554-1559.

Montag C, Weber B, Fliessbach K, Elger C, Reuter M. The BDNF val⁶⁶met polymorphism impacts parahippocampal and amygdala volume in healthy humans: incremental support for a genetic risk for depression. *Psychological Med* 2009; **39**: 1831-1839

Monteggia LM, Luikart B, Barrot M, Theobald D, Malkovska I, Nef S et al. Brain-derived neurotrophic factor conditional knockouts show gender differences in depression-related behaviors. *Biol Psychiatry* 2007; **61**: 187-197.

Monteleone P, Artini PG, Simi G, Cela V, Casarosa E, Begliuomini S et al. Brain derived neurotrophic factor circulating levels in patients undergoing IVF. *J Assist Reprad Genet* 2007; **24**: 477-480.

Monteleone P, Serritella C, Martiadis V, Maj M. Decreased concentrations of BDNF in both depressed and euthymic patients with unipolar depression and in euthymic patients with bipolar I and II disorders. *Bipolar Disord* 2008; **10**: 95-100.

Monteleone P, Fabrazzo M, Martiadis V, Serritella C, Pannuto M, Maj M. Circulating BDNF is decreased in women with anorexia and bulimia nervosa but not in women with binge eating disorder: relationships to co-morbid depression, psychopathology and hormonal variables. *Psychol Med* 2005; **35**: 897-905.

Montgomery SA, Åsberg MA. A new depression scale designed to be sensitive to change. *Br J Psychiatry* 1979; **134**: 382-389.

Mossner R, Mikova O, Koutsilieri E, Saoud M, Ehli AC, Muller N et al. Consensus paper of the WFSBP Task Force on Biological Markers: biological markers in depression. *World J Biol Psychiatry* 2007; **8**: 141-174.

Mosteller F, Colditz GA. Understanding research synthesis. *Annu Rev Public Health* 1996; **17**: 1-23.

Murakami S, Imbe H, Morikawa Y, Kubo C, Senba E. Chronic stress, as well as acute stress, reduces BDNF mRNA expression in the rat hippocampus but less robustly. *Neurosci Res* 2005; **53**: 129-139

Munafo MR, Flint J. Meta-analysis of genetic association studies. *Trends Gen* 2004; **20**: 439-444.

Murad MH, Montori VM. Synthesizing evidence: shifting the focus from individual studies to the body of evidence. *JAMA Psychiatry* 2013; **309**: 2217-2218.

Murer MG, Yan Q, Raisman-Vosari R. BDNF in the control human brain, and in Alzheimer's disease and Parkinson's disease. *Prog Neurobiology* 2001; **63**: 71-124.

Murphy GM, Sarginson JE, Ryan HS, O'Hara R, Schatzberg AF, Lazzeroni LC. BDNF and CREB1 genetic variants interact to affect antidepressant treatment outcomes in geriatric depression. *Pharmacogenet Genomics* 2013; **23**: 301-313.

N

Nagahara AH, Tuszynski MH. Potential therapeutic uses of BDNF in neurological and psychiatric disorders. *Nat Rev Drug Discovery* 2011; **10**: 209-219.

Nakahashi T, Fujimura H, Altar CA, Li J, Kambayashi J, Tandon NN et al. Vascular endothelial cells synthesize and secrete brain-derived neurotrophic factor. *FEBS Lett* 2000; **470**: 113-117.

Nakazato M, Hashimoto K, Shimizu E, Kumakiri C, Koizumi H, Okamura N et al. Decreased concentrations of serum BDNF in female patients with eating disorders. *Biol Psychiatry* 2003; **54**: 485-490.

Nederhof E, Bouma, EM, Oldewinkel AJ, Ormel J. Interaction between childhood adversity, BDNF val/met and serotonin transporter promoter polymorphism on depression: the TRAILS study. *Biol Psychiatry* 2010; **68**: 209-212

Nemeroff CB, Owens MJ. Treatment of mood disorders. *Nat Neurosci* 2002; **6**: 1068-1070.

Nemoto K, Fukamachi K, Nemoto F, Miyata S, Hamada M, Nakamura Y et al. Gene expression of neurotrophins and their receptors in cultured rat vascular smooth muscle cells. *Biochem Biophys Res Commun* 1998; **245**: 284-288.

Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. *Neuron* 2000; **34**: 13-25.

Nestler EJ, Gould E, Manji H, Bucan M, Duman RS, Gershenfeld HK et al. Preclinical models: status of basic research in depression. *Biol Psychiatry* 2002; **52**: 503-528.

Neumeister A, Pirker W, Willeit M, Praschak-Rieder N, Asenbaum S, Brucke T et al. Seasonal variation of availability of serotonin transporter binding sites in healthy female subjects as measured by [123I]-2 beta-carbomethoxy-3 beta-(4-iodophenyl)tropane and single photon emission computed tomography. *Biol Psychiatry* 2000; **47**: 158-160.

Nibuya M, Morinobu S, Duman RS. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci* 1995; **15**: 7539-7547.

Nowak J, Murray JJ, Oates JA, Fitzgerald GA. Biochemical evidence of a chronic abnormality in platelet and vascular function in healthy individuals who smoke cigarettes. *Circulation* 1987; **76**: 6-14.

Nugraha B, Korallus C, Gutenbrunner G. Serum level of BDNF in fibromyalgia syndrome correlates with depression but not anxiety. *Neurochem Int* 2013; **62**: 281-286.

O

Okamoto T, Yoshimura R, Ikenouchi-Sugita A, Hori H, Umene-Nakano W, Inoue Y et al. Efficacy of Electroconvulsive therapy is associated with changing blood levels of homovanilic acid and BDNF in refractory depressed patients: a pilot study. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; **32**: 1185-1190.

Oral E, Canpolat S, Yildirim S, Gulec M, Aliyev E, Aydin N et al. Cognitive functions and serum levels of BDNF in patients with major depressive disorder. *Brain Res Bull* 2012; **88**: 454-459.

Ozan E, Okur H, Eker C, Eker DE, Gönül AS, Akarsu N. The effect of depression, BDNF gene val⁶⁶met polymorphism and gender on serum BDNF levels. *Brain Res Bull* 2010; **81**: 61-65.

P

Pan W, Banks WA, Fasold MB, Bluth J, Kastin AJ. Transport of BDNF across the blood-brain barrier. *Neuropharmacology* 1998; **37**: 1553-1561.

Papakostas GI, Shelton RC, Kinrys G, Henry ME, Bakow BR, Lipkin SH et al. Assessment of multi-assay, serum-based biological diagnostic test for major depressive disorder: a pilot and replication study. *Mol Psychiatry* 2013; **18**: 332-339.

Pariente CM, Miller AH. Glucocorticoid receptors in major depression: relevance to pathophysiology and treatment. *Biol Psychiatry* 2001; **49**: 391-404.

Park H, Poo M-M. Neurotrophin regulation of neuronal circuit development and function. *Nat Rev Neurosci* 2013; **14**: 7-23.

Paykel ES. The clinical interview for depression development, reliability and validity. *J Affect Disord* 1985; **8**: 85-96.

Paykel ES, Ramana R, Cooper Z, Hayhurst H, Kerr J, Barocka A. Residual symptoms after partial remission: an important outcome in depression. *Psychol Med* 1995; **25**: 1171-1180.

Peen J, Schoevers RA, Beekman AT, Dekker J. The current status of urban-rural differences in psychiatric disorders. *Acta Psychiatr Scand* 2009; **121**: 84-93.

Penninx BWJH, Beekman AT, Smit JH, Nolen WA, Spinhoven P, Cuijpers P et al. The Netherlands Study of Depression and Anxiety (NESDA): rational, objectives and methods. *Int J Methods Psychiatr Res* 2008; **17**: 121-140.

Penninx BWJH, Milaneschi Y, Lamers F, Vogelzangs N. Understanding the somatic consequences of depression: biological mechanisms and the role of depression symptom profile. *BMC Med* 2013; **11**: 129-138.

Perlis R. A clinical stratification tool for predicting treatment resistance in major depressive disorder. *Biol Psychiatry* 2013; **74**: 7-14.

Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Performance of the trim and fill method in the presence of publication bias and between-study heterogeneity. *Statist Med* 2007; **26**: 4544-4562.

Petryshen TL, Sabeti PC, Aldinger KA, Fry B, Fan JB, Schaffner SF et al. Population genetic study of the Brain-Derived Neurotrophic Factor (BDNF) gene. *Mol Psychiatry* 2010; **15**: 810-815

Pezawas L, Verchinski BA, Mattay VS, Callicot JH, Kolochana BS, Straub RE et al. the BDNF val⁶⁶met polymorphism and variation in human cortical morphology. *J Neurosci* 2004; **24**: 10099-10102.

Piccinni A, Marazziti D, Del DA, Bianchi C, Roncaglia I, Mannari C et al. Diurnal variation of plasma brainderived neurotrophic factor (BDNF) in humans: an analysis of sex differences. *Chronobiol Int* 2008; **25**: 819-826.

Piccinni A, Marazziti D, Catena M, Domenici L, Del Debbio A, Bianchi C et al. Plasma and serum brain-derived neurotrophic factor (BDNF) in depressed patients during 1 year of antidepressant treatment. *J Affect Disord* 2008; **105**: 279-283.

Pillai A, Kale A, Joshi S, Naphade N, Raju MS, Nasrallah H et al. Decreased BDNF levels in CSF of drug-naïve first episode psychotic subjects : correlation with plasma BDNF and psychopathology. *Int J Neuropsychopharmacol* 2010; **13**: 535-539.

Pillai A, Bruno D, Sarreal SS, Hernando RT, Saint-Louis LA, Nierenberg J et al. Plasma BDNF levels vary in relation to body weight in females. *PLoS ONE* 2012; **7**: doi: 10.1371/journal.pone.0039358.

Pittenger C, Duman RS. Stress, depression, and neuroplasticity: a convergence of mechanisms. *Neuropsychopharmacol* 2008; **33**: 88-109.

Poo MM. Neurotrophins as synaptic modulators. *Nat Rev Neurosci* 2001; **2**: 24-32.

Post RM. Kindling and sensitization as models for affective episode recurrence, cyclicity, and tolerance phenomena. *Neurosci Biobehav Rev* 2007; **31**: 858-873.

Postolache TT, Mortensen PB, Tonelli LH, Jiao X, Frangakis C, Soriano JJ et al. Seasonal spring peaks of suicide in victims with and without prior hospitalization for mood disorders. *J Affect Disord* 2010; **121**: 88-93.

Prasad V, Jena AB. Prespecified falsification end points: can they validate true observational associations? *JAMA* 2013; **209**: 241-242.

Praschak-Rieder N, Willeit M, Wilson AA, Houle S, Meyer JH. Seasonal variation in human serotonin transporter binding. *Arch Gen Psychiatry* 2008; **65**: 1072-1078.

Prendergast BJ, Kay LM. Affective and adrenocorticotrophic responses to photoperiod in Wistar rats. *J Endocrinol* 2008; **20**: 261-267.

Prickaerts J, van den Hove DL, Fierens FL, Kia HK, Lenaerts I, Steckler T. Chronic corticosterone manipulations in mice affect brain cell proliferation rates, but only partly affect BDNF protein levels. *Neurosci Lett* 2006; **396**: 12-16.

Pruunsild P, Kazantseva A, Aid T, Palm K, Timmusk T. Dissecting the human BDNF locus: bidirectional transcription, complex splicing, and multiple promoters. *Genomics* 2007; **90**: 397-406.

Pyter LM, Reader BF, Nelson RJ. Short photoperiods impair spatial learning and alter hippocampal dendritic morphology in adult male white-footed mice (*Peromyscus leucopus*). *J Neurosci* 2005; **25**: 4521-4526.

Pyter LM, Nelson RJ. Enduring effects of photoperiod on affective behaviors in Siberian hamsters (*Phodopus sungorus*). *Behav Neurosci* 2006; **120**: 125-134.

Q

Quilty LC, Meusel LA, Bagby RM. Neuroticism as a mediator of treatment response to SSRIs in major depressive disorder. *J Affect Disord* 2008; **111**: 67-73.

R

Rakofsky JJ, Ressler KJ, Dunlop BW. BDNF function as a potential mediator of bipolar disorder and post-traumatic stress disorder comorbidity: *Mol Psychiatry* 2012; **17**: 22-35

Rasmusson AM, Shi L, Duman R. Downregulation of BDNF mRNA in the hippocampal dentate gyrus after re-exposure to cues previously associated with footshock. *Neuropsychopharmacol* 2002; **27**: 133-142.

Reichardt LF. Neurotrophin-regulated signaling pathways. *Phil Trans R Soc B* 2006; **361**: 1545-1564.

Reif A, Fritzen S, Finger M, Strobel A, Lauer M, Schitt A et al. Neural stem cell proliferation is decreased in schizophrenia, but not in depression. *Mol Psychiatry* 2006; **11**: 514-522.

Richter-Schmidinger T, Alexopoulos P, Horn M, Maus S, Reichel M, Rhein C et al. Influence of BDNF and apolipoprotein E genetic variants on hippocampal volume and memory performance in healthy young adults. *J Neural Trans* 2011; **118**: 249-257.

Roceri M, Cirulli F, Pessina C, Peretto P, Racagni G, Riva MA. Postnatal repeated maternal deprivation produces age-dependent changes of brain-derived neurotrophic factor expression in selected rat brain regions. *Biol Psychiatry* 2004; **55**: 708-714

Rojas-Vega S, Struder HK, Wahrmann BV, Schmidt A, Bloch W, Hollmann W. Acute BDNF and cortisol response to low intensity exercise and following ramp incremental exercise to exhaustion in humans. *Brain Res* 2006; **1121**: 59-65.

Rojas PS, Fritsch R, Rojas RA, Jara P, Fiedler JL. Serum BDNF and glucocorticoid receptor levels in lymphocytes as markers of antidepressant response in major depressive patients: a pilot study. *Psychiatry Res* 2011; **189**: 239-245.

Rosenfeld RD, Zeni L, Haniu M, Talvenheimo J, Radka SF, Bennett L et al. Purification and identification of brain-derived neurotrophic factor from human serum. *Protein Expr Purif* 1995; **6**: 465-471.

Rosenthal NE, Sack DA, Gillen JC, Lewy AJ, Goodwin FK, Davenport Y et al. Seasonal affective disorder: a description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry* 1984; **41**: 72-80.

Rush AJ, Gullion CM, Basco MR, Jarrett RB, Trivedi MH. The inventory of depressive symptomatology: psychometric properties. *Psychol Med* 1996; **26**: 477-486.

Rush AJ, Trivedi MH, Wisniewski SR, Nierenberg AA, Stewart JW, Warden D et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry* 2006; **163**: 1905-1917.

Rush AJ. The varied clinical presentations of major depressive disorders. *J Clin Psychiatry* 2007; **68**: S4-S10.

Russo-Neustadt AA, Ha T, Ramirez R, Kessler JP. Physical activity-antidepressant treatment combination: impact on brain-derived neurotrophic and behavior in an animal model. *Beh Brain Res* 2001; **120**: 87-95.

Rybakowski A, Czerski PM, Skibinska M, Hauser J. Polymorphisms on the BDNF gene and performance on a cognitive prefrontal test in bipolar patients. *Bipolar Disord* 2003; **5**: 468-472.

S

Sanderson S, Tatt ID, Higgs JPT. Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography. *Int J Epidemiol* 2007; **36**: 666-676.

Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, Dulawa S et al. Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 2003; **301**: 805-809.

Sapolsky RM, Plotsky PM. Hypercortisolism and its possible neural bases. *Biol Psychiatry* 1990; **27**: 937-952.

Sapolsky RM. Why stress is bad for your brain. *Science* 1996; **273**: 749-750.

Sairanen M, Lucas G, Ernfors P, Castren M, Castren E. BDNF and antidepressant drugs have differentiated but coordinated effects on neuronal turnover, proliferation, and survival in the adult dentate gyrus. *J Neurosci* 2005; **25**: 1089-1094.

Sarrias MJ, Artigas F, Martinez E, Gelpi E. Seasonal changes of plasma serotonin and related parameters: correlation with environmental measures. *Biol Psychiatry* 1989; **26**: 695-706.

Sartorius A, Hellweg R, Litzke J, Vogt M, Dormann C, Vollmayr B et al. Correlations and discrepancies between serum and brain tissue concentrations of neurotrophins after electroconvulsive treatment in rats. *Pharmacopsychiatry* 2009; **42**: 270-276.

Sasaki T, Niitsu T, Hashimoto T, Kanahara N, Shiina A, Hasegawa T et al. Decreased levels of brain-derived neurotrophic factor in male pediatric patients with depression. *Open Clin Chem J* 2011; **4**: 28-33.

Satohmura E, Baba H, Nakano Y, Maeshima H, Suzuki T, Arai H. Correlations between BDNF and clinical symptoms in medicated patients with major depression. *J Affect Disord* 2011; **135**: 332-335.

Scharfman HE, MacLusky NJ. Similarities between actions of estrogen and BDNF in the hippocampus: coincidence or clue? *Trends Neurosci* 2004; **28**: 79-85.

Schildkraut JJ. The catecholamine hypothesis of affective disorders: a review of supporting evidence. *J Neuropsychiatry Clin Neurosci* 1965; **7**: 524-533.

Schmidt HD, Duman RS. Peripheral BDNF produces anti-depressant-like effects in cellular and behavioral models. *Neuropsychopharmacol* 2010; **35**: 2378-2391.

Schmidt HD, Shelton RC, Duman RS. Functional biomarkers of depression: diagnosis, treatment, and pathophysiology. *Neuropsychopharmacol* 2011; **36**: 2375 – 2394.

Schofield PR, Williams LM, Paul RH, Gatt JM, Brown K, Luty A et al. Disturbances in selective information processing associated with the BDNF val⁶⁶met Polymorphism: evidence from cognition, the P300 and fronto-hippocampal systems. *Biol Psychology* 2009; **80**: 176-188.

Seeman MV. Psychopathology in women and men: focus on female hormones. *Am J Psychiatry* 1997; **154**: 1641 – 1647.

Sen S, Duman R, Sanacora G. Serum brain-derived neurotrophic factor, depression, and antidepressant medications: meta-analyses and implications. *Biol Psychiatry* 2008; **64**: 527-532.

Sharp T, Cowen PJ. 5-HT and depression: is the glass half-full? *Curr Opin Pharmacol* 2011; **11**: 45-51.

Shimizu E, Hashimoto K, Okamura N, Koike K, Komatsu N, Kumakiri C et al. Alterations of serum concentrations of BDNF in depressed patients with or without antidepressants. *Biol Psychiatry* 2003; **54**: 70–75.

Simon D, Boring III Jr. Sensitivity, specificity, and predictive value. In *Clinical Methods: the history, physical, and laboratory examinations*, Walker HK, Hall WD, Hurst JW eds. Butterworths 3rd edition, 1990.

Simon GE, von Korff M, Saunders K, Miglioretti DL, Crane PK, van Belle G et al. Association between obesity and psychiatric disorders in the US adult population. *Arch Gen Psychiatry* 2006; **63**: 824–830.

Siuciak JA, Lewis DR, Wiegand SJ, Lindsay RM. Antidepressant-like effect of BDNF. *Pharmacol Biochem Behav* 1996; **56**: 131-173.

Smith MA, Makino S, Kvetnansky R, Post RM. Stress and glucocorticoids affect the expression of brain-derived neurotrophic factor and neurotrophin-3 mRNAs in the hippocampus. *J Neurosci* 1995; **15**: 1768–1777.

Smith GD, Ebrahim S. Data dredging, bias, or confounding: they can all get you into the *BMJ* and the Friday papers. *BMJ* 2002; **325**: 1437-1438.

Sohrabji F, Lewis DK. Estrogen-BDNF interactions: implications for neurodegenerative diseases. *Front Neuroendocrinol* 2006; **27**: 404-414.

Song L, Che W, Min-Wei W, Murakami Y, Matsumoto K. Impairment of the spatial learning and memory induced by learned helplessness and chronic mild stress. *Pharmacol Biochem Behav* 2006; **83**: 186–193.

Song XY, Li F, Zhang FH, Zhong FH. Peripherally-derived BDNF promotes regeneration of ascending sensory neurons after spinal cord injury. *PLoS ONE* 2008; **3**: e1707.

Sözeri-Varma G, Enli Y, Toker-Uğurlu T, Alaçam H, Kalkan-Oğuzhanoglu N. Decreased serum BDNF levels in major depressive patients. *Neurol Psychiatry Brain Res* 2011; **17**: 84-88.

Spalding KL, Bergmann O, Alkass K, Bernard S, Salehpour M, Huttner HB et al. Dynamics of hippocampal neurogenesis in adult humans. *Cell* 2013; **153**: 1219-1227.

Spencer JL, Waters EM, Milner TA, Lee FS, McEwen BS. BDNF variant val⁶⁶met interacts with estrous cycle in the control of hippocampal function. *Proc Natl Acad Sci USA* 2010; **107**: 4395-4400.

Spinhoven P, Elzinga BM, Hovens JGFM, Roelofs K, Zitman FG, van Oppen P et al. The specificity of childhood adversities and negative life events across the life span to anxiety and depressive disorders. *J Affect Disord* 2010; **126**: 103-112.

Stanek K, Gunstad J, Leahy T, Glickman E, Alexander T, Spitznagel MB et al. Serum brain-derived neurotrophic factor is associated with reduced appetite in healthy older adults. *J Health Aging* 2008; **12**: 183-185.

Stern AJ, Savostyanova AA, Goldman A, Barnett AS, van der Veen JWC, Caillicot JH et al. Impact of the BDNF val⁶⁶met polymorphism on levels of N-Acetyl-Aspartate assessed by magnetic resonance spectroscopic imaging at 3 Tesla. *Biol Psychiatry* 2008; **64**: 856-862.

Sterne JAC, Smith GD. Sifting the evidence – what’s wrong with significance tests? *BMJ* 2001; **322**: 226-231.

Sterne JAC, Egger M, Smith GD. Investigating and dealing with publication and other biases in meta-analysis. *BMJ* 2001; **323**: 101-105.

Strohle A, Stoy M, Graetz B, Scheel M, Wittman A, Galinat J et al. Acute exercise ameliorates reduced BDNF in patients with panic disorder. *Psychoneuroendocrinol* 2010; **35**: 364–368.

Su SC, Sun MT, Wen MJ, Lin CJ, Chen YC, Hung YJ. BDNF, adiponectin, and proinflammatory markers in various subtypes of depression in young men. *Int J Psychiatry Med* 2011; **42**: 211-226.

Sutton MA, Schuman EM. Dendritic protein synthesis, synaptic plasticity, and memory. *Cell* 2006; **127**: 49–58.

Suwa M, Kishimoto H, Nofuji Y, Sasaki H, Radak Z, Kumagai S. Serum BDNF level is increased and associated with obesity in newly diagnosed female patients with type 2 diabetes mellitus. *Metabolism* 2006; **55**: 852–857.

Szeszko PR, Lipsky R, Mentschel C, Robinson D, Gunduz-Bruce H, Sevy S et al. BDNF val⁶⁶met polymorphism and volume of hippocampal formation. *Mol Psychiatry* 2005; **10**: 631-636.

T

Taliaz D, Stall N, Dar DE, Zangen A. Knockdown of brain-derived neurotrophic factor in specific brain sites precipitates behaviors associated with depression and reduces neurogenesis. *Mol Psychiatry* 2010; **15**: 80-92.

Taliaz D, Nagaraj V, Haramati S, Chen A, Zangen A. Altered BDNF expression in the ventral tegmental area but not in the hippocampus, is essential to antidepressant-like effects of electroconvulsive therapy. *Biol Psychiatry* 2012 AOP: 10.1016/j.biopsych.2012.07.025

Tang SW, Chu E, Hui T, Helmeeste D, Law C. Influence of exercise on serum BDNF concentrations in healthy human subjects. *Neurosci Lett* 2008; **431**: 62-65.

Takahashi T, Suzuki M, Tsunoda M, Kawamura Y, Takahashi N, Tsuneki H et al. Association between the BDNF val⁶⁶met polymorphism and brain morphology in a Japanese sample of schizophrenia and healthy comparisons. *Neurosci Lett* 2008; **435**: 34-39.

Tapia-Arancibia L, Aliaga E, Sihol M, Arancibia S. New insights into BDNF function in normal ageing and Alzheimer disease. *Brain Res Rev* 2008; **59**: 201-220.

Teixera AL, Barbosa IG, Diniz BS, Kummer A. Circulating levels ofBDNF: correlation with mood, cognition and motor function. *Biomark Med* 2010; **4**: 871-887.

Terracciano A, Lobina M, Piras MG, Cannas A, Meirelles O et al. Lower serum BDNF associated with higher neuroticism and depressive symptoms. *Psychosom Med* 2011; **74**: 638-642.

Terracciano A, Pinas MG, Lobina M, Mulas A, Meirelles O, Sutin AR et al. Genetics of serum BDNF: meta-analysis of the BDNF val⁶⁶met and genome-wide association study. *W J Biol Psychiatry* 2011; AOP: 10.3109.15622975.2011.616533.

The Cochrane Collaboration Handbook. www.cochrane.org (accessed 29 June 2013).

Thomaes K, Dorrepaal E, Draijer NPJ, de Ruiter MB, Elzinga BM, van Balkom AJ et al. Increased activation of the left hippocampus region in complex PTSD during encoding and recognition of emotional words: a pilot study. *Psychiatric Res: Neuroimaging* 2009; **171**: 44-53.

Thompson Ray M, Weickert CS, Wyatt E, Webster MJ. Decreased BDNF, trkB-TK+ and GAD mRNA expression in the hippocampus of individuals with schizophrenia and mood disorders. *J Psychiatry Neurosci* 2011; **36**: 195-203.

Toro R, Chupin M, Garnero L, Leonard G, Perron M, Pike B et al. Brain volumes and val⁶⁶met polymorphism of the BDNF gene: local or global effects? *Brain Struct Funct* 2009; **213**: 501-509.

Toups MSP, Greer TL, Kurian BT, Grannemann BD, Carmody TJ, Huebinger R et al. Effects of serum brain-derived neurotrophic factor on exercise augmentation treatment of depression. *J Psychiatric Res* 2011; **45**: 1301-1306.

Trajkovska V, Marcussen AB, Vinberg M, Hartvig P, Aznar S, Knudsen GM. Measurements of brain-derived neurotrophic factor: methodological aspects and demographical data. *Brain Res Bull* 2007; **73**: 143-149.

Trajkovska V, Vinberg M, Aznar S, Knudsen GM, Kessing LV. Whole blood BDNF concentrations in healthy twins discordant for affective disorder: association to life events and neuroticism. *J Affect Disord* 2008; **108**: 165-169.

Tripp A, Oh H, Guilloux J-P, Martinowich K, Lewis D, Sibille E. Brain-derived neurotrophic factor signaling and subgenual anterior cingulate cortex dysfunction in major depressive disorder. *Am J Psychiatry* 2012; **169**: 1194-1202.

Turner EH, Matthews AM, Linardatos E, Tell RA, Rosenthal R. Selective publication of antidepressant trials and its influence on apparent efficacy. *N Eng J Med* 2008; **358**: 252-260.

U

Ullrich A, Schlessinger J. Signal transduction by receptors with tyrosine kinase activity. *Cell* 1990; **61**: 203-212.

Umene-Nakano W, Yoshimura R, Ikenouchi-Sugita A, Hori H, Hayashi K, Ueda N et al. Serum levels of BDNF in comorbidity of depression and alcohol dependence. *Hum Psychopharmacol* 2009a; **24**: 409-413.

V

van Harmelen A-L, van Tol M-J, van der Wee NJA, Veltman DJ, Aleman A, Spinhoven et al. Reduced medial prefrontal cortex volume in adults reporting childhood emotional maltreatment. *Biol Psychiatry* 2010; **68**: 832-838.

van Tol M-J, van der Wee NJ, van den Heuvel OA, Demenescu LR, Nielen MM, Renken R et al. Regional brain volume in depression and anxiety disorders. *Arch Gen Psychiatry* 2010; **67**: 1002-1011.

van Tol M-J, Demenescu LR, van der Wee NJA, Korteas R, Nielen MMA, Den Boer JA et al. Emotional word encoding and recognition in depression and anxiety disorders. *Biol Psychiatry* 2012; **71**: 593-602.

Verhagen M, van der Meij A, van Deurzen PAM, Janzing JGE, Arias-Vasquez A et al. Meta-analysis of the BDNF val⁶⁶met polymorphism in major depressive disorder: effects of gender and ethnicity. *Mol Psychiatry* 2010; **15**: 260-271.

Videbech P, Ravnkilde B. Hippocampal volume and depression: a meta-analysis of MRI studies. *Am J Psychiatry* 2004; **161**: 1957-1966.

Vinberg M, Trajkovska V, Bennike B, Knorr U, Knudsen GM, Kessing LV. The BDNF val⁶⁶met polymorphism: relation to familial risk of affective disorder, BDNF concentrations and salivary cortisol. *PNEC* 2009; **34**: 1380-1389.

Vinberg M, Miskowiak K, Kessing LV. Brain-derived neurotrophic factor levels as possible predictor of psychopathology in healthy twins at high and low risk for affective disorder. *Psychoneuroendocrinol AOP*; dx.doi.org/10.1016/j.psyneuen.2013.09.007.

Voineskos AN, Lerch JP, Felsky D, Shaikh S, Rajji TK, Miranda D et al. The brain-derived neurotrophic factor val⁶⁶met polymorphism and prediction of neural risk for Alzheimer disease. *Arch Gen Psychiatry* 2011; **68**: 198-206.

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; for the STROBE initiative. The strengthening the Reporting of Observational Studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007; **370**: 1443-1457.

Voorhuis M, Broekmans FJ, Fauser BCJM, Onland-Moret NC, van der Schouw YT. Genes involved in initial follicle recruitment may be associated with age at menopause. *J Clin Endocrinol Metab* 2011; **96**: e473-e479.

W

Wacker HR, Battegay R, Mulleijans R, Schlosser C (2006). Using the CIDI-C in the general population. In: Stefanis CN, Rabavilas AD, Soldatos CR editors. *Psychiatry: a world perspective*. Amsterdam, The Netherlands: Elsevier Science Publishers.

Walf AA, Frye CA. A review and update of mechanisms of estrogen in the hippocampus and amygdala for anxiety and depression behavior. *Neuropsychopharmacol* 2006; **31**: 1097-1111.

Walton JC, Weil ZM, Nelson RJ. Influence of photoperiod on hormones, behavior, and immune function. *Front Neuroendocrinol* 2011; **32**: 303-319.

Wehr TA, Rosenthal NE. Seasonality and affective illness. *Am J Psychiatry* 1989; **146**: 829-839.

Wehr TA, Duncan WC, Sher L, Aeschbach D, Schwartz PJ, Turner EH et al. A circadian signal of change of season in patients with seasonal affective disorders. *Arch Gen Psychiatry* 2001; **58**: 1108-1114.

Wells GA, Shea B, O'Connell D. The Newcastle-Ottawa Scale (NOS) for assessing the quality non-randomised studies in meta-analyses. Ottawa Healthcare Institute. www.ohri.ca (accessed 12 April 2012 and 7 February 2013).

Werner NS, Meindl T, Engel RR, Rosner R, Riedel M, Reiser M et al. Hippocampal function during associative learning in patients with posttraumatic stress disorder. *J Psychiatric Res* 2009; **43**: 309-318.

Wichers M, Kenis G, Jacobs N, Mengelers R, Derom C, Vlietinck R et al. The BDNF val⁶⁶met x 5-HTTLPR and child adversity interaction and depressive symptoms: an attempt at replication. *Am J Med Genet Part B* 2008; **147B**: 120-123.

Wichers M, Kenis G, Jacobs N, Myin-Germeyns I, Schruers K, Mengelers R et al. The psychology of psychiatric genetics: evidence that positive emotions in females moderate genetic sensitivity to social stress associated with the BDNF val⁶⁶met polymorphism. *J Abn Psychology* 2008; **117**: 699-704.

Wiersma JE, Hovens JGFM, van Oppen P, Giltay EJ, van Schaik JF, Beekman ATF et al. The importance of childhood trauma and childhood life events for chronicity of depression in adults. *J Clin Psychiatry* 2009; **70**: 983-989.

Winthorst WH, Post WJ, Meesters Y, Penninx BWJH, Nolen WA. Seasonality in depressive and anxiety symptoms among primary care patients and in patients with depressive and anxiety disorders; results from the Netherlands study of depression and anxiety. *BMC Psychiatry* 2011; **11**: 198-216.

Wittchen HU. Reliability and validity studies of the WHO-Composite International Diagnostic Interview (CIDI): a critical review. *J Psychiatr Res* 1994; **28**: 57-84.

Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier D. Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The multicenter WHO/ADAMHA field trials. *Br J Psychiatry* 1991; **159**: 645-653.

Wojnar M, Brower KJ, Strobbe S, Ilgen M, Matsumoto H, Nowosad I et al. Association between val⁶⁶met BDNF gene polymorphism and post-treatment relapse in alcohol dependence. *Alcohol Clin Exp Res* 2009; **33**: 693-702.

Wolkowitz OM, Wolf J, Shelly W, Rosser R, Burke HM, Lerner GK et al. Serum BDNF levels before treatment predict SSRI response in depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; **15**: 1623-1630.

World Health Organization (WHO). Global Burden of Disease Report: 2004 update. 2008; www.who.int (accessed March 2010, April 2010, and February 2013).

World Health Organization (WHO). Collaborating Centre for Drug Statistics Methodology. 2008; www.whocc.no (accessed March 2010, April 2010, March 2012, and January 2011).

World Meteorological Organization (WMO). Guide to meteorological instruments and methods of observation. WMO: Geneva, 1996.

Workman JL, Bower SL, Nelson RJ. Enrichment and photoperiod interact to affect spatial learning and hippocampal dendritic morphology in white-footed mice (*Peromyscus leucopus*). *Eur J Neurosci* 2009; **29**: 161-170.

X

Y

Yasutake C, Kuroda K, Yanagawa T, Okamura T, Yoneda H. Serum BDNF, TNF-alpha and IL-1beta levels in dementia patients: comparison between Alzheimer's disease and vascular dementia. *Eur. Arch. Psychiatry Clin Neurosci* 2006; **256**: 402-406.

Young E, Korszun A. Sexual trauma, stress hormones and depression. *Mol Psychiatry* 2010; **15**: 23-28.

Yoshida T, Ishikawa M, Masaomi I, Hashimoto K. Serum levels of mature BDNF and its precursor in healthy subjects. *Open Clin Chem* 2012; **5**: 7-12.

Yoshida T, Ishikawa M, Niitsu T, Nakazato M, Watanabe H, Shiraishi T et al. Decreased serum levels of mature BDNF, but not its precursor, in patients with major depressive disorder. *PLoS ONE* 2012; **7**: e42676.

Yoshimura R, Mitoma M, Sugita A, Hori H, Okamoto T, Umene W et al. Effects of paroxetine or milnacipran on serum BDNF in depressed patients. *Prog Neuropsychopharmacol Biol Psychiatry* 2007; **31**: 1034-1037.

Yoshimura R, Sugita-Ikenouchi A, Hori H, Umene-Nakano W, Hayasi K, Katsuki A et al. A close correlation between plasma and serum levels of BDNF in healthy volunteers. *Int J Psychiatry Clin Prac* 2010; **14**: 220-222.

Yoshimura R, Kishi T, Suzuki A, Umene-Nakano W, Ikenouchi-Sugita A, Hori H et al. The BDNF polymorphism val⁶⁶met is associated with neither serum BDNF level nor response to selective serotonin reuptake inhibitors in depressed Japanese patients. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; **35**: 1022-1025.

Z

Zanardini R, Gazzoli A, Ventriglia M, Perez J, Bignotti S, Rossini PM et al. Effect of repetitive transcranial magnetic stimulation on serum BDNF in drug resistant depressed patients. *J Affect Disord* 2006; **91**: 83-86.

Zhao C, Deng W, Gage FH. Mechanisms and functional implications of adult neurogenesis. *Cell* 2008; **132**: 645-660.

Zhou Z, Lu T, Xu G, Yue X, Zhu W, Ma M et al. Decreased serum BDNF is associated with post-stroke depression but not with BDNF gene val⁶⁶met polymorphism. *Clin Chem Lab Med* 2011; **49**: 185-189.

Zhou L, Xiong J, Lim Y, Ruan Y, Huang C, Zhu Y et al. Upregulation of blood proBDNF and its receptors in major depression. *J Affect Disord*. AOP, doi: 10.1016/j.jad.2013.03.002.

Ziegenhorn AA, Schulte-Hebruggen O, Danker-Hopfe H, Malbranc M, Hartung H-D, Anders D. Serum neurotrophins - a study on the time course and influencing factors in a large old sample. *Neurobiol Aging* 2007; **28**: 1436-1445.