



Universiteit
Leiden
The Netherlands

Recent innovations in minimally invasive anterior and posterior lamellar keratoplasty

Parker, J.

Citation

Parker, J. (2017, July 4). *Recent innovations in minimally invasive anterior and posterior lamellar keratoplasty*. Retrieved from <https://hdl.handle.net/1887/50484>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/50484>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/50484> holds various files of this Leiden University dissertation

Author: Parker, Jack

Title: Recent innovations in minimally invasive anterior and posterior lamellar keratoplasty

Issue Date: 2017-07-04

Chapter 6

Descemet Membrane Endothelial Keratoplasty (DMEK): A Review

Jack Parker¹⁻³, John S. Parker³, and Gerrit R.J. Melles^{1,2,4}

¹Netherlands Institute for Innovative Ocular Surgery, Rotterdam, The Netherlands; ²Melles Cornea Clinic Rotterdam, The Netherlands; ³UAB Callahan Eye Hospital, Birmingham, Alabama; ⁴Amnitrans EyeBank Rotterdam, The Netherlands.

ABSTRACT

Descemet Membrane Endothelial Keratoplasty (DMEK) is the most recent step forward in the evolution of endothelial keratoplasty toward thinner grafts and more natural, anatomic corneal restoration. Offering unprecedented visual results and requiring no special or expensive equipment, DMEK has the potential to become the first line treatment for corneal endothelial disorders. The surgery's perceived shortcomings (primarily technical difficulty) have mostly been addressed by new "no-touch" procedures for both graft preparation and graft unfolding in the recipient eye. And as a result, DMEK has been gaining traction with ophthalmologists the world over. Now, in its most recent formulation, DMEK is ready for the typical corneal surgeon, in any clinical setting, and at low cost.

KEYWORDS: Descemet membrane endothelial keratoplasty, posterior lamellar keratoplasty, corneal transplantation, endothelium, surgical technique

INTRODUCTION

For almost 100 years, Penetrating Keratoplasty (PK) was the mainstay of therapy for patients with corneal endothelial disorders.¹ That changed in 1998 with the introduction of Posterior Lamellar Keratoplasty (PLK),²⁻⁴ later popularized in the United States as Deep Lamellar Endothelial Keratoplasty (DLEK).⁵⁻⁷ Selectivity was the new technique's primary advantage. By replacing only the inner aspect of the cornea, many of the suture, astigmatism, and wound healing problems of PK disappeared. But while effective, DLEK ultimately proved too technically challenging for widespread adoption. So, the surgery was simplified, giving rise to Descemet Stripping (Automated) Endothelial Keratoplasty (DS(A)EK).⁸⁻¹¹ And within five years, this modified technique became the global treatment of choice for corneal endothelial disorders. Still, few patients after DS(A)EK achieved best corrected visual acuities (BCVAs) exceeding 20/25. Probably, the graft's layer of attached stroma was to blame, which thickened the cornea and seemed to undermine its optical performance.¹²⁻¹⁶

A stroma-less graft was the solution, arriving in 2006 in the form of Descemet Membrane Endothelial Keratoplasty (DMEK).¹⁷⁻¹⁹ With a transplant composed solely of isolated Descemet membrane (and its endothelium), DMEK slashed graft thickness by 75% compared to DS(A)EK, from 80 microns down to 20. The results were dramatic: almost 80% of patients reached $\geq 20/25$ within 6 months after surgery.^{12,20,21}

Recently, DMEK has been refined into a standardized "no-touch" procedure, ready for the typical corneal surgeon in any clinical setting and at low cost.²² Compared to its predecessors (DSEK, DLEK, and their variations), DMEK provides better and faster visual recovery, usually with no additional complications. It is therefore poised to become the first-line option for corneal endothelial disorders worldwide.²³

PREOPERATIVE PREPARATION OF THE DMEK GRAFT

Ideally, DMEK grafts are prepared in an eye bank, 1-2 weeks before surgery. There, the tissue undergoes several rounds of additional screening. Principally, this consists of evaluating the cell density and morphology of the donor endothelium. Grafts which appear abnormal under the microscope – those with scarce or atypical cells, suspicious for being dysfunctional – are discarded, raising the quality of the pool of tissue for transplant. Preparing the grafts weeks in advance also adds convenience: it saves time and safeguards against unexpected tissue shortage on the day of surgery.²⁴

On the other hand, some ophthalmologists may prefer to create the grafts themselves, in the operating room, just before surgery.²⁵ This is especially true in the United States, where few eye banks currently supply ready-to-use DMEK tissue. Each graft takes 30

minutes to prepare, and all the steps are the same, whether in the operating room or the eye bank.

The initially described DMEK graft harvesting technique consisted of stripping Descemet membrane from a corneo-scleral rim submerged in saline. This method was proven safe and reproducible, with <5% tissue loss due to inadvertent tearing, and – surprisingly – no significant endothelial cell damage.²⁴⁻²⁸ Recently, the process was upgraded to a “no-touch” procedure, making the preparation both safer and easier.²⁹ As a bonus, the anterior portion of the corneas left over from creating the DMEK grafts (with the Descemet membrane stripped off, but otherwise intact) can be used for Deep Anterior Lamellar Keratoplasty (DALK). This added benefit applies only to DMEK, because DS(A)EK preparation – by incorporating some of the posterior stroma into the graft – mangles the corneal remains, leaving them less suitable for transplant.²⁹⁻³¹

DMEK SURGICAL TECHNIQUE

The standardized no-touch technique for DMEK was published by Dapena et al in 2011.²² In brief, a 3.0mm clear-cornea tunnel incision is made at the 12 o'clock position with a slit knife, followed by the creation of three side-ports using a surgical knife at 10:30, 1:30, and 7:30 (right eye) or 4:30 (left eye). Under air, the recipient's Descemet membrane is first scored 360 degrees then stripped from the posterior stroma using a reversed Sinsky hook (Catalogue no 50.1971B, D.O.R.C. International, Zuidland, The Netherlands). The DMEK graft is thoroughly rinsed with balanced salt solution (BSS, Alcon Nederland BV, Gorinchem, The Netherlands) and stained twice with trypan blue 0.06% (Catalogue no VBL.105.USA, Vision blue™; D.O.R.C. International) to enhance its visibility in the recipient anterior chamber. Already curled into a roll due to the inherent elastic properties of the membrane itself, the graft may be nudged into a “double roll” configuration by applying a flow of BSS directly across its surface.²²

After staining, the DMEK double-roll is sucked into a custom-made glass pipette (D.O.R.C. International), then injected into the recipient anterior chamber through the 12 o'clock incision “hinge down” so that the double roll faces upward. Once the graft has been inserted, its orientation can be checked (and verified as properly “hinge down”) through the use of the Moutsouris sign, whereby the tip of a 30G cannula, positioned atop the edge of the graft, will turn blue if it is embraced by an upward facing roll. If the tip does not turn blue, then the roll must be facing down, and therefore the graft is upside down, which can be corrected by gently flushing it within the anterior chamber (**Figure 1**).²²

With the graft properly oriented, it may be unfolded by injecting a small air bubble in between the double rolls, then stroking the surface of the cornea to move the bubble

and spread out the graft (Dapena technique). Once it has been fully unfolded, the graft is fixed against the recipient posterior stroma by completely filling the anterior chamber with air for a period of one hour. Afterwards, the air fill is reduced to 30-50%, and the patient is instructed to remain supine for 24 hours.²²




Figure 1. Artist rendering of the *Moutsouris sign*. (A and B) When the DMEK-graft is oriented correctly within the anterior chamber (double roll upward), the tip of the cannula can be positioned 'inside' a peripheral curl, so that the tip appears blue (arrows) because of the overlying blue tinted donor tissue (Moutsouris sign positive). (C and D) When the graft is positioned 'upside-down' (double roll downward), the tip of the cannula does not 'find' the curls, so the tip will not change in color (Moutsouris sign negative). [This figure has been published previously in Dapena et al. Arch Ophthalmol. 2011;129(1):88-94]

Variations on DMEK surgery do exist, however, with DMAEK and DMEK-S being the most prominent examples.³²⁻³⁵ These differ from regular DMEK in that a stromal rim is left attached to the periphery of the graft during preparation, which allows grasping and a "drag-and-drop" insertion method. Otherwise, the surgery is the same.

RESULTS

Visual Acuity

After DMEK, 77% of eyes may achieve a BCVA $\geq 20/25$ at 6 months, with 50% $\geq 20/20$. Visual rehabilitation is frequently fast, not uncommonly rebounding to 20/20 within the first post-operative week, and with most patients reaching their final BCVA within 1-3 months.^{19,22,23,26} (DMAEK and DMEK-S, likewise, seem to offer similarly good results.^{35,36})

No other form of corneal transplantation offers comparable outcomes. After PK, less than 50% of patients achieve visions of $\geq 20/40$, and then only at 1 year.³⁷ Following DS(A)EK, the average vision at 6 months is 20/40, rarely reaching 20/25 or better.¹²⁻¹⁶ Tellingly, in those patients with poor vision after DSEK, many dramatically improve with a re-operation to replace their DSEK graft with a DMEK (**Figure 2**).³⁸ Moreover, in people with one eye operated with each technique – one eye DSEK, one eye DMEK – overwhelmingly, they prefer the vision in their DMEK eye.³⁹




Figure 2. (A) Slit-lamp photograph 1 year after DSEK. Despite complete corneal clearance and minimal interface opacity, the patient's BCVA never improved beyond 20/100. Image (B) shows the same eye following a secondary DMEK for reasons of low visual acuity. After DMEK, vision improved to 20/25 at 1 month post-operatively.

Refractive Change and Stability

After DMEK, both the spherical equivalent (SE) and cylindrical error are frequently within 1.0D of the pre-operative refractive error. Pachymetric and refractive data show that the transplanted cornea stabilizes 3 months after surgery, at which point new glasses may be prescribed. Until then, most patients are able to wear their current prescription.²³

Endothelial Cell Density

Most DMEK grafts show a $\pm 30\%$ reduction in cell density 6 months after surgery. Thereafter, cell density falls at a steady, predictable rate – at about 10% per year.^{26,40-42} Interestingly, the transition to an entirely no-touch technique has had no effect on the measured “cell-loss” after DMEK.²² The strong implication is that mechanical damage during transplantation cannot be the cause. More likely, the rapid fall in cell density after surgery reflects a decline in cellular concentration – not number – as the endothelial cells migrate out from the graft onto peripheral parts of the patient’s posterior stroma.

Cell density measurements after DS(A)EK are almost identical, with a sharp $\pm 30\%$ drop-off in the first 6 months, followed by a regular decline of nearly 10% per year.⁴³⁻⁴⁵

A much larger decline is evident after PK, however, in which grafts commonly lose upwards of 40-55% within the first post-operative year. In addition, the rate of decline never appears to stabilize at a lower level, as with DS(A)EK and DMEK.⁴⁶⁻⁴⁸

COMPLICATIONS

Graft Detachment

Graft detachment is the most common complication following all forms of endothelial keratoplasty. With DS(A)EK, this may occur in 0-82% of surgeries.^{11,49-51} Similarly, detachment rates of 20-60% have been reported after DMEK, although many of these cases do not appear to be clinically significant.^{22,35,52-54} Frequently, DMEK detachments are small, peripheral, and temporary. And even when the detached areas are both large and central, some patients nevertheless achieve BCVAs $\geq 20/40$. In our own series, clinically significant detachments – those which reduced the patient’s vision and/or required re-intervention – occurred in 10% of eyes. Risk factors might include surgical inexperience, failing to completely unfold the donor membrane during surgery, implanting the graft upside down, the use of intra-ocular viscoelastics, use of plastic materials (rather than glass) to inject the tissue into the recipient anterior chamber, insufficient air-bubble support after surgery, and the use of Optisol rather than organ culture medium for graft storage pre-operatively.⁵²⁻⁵⁵

Management depends on the size of the detachment. Small detachments (less than one-third of the graft area) resolve spontaneously and rarely, if ever, require re-

intervention. Larger detachments, however, have more variable outcomes, complicating the management decision tree. In general, even with large detachments (greater than one-third of the graft area), most corneas eventually clear, although over a longer time period and then only 50% of patients achieve vision $\geq 20/40$. Because a satisfactory visual result may occur half the time after a large detachment without any subsequent intervention, reoperation – either with re-grafting or re-bubbling – ought to be an individualized decision, tailored to the patient’s preferences (i.e. for more surgery, in light of the possibility of better vision).⁵²⁻⁵⁵

Allograft rejection

Two years after DMEK, the allograft rejection rate is $\leq 1\%$. This is considerably lower than the reported rate after PK (5-15% in “low-risk” cases), and also lower than after DS(A)EK (10%).^{23,37,56-58} Likely, the explanation lies in DMEK’s thinner, stroma-less graft, which may be less immunogenic because it presents fewer antigens to the recipient’s immune system.^{23,57}

Secondary glaucoma

Because runaway pressures threaten both the survival of the graft and the health of the optic nerve, glaucoma is among the most important potential complications of any form of corneal transplantation. Reported rates after PK and DS(A)EK commonly range from 15-35%, but sometimes as high as 60% depending on the patient population and the steroid regimen.⁵⁹⁻⁶² Because the risk of allograft rejection after DMEK is relatively low, a lighter, less intense, steroid schedule is possible. (Specifically, we use 0.1% topical dexamethasone for just the first postoperative month, then switch to fluoromethalone thereafter.) Perhaps as a consequence, the reported rate of glaucoma is small – just 6.5% at 2 years. Most cases arise in eyes with a pre-existing history of pressure trouble, with relatively few “new” cases after surgery.⁶³

Two additional factors may contribute to DMEK’s low rate of secondary glaucoma. First, most patients receiving a DMEK for Fuchs Dystrophy are Caucasian, a population thought to be at lower risk. Second, one week prior to surgery, a peripheral iridotomy is made at the 12 o’clock position to prevent the development of a pupillary block glaucoma.⁶³

DMEK IN PHAKIC EYES

DMEK is safe for phakic eyes, although several additional protective steps are required. Just prior to transplant, the pupil should be constricted with 2% pilocarpine to protect the lens from accidental damage during surgery, either from air-bubble or instrument

induced trauma. Even so, 25% of phakic eyes may present with mild anterior subcapsular lens opacities or a Vossius ring (iris pigment imprint on the outer lens capsule). Usually, these pigment deposits disappear with time and do not affect final visual acuity. The rate of iatrogenic cataract formation necessitating phacoemulsification is reported at 4% at 2 years.^{64,65}

As a precaution, the size of the air bubble left behind in the anterior chamber after DMEK surgery ought to be reduced in phakic eyes, from 50% down to 30%. This may help prevent a mechanical angle closure glaucoma from developing (arising when a large air bubble presses against the lens, causing the lens to tilt forward and compress the angle).⁶⁵

FUTURE DIRECTIONS

Steadily, reports have been accumulating of corneas with detached grafts (after both DMEK and DS(A)EK) that nevertheless clear.^{66,67} When these corneas are viewed with specular and confocal microscopy, endothelial cells are clearly visible populating the recipient's posterior stroma (**Figure 3**). The prevailing speculation is that endothelial migration is responsible for this phenomenon, either by the donor cells, or host cells, or both.⁶⁸⁻⁷⁰ If widespread cell migration does indeed occur, then a simplified procedure, tentatively named "free-DMEK" or "Descemet Membrane Endothelial Transfer" (DMET) – in which the donor tissue is merely injected into the recipient anterior chamber after descemetorhexis – could be effective in the management of corneal endothelial disease.⁷¹ The advantages of this surgery, even over DMEK, would be enormous: perfect anatomical restoration, complete visual recovery, elimination of virtually all intra- and post-operative complications associated with endothelial keratoplasty, and an enormous reduction in the required surgical skills. Pending further study, DMET has the potential to become the preferred "no-keratoplasty" treatment for corneal endothelial disorders.




Figure 3. Slit-lamp photographs after DMEK (A, B) showing a clear cornea (yellow arrows) above a large centrally detached graft (green arrows). OCT demonstrates normal corneal thickness above the detachment (C), and confocal (D) and specular microscopy (E) reveal the presence of endothelial cells populating the recipient's posterior stroma in the detached area.

REFERENCES

1. Terry MA, "Endothelial Keratoplasty: Why aren't we all doing Descemet membrane endothelial keratoplasty?" *Cornea*, 2012 Feb 23. [Epub ahead of print].
2. Melles GRJ, Eggink FAGJ, Lander F, et al, A surgical technique for posterior lamellar keratoplasty, *Cornea*, 1998;17:618-26.
3. Melles GR, Lander F, Beekhuis WH, et al, Posterior lamellar keratoplasty for a case of pseudophakic bullous keratopathy, *Am J Ophthalmol*, 1999;127:340-1.
4. Melles GR, Lander F, Nieuwendaal C, Sutureless, posterior lamellar keratoplasty: A case report of a modified technique, *Cornea*, 2002;21:325-7.
5. Terry MA, Ousley PJ, Deep lamellar endothelial keratoplasty in the first United States patients: early clinical results, *Cornea*, 2001;20:239-43.
6. Terry MA, Ousley PJ, Deep lamellar endothelial keratoplasty visual acuity, astigmatism, and endothelial survival in a large prospective series, *Ophthalmology*, 2005;112:1541-8.
7. Ousley PJ, Terry MA, Stability of vision, topography, and endothelial cell density from 1 year to 2 years after deep lamellar endothelial keratoplasty surgery, *Ophthalmology*, 2005;112:50-7.
8. Melles GR, Wijdh RH, Nieuwendaal CP, A technique to excise the Descemet membrane from a recipient cornea (descemetorhexis), *Cornea*, 2004;23:286-8.
9. Price FW Jr, Price MO, Descemet's stripping with endothelial keratoplasty in 50 eyes: a refractive neutral corneal transplant, *J Refract Surg*, 2005;21:339-45.
10. Azar DT, Jain S, Sambursky R, Strauss L, Microkeratome-assisted posterior keratoplasty, *J Cataract Refract Surg*, 2001;27:353-6.
11. Gorovoy MS, Descemet-stripping automated endothelial keratoplasty, *Cornea*, 2006;25:886-9.
12. Dapena I, Ham L, Melles GRJ, Endothelial Keratoplasty: DSEK/DSAEK or DMEK – the thinner the better? *Curr Opin Ophthalmol*, 2009;20:299-307.
13. Bahar I, Kaiserman I, McAllum P, et al, Comparison of posterior lamellar keratoplasty techniques to penetrating keratoplasty, *Ophthalmology*, 2008;115:1525-33.
14. Chen ES, Terry MA, Shamie N, et al, Descemet-stripping automated endothelial keratoplasty: six month results in a prospective study of 100 eyes, *Cornea*, 2008;27:514-20.
15. Mearza AA, Qureshi MA, Rostron CK, Experience and 12-month results of Descemet-stripping endothelial keratoplasty (DSEK) with a small-incision technique, *Cornea*, 2007;26:279-83.
16. Terry MA, Shamie N, Chen ES, et al, Precut tissue for Descemet's stripping automated endothelial keratoplasty: Vision, astigmatism, and endothelial survival, *Ophthalmology*, 2009;116:248-56.
17. Melles GR, Lander F, Rietveld FJ, Transplantation of Descemet's membrane carrying viable endothelium through a small scleral incision, *Cornea*, 2002;21:415-8.
18. Melles GR, Ong TS, Ververs B, van der Wees J, Descemet membrane endothelial keratoplasty (DMEK), *Cornea*, 2006;25:987-90.
19. Melles GR, Ong TS, Ververs B, van der Wees J, Preliminary clinical results of Descemet membrane endothelial keratoplasty, *Am J Ophthalmol*, 2008;145:222-7.
20. Melles GR, Posterior lamellar keratoplasty: DLEK to DSEK to DMEK, *Cornea*, 2006;25:879-81.
21. Fernandez MM, Afshari NA, Endothelial Keratoplasty: From DLEK to DMEK, *Middle East Afr J Ophthalmol*, 2010;17:5-8.
22. Dapena I, Moutsouris K, Droutsas K, et al, Standardized 'no touch' technique for Descemet membrane endothelial keratoplasty (DMEK), *Arch Ophthalmol*, 2011;129:88-94.

23. Dirisamer M, Ham L, Dapena I, et al, Efficacy of Descemet membrane endothelial keratoplasty (DMEK): Clinical outcome of 200 consecutive cases after a 'learning curve' of 25 cases, *Arch Ophthalmol*, 2011;129:1435-43.
24. Lie JT, Birbal R, Ham L et al, Donor tissue preparation for Descemet membrane endothelial keratoplasty, *J Cataract Refract Surg*, 2008;34:1578-83.
25. Kruse FE, Laaser K, Cursiefen C, et al, A stepwise approach to donor preparation and insertion increases safety and outcome of Descemet membrane endothelial keratoplasty, *Cornea*, 2011;30:580-7.
26. Price MO, Giebel AW, Fairchild KM, Price FW Jr, Descemet's membrane endothelial keratoplasty: prospective multicenter study of visual and refractive outcomes and endothelial survival, *Ophthalmology*, 2009; 116:2361-8.
27. Zhu Z, Rife L, Yiu S, et al, Technique for preparation of the corneal endothelium-Descemet membrane complex for transplantation, *Cornea*, 2006;25:705-8.
28. Ignacio TS, Nguyen TT, Sarayba MA, et al, A technique to harvest Descemet's membrane with viable endothelial cells for selective transplantation, *Am J Ophthalmol*, 2005;139:325-30.
29. Groeneveld-van Beek E, Lie J, van der Wees J, et al., Standardized 'no-touch' donor tissue preparation for DALK and DMEK: Harvesting undamaged anterior and posterior transplants from the same donor cornea, *Acta Ophthalmologica* 2012; In press.
30. Heindl LM, Riss S, Laaser K, et al, Split cornea transplantation for 2 recipients - review of the first 100 consecutive patients, *Am J Ophthalmol*, 2011;152:523-32.
31. Lie JT, Groeneveld-van Beek EA, Ham L, et al, More efficient use of donor corneal tissue with Descemet membrane endothelial keratoplasty (DMEK): two lamellar keratoplasty procedures with one donor cornea, *Br J Ophthalmol*, 2010;94:1265-6.
32. Kymionis GD, Yoo SH, Diakonis VF, et al, Automated donor tissue preparation for Descemet membrane automated endothelial keratoplasty (DMAEK): An Experimental Study, *Ophthalmic Surg Lasers Imaging*, 2011;42:158-61.
33. McCauley MB, Price FW, Price MO, Descemet membrane automated endothelial keratoplasty: hybrid technique combining DSAEK stability with DMEK visual results, *J Cataract Refract Surg*, 2009;35:1659-64.
34. Studeny P, Farkas A, Vokrojova M, et al, Descemet membrane endothelial keratoplasty with a stromal rim (DMEK-S), *Br J Ophthalmol*, 2010;94:909-14.
35. Guerra FP, Anshu A, Price MO, et al, Descemet's membrane endothelial keratoplasty prospective study of 1-Year visual outcomes, graft survival, and endothelial cell loss, *Ophthalmology*, 2011; 118:2368-73.
36. McCauley MB, Price MO, Fairchild KM, et al, Prospective study of visual outcomes and endothelial survival with Descemet membrane automated endothelial keratoplasty, *Cornea*, 2011;30:315-9.
37. Williams KA, Muehlberg SM, Lewis RF, et al, How successful is corneal transplantation? A report from the Australian Corneal Graft Register, *Eye*, 1995;9:219-27.
38. Ham L, Dapena I, van der Wees J, Melles GR, Secondary DMEK for poor visual outcome after DSEK: donor posterior stroma may limit visual acuity in endothelial keratoplasty, *Cornea*, 2010;29:1278-83.
39. Guerra FP, Anshu A, Price MO, Price FW, Endothelial keratoplasty: fellow eyes comparison of Descemet stripping automated endothelial keratoplasty and Descemet membrane endothelial keratoplasty, *Cornea*, 2011;30:1382-6.
40. Dapena I, Dapena L, Dirisamer M, et al, Visual acuity and endothelial cell density following Descemet membrane endothelial keratoplasty (DMEK), *Arch Soc Esp Oftalmol*, 2011;86:395-401.

41. Ham L, Dapena I, van der Wees J, Melles GR, Endothelial cell density after Descemet membrane endothelial keratoplasty: 1- to 3-year follow-up, *Am J Ophthalmol*, 2010;149:1016-7.
42. Parker J, Dirisamer M, Naveiras M, et al, Endothelial cell density after Descemet membrane endothelial keratoplasty: 4-year follow-up, *Am J Ophthalmol*, 2011;151:1107.
43. Price MO, Fairchild KM, Price DA, et al, Descemet's stripping endothelial keratoplasty five-year graft survival and endothelial cell loss, *Ophthalmology*, 2011;118:725-9.
44. Price FW Jr, Price MO, Does endothelial cell survival differ between DSEK and standard PK?, *Ophthalmology*, 2009;116:367-8.
45. Price MO, Price FW Jr, Endothelial cell loss after Descemet stripping with endothelial keratoplasty influencing factors and 2-year trend, *Ophthalmology*, 2008;115:857-65.
46. Borderie VM, Boëlle PY, Touzeau O, et al, Predicted long-term outcome of corneal transplantation, *Ophthalmology*, 2009;116:2354-60.
47. Patel SV, David O, Hodge MS, Bourne WM, Corneal endothelium and postoperative outcomes 15 years after penetrating keratoplasty, *Am J Ophthalmol*, 2005;139:311-9.
48. Ing JJ, Ing HH, Nelson LR, et al, Ten-year postoperative results of penetrating keratoplasty, *Ophthalmology*, 1998;105:1855-65.
49. Busin M, Bhatt PR, Scorcia V, A modified technique for Descemet membrane stripping automated endothelial keratoplasty to minimize endothelial cell loss, *Arch Ophthalmol*, 2008;126:1133-7.
50. Koenig SB, Covert DJ, Early results of small-incision Descemet stripping and automated endothelial keratoplasty, *Ophthalmology*, 2007;114:221-6.
51. Terry MA, Shamie N, Chen ES, et al, Endothelial keratoplasty a simplified technique to minimize graft dislocation, iatrogenic graft failure, and pupillary block, *Ophthalmology*, 2008;115:1179-86.
52. Dapena I, Moutsouris K, Ham L, Melles GR, Graft detachment rate, *Ophthalmology*, 2010;117:847.
53. Dirisamer M, van Dijk K, Dapena I, et al, Prevention and management of graft detachment in Descemet membrane endothelial keratoplasty (DMEK), *Arch Ophthalmol*, 2012;130:280-91.
54. Dapena I, Ham L, Droutsas K, et al, Learning curve in Descemet's membrane endothelial keratoplasty: First series of 135 consecutive cases, *Ophthalmology*, 2011;118:2147-54.
55. Laaser K, Bachmann BO, Horn FK, et al, Donor tissue culture conditions and outcome after descemet membrane endothelial keratoplasty, *Am J Ophthalmol*, 2011;151:1007-18.
56. Anshu A, Price MO, Price FW Jr, Risk of corneal transplant rejection significantly reduced with Descemet's membrane endothelial keratoplasty, *Ophthalmology*, 2012;119:536-40.
57. Dapena I, Ham L, Netukova M, et al, Incidence of early allograft rejection following Descemet membrane endothelial keratoplasty (DMEK), *Cornea*, 2011;30:1341-45.
58. Li JY, Terry MA, Goshe J, et al, Graft rejection after Descemet's stripping automated endothelial keratoplasty: graft survival and endothelial cell loss, *Ophthalmology*, 2012;119:90-4.
59. Vajaranant TS, Price MO, Price FW, Visual acuity and intraocular pressure after Descemet's stripping endothelial keratoplasty in eyes with and without preexisting glaucoma, *Ophthalmology*, 2009;116:1644-50.
60. Erdurmus M, Cohen EJ, Yildiz EH, et al, Steroid-induced intraocular pressure elevation or glaucoma after penetrating keratoplasty in patients with keratoconus or Fuchs dystrophy, *Cornea*, 2009;28:759-64.
61. Greenlee EC, Kwon YH, Graft failure: III. glaucoma escalation after penetrating keratoplasty, *Int Ophthalmol*, 2008;28:191-207.
62. Allen MB, Lieu P, Mootha VV, et al, Risk factors for intraocular pressure elevation after Descemet stripping automated endothelial keratoplasty, *Eye Contact Lens*, 2010;36:223-7.

63. Naveiras M, Dirisamer, Parker J, et al, Causes of glaucoma after Descemet membrane endothelial keratoplasty (DMEK), *Am J Ophthalmol*, 2012, Epub
64. Dapena I, Ham L, Tabak S, et al, Phacoemulsification after Descemet membrane endothelial keratoplasty, *J Cataract Refract Surg*, 2009;35:1314-5.
65. Parker J, Dirisamer M, Naveiras M, et al, Outcomes of Descemet membrane endothelial keratoplasty in phakic eyes, *J Cataract Refract Surg*, 2012, in Press.
66. Price FW Jr, Price MO, Comment on 'Spontaneous corneal clearance despite graft detachment after descemet membrane endothelial keratoplasty', *Am J Ophthalmol*, 2010;149:173-4; author reply 174-5.
67. Zafirakis P, Kymionis GD, Grentzelos MA, Livir-Rallatos G, Corneal graft detachment without corneal edema after Descemet stripping automated endothelial keratoplasty, *Cornea*, 2010;29:456-8.
68. Jacobi C, Zhivov A, Korbmacher J, et al, Evidence of endothelial cell migration after Descemet membrane endothelial keratoplasty, *Am J Ophthalmol*, 2011;152:537-42.
69. Lagali N, Stenevi U, Claesson, M, et al, Donor and recipient endothelial cell population of the transplanted human cornea: a two-dimensional imaging study, *Invest Ophthalmol Vis Sci*, 2010;51:1898-1904.
70. Stewart RM, Hiscott PS, Kaye SB, Endothelial migration and new Descemet membrane after endothelial keratoplasty, *Am J Ophthalmol*, 2010;149:683; author reply 683-4.
71. Dirisamer M, Yeh RY, van Dijk K, et al, First series of Descemet membrane endothelial transfer (DMET) suggests recipient endothelium contributes to corneal clearance, *Am J Ophthalmol*; In press.

