



Universiteit
Leiden
The Netherlands

The role of the Arabidopsis AHL15/REJUVENATOR gene in developmental phase transitions

Karami, O.; Karami O.

Citation

Karami, O. (2017, September 5). *The role of the Arabidopsis AHL15/REJUVENATOR gene in developmental phase transitions*. Retrieved from <https://hdl.handle.net/1887/54683>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/54683>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/54683> holds various files of this Leiden University dissertation

Author: Karami, Omid

Title: The role of the Arabidopsis AHL15/REJUVENATOR gene in developmental phase transitions

Issue Date: 2017-09-05

Chapter 4

Rapid chromatin decondensation by the Arabidopsis AT-Hook domain protein AHL15 leads to long-term repression of the ageing pathway

Omid Karami¹, Christiaan Henkel ² and Remko Offringa¹

¹ Plant Developmental Genetics, and ² Bioinformatics and Genomics, Institute of Biology Leiden, Leiden University, Sylviusweg 72, 2333 BE Leiden, The Netherlands

Abstract

In the previous chapters, we documented that overexpression of the *Arabidopsis* nuclear protein AHL15 leads to reprogramming of somatic cells to embryonic cells and to suppression of plant ageing. Here we show that transient (4 hours) activation of overexpressed AHL15-GR in *Arabidopsis* seedlings has long-term effects on plant development. RNA sequencing analysis detected an extensive reprogramming of the transcriptome 4 hours after AHL15-GR activation, with respectively 540 and 1107 genes showing more than 2-fold up- and down-regulation. AHL15 seemed to act in a transcription level-dependent manner, activating predominantly low expressed genes and repressing mostly highly expressed genes. Rapid decondensation of heterochromatin was observed after AHL15 activation in leaf primordia and axillary meristems, indicating that the global reprogramming of the transcriptome by transient activation of this AT-Hook domain protein might at least in part be caused by extensive modulation of the chromatin configuration. Co-activated or co-repressed genes were often physically linked in small chromosomal clusters, which is in line with regulation at the chromatin level. More detailed analysis of down-regulated genes indicated that AHL15 represses plant ageing by targeting several components of the ageing pathway, including the *SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE* (*SPL*) genes, GA biosynthesis and photosynthesis-dependent sugar production.

Keywords: AHL15, *Arabidopsis*, heterochromatin decondensation, RNA sequencing, transcriptome

Introduction

In flowering plants, ageing is defined by a series of developmental phase transitions that starts with the embryonic to vegetative phase change during germination, and is followed by the juvenile to adult (vegetative) phase change, and the vegetative to reproductive phase change, ultimately culminating in gametogenesis, embryogenesis and seed production. Each developmental phase is characterized by a unique set of morphological traits and the production and growth of specific organs (Huijser and Schmid, 2011). During the past few years, molecular genetic studies have demonstrated that the timing of these developmental phase transitions is orchestrated by specific sets of key regulators, which the SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) proteins play a central role in promoting the vegetative (juvenile to adult)- and the vegetative to reproductive phase transition. Early during development, high levels of *miR156* repress the production of the SPL proteins, thereby maintaining plants in the juvenile phase. A gradual decrease in *miR156* expression during the progression of development leads to increased SPL levels, which promote the progression of development through the subsequent phase transitions (Wu and Poethig, 2006; Xie et al., 2006; Saeteurn K, 2007; Wu et al., 2009; Andrés and Coupland, 2012).

In Chapters 2 and 3 we have shown that the AT-HOOK MOTIF NUCLEAR LOCALIZED protein AHL15 and other members of the plant-specific AHL protein family redundantly repress the ageing pathway by delaying the juvenile to adult and the vegetative to reproductive phase transition. In fact, overexpression of *AHL15* was able to reverse developmental phase transitions, resulting in the hormone-independent induction of somatic embryos on immature zygotic embryos or seedlings, and converting the monocarpic annual *Arabidopsis thaliana* into a polycarpic plant. How AHL15 can have such a dramatic effect on developmental phase transitions is one of the key questions to be addressed.

The paradigm for transcription factors is that they are nuclear proteins that bind to upstream regulatory DNA sequences and activate and/or repress transcription of target genes by respectively facilitating or inhibiting the recruitment of RNA polymerase to the transcription start site. By contrast, proteins of the plant-specific AHL family are a specific class of nuclear proteins that unlike most transcription factors, do not bind the major groove of the DNA helix, but instead interact with the narrow minor groove of DNA (Matsushita et al., 2007; Ng et al., 2009; Zhao et al., 2013). AHL proteins contain at least one AT-hook DNA binding motif and a Plant and Prokaryote Conserved (PPC) domain. Based on mutants and protein-protein interaction studies, the AHL family members have been proposed to bind AT-rich DNA regions as hetero-multimeric complexes that recruit other transcription factors through their interacting PPC domains (Zhao et al., 2013). In addition, it has been shown that AHL proteins repress transcription of several key developmental regulatory genes, possibly through modulation of the epigenetic code in the vicinity of its DNA binding regions (Lim et al., 2007; Ng et al., 2009; Yun et al., 2012). Some evidence has been obtained that AHL proteins function by altering the organization of the chromatin structure (Lim et al., 2007; Ng et al., 2009; Yun et al., 2012; Xu et al., 2013). Thus, different modes of transcription regulation by the AHL proteins have been described, but since this plant specific class of nuclear proteins is not well studied, their exact mode of action is still elusive.

Many AT-rich sequences in the DNA function as matrix attachment regions (MARs), which are well known to interact with the nuclear matrix, a fibrillar network of proteins inside the nucleus. Although, MARs are widely distributed in the genome, they are commonly found at the boundaries of transcription units. MARs play important roles in the higher-order organization of chromatin structure, thereby regulating gene expression (Heng et al., 2004; Girod et al., 2007; Chavali et al., 2011; Wilson and Coverley, 2013). The majority of the animal AT-hook motif containing DNA-binding proteins are localized to MARs where they are associated with proteins that modulate chromatin architecture (Fusco and Fedele, 2007). Also for the plant-specific AHL proteins it has been shown that they preferentially bind to the AT-rich DNA sequences of MARs (Morisawa et al., 2000; Fujimoto et al., 2004; Lim et al., 2007; Ng et al., 2009; Xu et al., 2013). However, a correlation between the function of AHL proteins and their localization to MARs has not been documented yet.

Here we observed that short-term activation (4 hours) of 35S::AHL15-GR seedlings, overexpressing a fusion between AHL15 and the dexamethasone (DEX) responsive domain of the glucocorticoid receptor, resulted in long-term effects on plant development, such as delayed flowering, enhanced branching and the recurrent formation of aerial rosettes, converting monocarpic *Arabidopsis* into a polycarpic plant. In order to understand the AHL15 mode of action, we compared the chromatin configuration and transcriptome of mock- or DEX-treated 35S::AHL15-GR seedlings by respectively nuclear staining and high throughput next generation sequencing of transcripts (Illumina RNA-Seq) (Mortazavi et al., 2008). Rapid decondensation of heterochromatin was observed after AHL15-GR activation in leaf primordia and axillary meristems, indicating that the observed global reprogramming of the transcriptome by transient activation of this AT-Hook domain protein might at least in part be caused by extensive modulation of the chromatin configuration. More detailed analysis of down-regulated genes indicated that AHL15 represses plant ageing by targeting several components of the ageing pathway, including the *SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE* (SPL) genes, GA biosynthesis and photosynthesis-dependent sugar production.

Results

Short-term activation of AHL15-GR has long-term effects on plant development

In Chapter 3 we showed that *Arabidopsis* plants that constitutively express AHL15 (35S::AHL15) or plants that express a dexamethasone (DEX) activatable version of AHL15 (35S::AHL15-GR) and were subjected to continuous DEX treatment both showed a strong delay in the juvenile-to-adult transition and flowering time. In contrast, *ahl* loss-of-function mutant plants or plants expressing a dominant negative AHL15-GUS fusion showed a premature vegetative phase change and early flowering. Moreover, AHL15 overexpression induced rejuvenation of axillary meristems, resulting in enhanced branching and in the production of aerial rosettes, converting monocarpic *Arabidopsis* into a polycarpic plant. For typical transcription factors it has been shown that they act transiently, whereas other nuclear factors have a more long-lasting effect on the gene expression profile of a cell, as they act on the chromatin structure by inducing epigenetic changes (Bratzel and Turck,

2015). In order to analyse the mode of action of AHL15, 5-day-old *35S::AHL15-GR* seedlings were submerged in water with 20 μ M DEX or without DEX (mock) for 4 and 8 hours, and subsequently DEX was removed by washing the seedlings in water and transferring them to soil. DEX-incubated seedlings developed much slower and the resulting plants showed a significant delay in flowering compared to the plants derived from the mock-treated seedlings (Fig. 1A). DEX-treatment for 8 hours enhanced the phenotypes observed in the 4 hour DEX-treated seedlings and derived plants (not shown). Also, 35-day-old flowering *35S::AHL15-GR* plants that were only sprayed once with 20 μ M DEX developed aerial rosettes from their axillary meristems 7-10 days after spraying (Fig. 1B), and new aerial rosettes continued to develop for at least 4 months after spraying (Fig. 1C-E). These data indicates that short-term activation of AHL15 leads to long-term changes in development, which would be in line with a function for AHL15 in chromatin remodeling.




Figure 1 Short term activation of AHL15 has long term effects on plant development. (A) Phenotype (upper panel) and quantification of the number of rosette leaves (lower panel) of 35 day old flowering *35S::AHL15-GR* plants that were grown from 5 day old seedlings that were submerged for 4 hours in water (4h_mock) or in 20 μ M DEX solution (4h_DEX). Asterisk in the graph indicates a significant difference between mock- and DEX-treated plants (Student's *t*-test, $p < 0.01$), and error bars indicate standard error of the mean ($n = 20$). (B-E) Aerial rosettes developing from axillary meristems in *35S::AHL15-GR* plants 10 days (B), or one (C), two (D) or four months (E) after spraying 1 month old flowering *35S::AHL15-GR* plants with 20 μ M DEX solution. Plants were grown under LD.

AHL15 extensively reprograms the *Arabidopsis* transcriptome

In order to uncover the molecular basis of these long term developmental changes induced by AHL15, the genome-wide expression changes were compared between DEX-treated and untreated 35S::AHL15-GR seedlings. As the major developmental changes induced by AHL15 overexpression were observed in the shoot (Chapter 3, Figure 1), we isolated RNA from the shoot part (i.e. shoot apex, cotyledons and top part of hypocotyl) of 5-day-old 35S::AHL15-GR seedlings submerged for 4 hours in mock or in 20 μ M DEX, or for 8 hours in 20 μ M DEX. RNA sequencing was performed on RNA isolated from three biological replicates for each treatment. A slight but significant reduction of AHL15-GR expression was observed in DEX-treated seedlings compared to untreated samples (Table 1), suggesting that AHL15 modulates the 35S promoter activity in this time period.

Table 1. Overview of raw data obtained by Illumina sequencing on RNA isolated from the shoot part of 5-day-old 35S::AHL15-GR seedlings that were treated with water for 4 hours (4h_mock)) or with 20 μ M DEX for 4 or 8 hours (resp. 4h_DEX or 8h_DEX).

Sample #	Treatment	Raw sequencing reads	Expression level of AHL15 ¹	
1	4h_mock	14285557	678.13	687.06 $\pm$ SE
2	4h_mock	17597921	636.59	
3	4h_mock	27463033	746.46	
4	4h_DEX	32465125	519.36	519.70 $\pm$ SE *
5	4h_DEX	21100996	526.87	
6	4h_DEX	28328909	512.89	
7	8h_DEX	30340913	519.13	532.90 $\pm$ SE *
8	8h_DEX	26594329	538.09	
9	8h_DEX	30496134	541.48	

¹ Normalized Reads Per Kilobase Million (RPKM) values. Right column indicates the average value of the three biological replicates $\pm$ standard error of the mean.

* Significantly different from 4h_mock (Student's *t*-test, $p < 0.001$)

Annotation of the reads using the *Arabidopsis* Information Resource (www.arabidopsis.org) showed that the expression of 22570 genes could be detected in the 4 hours mock samples, which is 83% of the total gene number (Table S1). Statistical analysis showed that 13483 genes were differentially expressed in 4 hours DEX-treated compared to mock treated 35S::AHL15-GR shoot organs (Table S2). However, only 1663 genes showed a fold change of ≥ 2 (and $P \leq 0.05$) after 4 hours DEX treatment (Fig. 2A, B, Table S3 and S4). Verification of the change in expression for 4 selected genes by qRT-PCR analysis showed a good agreement with the RNA-sequencing results (Table 2).

Comparison of the up- and downregulated gene sets showed a considerable overlap between 4 and 8 hours DEX treatment (Fig. 2A,B). Still, however, the number of up- (Table S5) and down-regulated (Table S6) genes increased significantly after 8 hours DEX treatment (Fig. 2A,B). Several of the genes that did not show a ≥ 2 fold change after 4 hours DEX treatment reached this level after 8 hours treatment (Fig. 2A,B). Conversely, some genes that reached ≥ 2 fold change after 4 hours, did not appear anymore in the ≥ 2 fold gene set after 8

hours (Fig. 2A,B), indicating time-dependent transcriptome changes by AHL15-GR. The latter result would be more in line with a role for AHL15 as a gene-specific transcription factor.

Table 2. Confirmation of RNA-seq data by qRT-PCR analysis.

Locus	Name and short description	Fold change found by RNA Seq	P Value	Fold change detected by qRT-PCR ¹	P Value
AT5G59490	Haloacid dehalogenase-like hydrolase (HAD) superfamily protein	135.5	2.07E-145	78.30 ± 7.80	0.003
AT4G22770	AT hook motif DNA-binding family protein 2	15.19	8.02E-139	9.32 ± 1.45	0.005
AT2G17740	Cysteine/Histidine-rich C1 domain family protein	0.002	1.21E-62	0.008 ± .002	0.0003
AT1G78580	Trehalose-6-phosphate synthase 1	0.35	2.11E-54	0.43 ± 0.06	0.021

¹Shown is the fold changes ± standard error of the mean (n=3). P value: Student's t-test.

Notably, we found that the down-regulated genes were generally expressed at higher levels in the 4h_mock control sample, whereas the up-regulated genes were expressed at relatively low levels in the 4h_mock control sample (Fig. 2C). This data suggests that AHL15 reverses phase changes by modulating gene activity in a transcription level-dependent manner, repressing genes that are highly expressed and activating genes that are low expressed during a specific developmental phase. Global prediction of the transcriptome changes using gene ontology (GO) examination (Du et al., 2010) (<http://bioinfo.cau.edu.cn/agriGO/>) and TAIR as reference for the annotation of *Arabidopsis* genes indicated that up- and down-regulated gene sets grouped in many different biological categories (Tables S7 and S8). These results suggested that, beside its action as regulator of individual genes, AHL15 contributes to a more global reprogramming of biological processes by inducing global changes in gene transcription.

Whereas most gene families represented in the transcriptome profiles (Table S2) had gene members that were either up- or down-regulated, members of the *CYSTEIN-RICH RECEPTOR-LIKE KINASE* (CRK) gene family only showed down-regulation of gene expression by AHL15 (Table 3). Notably, several of the down-regulated CRKs by AHL15 are neighboring genes that are located in a tandem arrays on chromosome 4 (Fig. 3A). This triggered us to look into the localization of up- or down-regulated genes, and surprisingly, a high rate (~75%) of co-activation or co-repression of neighboring genes by AHL15 was observed across the genome (Tables S9, and Fig. 3B). These observations suggest that AHL15 modulates gene expression in a chromosomal position- rather than a gene-specific manner.




Figure 2 Changes in the transcriptome of *Arabidopsis* 35S::AHL15-GR seedling shoots after DEX treatment. (A, B) Venn diagrams indicating the number of (overlapping) down-regulated (A) or up-regulated (B) genes in 35S::AHL15-GR seedling shoots with a ≥ 1.5 fold or ≥ 2 fold change in gene expression following 4 or 8 hours of DEX treatment. (D) Graph showing the percentage of genes having a specific expression level (in RPKM) in mock treated 35S::AHL15-GR seedling shoots that are either ≥ 2 fold up- (red) or ≥ 2 fold down- (blue) regulated by 4 hours DEX treatment.

Table 3. Members of the *Cysteine-rich receptor-like kinase* gene family were all down-regulated in 35S::AHL15-GR seedling shoots after 4 hours DEX treatment

Locus	Name and short description	Fold change	P Value
AT4G23180	cysteine-rich RLK (RECEPTOR-like protein kinase) 10	0.34	5.49E-29
AT4G23190	cysteine-rich RLK (RECEPTOR-like protein kinase) 11	0.32	4.58E-39
AT4G23200	cysteine-rich RLK (RECEPTOR-like protein kinase) 12	0.04	1.92E-87
AT4G23210	cysteine-rich RLK (RECEPTOR-like protein kinase) 13	0.08	1.28E-44
AT4G23260	cysteine-rich RLK (RECEPTOR-like protein kinase) 18	0.138	9.12E-165
AT4G23270	cysteine-rich RLK (RECEPTOR-like protein kinase) 19	0.82	0.0098589
AT1G70520	cysteine-rich RLK (RECEPTOR-like protein kinase) 2	0.40	5.80E-33
AT4G23290	cysteine-rich RLK (RECEPTOR-like protein kinase) 21	0.11	6.19E-110
AT4G23300	cysteine-rich RLK (RECEPTOR-like protein kinase) 22	0.17	1.37E-106
AT4G21400	cysteine-rich RLK (RECEPTOR-like protein kinase) 28	0.77	7.47E-05
AT4G21410	cysteine-rich RLK (RECEPTOR-like protein kinase) 29	0.94	0.33050
AT1G70530	cysteine-rich RLK (RECEPTOR-like protein kinase) 3	0.40	7.76E-26
AT4G11460	cysteine-rich RLK (RECEPTOR-like protein kinase) 30	0.31	5.07E-19
AT4G04490	cysteine-rich RLK (RECEPTOR-like protein kinase) 36	0.14	1.29E-53
AT4G04570	cysteine-rich RLK (RECEPTOR-like protein kinase) 40	0.30	4.74E-63
AT4G00970	cysteine-rich RLK (RECEPTOR-like protein kinase) 41	0.09	2.31E-62
AT5G40380	cysteine-rich RLK (RECEPTOR-like protein kinase) 42	0.25	1.70E-23
AT4G23130	cysteine-rich RLK (RECEPTOR-like protein kinase) 5	0.19	5.63E-41




Figure 3 AHL15 modulates gene expression in a chromosomal region-dependent manner. (A) Schematic representation of *Arabidopsis* chromosome 4 showing the tandem arrays of CRKs genes on this chromosome. RNA Seq analysis on 5 day old 35S::AHL15-GR seedlings shoots treated for 4 hours with 20 μ M DEX shows that some neighboring CRK genes are co-repressed (indicated by the red line). (B) Schematic representation of *Arabidopsis* chromosome 1 showing for one region that neighboring genes are co-repressed (red line) or co-induced (green line) in 5 day old 35S::AHL15-GR seedling shoots following 4 hours DEX treatment.

AHL15 overexpression results in reduced heterochromatin condensation

Based on previous analysis in animals that AT-hook proteins are able to change the higher order compaction of chromatin organization (Catez et al., 2004; Kishi et al., 2012; Postnikov and Bustin, 2016), we compared the chromatin organization in leaf cells of two week old 35S::AHL15 seedlings with that in wild-type leaf cells. By imaging DAPI stained nuclei or nuclei of leaf cells expressing GFP-tagged histone H2B, a considerable reduction of heterochromatin could be detected in 35S::AHL15 leaf cells compared to wild-type cells (Fig. 4A). This reduction in heterochromatin by AHL15 was even more clearly observed by imaging GFP-H2B in leaf primordia produced from axillary meristems on inflorescences (Fig. 4B). Time-lapse imaging of GFP signals revealed a rapid gradual reduction in heterochromatin condensation in DEX-treated compared to mock-treated 35S::AHL15-GR leaf primordia (Fig. 4B), indicating that the AHL15-induced heterochromatin opening occurs in a time-dependent manner and within the time frame of the transcriptome analysis. These data support the view that the observed global reprogramming of the transcriptome by AHL15 might be caused by extensive modulation of the chromatin configuration.




Figure 4 AHL15 overexpression induces rapid heterochromatin decondensation. (A) Visualization of heterochromatin using DAPI-staining or H2B-GFP labelling of nuclei in leaf primordia or fully developed leaf cells in 2-week-old wild-type (upper row) or 35S::AHL15 (lower row) plants. (B) Heterochromatin labelling by H2B-GFP in nuclei of 35S::AHL15-GR axillary leaf primordium cells at 4, 8, 24, or 48 hours after mock (upper row) or DEX treatment (lower row).

AHL15 represses ageing genes

Since *AHL15* expression delays and even reverses plant ageing (Chapter 2 and 3), we searched for age regulatory genes among the differentially expressed genes in the transcriptome profiles. The first gene family we looked for were the *SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE* (*SPL*) genes encoding transcription factors that promote the juvenile-to-adult transition in *Arabidopsis* (Wu and Poethig, 2006; Xie et al., 2006; Saeteurn K, 2007; Wu et al., 2009). *SPL* mRNAs are targeted by MicroRNA156 (*miR156*), and the juvenile to adult transition is attributed to the down-regulation of *miR156* gene expression and the subsequent increase in *SPL* protein abundance (Wu and Poethig, 2006; Xie et al., 2006; Saeteurn K, 2007; Wu et al., 2009). Our transcriptome data revealed that *SPL4*, *SPL9*, *SPL13* and *SPL15* were significantly down regulated after *AHL15-GR* activation (Fig 5A), whereas the expression of *SPL2*, *SPL11*, *SPL10* was not significantly changed (Fig 4A). The expression of *SPL3* and *SPL5* was undetectable in the RNA seq-based transcriptome profile (Fig 5A), which could be related to the high levels of degradation of *SPL3* and *SPL5* mRNAs by *miR165* in young seedlings. This RNA seq-based result triggered us to further study the genetic regulation *SPL* genes by *AHL15*. Quantitative RT-PCR (qPCR) analysis confirmed that the expression of *SPL3*, *SPL9*, and *SPL15* was significantly down-regulated in 4 hour DEX-treated *35S::AHL15-GR* seedlings (Fig. 5B). Time-lapse GUS staining of *pSPL3::GUS-SPL3* plants (Yang et al., 2011) revealed a gradual reduction of *SPL3* expression in response to continuous DEX activation of *AHL15-GR* (Fig. 5C). These results indicate that the *SPL* genes are gradually suppressed by *AHL15*.

Consistent with maximum activity of the *SPL* genes in inflorescence stems (Cardon et al., 1999), we analyzed the effect of *AHL15* overexpression on *SPL* expression levels in inflorescence nodes. Compared to wild-type inflorescence nodes, qPCR detected only low expression of *SPL3*, *SPL9* and *SPL15* in *35S::AHL15* inflorescence nodes (Fig. 5D). These results displayed that *AHL15* overexpression strongly suppresses *SPL* genes in the reproductive phase, which is likely to maintain axillary meristems in *35S::AHL15* plants in the vegetative state during flowering, thereby resulting in the production of areal rosettes and enhanced branching (Chapter 3).

To determine whether down-regulation of *SPL* genes by *AHL15* is dependent on *miR156* function, two *miR156*-insensitive reporters, *pSPL9::rSPL9-GUS* and *pSPL3::GUS-rSPL3* (Yang et al., 2011), were introduced into *35S::AHL15-GR* plants. The expression of *pSPL9::rSPL9-GUS* and *pSPL3::GUS-rSPL3* was significantly down-regulated in DEX-treated *35S::AHL15-GR* seedlings compared with mock-treated seedlings (Fig. 5E). These results indicated that *AHL15* overexpression suppresses *SPL* genes in a *miRNA156*-independent manner.




Figure 5 AHL15 represses *SPL* expression in a miRNA-independent manner. (A) The relative expression level (in RPKM) of different *SPL* genes detected by RNA Seq analysis on 5 day old 35S::AHL15-GR seedlings shoots following 4 hours water (mock) or 20 μ M DEX treatment. (B) The relative expression level of *SPL* genes by qPCR analysis on shoot parts of 2 week old 35S::AHL15-GR seedlings incubated for 1 day on medium without (mock) or with 20 μ M DEX (DEX). (C) Histochemical staining for GUS activity on 3 week old GUS-*SPL3* 35S::AHL15-GR seedlings incubated on medium without (mock, right) or with 20 μ M DEX (DEX, left) for 1, 2 or 3 days (respectively D1, D2 or D3). (D) Relative expression level of *SPL3*, *SPL9* and *SPL15* by qPCR analysis on the base regions of 1 week old inflorescences of wild-type (Col) or 35S::AHL15 #1 or #2 plants. (E) Histochemical staining for GUS activity on 2 week old water (mock) or 20 μ M DEX (DEX) treated 35S::AHL15-GR plants expressing miR156-resistant -*SPL3*-r*SPL3*-GUS or *SPL9*-r*SPL9*-GUS reporters. Asterisks in A, B and D indicate significant difference from mock-treated (A,B) or wild-type (D) plants (Student's *t*-test, $p < 0.01$). Error bars indicate standard error of the mean ($n = 3$).

One of the plant hormones involved in ageing is gibberellic acid (GA), as it promotes the juvenile-to-adult or the vegetative-to-reproductive phase transitions (Telfer et al., 1997; Mutasa-Göttgens and Hedden, 2009; Poethig, 2013). In our transcriptome profile we observed that the GA biosynthesis gene *GIBBERELLIN 20-OXIDASE 2* (*GA20OX2*) was highly down-regulated in 4 hour DEX-treated *35S::AHL15* seedlings (Fig 6A). *GA20OX2* is a rate limiting enzyme in the last steps of the GA biosynthetic pathway (Huang et al., 1998; Rieu et al., 2008; Andrés et al., 2014). In line with this down-regulation, meristem rejuvenation in DEX treated flowering *35S::AHL15-GR* plants was remarkably reduced when these plants were treated with GA 3 days after DEX (Fig 6B). We conclude that the repression of GA biosynthesis, next to a decrease in *SPL* gene expression, plays an important role in the AHL15-mediated suppression of phase transitions and axillary meristem rejuvenation.



Figure 6 AHL15-induced developmental changes by reduced GA biosynthesis. (A) The relative expression level of the *GA20OX2* gene detected by RNA Seq analysis on 5 day old *35S::AHL15-GR* seedlings submerged for 4 hours in water (4h_mock) or 20 μ M DEX solution (4h_DEX). Asterisk indicates significant difference from mock seedlings (Student's *t*-test, $p < 0.001$). Error bars indicate standard error of the mean ($n = 3$). (B) Shoot phenotypes of two month old *35S::AHL15-GR* plants derived from 35 day old DEX sprayed plants that were subsequently sprayed 3 days later with water (-GA) or with 15 μ M GA4 (+GA).

Reduced photosynthesis in *35S::AHL15* plants causes sucrose-dependent seedling growth

In the transcriptome profile of DEX-treated *35S::AHL15-GR* seedlings we detected down-regulation of several photosynthesis-related genes (Table 4). The MapMan hierarchical ontology software (Thimm et al., 2004) showed that several of the down regulated genes encode for components of photosystem I and II, such as the light harvesting complexes, and the reaction centers (Fig. 7A, Table 4). These are among the most highly expressed genes in plants, and the fact that they are repressed by AHL15 is in line with the previously proposed transcription level-dependent manner of regulating gene expression by AHL15.

Recent studies suggest that the increasing photosynthesis efficiency and sugar concentration in shoot organs of the developing young plant promote the vegetative phase change (Matsoukas et al., 2013; Yang et al., 2013; Yu et al., 2013). The suppression of

ageing in *35S::AHL15* seedlings could therefore at least in part be caused by a delayed increase in the photosynthesis efficiency. To confirm that *AHL15* overexpression reduces the photosynthesis efficiency and thus the sugar concentration, we germinated wild-type or *35S::AHL15* seedlings on medium with or without sucrose. On medium with sucrose, both wild-type and *35S::AHL15* seedlings developed, albeit that development of *35S::AHL15* seedlings was retarded, as observed previously (Chapter 3). On medium without sugar wild-type seedlings also developed, but generally much slower than sucrose grown seedlings, suggesting that photosynthesis under these *in vitro* conditions is rate limiting (Fig. 7B). In contrast, *35S::AHL15* seeds did germinate on medium without sucrose but seedling development was completely arrested (Fig. 7B). The results indicate that the germinating seedlings lack endogenous sugars, and therefore are completely dependent for their development on externally provided sucrose, most likely due to *AHL15*-mediated repression of photosynthesis during germination.

Table 4. List of photosynthesis-related genes that were down-regulated in *35S::AHL15-GR* seedling shoots after 8 hours DEX induction

Locus	Name and descriptions	Expression level in 4h_water	Fold change	P Value
AT4G10340	light harvesting complex of photosystem II 5	3485	0.37	1.58E-44
AT5G01530	light harvesting complex photosystem II	2405	0.45	3.93E-35
AT3G08940	light harvesting complex photosystem II	1043	0.34	2.54E-37
AT1G15820	light harvesting complex photosystem II subunit 6	2323	0.44	1.29E-32
AT5G54270	light-harvesting chlorophyll B-binding protein 3	3520	0.18	5.28E-78
AT3G47470	light-harvesting chlorophyll-protein complex I subunit A4	3885	0.18	2.13E-93
AT2G34430	light-harvesting chlorophyll-protein complex II subunit B1	2402	0.08	1.22E-118
AT3G54890	photosystem I light harvesting complex gene 1	3839	0.23	2.29E-103
AT1G61520	photosystem I light harvesting complex gene 3	2957	0.39	8.23E-42
AT2G05100	photosystem II light harvesting complex gene 2.1	1773	0.04	2.11E-109
AT2G05070	photosystem II light harvesting complex gene 2.2	1495	0.05	7.91E-101
AT3G27690	photosystem II light harvesting complex gene 2.3	362	0.02	8.19E-97
AT2G34420	photosystem II light harvesting complex gene B1B2	5162	0.36	2.86E-51
AT1G03130	photosystem I subunit D-2	612	0.36	1.03E-32
AT1G31330	photosystem I subunit F	3256	0.41	1.78E-40
AT3G50820	photosystem II subunit O-2	713	0.41	4.87E-38
AT1G08380	photosystem I subunit O	3446	0.37	2.84E-58
AT5G64040	photosystem I reaction center subunit PSI-N, chloroplast	2222	0.37	2.58E-37
AT4G28750	Photosystem I reaction center subunit IV / PsaE protein	1711	0.49	2.94E-26
AT2G30570	photosystem II reaction center W	2243	0.48	5.79E-26




Figure 7 Repression of photosynthesis by AHL15 results in sucrose-dependent seedling growth. (A) A simplified scheme generated by the MAPMAN software (Thimm et al., 2004) showing that the expression of most genes encoding Photosystem I and II components is repressed in 35S::AHL15-GR seedling shoots following 8 hours of DEX treatment. Blue squares indicate up-regulated genes, red squares indicate down-regulated genes, and gray dots indicate genes for which the expression did not change. (B) Phenotypes of 4 week old wild-type (Col-0, top) and 35S::AHL15 plants (bottom) germinated and grown on medium containing 1% sucrose (left) or no sucrose (right).

Discussion

The *Arabidopsis* AT-hook motif nuclear-localized protein AHL15 delays phase transitions during plant development. In fact, overexpression of this protein can even reverse these phase transitions, resulting in the 2,4-D-independent induction of somatic embryos on cotyledons of immature zygotic embryos or seedlings, or in the appearance of juvenile aerial rosettes from axillary meristems of flowering plants. In addition to these initial results (Chapters 2 and 3 of this thesis), we observed that short transient activation of a constitutively expressed AHL15-GR fusion led to long-term effects on plant developmental timing and ageing. Activation of AHL15-GR by DEX treatment of seedlings resulted in a significant delay of development and flowering for up to a month after treatment, whereas spraying flowering plants with DEX resulted in the development of aerial rosettes from axillary meristems for up to 4 months after treatment. These results suggested that AHL15 establishes a long-term molecular memory, which made us wonder about the mode of action of this plant-specific class of nuclear DNA binding proteins.

Genome-wide analysis of transcriptome changes following transient activation of AHL15-GR showed that AHL15 modulates the expression level of a large number of genes participating in various biological processes. Some genes were induced after 4 hours but showed basal expression levels again after 8 hours, suggesting the direct transient activation of target genes that is typical for normal transcription factors. On the other hand, the large number of genes with a changed expression profile suggested a more global reprogramming of cellular processes. Previous studies have demonstrated that the dynamics of higher-order chromatin organization plays a critical role in both the global regulation of the transcriptome (Ho and Crabtree, 2010; Li and Reinberg, 2011; Pombo and Dillon, 2015) and the establishment of long-term molecular memory (He and Amasino, 2005; Jarillo et al., 2009; Harmston and Lenhard, 2013).

An additional result supporting a regional rather than single gene regulation mode was the observation that neighboring genes are co-activated or co-suppressed by AHL15. Previous analyses of genome-wide gene expression datasets using bioinformatics approaches have shown that neighboring genes in *Arabidopsis* are more co-expressed than random gene pairs (Williams and Bowles, 2004; Zhan et al., 2006; Chen et al., 2010; Wada et al., 2012; Yeaman, 2013; Kundu et al., 2017). Several mechanisms have been proposed to explain this phenomenon, such as gene duplications, shared promoters or common transcription factor binding motifs, but chromatin architecture has always been considered as the major source of co-expression of neighboring genes (Grob et al., 2014; Pombo and Dillon, 2015; Quintero-cadena and Sternberg, 2016). A recent genome-wide data analysis has excluded that co-regulation of neighboring genes in *Arabidopsis* is caused by gene duplications or the presence of common promoter motifs in neighboring genes (Kundu et al., 2017). Instead, co-regulation could be clearly correlated with local rearrangement of chromatin configuration (Kundu et al., 2017). Therefore, we suggest that the co-activation or co-repression of neighboring genes across genome by AHL15 is most likely also caused by extensive modulation of the chromatin configuration.

In animals, the contribution of higher-order chromatin organization to ageing processes has been well documented (Moraes, 2014; Chandra and Kirschner, 2016; Gorbunova and

Seluanov, 2016). In contrast, the role of global organization of chromatin architecture in plant Ageing is not described yet. *Arabidopsis* adult leaves display a visible increase of chromatin compaction compared to juvenile leaves (Exner and Hennig, 2008), but the actual involvement of this chromatin compaction in the juvenile-to-adult transition has not been reported yet. Our results suggest that the global changes in gene expression after 4 hours of DEX-induced AHL15-GR activation coincide with rapid de-condensation of heterochromatin, thereby clearly linking plant ageing or tissue rejuvenation to respectively an increase or a reduction in the heterochromatin. Our results suggest that the extensive reprogramming of the transcriptome and the observed establishment of a long-term molecular memory following AHL15-GR activation might at least in part be caused by an extensive reorganization of the chromatin configuration.

Previously, it has been shown that the juvenile-to-adult vegetative transition is mediated mainly by SPL proteins, which are central components of the ageing pathway in several plant species (Wu and Poethig, 2006; Xie et al., 2006; Saeteurn K, 2007; Wang et al., 2011; Zhang et al., 2011; Fu et al., 2012; Xie et al., 2012). *SPL* genes are well known targets of miR156 (Wu and Poethig, 2006; Xie et al., 2006;; Wu et al., 2009), and a steep decline in miR156 levels over 1-2 weeks of shoot development results in an increase in SPL protein levels (Xu et al., 2016). This pushes plant development to the adult vegetative phase, and from there to the induction of flowering. Our expression- and genetic analyses showed that AHL15 represses the expression of several *SPL* genes, independent of miR156. In both *Arabidopsis* wild-type and *AHL15* overexpression plants this delays both the juvenile to adult and the vegetative to reproductive phase transitions. However, we also found evidence that AHL15 represses *GA20ox2*, a rate limiting enzyme in GA biosynthesis pathway (Huang et al., 1998; Rieu et al., 2008; Andrés et al., 2014). In several plant species GA was shown to promote the vegetative phase transition (Telfer et al., 1997; Park et al., 2017). Moreover, we found that AHL15 represses several genes encoding components of the photosynthesis machinery. Recent studies have shown that the gradual increase in the sugar levels as a result of enhanced photosynthetic efficiency or an increase in leaf numbers promotes the juvenile-to-adult transition in *Arabidopsis*, tobacco, and tomato (Matsoukas et al., 2013; Yang et al., 2013; Yu et al., 2013). In contrast, a mutation in the *Arabidopsis* *CAO* gene that causes a low photosynthetic efficiency, was found to prolong the juvenile vegetative phase (Espine et al., 1999; Yu et al., 2013). By combining all our findings we propose a model (Fig. 8) whereby AHL15 represses the *Arabidopsis* ageing pathway by inducing changes in higher-order chromatin organization that lead to repression of *SPL* genes, GA biosynthesis, and photosynthesis-mediated sugar production. Unfortunately, how AHL15 alters the chromatin structure remains unclear. Detailed studies on the chromatin configuration by new approaches such as chromosome conformation capture technologies (Dekker et al., 2013), and comparative analysis of the putative binding sites of AHL15 with our transcriptome data are directions of future research that might provide more insight into the mode of action of this plant-specific AT-motif protein.




Figure 8 Model for suppression of plant ageing by AHL15. Large-scale chromatin opening by AHL15 leads to repression of key age promoting factors such as the *SPL* genes, gibberellin (GA) biosynthesis, and photosynthesis leading to sugar production. Arrows indicate activation or induction, and blunted lines indicate repression. Note that AHL15-mediated *SPL* suppression is independent of miR156.

Methods

Plant material and RNA isolation and sequencing

The *Arabidopsis thaliana* (Col-0) plant line harboring the 35S::AHL15-GR construct is described in Chapter 3. The reporter lines *pSPL3::GUS-SPL3*, *pSPL3::rSPL3-GUS*, *pSPL9::rSPL9-GUS* (Yang et al., 2011) and *H2B::H2B-GFP* (Fang and Spector, 2005) have been described previously. Seeds were surface sterilized (30 sec 70% ethanol, 10 minutes 1% chlorine, followed by washes in sterile water) and germinated after three days incubation at 4°C on MA medium (Masson and Paszkowski, 1992) containing 1% or no sucrose, and 0.7% agar at 21 °C and a 16 hours photoperiod. Following germination, 14 days old seedlings were transferred to soil and grown at 21 °C, 65% relative humidity, and long-day (LD: 16 hours photoperiod) condition.

For the transcriptome analysis (specifically), MA plates with 5 day old seedlings were flooded with water containing 1 ml ethanol per liter (mock), or with water to which dexamethasone (DEX, Sigma-Aldrich) dissolved in 1 ml ethanol was added to a final

concentration of 20 μ M. After 4 and 8 hours treatment, the shoot part of seedlings, including the shoot apex and cotyledons, was carefully separated from the hypocotyl and immediately frozen in liquid nitrogen and stored at -80 °C for RNA isolation.

Total RNA was extracted using a Qiagen RNeasy Plant Mini Kit, and the quality of the isolated RNA was validated using a nanodrop spectrophotometer (NanoDrop, ND-1000, Life Science). The isolated RNAs were reverse transcribed and sequenced on an Illumina HiSeq 2000 (100 nucleotides single reads). Three biological replicates were performed.

To test the effect of GA on AHL15-GR activation, 35 day old flowering 35S::*AHL15-GR* plants were first sprayed with 20 μ M DEX, followed 3 days later by spraying with 15 μ M GA4 (Sigma-Aldrich).

Quantification of expression levels and differential expression analysis

The quality control of all sequencing samples was carried out using FASTQC (version 0.10.1: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). Reference sequences and annotations for the *Arabidopsis* genome (TAIR10) were obtained from www.arabidopsis.org. To obtain expression levels the reads were aligned to the *Arabidopsis* genome sequence using Tophat2 (version 2.0.10) (Kim et al., 2013), using Bowtie2 (version 2.1.0) (Langmead and Salzberg, 2012) as the short read aligner at ‘very sensitive’ settings. Secondary alignments, i.e. alignments that meet Tophat’s criteria but are less likely to be correct than simultaneously reported primary alignments, were removed from the BAM files using SAMtools (version 0.1.18) (Li et al., 2009). Fragment alignment counts per transcript were determined from SAM alignment files using the Python package HTSeq-count (version 0.5.3p9) (Anders et al., 2014) with ‘strict’ settings to exclude reads aligning ambiguously with respect to annotated gene structures. Counts were summarized at the level of annotated genes, resulting in between 12.992.327 and 29.972.677 aligned fragments per sample. Read counts per annotated gene were normalized across all samples using the DESeq-like robust scaling factor (Anders and Huber, 2010) on reads per kilobase per million mapped reads (RPKM) values. For 13496 genes which had an expression value ≥ 10 in at least two samples, differential expression statistics were calculated using the R package edgeR (version 3.2.4) (Robinson et al., 2010).

Gene ontology term analysis

Gene ontology (GO) term analysis for identification of enriched functional categories was performed using the agriGO single enrichment analysis tool (Du et al., 2010) (<http://bioinformatics.cau.edu.cn/easygo/>) with TAIR10 GO annotations. The MapMan software (Thimm et al., 2004) (<http://mapman.gabipd.org/>) was used to visualize pathways containing multiple genes with significant changes in expression.

Quantitative real-time PCR analysis

RNA isolation was performed using a RevertAidTM Kit (Thermo Scientific). For qRT-PCR (qPCR), 1 μ g of total RNA was used for cDNA synthesis with the iScriptTM cDNA Synthesis

Kit (BioRad). PCR was performed using the SYBR-Green PCR Master mix and amplification was run on a CFX96 thermal cycler (BioRad). The Pfaffl method was used to determine relative expression levels (Pfaffl, 2001). Expression was normalized using the β -*TUBULIN*-6 gene. Three biological replicates were performed, with three technical replicates each. The primers used are described in Table 5.

Table 5: Sequences of DNA primers used for qRT-PCR (from 5' to 3')

Name*	Sequence (5' to 3')	Purpose
q AT5G59490-F	TTCGAGAAAGCCTTCGACAT	qRT-PCR AT5G59490
q AT5G59490-R	ATAACTCGGGGAATGCCTCT	
q AT4G22770-F	TGCAGCCACTCCTATTCAAGT	qRT-PCR AT4G22770
q AT4G22770-R	GAAGGAAAAGACGGTGTCCAT	
q AT2G17740-F	GTCTGTGCCTGAGACCATGA	qRT-PCR AT2G17740
q AT2G17740-R	CTTCAGCAACGCATGAATGT	
q AT1G78580-F	GAAACTCAAGACGTCCTTCACCAG	qRT-PCR AT1G78580
q AT1G78580-R	TCTAGCATTGGTGCAGTACGAC	
qSPL3-F	CTCATGTTCGGATCTCTGGTC	qRT-PCR <i>SPL3</i>
qSPL3-R	TTTCCGCCTTCTCTCGTTGTG	
qSPL9-F	AACAATACATGGCGAGCTTCTT	qRT-PCR <i>SPL9</i>
qSPL9-R	ATTGCCGTGCCACTACTTATCT	
qSPL15-F	AATCCAGTTAGGGAAACCCATC	qRT-PCR <i>SPL15</i>
qSPL15-R	GAGTCGAAACCAGAAAGATGGTC	

*, F: forward; R: reverse

Histological staining and microscopy

Histochemical β -glucuronidase (GUS) staining of transgenic lines expressing GUS was performed as described previously (Anandalakshmi et al., 1998). Tissues were stained for 4 hours at 37°C, followed by rehydration by incubation for 10 minutes in a graded ethanol series (75, 50 and 25%). GUS stained tissues were observed and photographed using a LEICA MZ12 microscopy (Switzerland) equipped with a LEICA DC500 camera.

DAPI staining of nuclei was performed as described previously (Zanten et al., 2011). Samples were fixed in ice-cold Carnoy's fixative (1:3 acetic acid:ethanol), then treated with enzymatic cell wall degrading solution containing 0.5% cellulose Onozuka R10 (Duchefa), 0.25% macerozyme R10 (Duchefa), and 0.1% Triton X-100 for 1 h 30 min at room temperature. The samples were mounted with Vectashield (Vector laboratories) supplemented with 2 μ g/ml 4',6-diamidino-2-phenylindole (DAPI) before microscopic observation.

The heterochromatin phenotypes of the DAPI-stained leaf cells were recorded using a confocal laser scanning microscope (ZEISS-003-18533) using a 405 laser, a 350 nm LP excitation filter and a 425-475 nm BP emission filter. The H2B-GFP fusion protein was visualized using the same laser scanning microscope with a 534 laser, a 488 nm LP excitation filter and a 500-525 nm BP emission filters.

References

- Anandalakshmi, R., Pruss, G. J., Ge, X., Marathe, R., Mallory, A. C., Smith, T. H., and Vance, V.B. (1998). A viral suppressor of gene silencing in plants. *Proc. Natl. Acad. Sci.* **95**: 13079-13084.
- Anders, S. and Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biol.* **11**: R106.
- Anders, S., Pyl, P.T., and Huber, W. (2014). HTSeq – A Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 0–5.
- Andrés, F. and Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. *Nat. Rev. Genet.* **13**: 627–639.
- Andrés, F., Porri, A., Torti, S., Mateos, J., Romera-Branchat, M., García-Martínez, J.L., Fornara, F., Gregis, V., Kater, M.M., and Coupland, G. (2014). SHORT VEGETATIVE PHASE reduces gibberellin biosynthesis at the *Arabidopsis* shoot apex to regulate the floral transition. *Proc. Natl. Acad. Sci.* **111**: E2760-9.
- Bratzel, F. and Turck, F. (2015). Molecular memories in the regulation of seasonal flowering: from competence to cessation. *Genome Biol.* **16**: 192.
- Cardon, G., Höhmann, S., Klein, J., Nettekheim, K., Saedler, H., and Huijser, P. (1999). Molecular characterisation of the *Arabidopsis* *SBP-box* genes. *Gene* **237**: 91–104.
- Catez, F., Yang, H., Tracey, K.J., Reeves, R., Misteli, T., and Bustin, M. (2004). Network of dynamic interactions between histone H1 and high-mobility-group proteins in chromatin. *Mol. Cell. Biol.*: 4321–4328.
- Chandra, T. and Kirschner, K. (2016). Chromosome organisation during ageing and senescence. *Curr. Opin. Cell Biol.* **40**: 161–167.
- Chavali, P.L., Funa, K., and Chavali, S. (2011). Cis-regulation of microRNA expression by scaffold/matrix-attachment regions. *Nucleic Acids Res.* **39**: 6908–6918.
- Chen, W., Meaux, J. De, and Lercher, M.J. (2010). Co-expression of neighbouring genes in *Arabidopsis*: separating chromatin effects from direct interactions. *BMC Genomics* **11**: 2–8.
- Dekker, J., Marti-Renom, M. A., and Mirny, L.A. (2013). Exploring the three-dimensional organization of genomes: interpreting chromatin interaction data. *Nat. Rev. Genet.* **14**: 390–403.
- Du, Z., Zhou, X., Ling, Y., Zhang, Z., and Su, Z. (2010). agriGO: a GO analysis toolkit for the agricultural community. *Nucleic Acids Res.* **38**: 64–70.
- Espineda, C.E., Linford, A.S., Devine, D., and Brusslan, J.A. (1999). The *AtCAO* gene, encoding chlorophyll a oxygenase, is required for chlorophyll b synthesis in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* **96**: 10507–10511.
- Exner, V. and Hennig, L. (2008). Chromatin rearrangements in development. *Curr. Opin. Plant Biol.*: 64–69.
- Fang, Y. and Spector, D.L. (2005). Centromere positioning and dynamics in living *Arabidopsis* plants. *Genetics* **16**: 5710–5718.
- Fu, C., Sunkar, R., Zhou, C., Shen, H., Zhang, J.-Y., Matts, J., Wolf, J., Mann, D.G.J., Stewart, C.N., Tang, Y., and Wang, Z.-Y. (2012). Overexpression of *miR156* in switchgrass (*Panicum virgatum* L.) results in various morphological alterations and leads to improved biomass production. *Plant Biotechnol. J.* **10**: 443–452.
- Fujimoto, S., Matsunaga, S., Yonemura, M., Uchiyama, S., Azuma, T., and Fukui, K. (2004).

- Identification of a novel plant MAR DNA binding protein localized on chromosomal surfaces. *Plant Mol. Biol.* **56**: 225–239.
- Fusco, A. and Fedele, M.** (2007). Roles of HMGA proteins in cancer. *Nat. Rev. Cancer* **7**: 899–910.
- Girod, P., Nguyen, D., Calabrese, D., Puttini, S., Martinet, D., Regamey, A., Saugy, D., Beckmann, J.S., Bucher, P., and Mermoud, N.** (2007). Genome-wide prediction of matrix attachment regions that increase gene expression in mammalian cells. *Nat. Methods* **4**: 747–753.
- Gorbunova, V., and Seluanov, A.** (2016). DNA double strand break repair , aging and the chromatin connection. In *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis* (Elsevier B.V.), pp. 2–6.
- Grob, S., Schmid, M.W., and Grossniklaus, U.** (2014). Article Hi-C analysis in *Arabidopsis* identifies the KNOT , a atructure with similarities to the flamenco locus of drosophila. *Mol. Cell* **55**: 678–693.
- Harmston, N. and Lenhard, B.** (2013). Chromatin and epigenetic features of long-range gene regulation. *Nucleic Acids Res.* **41**: 7185–7199.
- He, Y. and Amasino, R.M.** (2005). Role of chromatin modification in flowering-time control. *Trends Plant Sci.* **10**: 30–35.
- Heng, H.H.Q., Goetze, S., Ye, C.J., Liu, G., Stevens, J.B., Bremer, S.W., Wykes, S.M., Bode, J., and Krawetz, S.** (2004). Chromatin loops are selectively anchored using scaffold/matrix-attachment regions. *J. Cell Sci.* **117**: 999–1008.
- Ho, L. and Crabtree, G.R.** (2010). Chromatin remodelling during development. *Nature* **463**: 474–484.
- Huang, S., Raman, A.S., Ream, J.E., Fujiwara, H., Cerny, R.E., and Brown, S.M.** (1998). Overexpression of *20-Oxidase* confers a gibberellin- overproduction phenotype in *Arabidopsis*. *Plant Physiol.* **118**: 773–781.
- Huijser, P. and Schmid, M.** (2011). The control of developmental phase transitions in plants. *Development* **138**: 4117–4129.
- Jarillo, J.A., Piñeiro, M., Cubas, P., and Martínez-zapater, J.** (2009). Chromatin remodeling in plant development. *Int. J. Dev. Biol.* **1596**: 1581–1596.
- Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L.** (2013). TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* **14**: R36.
- Kishi, Y., Fujii, Y., Hirabayashi, Y., and Gotoh, Y.** (2012). HMGA regulates the global chromatin state and neurogenic potential in neocortical precursor cells. *Nat. Neurosci.* **15**: 1127–1133.
- Kundu, S., Pendle, A., Irish, V.F., Shaw, P., Reimeg, J., Emanuelsson, O., Nakayama, N., and Sundstr, J.F.** (2017). Genome-wide identification of physically clustered genes suggests chromatin-level co-regulation in male reproductive development in *Arabidopsis thaliana*. *Nucleic Acids Res.* **45**: 3253–3265.
- Langmead, B. and Salzberg, S.L.** (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**: 357–359.
- Li, G., and Reinberg, D.** (2011). Chromatin higher-order structures and gene regulation. *Curr Opin Genet Dev.* **21**: 49–64.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.** (2009). The Sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078–2089.
- Lim, P.O., Kim, Y., Breeze, E., Koo, J.C., Woo, H.R., Ryu, J.S., Park, D.H., Beynon, J., Tabrett, A.,**

- Buchanan-Wollaston, V., and Nam, H.G.** (2007). Overexpression of a chromatin architecture-controlling AT-hook protein extends leaf longevity and increases the post-harvest storage life of plants. *Plant J.* **52**: 1140–1153.
- Masson, J., and Paszkowski, J.** (1992). The culture response of *Arabidopsis thaliana* protoplasts is determined by the growth conditions of donor plants. *Plant J.* **2**: 829–833.
- Matsoukas, I.G., Massiah, A.J., and Thomas, B.** (2013). Starch metabolism and antiflorigenic signals modulate the juvenile-to-adult phase transition in *Arabidopsis*. *Plant. Cell Environ.* **36**: 1802–1811.
- Matsushita, A., Furumoto, T., Ishida, S., and Takahashi, Y.** (2007). AGF1, an AT-hook protein, is necessary for the negative feedback of *AtGA3ox1* encoding GA 3-oxidase. *Plant Physiol.* **143**: 1152–1162.
- Moraes, S.** (2014). Gerontology & geriatric changes in chromatin structure along aging. *Gerontol. Geriatr. Res.* **3**: 1000157.
- Morisawa, G., Han-Yama, A., Moda, I., Tamai, A., Iwabuchi, M., and Meshi, T.** (2000). *AHM1*, a novel type of nuclear matrix-localized, MAR binding protein with a single AT hook and a J domain-homologous region. *Plant Cell* **12**: 1903–1916.
- Mortazavi, A., Williams, B.A., Mccue, K., Schaeffer, L., and Wold, B.** (2008). Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* **5**: 1–8.
- Mutasa-Göttgens, E. and Hedden, P.** (2009). Gibberellin as a factor in floral regulatory networks. *J. Exp. Bot.* **60**: 1979–1989.
- Ng, K.-H., Yu, H., and Ito, T.** (2009). AGAMOUS controls *GIANT KILLER*, a multifunctional chromatin modifier in reproductive organ patterning and differentiation. *PLoS Biol.* **7**: e1000251.
- Park, J., Oh, D., Dassanayake, M., Nguyen, K.T., Ogas, J., and Choi, G.** (2017). Gibberellin signaling requires chromatin remodeler PICKLE to promote vegetative growth and phase transitions. *Plant Physiol.* **173**: 1463–1474.
- Pfaffl, M.W.** (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**: e45–e45.
- Poethig, R.S.** (2013). Vegetative phase change and shoot maturation in plants. *Curr. Top. Dev. Biol.* **105**: 125–152.
- Pombo, A. and Dillon, N.** (2015). Three-dimensional genome architecture: players and mechanisms. *Nat. Rev. Mol. Cell Biol.* **16**: 245–257.
- Postnikov, Y. V and Bustin, M.** (2016). Functional interplay between histone H1 and HMG proteins in chromatin. *Biochim. Biophys. Acta* **1859**: 462–467.
- Quintero-cadena, P. and Sternberg, P.W.** (2016). Enhancer sharing promotes neighborhoods of transcriptional regulation across eukaryotes. *G3* **6**: 4167–4174.
- Rieu, I., Ruiz-Rivero, O., Fernandez-Garcia, N., Griffiths, J., Powers, S.J., Gong, F., Linhartova, T., Eriksson, S., Nilsson, O., Thomas, S.G., Phillips, A.L., and Hedden, P.** (2008). The gibberellin biosynthetic genes *AtGA20ox1* and *AtGA20ox2* act, partially redundantly, to promote growth and development throughout the *Arabidopsis* life cycle. *Plant J.* **53**: 488–504.
- Robinson, M.D., McCarthy, D.J., and Smyth, G.K.** (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.
- Saeteurn, K. H.S., and huck, G.C.A.** (2007). The heterochronic maize mutant *Corngrass1* results from overexpression of a tandem microRNA. *Nat. Genet.* **39**: 544–549.
- Telfer, A., Bollman, K.M., and Poethig, R.S.** (1997). Phase change and the regulation of trichome

- p>
distribution in
- Arabidopsis thaliana*
- .
- Development*
- 124**
- : 645–654.
- Thimm, O., Bläsing, O., Gibon, Y., Nagel, A., Meyer, S., Krüger, P., Selbig, J., Müller, L. a., Rhee, S.Y., and Stitt, M.** (2004). Mapman: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* **37**: 914–939.
- Wada, M., Takahashi, H., Nakamura, K., Hirai, M.Y., Ohta, D., and Kanaya, S.** (2012). Prediction of operon-like gene clusters in the *Arabidopsis thaliana* genome based on co-expression analysis of neighboring genes. *Gene* **503**: 56–64.
- Wang, J.-W., Park, M.Y., Wang, L.-J., Koo, Y., Chen, X.-Y., Weigel, D., and Poethig, R.S.** (2011). miRNA control of vegetative phase change in trees. *PLoS Genet.* **7**: e1002012.
- Williams, E.J.B. and Bowles, D.J.** (2004). Coexpression of neighboring Genes in the genome of *Arabidopsis thaliana*. *Genome Res.* **14**: 1060–1067.
- Wilson, R.H.C. and Coverley, D.** (2013). Relationship between DNA replication and the nuclear matrix. *Genes Cells* **18**: 17–31.
- Wu, G., Park, M.Y., Conway, S.R., Wang, J.-W., Weigel, D., and Poethig, R.S.** (2009). The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**: 750–759.
- Wu, G. and Poethig, R.S.** (2006). Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target *SPL3*. *Development* **133**: 3539–3547.
- Xie, K., Shen, J., Hou, X., Yao, J., Li, X., Xiao, J., and Xiong, L.** (2012). Gradual increase of miR156 regulates temporal expression changes of numerous genes during leaf development in rice. *Plant Physiol.* **158**: 1382–1394.
- Xie, K., Wu, C., and Xiong, L.** (2006). Genomic organization, differential expression, and interaction of *SQUAMOSA Promoter-Binding-Like* transcription factors and *microRNA156* in rice. *Plant Physiol.* **142**: 280–293.
- Xu, M., Hu, T., Zhao, J., Park, M.-Y., Earley, K.W., Wu, G., Yang, L., and Poethig, R.S.** (2016). Developmental functions of miR156-regulated *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE* (*SPL*) genes in *Arabidopsis thaliana*. *PLoS Genet.* **12**: e1006263.
- Xu, Y., Wang, Y., Stroud, H., Gu, X., Sun, B., Gan, E.-S., Ng, K.-H., Jacobsen, S.E., He, Y., and Ito, T.** (2013). A matrix protein silences transposons and repeats through interaction with retinoblastoma-associated proteins. *Curr. Biol.* **23**: 345–350.
- Yang, L., Conway, S.R., and Poethig, R.S.** (2011). Vegetative phase change is mediated by a leaf-derived signal that represses the transcription of *miR156*. *Development* **138**: 245–249.
- Yang, L., Xu, M., Koo, Y., He, J., and Poethig, R.S.** (2013). Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of *MIR156A* and *MIR156C*. *Elife* **2**: e00260.
- Yeaman, S.** (2013). Genomic rearrangements and the evolution of clusters of locally adaptive loci. *Proc. Natl. Acad. Sci.*
- Yu, S., Cao, L., Zhou, C.-M., Zhang, T.-Q., Lian, H., Sun, Y., Wu, J., Huang, J., Wang, G., and Wang, J.-W.** (2013). Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *Elife* **2**: e00269.
- Yun, J., Kim, Y.-S., Jung, J.-H., Seo, P.J., and Park, C.-M.** (2012). The AT-hook motif-containing protein AHL22 regulates flowering initiation by modifying *FLOWERING LOCUS T* chromatin in *Arabidopsis*. *J. Biol. Chem.* **287**: 15307–16.
- Zanten, M. Van, Koini, M.A., Geyer, R., Liu, Y., Brambilla, V., and Bartels, D.** (2011). Seed maturation in *Arabidopsis thaliana* is characterized by nuclear size reduction and increased chromatin condensation. *Proc. Natl. Acad. Sci.* **108**: 1–6.

- Zhan, S., Horrocks, J., and Lukens, L.N.** (2006). Islands of co-expressed neighbouring genes in *Arabidopsis thaliana* suggest higher-order chromosome domains. *Plant J.* **45**: 347–357.
- Zhang, X., Zou, Z., Zhang, J., Zhang, Y., Han, Q., Hu, T., Xu, X., Liu, H., Li, H., and Ye, Z.** (2011). Over-expression of *sly-miR156a* in tomato results in multiple vegetative and reproductive trait alterations and partial phenocopy of the *sft* mutant. *FEBS Lett.* **585**: 435–439.
- Zhao, J., Favero, D.S., Peng, H., and Neff, M.M.** (2013). *Arabidopsis thaliana* AHL family modulates hypocotyl growth redundantly by interacting with each other via the PPC/DUF296 domain. *Proc. Natl. Acad. Sci.* **110**: 4688–4697.

