



Universiteit
Leiden
The Netherlands

Obesity and type 2 diabetes : cardiovascular and cerebral aspects

Widya, R.L.; Widya R.L.

Citation

Widya, R. L. (2017, September 5). *Obesity and type 2 diabetes : cardiovascular and cerebral aspects*. Retrieved from <https://hdl.handle.net/1887/54684>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/54684>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/54684> holds various files of this Leiden University dissertation

Author: Widya, Ralph Lennart

Title: Obesity and type 2 diabetes : cardiovascular and cerebral aspects

Issue Date: 2017-09-05

Chapter 8

Increased amygdalar and hippocampal volumes in elderly obese individuals with or at risk of cardiovascular disease

Ralph L. Widya
Albert de Roos
Stella Trompet
Anton J.M. de Craen
Rudi G.J. Westendorp
Johannes W.A. Smit
Mark A. van Buchem
Jeroen van der Grond

for the PROSPER Study Group

American Journal of Clinical Nutrition 2011;93(6):1190-1195

ABSTRACT

Background

The basal ganglia, hippocampus, and thalamus are involved in the regulation of human feeding behavior. Recent studies have shown that obesity [body mass index (BMI; in kg/m^2) > 30] is associated with loss of gray and white matter. It is unknown whether the subcortical brain structures that are actually involved in feeding behavior also show volume changes in obesity. Therefore, the purpose of this study was to evaluate the volumes of the basal ganglia, hippocampus, and thalamus in obesity.

Methods

Three-dimensional T1-weighted magnetic resonance imaging scans of the brain were analyzed by using automatic segmentation to measure volumes of the nucleus accumbens, globus pallidus, amygdala, putamen, caudate nucleus, thalamus, and hippocampus in 471 subjects (mean age: 74.4 y; 56% men).

Results

Obese subjects had larger left ($P = 0.013$) and right ($P = 0.003$) amygdalar volumes and a larger left hippocampal volume ($P = 0.040$) than did normal-weight subjects (BMI < 25). None of the other subcortical structures differed in size between these groups. After correction for age, sex, smoking, hypertension, and pravastatin use, BMI was associated with left ($\beta = 0.175$, $P = 0.001$) and right ($\beta = 0.157$, $P = 0.001$) amygdalar volumes and with left hippocampal volume ($\beta = 0.121$, $P = 0.016$).

Conclusions

This study showed that the amygdala and hippocampus are enlarged in obesity. In consideration of the function of these structures, this finding may indicate that hedonic memories could be of major importance in the regulation of feeding. Because of the cross-sectional design, cause and effect could not be discriminated in this study.

INTRODUCTION

Obesity is a major public health issue strongly associated with chronic diseases, including diabetes mellitus and cardiovascular disease^{1,2}. It is well recognized that many brain structures are functionally involved in the regulation of food intake, including internal and external sensory inputs³. Regulation of food intake is controlled by multiple cognitive factors, including memorial representations of foods and their environmental context, and emotional and rewarding properties of such representations and their hedonic effects³. These factors are regulated by signaling molecules through the corticolimbic brain systems, in which the basal ganglia have an important role⁴.

In addition to homeostatic regulation of food intake driven by inputs of the basal ganglia, thalamus, and hippocampus, generic morphometric changes of the brain have been described as well. There is growing evidence that an increased body mass index (BMI) is independently associated with whole-brain atrophy^{5,6}. Other studies have shown an evident correlation between obesity and loss of cortical gray matter⁷⁻¹⁰. On the other hand, cortical gray and white matter volume loss has also been described in patients with a very low BMI, as in anorexia nervosa¹¹⁻¹⁵. Mechanisms behind these relations have not yet been clarified.

To our knowledge, the relation between obesity and the anatomy of subcortical structures that play a role in human food regulation have not been investigated previously. In the present study we aimed to investigate whether the basal ganglia, hippocampus, and thalamus are different between obese subjects and overweight and normal-weight subjects.

METHODS

Subjects

All subjects were derived from the magnetic resonance imaging (MRI) substudy of the PROspective Study of Pravastatin in the Elderly at Risk (PROSPER). PROSPER was a double-blind, randomized, placebo-controlled trial aimed at assessing the effect of therapy with 40 mg pravastatin on vascular events in 5804 men and women from Scotland, Ireland, and the Netherlands aged 70-82 y with vascular disease or at risk of vascular disease¹⁶. The PROSPER study was initiated in 1997. In the Netherlands, all consenting subjects were enrolled beginning in May 1998. The inclusion criteria for this study were as follows: men or women aged 70-82 y; total cholesterol of 4.0-9.0 mmol/L; stroke, transient ischemic attack, myocardial infarction, arterial surgery, or amputation for vascular disease > 6 mo before study entry; or one or more of the following risk factors for vascular disease: current smoking, hypertension, current drug treatment, known diabetes mellitus, or fasting blood glucose > 7 mmol/L. The institutional ethics review boards of all centers approved the protocol, and all participants gave written informed consent. The protocol was consistent with the Declaration of Hel-



sinki. All subjects had a Mini-Mental State Examination score of ≥ 24 . Effects of dementia and/or Alzheimer disease on the volumes of the basal ganglia were hereby automatically eliminated^{16,17}. Weight and height data for all participants were collected. The results were compared between subjects on the basis of BMI (in kg/m^2) groups: normal-weight subjects ($\text{BMI} < 25$), overweight subjects ($\text{BMI} = 25\text{-}30$), and obese subjects ($\text{BMI} > 30$).

Magnetic resonance acquisition

Of the 1100 Dutch participants in PROSPER, 494 subjects underwent a high-resolution three-dimensional (3D) T1-weighted (T1-w) MRI. All imaging was performed on an MR system operating at a field strength of 1.5 Tesla (Philips Medical Systems, Best, Netherlands). Three-dimensional T1-w scans were obtained by using the following parameters: repetition time = 30 ms, echo time = 4.6 ms, flip angle = 30° , slice thickness = 1.5 mm, 120 slices, no interslice gap, field of view = 220×220 mm, matrix = 256×256 , and in-plane resolution = 1.12×1.12 mm.

MRI processing

Segmentation of the various subcortical structures was performed on the original data set ($n = 494$) and on the same data set of which the left and right hemispheres were mirrored in the lateral plane. This approach was chosen to overcome potential systematic lateralization bias in preferential volume determinations by the algorithm FIRST (FMRIB's Integrated Registration and Segmentation Tool). To calculate the individual left and right volumes of the separate basal ganglia, the original volume and the corresponding mirrored volume of that particular intracranial structure were averaged.

FIRST was applied to estimate the volumes of left and right structures separately in 7 subcortical regions: nucleus accumbens, globus pallidus, amygdala, putamen, caudate nucleus, thalamus, and hippocampus. FIRST is part of FSL (FMRIB's Software Library) and performs both registration and segmentation of the mentioned subcortical regions^{18,19}. During registration, the input data (3D T1 images) are transformed to Montreal Neurological Institute (MNI) 152 standard space. For intrasubject comparison studies, the algorithm uses 12 df to transform the image (i.e., 3 translations, 3 rotations, 3 scalings, and 3 skews). After registration, a subcortical mask was applied to locate the different subcortical structures, followed by segmentation based on voxel intensities. Absolute volumes of subcortical structures were calculated, taking into account the transformations made in the first stage. The software was set to archive all single-structure segmentations. Border correction was used to increase the accuracy of the volumetric measurements²⁰. Border-corrected segmentations were archived as well. Volumetric data were obtained from the archived segmentations and normalized to MNI space. Incorrect segmentation of individual structures (volume greater or less than the mean group volume $\pm 3 \times \text{SD}$, usually with the value of "0") was excluded for analysis based on Tukey's rule²¹.

The single time point measurement technique SIENAX was performed to obtain estimates of gray and white matter volumes as well as brain volume. SIENAX starts by extracting brain and skull images from the single whole-head input data. The brain image is then affine-registered to MNI 152 space (by using the skull image to determine the registration scaling), done primarily to obtain the volumetric scaling factor to be used as normalization for head size. Next, tissue-type segmentation with partial volume estimation is carried out to calculate total volume of brain tissue (including separate estimates of volumes of gray matter, white matter, peripheral gray matter, and ventricular cerebrospinal fluid)¹⁹. Atrophy was defined as intracranial volume: parenchymal volume / intracranial volume \times 100%. All volumetric data were normalized to standard space, thereby correcting for interindividual differences in brain size. All MRI scans were reviewed by M.A.v.B. (> 15 y of neuroradiologic experience) for incidental findings.

Statistical analysis

Statistical analysis was performed by using SPSS for Windows (version 16.0.2; SPSS, Chicago, Ill). All descriptive data were expressed as means \pm SDs or *n* (%). To compare group data, a one-factor analysis of variance was used, followed by a Scheffe's post hoc test to correct for multiple comparisons. For categorical variables, the chi-square test or Fisher's exact test was used. The association between basal ganglia volumes and BMI was determined by using a multiple linear regression analysis, with age, sex, smoking, hypertension, and pravastatin use as covariates. β regression coefficients and *P* values are reported. *P* values < 0.05 were considered significant.

RESULTS

Three-dimensional T1-w MRI of the brain was performed in 494 subjects. Except for age-related changes, none of our subjects had unexpected findings on MRI. After exclusion of cases with poor segmentation, 471 subjects (mean age: 74.4 y; 56% men) remained for volume calculation. The BMI of our sample population was 26.8 ± 3.6 , with a minimum of 15.8 and a maximum of 43.5. The distribution of all BMIs is shown in **Figure 1**. On the basis of BMI, 140 subjects were classified as normal-weight (BMI = 23.2 ± 1.6), 256 subjects as overweight (BMI = 27.0 ± 1.4), and 75 subjects as obese (BMI = 32.8 ± 2.9). Demographic profiles for each group are shown in **Table 1**.




Figure 1. BMI distribution for all subjects.



Table 1. Demographics and group characteristics

Variable	Normal-weight	Overweight	Obese	P value		
	(n = 140)	(n = 256)	(n = 75)	Normal-weight vs. overweight	Overweight vs. obese	Normal-weight vs. obese
Male sex [n (%)] [†]	82 (59)	152 (59)	29 (39)	0.961	0.002*	0.008*
Age (y)	74.8 ± 3.3	74.3 ± 3.1	73.8 ± 3.0	0.277	0.469	0.071
Weight (kg)	67.7 ± 8.4	78.7 ± 8.2	92.6 ± 11.5	0.000*	0.000*	0.000*
Height (cm)	170.7 ± 7.9	170.5 ± 8.4	167.9 ± 9.3	0.978	0.062	0.068
BMI (kg/m ²)	23.2 ± 1.6	27.0 ± 1.4	32.8 ± 2.9	0.000*	0.000*	0.000*
MMSE score (points)	28.4 ± 1.3	28.4 ± 1.4	28.1 ± 1.6	0.993	0.288	0.397
History of vascular disease [n (%)] [†]	60 (43)	109 (43)	29 (39)	0.958	0.637	0.653
History of MI [n (%)] [†]	7 (5)	34 (13)	13 (17)	0.083	0.486	0.079
History of stroke/TIA [n (%)] [†]	30 (21)	30 (12)	13 (17)	0.121	0.281	0.592
Current smokers [n (%)] [†]	37 (26)	47 (18)	10 (13)	0.080	0.401	0.042*
Hypertension [n (%)] [†]	77 (55)	159 (62)	58 (77)	0.203	0.021*	0.002*
DM [n (%)] [†]	14 (10)	50 (20)	17 (23)	0.047*	0.666	0.020*
Insulin use [n (%)] [‡]	1 (1)	6 (2)	3 (4)	0.429	0.428	0.123
Fasting glucose (mmol/L)	5.4 ± 1.1	5.9 ± 1.7	6.0 ± 1.8	0.003*	0.883	0.013*
Systolic BP (mmHg)	155.6 ± 21.9	158.8 ± 21.9	158.2 ± 18.2	0.375	0.982	0.694
Diastolic BP (mmHg)	84.7 ± 10.6	86.3 ± 10.7	87.7 ± 10.9	0.327	0.634	0.143
Total cholesterol (mmol/L)	5.8 ± 0.9	5.7 ± 0.8	5.7 ± 0.8	0.832	0.968	0.793
LDL cholesterol (mmol/L)	3.9 ± 0.7	3.9 ± 0.7	3.9 ± 0.7	0.880	0.997	0.957
HDL cholesterol (mmol/L)	1.3 ± 0.4	1.2 ± 0.3	1.2 ± 0.3	0.000*	0.509	0.000*
Triglycerides (mmol/L)	1.3 ± 0.5	1.6 ± 0.7	1.6 ± 0.6	0.000*	0.845	0.003*

MMSE = Mini-Mental State Examination, MI = myocardial infarction, TIA = transient ischemic attack, DM = diabetes mellitus, BP = blood pressure. Data were compared by one-factor ANOVA with Scheffé's post hoc analysis.

* $P < 0.05$.

[†] Chi-square test.

[‡] Fisher's exact test.

Obese subjects had larger left ($P = 0.013$) and right ($P = 0.003$) amygdalar volumes and a larger left hippocampal volume ($P = 0.040$) than did normal-weight subjects. None of the other basal ganglia or the thalamus differed significantly in size between groups (**Table 2**).

Multiple linear regression analysis examined the association between BMI and the basal ganglia, thalamus, and hippocampus, the left and right sides of which were assessed separately. After correction for age, sex, smoking, hypertension, and pravastatin use, a significant association was observed between BMI and normalized left ($\beta = 0.175$, $P = 0.001$) and right ($\beta = 0.157$, $P = 0.001$) amygdalar volumes, and normalized left hippocampal volume ($\beta = 0.121$, $P = 0.016$) (**Figure 2**). This indicates that a high BMI is associated with large volumes

Table 2. Magnetic resonance imaging characteristics of normal-weight, overweight, and obese subjects

	Normal-weight		Overweight		Obese		P value		
	n	Volume cm ³	n	Volume cm ³	n	Volume cm ³	Normal-weight vs. overweight	Overweight vs. obese	Normal-weight vs. obese
L nuc acc	86	0.58 ± 0.16	156	0.60 ± 0.16	45	0.61 ± 0.18	0.728	0.879	0.577
R nuc acc	68	0.54 ± 0.16	164	0.54 ± 0.16	38	0.51 ± 0.16	1.000	0.779	0.814
L glob pal	117	1.96 ± 0.46	214	1.95 ± 0.40	65	1.97 ± 0.43	0.978	0.939	0.986
R glob pal	119	1.97 ± 0.45	213	1.95 ± 0.34	64	1.99 ± 0.41	0.864	0.733	0.948
L amygy	126	1.95 ± 0.36	226	2.02 ± 0.33	66	2.10 ± 0.33	0.144	0.253	0.013*
R amygy	130	1.93 ± 0.32	224	2.01 ± 0.33	68	2.10 ± 0.35	0.151	0.097	0.003*
L putam	120	5.16 ± 0.61	226	5.18 ± 0.61	69	5.17 ± 0.75	0.952	0.996	0.988
R putam	123	5.34 ± 0.61	220	5.33 ± 0.61	67	5.18 ± 0.61	1.000	0.184	0.235
L caud nuc	126	3.57 ± 0.50	222	3.58 ± 0.51	69	3.51 ± 0.53	0.966	0.611	0.772
R caud nuc	127	4.05 ± 0.57	225	3.98 ± 0.63	68	3.84 ± 0.63	0.560	0.277	0.077
L thal	129	8.01 ± 0.65	224	8.09 ± 0.64	69	8.09 ± 0.74	0.481	0.998	0.707
R thal	128	8.04 ± 0.66	224	8.13 ± 0.68	69	8.13 ± 0.74	0.506	0.998	0.656
L hipp	124	4.52 ± 0.68	223	4.63 ± 0.58	64	4.77 ± 0.70	0.309	0.301	0.040*
R hipp	124	4.60 ± 0.63	219	4.66 ± 0.59	64	4.77 ± 0.57	0.600	0.476	0.177
Gray matter	143	5.93 ± 0.50 x 10 ²	252	5.91 ± 0.41 x 10 ²	76	5.93 ± 0.44 x 10 ²	0.953	0.923	0.990
White matter	143	7.71 ± 0.41 x 10 ²	252	7.66 ± 0.36 x 10 ²	76	7.66 ± 0.41 x 10 ²	0.495	0.999	0.665
Gray + white	143	13.64 ± 0.71 x 10 ²	252	13.57 ± 0.62 x 10 ²	76	13.60 ± 0.69 x 10 ²	0.667	0.970	0.911

Values are means ± SDs unless otherwise indicated. L = left, R = right, nuc acc = nucleus accumbens, glob pal = globus pallidus, amygy = amygdala, putam = putamen, caud nuc = caudate nucleus, thal = thalamus, hipp = hippocampus, Gray matter = volume of normalized gray matter, White matter = volume of normalized white matter, Gray + white = total volume of white and gray matter. Data were compared by one-factor ANOVA with Scheffe's post hoc analysis.

* $P < 0.05$.



of amygdala and left hippocampus. BMI showed no association with other basal ganglia or thalamus volumes (**Table 3**).

The subjects included in this study might have had one or more risk factors for vascular disease, namely current smoking, hypertension, current drug treatment, known diabetes mellitus, or fasting blood glucose > 7 mmol/L. We found no association between volumes of basal ganglia and any of these cardiovascular disease risk factors.




Figure 2. Bivariate relations between amygdalar and hippocampal volumes and BMI. Reported β and P values were obtained by multiple linear regression analysis with correction for age, sex, smoking, hypertension, and pravastatin use.

DISCUSSION

The main findings of this study were enlargements of the amygdala and hippocampus in the obese subjects. In addition, an association of amygdalar and hippocampal enlargement with BMI corrected for age, sex, smoking, hypertension, and pravastatin use was observed. Both brain structures play a crucial role in the process of feeding behavior. Interestingly, none of the other investigated brain structures showed a significant association with BMI.

Table 3. Association between BMI and volumes of subcortical structures

	Crude		Adjusted	
	β	<i>P</i> value	β	<i>P</i> value
L nuc acc	0.036	0.543	0.044	0.455
R nuc acc	-0.012	0.843	0.002	0.968
L glob pal	0.023	0.650	0.021	0.683
R glob pal	0.021	0.679	0.020	0.707
L amyg	0.164	0.001*	0.175	0.001*
R amyg	0.151	0.002*	0.157	0.001*
L putam	-0.011	0.818	0.020	0.667
R putam	-0.068	0.171	-0.050	0.300
L caud nuc	-0.050	0.308	-0.027	0.577
R caud nuc	-0.079	0.109	-0.045	0.353
L thal	0.022	0.658	0.056	0.229
R thal	0.037	0.446	0.064	0.180
L hipp	0.118	0.016*	0.121	0.016*
R hipp	0.063	0.202	0.069	0.176
Gray matter	0.005	0.921	-0.054	0.202
White matter	-0.060	0.190	-0.057	0.226
Gray + white	-0.032	0.485	-0.070	0.119

Results are from a multiple linear regression analysis. L = left, R = right, nuc acc = nucleus accumbens, glob pal = globus pallidus, amyg = amygdala, putam = putamen, caud nuc = caudate nucleus, thal = thalamus, hipp = hippocampus, Gray matter = volume of normalized gray matter, White matter = volume of normalized white matter, Gray + white = total volume of white and gray matter.

* Significant β values before and after adjustment for age, sex, smoking, hypertension, and pravastatin use.

The amygdala plays an important role in the coordination of appetitive behavior. It is part of a complex neural system that is responsible for the evaluation of food. The electrophysiological responses on taste and picture-odor contingencies in the amygdala suggest a sensory function^{22, 23}. Furthermore, rat studies suggest that the amygdala is critical to learning representations of specific experiences with food and using them to guide appetitive behavior²⁴. Enlargement of the amygdala with respect to BMI has not been described before. It is known that enlargement can occur in patients with depression^{25, 26}, psychosis²⁷, autism²⁸, and bipolar disorder^{29, 30}. A slight enlargement of the hippocampus has also been reported in bipolar disorder³⁰.

Although we found that volume changes were more pronounced in the right hemisphere of the amygdala and in the left hemisphere of the hippocampus, we believe that these differences are statistically not convincing enough to speculate on lateralization.

A possible explanation for amygdalar enlargement is higher glucose metabolism and cerebral blood flow³¹. However, a direct relation with amygdalar enlargement remains unclear, because cerebral blood flow and metabolism might be affected by multiple physi-



ologic events, e.g., dynamic changes in neurotransmitter-neuroreceptor function²⁶. Other proposed mechanisms of amygdalar enlargement are an increase in size or number of neurons or glial cells, increased connective tissue, and increased intercellular fluid³¹. Rat studies have shown that chronic stress induces dendritic remodeling of the hippocampus and amygdala, which leads to atrophy and hypertrophy, respectively³². Hypermetabolism of the amygdala may also result from an increased afferent glutamatergic transmission between and within amygdala nuclei³³. Glutamatergic transmission is part of the endocannabinoid system, which is related to obesity when dysregulated³⁴. On the basis of this principle, pharmacologic agents, particularly cannabinoid receptor antagonists, are currently used as antiobesity drugs³⁵.

The amygdala and hippocampus relate memory to pleasure, respectively, by establishing the reward value of food and food-related stimuli or situations and by consolidating pleasure memories. On the basis of the previously described mechanisms, we assume that our findings are in part explained by an increased activity and possibly an elevated metabolism in the amygdala and hippocampus. Statements about the order of events (excessive feeding leading to or resulting from enlargement) are limited because this study was cross-sectional. However, it is just as likely that the described changes are the effect of enjoying food on hedonic centers, rather than hedonic center changes leading to eating more food.

Studies have shown that smoking history is associated with brain atrophy³⁶⁻³⁹. Therefore, we adjusted for smoking in the linear regression analysis. Hypertension has also been recognized as a risk factor for brain atrophy^{40, 41}, and moreover, hippocampal atrophy⁴². Hence, we used hypertension as a covariate in the linear regression analysis.

We realize that in an explorative study, significant results may occur by coincidence. In the current study, we performed a one-factor analysis of variance with subsequent Scheffé's post hoc analysis, correcting for multiple comparisons between groups (Table 2). We did not correct a priori for the number of structures (Tables 2 and 3). Because the amygdala is associated with BMI (Tables 2 and 3) in both hemispheres, we believe that our data are not driven by coincidence. Still, it must be emphasized that significant findings that are just below the threshold value ($P = 0.05$) should, in general, be treated with care.

No association was found between volumes of basal ganglia and cardiovascular disease risk factors. Because only cross-sectional data were available, no statements can be made about the influence of the duration of cardiovascular disease risk factors. For the same reason, cause and effect cannot be distinguished.

The obese group consisted of more women than men. Estrogen use has been shown to increase hippocampal volume relative to estrogen nonuse and findings in males⁴³. Estrogen therapy was not investigated in this study. Therefore, sex or estrogen therapy could have been confounders in the relation between BMI and hippocampal volume.

We used the algorithm FIRST to estimate volumes. Existing and upcoming evidence indicate that different segmentation tools show structural differences between them. Com-

parison of data assessed with different segmentation techniques is therefore discouraged. A final limitation of this study was that it was performed in elderly individuals. Therefore, our results cannot be generalized to the population aged < 70 y.

In conclusion, this explorative study showed that the amygdala and hippocampus, which are very important structures in regulating feeding behavior, are enlarged in obesity. Furthermore, an association between BMI and enlargement was seen. The finding that these anatomic changes only occurred in the amygdala and hippocampus, and not in other brain structures that play a role in feeding behavior, suggests that cognitive aspects may be a major issue in the regulation of feeding. To our knowledge, this is the first study to have explored the relation between obesity and the basal ganglia specifically. Additional studies are required to examine the mechanisms behind enlargement of the amygdala and hippocampus in obesity.



REFERENCES

1. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 2010;303:235-241.
2. Flegal KM, Graubard BI, Williamson DF, Gail MH. Sources of differences in estimates of obesity-associated deaths from first National Health and Nutrition Examination Survey (NHANES I) hazard ratios. *Am J Clin Nutr* 2010;91:519-527.
3. Berthoud HR. Interactions between the "cognitive" and "metabolic" brain in the control of food intake. *Physiol Behav* 2007;91:486-498.
4. Adan RA, Vanderschuren LJ, la Fleur SE. Anti-obesity drugs and neural circuits of feeding. *Trends Pharmacol Sci* 2008;29:208-217.
5. Ward MA, Carlsson CM, Trivedi MA, Sager MA, Johnson SC. The effect of body mass index on global brain volume in middle-aged adults: a cross sectional study. *BMC Neurol* 2005;5:23.
6. Gunstad J, Paul RH, Cohen RA, Tate DF, Spitznagel MB, Grieve S, Gordon E. Relationship between body mass index and brain volume in healthy adults. *Int J Neurosci* 2008;118:1582-1593.
7. Pannacciulli N, Del Parigi A, Chen K, Le DS, Reiman EM, Tataranni PA. Brain abnormalities in human obesity: a voxel-based morphometric study. *Neuroimage* 2006;31:1419-1425.
8. Taki Y, Kinomura S, Sato K, Inoue K, Goto R, Okada K, Uchida S, Kawashima R, Fukuda H. Relationship between body mass index and gray matter volume in 1,428 healthy individuals. *Obesity (Silver Spring)* 2008;16:119-124.
9. Walther K, Birdsill AC, Glisky EL, Ryan L. Structural brain differences and cognitive functioning related to body mass index in older females. *Hum Brain Mapp* 2010;31:1052-1064.
10. Raji CA, Ho AJ, Parikshak NN, Becker JT, Lopez OL, Kuller LH, Hua X, Leow AD, Toga AW, Thompson PM. Brain structure and obesity. *Hum Brain Mapp* 2010;31:353-364.
11. Katzman DK, Lambe EK, Mikulis DJ, Ridgley JN, Goldbloom DS, Zipursky RB. Cerebral gray matter and white matter volume deficits in adolescent girls with anorexia nervosa. *J Pediatr* 1996;129:794-803.
12. Frank GK, Bailer UF, Henry S, Wagner A, Kaye WH. Neuroimaging studies in eating disorders. *CNS Spectr* 2004;9:539-548.
13. Wagner A, Greer P, Bailer UF, Frank GK, Henry SE, Putnam K, Meltzer CC, Ziolkowski SK, Hoge J, McConaha C, Kaye WH. Normal brain tissue volumes after long-term recovery in anorexia and bulimia nervosa. *Biol Psychiatry* 2006;59:291-293.
14. Muhlau M, Gaser C, Ilg R, Conrad B, Leibl C, Cebulla MH, Backmund H, Gerlinghoff M, Lommer P, Schnebel A, Wohlschlagel AM, Zimmer C, Nunnemann S. Gray matter decrease of the anterior cingulate cortex in anorexia nervosa. *Am J Psychiatry* 2007;164:1850-1857.
15. Castro-Fornieles J, Bargallo N, Lazaro L, Andres S, Falcon C, Plana MT, Junque C. A cross-sectional and follow-up voxel-based morphometric MRI study in adolescent anorexia nervosa. *J Psychiatr Res* 2009;43:331-340.
16. Shepherd J, Blauw GJ, Murphy MB, Bollen EL, Buckley BM, Cobbe SM, Ford I, Gaw A, Hyland M, Jukema JW, Kamper AM, Macfarlane PW, Meinders AE, Norrie J, Packard CJ, Perry IJ, Stott DJ, Sweeney BJ, Twomey C, Westendorp RG, on behalf of the PROSPER study group. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet* 2002;360:1623-1630.
17. Folstein MF, Folstein SE, McHugh PR. "Mental state": A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-198.
18. de Jong LW, van der Hiele K, Veer IM, Houwing JJ, Westendorp RG, Bollen EL, de Bruin PW, Middelkoop HA, van Buchem MA, van der Grond J. Strongly reduced volumes of putamen and thalamus in Alzheimer's disease: an MRI study. *Brain* 2008;131:3277-3285.
19. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H,

- Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 2004;23(Suppl 1):S208-219.
20. Patenaude B. Bayesian shape and appearance models. Technical report TR07BP1. FMRIB Center, University of Oxford. 2007. Published 2007.
 21. Tukey JW. Exploratory data analysis. Reading, MA: Addison-Wesley Publishing Co., 1977.
 22. Rolls ET, Critchley HD, Browning AS, Hernadi I, Lenard L. Responses to the sensory properties of fat of neurons in the primate orbitofrontal cortex. *J Neurosci* 1999;19:1532-1540.
 23. Gottfried JA, O'Doherty J, Dolan RJ. Encoding predictive reward value in human amygdala and orbitofrontal cortex. *Science* 2003;301:1104-1107.
 24. Hatfield T, Han JS, Conley M, Gallagher M, Holland P. Neurotoxic lesions of basolateral, but not central, amygdala interfere with Pavlovian second-order conditioning and reinforcer devaluation effects. *J Neurosci* 1996;16:5256-5265.
 25. Tebartz van Elst L, Woermann F, Lemieux L, Trimble MR. Increased amygdala volumes in female and depressed humans. A quantitative magnetic resonance imaging study. *Neurosci Lett* 2000;281:103-106.
 26. Frodl T, Meisenzahl EM, Zetsche T, Born C, Jäger M, Groll C, Bottlender R, Leinsinger G, Möller HJ. Larger amygdala volumes in first depressive episode as compared to recurrent major depression and healthy control subjects. *Biol Psychiatry* 2003;53:338-344.
 27. Velakoulis D, Wood SJ, Wong MT, McGorry PD, Yung A, Phillips L, Smith D, Brewer W, Proffitt T, Desmond P, Pantelis C. Hippocampal and amygdala volumes according to psychosis stage and diagnosis: a magnetic resonance imaging study of chronic schizophrenia, first-episode psychosis, and ultra-high-risk individuals. *Arch Gen Psychiatry* 2006;63:139-149.
 28. Mosconi MW, Cody-Hazlett H, Poe MD, Gerig G, Gimpel-Smith R, Piven J. Longitudinal study of amygdala volume and joint attention in 2- to 4-year-old children with autism. *Arch Gen Psychiatry* 2009;66:509-516.
 29. Altshuler LL, Bartzokis G, Grieder T, Curran J, Mintz J. Amygdala enlargement in bipolar disorder and hippocampal reduction in schizophrenia: an MRI study demonstrating neuroanatomic specificity. *Arch Gen Psychiatry* 1998;55:663-664.
 30. Strakowski SM, DelBello MP, Sax KW, Zimmerman ME, Shear PK, Hawkins JM, Larson ER. Brain magnetic resonance imaging of structural abnormalities in bipolar disorder. *Arch Gen Psychiatry* 1999;56:254-260.
 31. Altshuler LL, Bartzokis G, Grieder T, Curran J, Jimenez T, Leight K, Wilkins J, Gerner R, Mintz J. An MRI study of temporal lobe structures in men with bipolar disorder or schizophrenia. *Biol Psychiatry* 2000;48:147-162.
 32. Vyas A, Mitra R, Shankaranarayana Rao BS, Chattarji S. Chronic stress induces contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. *J Neurosci* 2002;22:6810-6818.
 33. Drevets WC, Price JL, Bardgett ME, Reich T, Todd RD, Raichle ME. Glucose metabolism in the amygdala in depression: relationship to diagnostic subtype and plasma cortisol levels. *Pharmacol Biochem Behav* 2002;71:431-447.
 34. Mouslech Z, Valla V. Endocannabinoid system: An overview of its potential in current medical practice. *Neuro Endocrinol Lett* 2009;30:153-179.
 35. Rodriguez de Fonseca F, Del Arco I, Bermudez-Silva FJ, Bilbao A, Cippitelli A, Navarro M. The endocannabinoid system: physiology and pharmacology. *Alcohol Alcohol* 2005;40:2-14.
 36. Swan GE, Lessov-Schlaggar CN. The effects of tobacco smoke and nicotine on cognition and the brain. *Neuropsychol Rev* 2007;17:259-273.
 37. Swan GE, Decarli C, Miller BL, Reed T, Wolf PA, Carmelli D. Biobehavioral characteristics of



- nondemented older adults with subclinical brain atrophy. *Neurology* 2000;54:2108-2114.
38. Longstreth WT, Jr., Diehr P, Manolio TA, Beauchamp NJ, Jungreis CA, Lefkowitz D, for the Cardiovascular Health Study Collaborative Research Group. Cluster analysis and patterns of findings on cranial magnetic resonance imaging of the elderly: the Cardiovascular Health Study. *Arch Neurol* 2001;58:635-640.
 39. Brody AL, Mandelkern MA, Jarvik ME, Lee GS, Smith EC, Huang JC, Bota RG, Bartzokis G, London ED. Differences between smokers and nonsmokers in regional gray matter volumes and densities. *Biol Psychiatry* 2004;55:77-84.
 40. Gianaros PJ, Greer PJ, Ryan CM, Jennings JR. Higher blood pressure predicts lower regional grey matter volume: Consequences on short-term information processing. *Neuroimage* 2006;31:754-765.
 41. Wiseman RM, Saxby BK, Burton EJ, Barber R, Ford GA, O'Brien JT. Hippocampal atrophy, whole brain volume, and white matter lesions in older hypertensive subjects. *Neurology* 2004;63:1892-1897.
 42. Korf ES, White LR, Scheltens P, Launer LJ. Midlife blood pressure and the risk of hippocampal atrophy: the Honolulu Asia Aging Study. *Hypertension* 2004;44:29-34.
 43. Lord C, Buss C, Lupien SJ, Pruessner JC. Hippocampal volumes are larger in postmenopausal women using estrogen therapy compared to past users, never users and men: a possible window of opportunity effect. *Neurobiol Aging* 2008;29:95-101.

