



Universiteit
Leiden
The Netherlands

Pharmacogenetics of sunitinib in metastatic renal cell carcinoma

Diekstra, M.H.M.; Diekstra M.H.M.

Citation

Diekstra, M. H. M. (2017, September 13). *Pharmacogenetics of sunitinib in metastatic renal cell carcinoma*. Retrieved from <https://hdl.handle.net/1887/55944>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/55944>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden




The handle <http://hdl.handle.net/1887/55944> holds various files of this Leiden University dissertation

Author: Diekstra, Meta

Title: Pharmacogenetics of sunitinib in metastatic renal cell carcinoma

Date: 2017-09-13



CHAPTER 1

General introduction

Renal cell carcinoma: epidemiology, pathology and risk factors

The most common type of kidney cancer (>90%) is renal cell carcinoma (RCC) which comprises about 2-3% of all malignancies in adults.^{1,2} RCC typically occurs in people aged between 60 and 70 years old and is more common in men than in women.¹⁻³ In 2013, worldwide 350,000 people were diagnosed with RCC with more than 140,000 deaths. In the Netherlands 2,327 new cases were diagnosed in that year.^{3,4} The age-adjusted incidence of RCC seems to have increased in the last couple of years with a higher incidence in developed countries compared to developing countries. The higher RCC incidence could be explained by enhanced use of imaging techniques such as ultrasonography, computed tomography (CT) and magnetic resonance imaging (MRI).^{2,3} In most cases tumors are small and non-metastatic at first presentation, but approximately 17% of RCCs are metastasized at first diagnosis. Preferred sites of metastasis are lung, liver, bone, brain, and only rarely adrenal glands or the contralateral kidney.¹⁻³

RCC is a heterogeneous tumor encompassing a wide variety of tumor types with clear cell RCC as the most common histological subtype ($\approx 75\%$). Less common are the non-clear cell types: papillary type I (5%), papillary type II (10%), chromophobes (7%), medullary, collecting duct, and translocation renal tumors (**Figure 1**).¹⁻³ Moreover, multiple subtypes can be distinguished within a tumor. Non-clear cell subtypes have a poor prognosis in contrast to clear cell RCCs that show improved survival outcomes.⁵

Risk factors for RCC are smoking, obesity, hypertension or hereditary causes.¹⁻³ A mutation in the von Hippel-Lindau (VHL) gene is the most frequent hereditary cause (>80% of the clear cell RCCs). This causes a dysfunction of the VHL protein which prevents VHL to form a complex with hypoxia inducible factor α (HIF- α) so that HIF- α cannot be destroyed by proteasome degradation. Consequently, HIF- α accumulates in the cell inducing the secretion of proangiogenic factors, including the vascular endothelial growth factor (VEGF), that allow tumors to grow easily through oxygen supply by blood vessel formation.^{1,3,6}

RCC treatment options in the era of targeted therapy

First and foremost, before any treatment is being considered and if RCC is discovered at an early stage, surgical removal is the standard of care with the aim to cure the disease, provided that the patient has a good performance status. Minimally invasive techniques such as nephron sparing surgery (partial nephrectomy), exposure to extreme temperatures (cryoablation or radiofrequency ablation) are used rather than radical nephrectomy (removal of the entire kidney) to prevent chronic kidney disease and other morbidity.¹⁻³ Elderly patients with a poor prognosis or comorbidity and diagnosed with (localized) RCC with a tumor size <40 mm, should be under active surveillance by regularly imaging. Especially in asymptomatic patients, this period of observation before starting treatment is essential. In about 20% of the cases, small renal tumors may turn

out to be benign; slowly growing and self-limiting tumors, and only in 1 to 2% of the cases metastatic disease was reported.¹⁻³

In case of metastases or if surgery is not deemed feasible, treatment options will be more palliative than curative. Radiotherapy can offer pain relief in bone or brain metastases and chemotherapy has little or no effect.¹⁻³ The current drug arsenal against metastatic renal cell carcinoma (mRCC) is large and continues to expand. For a long time cytokines (interleukin-2 and interferon-alpha) were the only available treatment options with limited efficacy and substantial side effects. Since 2005 the number of drugs against kidney cancer has been growing with the arrival of targeted therapies that act on small proteins present in signal transduction pathways in and around the tumor. Targeted therapies that act in the tumor cell and inhibit the mammalian target of rapamycin (mTOR) are the so-called mTOR inhibitors everolimus and temsirolimus and cause a decreased expression of HIF- α . Temsirolimus is used in patients with a poor prognosis but is less and less prescribed, while everolimus is indicated as second-line treatment after VEGF-targeted therapy for good or intermediate prognosis. An alternative treatment option is the monoclonal antibody bevacizumab used in combination with interferon-alpha, which functions in between endothelial cells and tumor cells targeting VEGF and preventing it from binding to its receptor. Especially the (multi)targeted tyrosine kinase inhibitors (TKIs) showed to be very successful and target receptors on the endothelial cell with an anti-angiogenic effect (**Figure 1**).

Sorafenib was the first TKI approved for the treatment of RCC.⁷ Sunitinib entered the market in 2006. In the pivotal phase III trial of sunitinib in mRCC, it was shown that sunitinib significantly improves progression-free survival (PFS) and objective response rate compared to the former standard choice of therapy interferon-alpha ($P<0.001$) and an almost significant improvement in overall survival (OS) was seen for sunitinib ($P=0.051$).⁸ Later, in 2009, pazopanib was approved for mRCC treatment and meanwhile many more TKIs were being investigated, such as axitinib.^{9,10} The targeted therapies as mentioned above have been proven effective in clear cell RCC. For non-clear cell RCC however, less is known about the efficacy of anti-VEGF therapies and mTOR-inhibitors.⁵

Sunitinib pharmacology

The work described in this thesis focuses on sunitinib. Sunitinib is administered orally, most often given as 50 mg in a 4 week on/2 week off schedule. Dose reduction can be applied in steps of 12.5 mg where a continuous regimen of 37.5 mg is also applied. The main pharmacodynamic targets of sunitinib are VEGF-receptors type 1, 2 and 3, platelet-derived growth factor receptors (PDGFRs) α and β , Fms-related tyrosine kinase 3 (FLT3), and cytokine receptor Kit (c-KIT).⁸ Sunitinib is mainly metabolized by cytochrome P450 (CYP)3A4 to its active metabolite SU12662, and inactive metabolites. Both sunitinib and




Figure 1: Overview of RCC risk factors, tumor types, and treatment options with a focus on sunitinib and pharmacogenetics of sunitinib in mRCC.

SU12662 embody the total amount of active drug in plasma.¹¹⁻¹³ Furthermore, CYP3A5, CYP1A1 and CYP1A2 could be involved in sunitinib metabolism. The pregnane X receptor (NR1I2), the constitutive androstane receptor (NR1I3) and P450 oxidoreductase (POR) can influence CYP3A4 expression or activity.^{14,15} Other important modulators of sunitinib pharmacokinetics (PK) are the efflux transporters ATP-binding cassette (ABCB1) and the breast cancer resistance protein (ABCG2).¹⁶ Patients show a high variability in PK of sunitinib and SU12662 and respond very differently to drugs with large differences in toxicity and efficacy.^{12,13,17-19} Most common side effects are fatigue, hand-foot syndrome (HFS), hypertension, mucositis, hair or skin discoloration and hypothyroidism. About 30% of RCC patients on sunitinib required a dose reduction for any of these adverse events.^{13,17-19}

Despite successful advances in RCC treatment since the advent of targeted therapy, we have not yet reached our goal. Interpatient variability in response is large and the great difficulty is that we cannot predict an individual's response to sunitinib, which is highly essential in order to give patients an optimally effective and safe treatment and to efficiently use the available drug arsenal without adjustments afterwards. For that reason, we need specific biomarkers that can predict the individual treatment response with regard to efficacy and toxicity of sunitinib.

Pharmacogenetics of sunitinib

It is plausible that genetic variability in genes involved in the PK or pharmacodynamics (PD) of a drug are important determinants of a person's response to the drug with respect to adverse events and efficacy. Pharmacogenetics thoroughly explores this hypothesis. Previous pharmacogenetic studies used the candidate gene approach to select small genetic variations, namely Single Nucleotide Polymorphisms (SNPs), located in genes thought to be related to PK and PD of sunitinib. PK related genetic variants in genes coding for CYP enzymes, drug efflux transporters or influencers of CYP3A activity have been identified to be associated with toxicity or efficacy of sunitinib in mRCC.^{17-19,20-21} SNPs in *CYP1A1*, *CYP3A5*, *ABCB1*, *ABCG2*, and *NR1I3* showed an association directly with toxicity or with dose reductions due to toxicity.^{17,18} Furthermore, SNPs in *CYP3A5*, *ABCB1*, *NR1I2* and *NR1I3* have been associated with PFS on sunitinib.¹⁹ Regarding PD, SNPs associated with sunitinib toxicity or efficacy were located in genes encoding receptors on the endothelial cells or endothelial nitric oxide synthase (eNOS). SNPs in *VEGF-A*, *VEGFR-1*, *VEGFR-2*, *VEGFR-3*, *FLT3*, and *eNOS* have been associated with adverse events (particularly hypertension), PFS or OS on sunitinib (**Figure 1**).^{19,20,21} Yet, no genetic biomarker has been implemented for clinical use and little is known about the mechanisms underlying possible SNP effects. Therefore, the objective of this thesis is to investigate whether pharmacogenetic testing of SNPs prior to treatment initiation may possibly predict the individual response to sunitinib in mRCC and therefore benefit the patient's wellbeing.

Aim and outline of thesis

The general aim of this thesis was to study the predictive ability of germline pharmacogenetic markers on toxicity and efficacy outcomes of sunitinib in mRCC patients. This research is part of a European collaborative project on "Targeted therapy in Renal cell cancer: GEnetic and Tumor related biomarkers for response and toxicity" (EuroTARGET) funded by the European Commission under the Seventh Framework Programme (FP7).²²

Chapter 2 systematically reviews studies on SNPs (pharmacogenetics) but also gene expression or epigenetic biomarkers (pharmacogenomics) as predictors of TKI treatment

outcome in mRCC containing studies from 2005 to 2015. The quality of included studies is discussed and the prospect for clinical implementation of study results or the need for follow-up studies.

Previously, SNPs have been associated with toxicity or efficacy outcomes on sunitinib, but it is not clear whether these SNPs are also causally related to PK of sunitinib.¹⁷⁻¹⁹ In **Chapter 3** it is studied whether previously associated SNPs located in genes involved in the metabolism or absorption of sunitinib influence the clearance of sunitinib and its active metabolite SU12662.

In **Chapter 4** a pooled dataset is formed by including patients from previous pharmacogenetics analyses on sunitinib performed in the United States, Spain and the Netherlands.^{17-19,20,21} The aim of this study is to investigate whether SNPs associated with toxicity and efficacy on sunitinib in earlier exploratory studies could be confirmed in a larger group of 333 mRCC patients. To this end, 22 significantly associated SNPs ($P < 0.05$) from previous exploratory studies are selected and retested in this study for association with sunitinib treatment outcomes.

At the American Society of Clinical Oncology (ASCO) annual meeting in 2012, Urun *et al.* reported that SNP rs4646437G>A in *CYP3A4* had a protective effect on sunitinib related toxicity as shown in a cohort of 159 mRCC patients.²³ In **Chapter 5** the relationship between rs4646437 in *CYP3A4* and sunitinib toxicity or efficacy is further investigated in a large cohort of mRCC patients.

A set of relatively new SNPs (reported from 2011 onwards) in *CYP3A4*, *NR1I2*, *POR*, *IL8*, *IL4-R*, *IL13*, *HIF1A* and *MET* were reported as potential prognostic or predictive markers for TKI treatment outcome.²⁴⁻³⁰ With an exploratory nature, **Chapter 6** evaluates whether these SNPs play a role in sunitinib toxicity or efficacy outcomes in a large cohort of sunitinib treated mRCC patients.

In **Chapter 7** SNP rs34231037 in *VEGFR-2* (*KDR*) is analyzed in a large series of clear cell mRCC patients treated with sunitinib for the possible effect on response and PFS. This study objective was based on a Genome Wide Association Study (GWAS) of Maitland *et al.* in which this SNP was identified as a possible predictor for variations of soluble VEGFR2 (sVEGFR2) in the blood and sVEGFR2 levels at baseline and after 4 weeks of pazopanib treatment.³¹

A better understanding of the relationships between sunitinib exposure, the pharmacological response, and the clinical outcomes is vital to find predictive biomarkers for treatment outcome. **Chapter 8** describes a nonlinear mixed-effects population PK/PD model that is built by means of blood sampling data of sunitinib treated patients diagnosed with either mRCC or colorectal carcinoma (CRC) and a group of 12 healthy volunteers.³² The aim of this study is to evaluate the predictive performance of this model including PK parameters, blood pressure, sVEGFR-2, sVEGFR-3, and SNP genotypes.

Subsequently, the applicability of the model to predict clinical outcomes on sunitinib treatment is explored.

Due to the rapid developments in genetic testing technologies, possibilities to test more than one SNP at a time become easily accessible. In **Chapter 9** about 1 million SNPs are tested for the association with sunitinib toxicity or efficacy in a GWAS that allows less biased hypothesis-free testing for associations as compared to the candidate gene approach used in the aforementioned chapters.

This thesis closes with a general discussion and future perspective on the topic in **Chapter 10** and a summary written in English and Dutch in **Chapter 11**.

REFERENCES

1. Rini BI, Campbell SC, Escudier B. Renal cell carcinoma. *Lancet*. 2009;373(9669):1119-1132.
2. Escudier B, Porta C, Schmidinger M, et al. Renal cell carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2016;27(Suppl 5):v58-v68.
3. Capitanio U, Montorsi F. Renal cancer. *Lancet*. 2016;387(10021):894-906.
4. Database with reliable objective data on the incidence, prevalence, survival and mortality of all cancer cases; The Dutch Cancer Registration (NKR), managed by the Dutch Integral Cancer Center (IKNL). URL: <http://www.cijfersoverkanker.nl/>.
5. Kroeger N, Xie W, Lee JL, et al. Metastatic non-clear cell renal cell carcinoma treated with targeted therapy agents: characterization of survival outcome and application of the International mRCC Database Consortium criteria. *Cancer*. 2013;119(16):2999-3006.
6. George DJ, Kaelin WG Jr. The von Hippel-Lindau protein, vascular endothelial growth factor, and kidney cancer. *N Engl J Med*. 2003;349(5):419-421.
7. Escudier B, Eisen T, Stadler WM, et al. Sorafenib in advanced clear cell renal-cell carcinoma. *N Engl J Med*. 2007;356(2):125-134.
8. Motzer RJ, Hutson TE, Tomczak P, et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2009;27(22):3584-3590.
9. Sternberg CN, Davis ID, Mardiak J, et al. Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial. *J Clin Oncol*. 2010;28(6):1061-1068.
10. Rini BI, Escudier B, Tomczak P, et al. Comparative effectiveness of axitinib versus sorafenib in advanced renal cell carcinoma (AXIS): a randomised phase 3 trial. *Lancet*. 2011;378(9807):1931-1939.
11. Faivre S, Delbaldo C, Vera K, et al. Safety, pharmacokinetic, and antitumor activity of SU11248, a novel oral multitarget tyrosine kinase inhibitor, in patients with cancer. *J Clin Oncol*. 2006;24(1):25-35.
12. Houk BE, Bello CL, Kang D, Amantea M. A population pharmacokinetic meta-analysis of sunitinib malate (SU11248) and its primary metabolite (SU12662) in healthy volunteers and oncology patients. *Clin Cancer Res*. 2009;15(7):2497-2506.
13. Houk BE, Bello CL, Poland B, Rosen LS, Demetri GD, Motzer RJ. Relationship between exposure to sunitinib and efficacy and tolerability endpoints in patients with cancer: results of a pharmacokinetic/pharmacodynamic meta-analysis. *Cancer Chemother Pharmacol*. 2010;66(2):357-371.
14. Lamba J, Lamba V, Strom S, Venkataramanan R, Schuetz E. Novel single nucleotide polymorphisms in the promoter and intron 1 of human pregnane X receptor/NR1I2 and their association with CYP3A4 expression. *Drug Metab Dispos*. 2008;36(1):169-181.

15. Elens L, Nieuweboer AJ, Clarke SJ, et al. Impact of POR*28 on the clinical pharmacokinetics of CYP3A phenotyping probes midazolam and erythromycin. *Pharmacogenet Genomics*. 2013;23(3):148-155.
16. Tang SC, Lagas JS, Lankheet NA, et al. Brain accumulation of sunitinib is restricted by P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) and can be enhanced by oral elacridar and sunitinib coadministration. *Int J Cancer*. 2012;130(1):223-233.
17. van Erp NP, Eechoute K, van der Veldt AA, et al. Pharmacogenetic pathway analysis for determination of sunitinib-induced toxicity. *J Clin Oncol*. 2009;27(26):4406-4412.
18. Garcia-Donas J, Esteban E, Leandro-García GJ, et al. Single nucleotide polymorphism associations with response and toxic effects in patients with advanced renal-cell carcinoma treated with first-line sunitinib: a multicentre, observational, prospective study. *Lancet Oncol*. 2011;12(12):1143-1150.
19. van der Veldt AA, Eechoute K, Gelderblom H, et al. Genetic polymorphisms associated with a prolonged progression-free survival in patients with metastatic renal cell cancer treated with sunitinib. *Clin Cancer Res*. 2011;17(3):620-629.
20. Eechoute K, van der Veldt AA, Oosting S, et al. Polymorphisms in endothelial nitric oxide synthase (eNOS) and vascular endothelial growth factor (VEGF) predict sunitinib-induced hypertension. *Clin Pharmacol Ther*. 2012;92(4):503-510.
21. Kim JJ, Vaziri SA, Rini BI et al. Association of VEGF and VEGFR2 single nucleotide polymorphisms with hypertension and clinical outcome in metastatic clear cell renal cell carcinoma patients treated with sunitinib. *Cancer*. 2012;118(7):1946-1954.
22. EuroTARGET: Targeted therapy in Renal cell cancer: GENetic and Tumour related biomarkers for response and toxicity. European collaborative project funded by the European Commission under the Seventh Framework Programme (FP7). URL: <http://www.eurotargetproject.eu/>.
23. Urun Y, Gray K, Signoretti S, et al. Pharmacogenetics as predictor of sunitinib and mTOR inhibitors toxicity in patients with metastatic renal cell carcinoma (mRCC). *ASCO Meeting Abstr* 4570. 2013;31. URL: <http://meetinglibrary.asco.org/content/113157-132>.
24. Elens L, Bouamar R, Hesselink DA et al. A new functional CYP3A4 intron 6 polymorphism significantly affects tacrolimus pharmacokinetics in kidney transplant recipients. *Clin Chem*. 2011;57(11):1574-1583.
25. Elens L, van Gelder T, Hesselink DA, Haufroid V, van Schaik RH. CYP3A4*22: promising newly identified CYP3A4 variant allele for personalizing pharmacotherapy. *Pharmacogenomics*. 2013;14(1):47-62.
26. Xu CF, Bing NX, Ball HA, et al. Pazopanib efficacy in renal cell carcinoma: evidence for predictive genetic markers in angiogenesis-related and exposure-related genes. *J Clin Oncol*. 2011;29(18):2557-2564.

27. de Jonge H, Metalidis C, Naesens M, Lambrechts D, Kuypers DR. The P450 oxidoreductase *28 SNP is associated with low initial tacrolimus exposure and increased dose requirements in CYP3A5-expressing renal recipients. *Pharmacogenomics*. 2011;12(9):1281-1291.
28. Oneda B, Crettol S, Jaquenoud Sirot E, Bochud M, Ansermot N, Eap CB. The P450 oxidoreductase genotype is associated with CYP3A activity in vivo as measured by the midazolam phenotyping test. *Pharmacogenet Genomics*. 2009;19(11):877-883.
29. Chu H, Wang M, Yan F, et al. Polymorphisms in the IL13 and IL-4R genes are associated with the development of renal cell carcinoma. *Ann Oncol*. 2012;23(8):2114-2121.
30. Schutz FA, Pomerantz MM, Gray KP, et al. Single nucleotide polymorphisms and risk of recurrence of renal-cell carcinoma: a cohort study. *Lancet Oncol*. 2013;14(1):81-87.
31. Maitland ML, Xu CF, Cheng YC, et al. Identification of a variant in KDR associated with serum VEGFR2 and pharmacodynamics of Pazopanib. *Clin Cancer Res*. 2015;21(2):365-372.
32. Lindauer A, Di Gion P, Kanefendt F, et al. Pharmacokinetic/pharmacodynamic modeling of biomarker response to sunitinib in healthy volunteers. *Clin Pharmacol Ther*. 2010;87(5):601-608.

