



Universiteit
Leiden
The Netherlands

Pharmacogenetics of sunitinib in metastatic renal cell carcinoma

Diekstra, M.H.M.; Diekstra M.H.M.

Citation


Diekstra, M. H. M. (2017, September 13). *Pharmacogenetics of sunitinib in metastatic renal cell carcinoma*. Retrieved from <https://hdl.handle.net/1887/55944>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/55944>

Note: To cite this publication please use the final published version (if applicable).





CHAPTER 2

A decade of pharmacogenomics research on tyrosine kinase inhibitors in metastatic renal cell cancer: a systematic review

Diekstra MH, Swen JJ, Gelderblom H, Guchelaar H-J.

Exp Rev of Mol Diagn. 2016;16(5):605-618.

ABSTRACT

Objective The individual response to targeted tyrosine kinase inhibitors (TKIs) in the treatment of metastatic renal cell cancer (mRCC) is highly variable. Outlined in this article are findings on potential biomarkers for TKI treatment outcome in mRCC and an evaluation of the status of clinical implementation.

Methods Articles were selected by two independent reviewers using a systematic search in five medical databases on renal cell carcinoma, TKIs, and pharmacogenetics.

Results Many researchers have focused on predictive biomarkers for treatment outcome of targeted therapies in mRCC patients. Attempts to explain differences in efficacy and toxicity of TKIs by use of genetic variants in genes related to the pharmacokinetics and pharmacodynamics of the drug have been successful.

Conclusion Most findings on potential biomarkers have not been validated and therefore biomarker testing to guide choice of therapy and dose in mRCC is not yet feasible.

INTRODUCTION

Using pharmacogenomics in our endeavor to personalize mRCC treatment

Renal cell cancer (RCC) represents ~3% of cancer in adults (approximately 170,000 people worldwide per year). RCC typically occurs in adults aged 50-70 years and is more common in men than in women. For many years, cytokine therapies (interleukin-2 and interferon-alpha) were the only available treatment options with limited efficacy and considerable toxicity.^{1,2} In the last decade, multiple targeted agents for the treatment of metastatic RCC (mRCC) have become available, namely: the multitargeted oral tyrosine kinase inhibitors (TKIs) with antiangiogenic activity. Of the first generation TKIs, sorafenib was approved for the treatment of mRCC in 2005, followed by sunitinib (2006) and pazopanib (2009). The first-generation TKIs target the vascular endothelial growth factor (VEGF) pathway by inhibition of the vascular endothelial growth factor receptors (VEGFR-1, 2, and 3) but also inhibit many other targets. Sorafenib, sunitinib, and pazopanib all target the platelet-derived growth factor receptor (PDGFR) α and/or β and the c-Kit protein (c-Kit). Additionally, sunitinib and sorafenib inhibit FMS-like tyrosine kinase 3 (Flt-3) and glial cell line-derived neurotrophic factor receptor (REarranged during Transfection) tyrosine kinases. Furthermore, sunitinib inhibits colony-stimulating factor 1 receptor, and sorafenib specifically inhibits the RAF kinases (C-RAF and B-RAF).³⁻⁶

Starting 2012, the second-generation TKIs emerged with an improved potency and selectivity, such as axitinib. Many of these second-generation TKIs were assessed for the treatment of mRCC, such as tivozanib, cediranib, and cabozantinib.^{2-5,7,8} Axitinib predominantly blocks the VEGF receptors, with a 50-450 times higher potency than the first-generation TKIs and less off-target activity, which could explain the larger therapeutic window and fewer side effects.^{2,7}

Increased exposure to sunitinib is associated with improved survival outcomes but also an increased risk for adverse events.^{3,5} Sunitinib revealed an improvement in progression-free survival (PFS) and objective response rate compared to interferon-alpha ($P < 0.001$) and showed a tendency towards an improved overall survival (OS) ($P = 0.051$).³ For several years, sunitinib has been considered the first-choice treatment for mRCC. However, based on the results of the PISCES trial (a randomized, controlled, double-blind, cross-over trial assessing treatment preference for pazopanib versus sunitinib in patients with metastatic renal cell carcinoma) and the larger COMPARZ trial (a phase 3 study of pazopanib versus sunitinib in the treatment of locally advanced and/or metastatic renal cell carcinoma)^{9,10} indicating noninferiority for pazopanib with regard to toxicity, patient preference, and quality of life, both pazopanib and sunitinib are prescribed as first-line treatment for mRCC.^{9,10}

The individual response to TKIs is highly variable in mRCC patients: some experience severe toxicities for which dose reductions or even cessation of therapy is needed, while others show no response at all. Previous studies identified single nucleotide polymorphisms (SNPs) in genes encoding enzymes or transporters related to pharmacokinetics (PK) and pharmacodynamics (PD) of TKIs which have been associated with toxicity and efficacy of TKIs.^{11,12} Also cytokines and antiangiogenic factors have been studied as potential predictors for TKI treatment outcome in mRCC patients. Ultimately, biomarker testing prior to start of therapy may result in providing the individual patient the most effective treatment with the least possible side effects. Several review articles on pharmacogenomics in mRCC have been published¹¹⁻¹⁴, but only one with a systematic approach.¹⁴ In our current systematic review article, we strictly focus on the predictive value of biomarkers (as opposed to the prognostic value), especially SNPs (pharmacogenetics biomarkers) and only in TKI treatment. Here, we outline the findings from currently available pharmacogenomics studies on potential predictive biomarkers for mRCC treatment outcome. In addition, we evaluate the quality of the selected studies, clarify the status on implementation of these predictive biomarkers into the clinic, and give recommendations as to what the focus of attention should be in future studies.

METHODS

Protocol

A systematic review was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.¹⁵ Prior to the literature search, a systematic review protocol was prepared following the PRISMA-P 17-item checklist (**Supplementary document 1**)¹⁵ and registered at the international prospective register of systematic reviews (PROSPERO; CRD42015016509).¹⁶

Systematic literature search

Together with an experienced librarian, we performed a systematic search in 5 medical databases: MEDLINE, EMBASE (OVID version), Web of Science, The Cochrane Library, and Cochrane Central Register of Controlled Trials (CENTRAL). This search comprised three topics: renal cell carcinoma, TKIs (e.g. sunitinib, sorafenib), and pharmacogenetics (**Supplementary document 2**). All databases were searched for keywords on these items taking into account the terminological and technical differences between these databases. Various synonyms and related terms for all subjects were used. The search was performed on 14 January 2015. Results were limited to articles in the English language and from the year 2004 onwards. Meeting abstracts were excluded. From

our initial search up to the moment of our final selection of articles, new studies may have appeared. To cover articles published after 14 January 2015, the initial search was repeated in MEDLINE on 31 May 2015.

Selection process and data extraction

Articles were first selected by title, then by abstract, and finally relevant data were extracted systematically from included full text articles. Selection of included articles was performed by two independent reviewers (MD, JS). In case of disagreement, a third reviewer (HJG) made the final decision. References were eligible when genetic biomarkers (i.e. SNPs, genetic variants, gene/protein expression) were investigated in relation to TKI treatment and mRCC. Review articles, preclinical studies (animal models and in vitro studies), phase I or II clinical trials, and papers discussing pathology, prognosis, tumor biology, or somatic mutations (e.g. Von Hippel-Lindau mutations) were excluded. Accordingly, articles were excluded if nontreatment-related biomarkers were studied or if no TKI response was assessed, because these generally consider prognostic biomarkers.

Extracted data were first author, year of publication, demographics (age, sample size, sex, ethnicity), sample material and origin (i.e. germline DNA from blood, serum, or other), methods, tumor type, drug information (i.e. drug name, treatment line, dose), evaluated biomarkers, endpoints, and statistical approach. Articles resulting from the secondary search on 31 May 2015 were subject to an identical selection process. Using cross-references of the included articles from our systematic search, relevant articles were added to our final selection. Clinical validity (the robustness of the statistical association between a pharmacogenetic variant and outcome of drug therapy) and clinical utility (a test's health-care value) of the included articles was evaluated in a descriptive manner.^{17,18}

RESULTS

Search results

Our primary search identified 968 articles (**Supplementary document 2**). The secondary search identified 30 additional articles. After selection by title, abstract, and full text, a total of 54 articles were available for evaluation in this review. **Figure 1** represents a flowchart on the study selection process. The extracted data from the included articles are presented in **Table 1** and **Supplementary document 3**.




Figure 1 Flowchart of study selection process. Our original search on the 14th of January 2015 resulted in a total of 968 articles (**Supplementary document 2**). For the first selection by title, there was a disagreement between the two independent reviewers on a number of 114 titles (12%). After consensus was reached with the last review author (HJG), this resulted in a number of 104 articles which was complemented with 33 additional articles of the search from January 14 to May 31, 2015. A total of 134 articles was available for the second selection step based on abstract reading. Consensus had to be reached for 20 of the 134 abstracts (15%) and resulted in exclusion of 67 articles. Consequently, a number of 67 articles underwent full text review and data extraction. Finally, 15 articles were excluded for data extraction because no treatment was evaluated ($n = 8$), prognostic biomarkers were investigated ($n = 5$) or these were no original articles but commentaries ($n = 2$) and using cross-references, we included two additional studies.^{37,69} Ultimately, a total of 54 articles was included in our systematic review.

SNPs associated with TKI treatment outcome in mRCC patients

From 2007 up till now, several genetic variants have been assessed for their possible association with the treatment outcome of sunitinib, pazopanib, sorafenib, or axitinib. These SNPs have been selected according to the candidate gene approach based on the knowledge on PK and PD of the drug. The reported findings in this field are outlined below and summarized in **Table 1** and **Figure 2**.¹⁹⁻⁴⁴

Table 1 Reported SNP associations ($P \leq 0.05$) with TKI treatment outcome in mRCC from 2009 to 2015.

First Author, year	TKI	No. of patients	Ethnicity	Associated SNPs	Study endpoint	Effect size	95%CI	P value
van Erp <i>et al.</i> , 2009 ¹⁹	SUN	219	white: 93.6%	<i>CYP1A1</i> rs1048943	Leukopenia	OR:6.24	1.20-32.42	$P=0.029$
				<i>FLT3</i> rs1933437		OR:0.36	0.17-0.77	$P=0.008$
				<i>NR1I3</i> (rs2307424, rs2307418, rs4073054) CAG haplotype		OR:1.74	1.02-2.96	$P=0.041$
				<i>VEGFR2</i> rs2305948	Any toxicity > grade 2	OR:2.39	1.02-5.60	$P=0.046$
				<i>ABCG2</i> (rs55930652, rs2622604)		OR:0.38	0.17-0.83	$P=0.016$
				<i>CYP1A1</i> rs1048943	Mucosal inflammation	OR:4.03	1.24-13.09	$P=0.021$
				<i>ABCB1</i> (rs1045642, rs1128503, rs2032582) TTT haplotype	HFS	OR:0.39	0.16-0.94	$P=0.035$
Garcia-Donas <i>et al.</i> , 2011 ²⁰	SUN	89 (efficacy)	mostly European	<i>VEGFR3</i> rs307826	PFS	HR:3.57	1.75-7.30	$P=0.0079$
				<i>VEGFR3</i> rs307821		HR:3.31	1.64-6.68	$P=0.014$
		95 (toxicity)		<i>CYP3A5</i> rs776746	dose reductions due to toxicity	HR:3.75	1.67-8.41	$P=0.022$
Beuselinck <i>et al.</i> , 2014 ²¹	SUN	96	95% CAU	<i>ABCB1</i> rs1125803	Time-to-dose-reduction (TTDR)	-	-	$P=0.012$
				<i>ABCB1</i> rs2032582		-	-	$P=0.024$
Mizuno <i>et al.</i> , 2010 ²²	SUN	5	JAP	<i>ABCG2</i> rs2231142	exposure and toxicity	-	-	$P<0.05$
Mizuno <i>et al.</i> , 2012 ²³	SUN	19	JAP	<i>ABCG2</i> rs2231142	Dose-adjusted AUC	-	-	$P=0.02$
Miura <i>et al.</i> , 2014 ²⁴	SUN	1	JAP	<i>ABCG2</i> rs2231142	grade 3 and 4 toxicity	-	-	-
Eechoute <i>et al.</i> , 2012 ²⁵	SUN	255	91% CAU	<i>VEGFA</i> (rs699947, rs833061, rs2010963) ACG haplotype	SBP	-	-	$P=0.014$
					MAP	-	-	$P=0.036$
					Grade 3 hypertension	OR:0.59	0.34-1.03	$P=0.031$
				<i>eNOS</i> rs2070744	Grade 3 hypertension	OR:2.62	1.08-6.35	$P=0.045$
Kim <i>et al.</i> , 2012 ²⁶	SUN	63	CAU	<i>VEGFA</i> rs2010963	prevalence of hypertension	OR:13.62	3.71-50.04	$P=0.05$
				<i>VEGFA</i> rs2010963	duration of hypertension	21%	12%-29%	$P=0.02$
				<i>VEGFA</i> rs3025039 + <i>VEGFR2</i> rs2305948	OS	HR:3.18	1.12-8.99	$P=0.03$
Kim <i>et al.</i> , 2013 ²⁷	SUN	65	Asian (Korea)	<i>ABCG2</i> rs2231142	grade 3 or 4 thrombocytopenia	OR:9.90	1.16-infinity	$P=0.04$
					grade 3 or 4 neutropenia	OR:18.20	1.49-222.09	$P=0.02$
					grade 3 or 4 HFS	OR:28.46	2.22-364.94	$P=0.01$
Takayoshi <i>et al.</i> , 2015 ²⁸	SUN	1	probably JAP	<i>ABCB1</i> (rs1045642, rs1128503, rs2032582)	Severe toxicity	-	-	-
Teo <i>et al.</i> , 2015 ²⁹	SUN	25	CHI = 86.1% (other Malay or Indian)	<i>ABCB1</i> rs1045642	sunitinib exposure	76.81 vs 56.55 ng/ml		$P=0.03$
					all-grade rash	RR:3.00	1.17-7.67	$P<0.05$
					all-grade mucositis	RR:1.60	1.10-2.34	$P<0.05$
					disease progression	RR:4.57	1.08-19.42	$P=0.05$

Table 1 Continued

First Author, year	TKI	No. of patients	Ethnicity	Associated SNPs	Study endpoint	Effect size	95%CI	P value
Diekstra <i>et al.</i> , 2015 ³⁰	SUN	333	96% white	CYP3A5 rs776746	dose reductions	OR:2.0	1.0-4.0	P=0.039
				ABCB1 (rs1128503, rs2032582, rs1045642)	PFS	HR:1.9	1.3-2.6	P=0.000275
van der Veldt <i>et al.</i> , 2011 ³¹	SUN	136	95.6% CAU	CYP3A5 rs776746	PFS	HR:0.27	0.08-0.89	P=0.032
				NR1I3 (rs2307424, rs2307418, rs4073054) CAT haplotype		HR:1.76	1.11-2.79	P=0.017
				ABCB1 (rs1045642, rs1128503, rs2032582) TCG haplotype		HR:0.52	0.29-0.95	P=0.033
Scartozzi <i>et al.</i> , 2013 ³²	SUN	84	CAU (Italy)	VEGFA rs833061	PFS	HR:0.71	-	P=0.0197
					OS	HR:0.69		P=0.0011
				VEGFA rs2010963	PFS	HR:0.19	-	P=0.0201
				VEGFR3 rs6877011	PFS	HR:0.35	-	P<0.0001
Motzer <i>et al.</i> , 2014 ³³	SUN	202	88% CAU	VEGFA rs699947	PFS, OS, objective response or time-to-tumor progression	no significant correlations		
				VEGFA rs1570360				
				VEGFR3 rs448012				
				VEGFR3 rs307821				
				VEGFR3 rs307826				
Beuselinck <i>et al.</i> , 2013 ³⁴	SUN	88	94% CAU	ABCB1 rs1128503	PFS	HR:0.464	0.234-0.918	P=0.027
				NR1I3 rs4073054		HR:1.864	1.082-3.210	P=0.025
				VEGFR3 rs307821		HR:1.981	1.060-3.702	P=0.032
				FGFR2 rs2981582		HR:2.669	1.094-6.511	P=0.031
				NR1I2 rs2276707		HR:2.978	1.012-8.761	P=0.047
				ABCB1 rs1128503	OS	HR:0.415	0.193-0.894	P=0.025
				NR1I3 rs4073054		HR:1.927	1.046-3.549	P=0.035
				VEGFR3 rs307821		HR:2.265	1.202-4.268	P=0.011
				NR1I3 rs2307424		HR:1.913	1.006-3.636	P=0.048
Beuselinck <i>et al.</i> , 2014 ³⁵	SUN	91	93% CAU	VEGFR1 rs9582036	Response Rate (RR)	0% vs 46%		P=0.028
					OS	HR:0.249	0.078-0.799	P=0.008
				VEGFR1 rs9554320	PFS	HR:2.713	1.321-5.575	P=0.005
Mizuno <i>et al.</i> , 2014 ³⁶	SUN	19	JAP	ABCG2 rs2231142	Oral clearance (CL/F)	Improvement of PK model: Δ OFV=8.59 (P=0.003)		
Diekstra <i>et al.</i> , 2014 ³⁷	SUN	114		CYP3A4 rs35599367	Sunitinib clearance	-22.5%	-	P=0.01
				CYP3A5 rs776746	SU12662 clearance	+28.0%	-	P=0.04
				ABCB1 rs2032582	Sunitinib clearance	+18.0%	-	P=0.02
Xu <i>et al.</i> , 2010 ³⁸	PAZ	116 (exploratory) 130 (replication)	white	UGT1A1 rs8175347 (*28) TA7 homozygotes	Bilirubin levels	OR:13.1	5.3-32.2	P=4.5×10 ⁻⁵
Xu <i>et al.</i> , 2011 ³⁹	PAZ	115 (exploratory) and 128 (replication)	white	HFE rs2858996 + rs707889 (in strong LD)	ALT levels	OR:39.7	2.2-703.7	P=0.006

Table 1 Continued

First Author, year	TKI	No. of patients	Ethnicity	Associated SNPs	Study endpoint	Effect size	95%CI	P value
Xu <i>et al.</i> , 2011 ⁴⁰	PAZ	397	84% white	<i>IL8</i> rs1126647	PFS	HR:1.8	1.2-2.7	<i>P</i> =0.009
				<i>HIF1A</i> rs11549467		HR:1.8	1.1-3.1	<i>P</i> =0.03
				<i>HIF1A</i> rs11549467	Response rate (RR)	RR:30% vs. 43%		<i>P</i> =0.02
				<i>NR1I2</i> rs3814055		RR:37% vs. 50%		<i>P</i> =0.03
				<i>VEGFA</i> rs833061		RR:33% vs. 51%		<i>P</i> =0.02
Motzer <i>et al.</i> , 2013 ⁴¹	SUN or PAZ	719 (SUN: N=350, PAZ: N=369)	Europe, North America, Asia, Australia	<i>UGT1A1</i> rs8175347 and rs4148323 (*28, *37 and *60)	Bilirubin levels on sunitinib	OR:4.51	1.26-16.11	<i>P</i> =0.024
					Bilirubin levels on pazopanib	OR:3.65	1.31-10.16	<i>P</i> =0.012
Xu <i>et al.</i> , 2015 ⁴²	PAZ or SUN	3 substudies: 186 pts (87% white) (PAZ) 690 pts (353 PAZ + 337 SUN) 98% white 88 (SUN)		<i>IL8</i> rs1126647	OS	HR:1.32	1.15-1.52,	<i>P</i> =8.8×10 ⁻⁵
Tsuchiya <i>et al.</i> , 2013 ⁴³	SOR	33	JAP	<i>ABCC2</i> rs717620	high-grade skin rash	35 vs 0%	-	<i>P</i> =0.032
				<i>HLA-A*24</i>		-	-	<i>P</i> =0.049
Escudier <i>et al.</i> , 2015 ⁴⁴	AXI and SOR	305 (AXI: N=159, SOR: N=146)	94.1% white	<i>VEGFR2</i> rs2071559	PFS sorafenib	2.223	1.268-3.897	<i>P</i> =0.0053
					OS sorafenib	2.584	1.390-4.803	<i>P</i> =0.0027

ABCB1: ATP-binding cassette member B1; ABCG2: ATP-binding cassette member G2; ALT: alanine aminotransferase; AST: aspartate aminotransferase; AXI: axitinib; CAU: Caucasian; CC: clear cell renal cell carcinoma; CHI: Chinese; GIST: gastrointestinal stromal tumor; JAP: Japanese; ΔOFV: delta Objective Function Value; OS: overall survival; PAZ: pazopanib; PFS: progression-free survival; PK: pharmacokinetic; SOR: sorafenib; SUN: sunitinib.

Pharmacogenetics to predict sunitinib toxicity

The first pharmacogenetic study on sunitinib tested for associations of genetic polymorphisms with toxicity endpoints ($P < 0.05$) in a cohort of 219 patients with mRCC (N=159), gastrointestinal stromal tumor (GIST) (N=50), or other tumors (N=10). Van Erp *et al.* observed an increased risk for leukopenia in *CYP1A1* rs1048943, and *FLT3* rs1933437 and absence of CAG in the *NR1I3* haplotype (rs2307424, rs2307418, rs4073054). *CYP1A1* rs1048943 was associated with mucosal inflammation, and presence of TTT in *ABCB1* (rs1045642, rs1128503, rs2032582) was associated with HFS. Presence of the TT copy in *ABCG2* (rs55930652, rs2622604) and *VEGFR2* rs2305948 was associated with any toxicity >grade 2.¹⁹

Later, in a group of 95 mRCC patients, Garcia-Donas *et al.* reported that the *CYP3A5* A-allele (*CYP3A5**1) of rs776746 was associated with dose reductions due to toxicity

(adjusted $P < 0.05$).²⁰ In a study of Beuselinck *et al.*, an increased time-to-dose reduction was seen for the TT genotype patients of rs1128503 and rs2032582 in *ABCB1* compared to C-allele and G-allele carriers, respectively.²¹

Mizuno *et al.*^{22,23} observed a case of severe haematological toxicity in an RCC patient with the AA genotype of rs2231142 in *ABCG2*. This prompted a study in 5 patients (1 variant homozygous, 3 heterozygous, and 1 wild-type homozygous) in which they observed an elevated sunitinib exposure for AA carriers.²² The variant allele A is common in Asians (~30%), but rare in Caucasians (~10%) and African Americans (~5%). In a subsequent study with 19 Japanese patients, only 1 patient had the AA genotype of rs2231142 in *ABCG2*.²³ It was reported that this SNP was associated with an increase in systemic exposure to sunitinib, possibly causing thrombocytopenia and hypertension.²³ Miura *et al.*²⁴ followed with a case report in 2014 describing a patient with mRCC who developed severe grade 3 and 4 adverse events on sunitinib, namely fever, thrombocytopenia, transaminase elevation, hypoxia, and pulmonary edema. Again, the AA genotype of rs2231142 in *ABCG2* was suggested to explain these toxicities in Asians.

Hypertension is a common adverse effect of sunitinib. Eechoute *et al.* identified genetic biomarkers by using the pathway approach using data from 255 patients with clear cell mRCC (N=167), non-clear cell mRCC (N=22), an unknown type of RCC (N=4), GIST (N=53), or another tumor type (N=9).²⁵ An ACG haplotype in *VEGF-A* (rs699947, rs833061, rs2010963) was associated with elevations in systolic blood pressure (SBP) and mean arterial pressure (MAP). The same haplotype in *VEGF-A* and presence of the C-allele of rs2070744 in *eNOS* showed a tendency to develop grade 3 hypertension.²⁵ Similarly, Kim *et al.*²⁶ evaluated 63 sunitinib-treated mRCC patients for the association of SNPs in *VEGF* or *VEGF-R* with hypertension and clinical outcome. The GG genotype of SNP rs2010963 in *VEGF* was associated with a greater chance for the occurrence and duration of hypertension.²⁶

Sunitinib-induced toxicities in relation to pharmacogenetic determinants were studied by Kim *et al.*²⁷ in a number of 65 Korean mRCC patients. Compared to C-allele carriers, the AA genotype of rs2231142 in *ABCG2* had an increased risk for grade 3 or 4 thrombocytopenia, neutropenia, and HFS.²⁷ In addition, a case report of Takayoshi *et al.*²⁸ reported a 65-year-old woman with mRCC who showed severe toxicities and was found to have the TTT haplotype on *ABCB1* (rs1045642, rs1128503, rs2032582). It was thought that the severe toxicities as observed in this patient could be associated with a high sunitinib exposure, probably explained by the *ABCB1* genetic polymorphism.²⁸

Teo *et al.*²⁹ reported an association analysis in a Chinese population of 25 patients to investigate the effect of SNPs on both PK and clinical outcome (i.e. toxicity and response). The variant CC genotype of rs1045642 in *ABCB1* was associated with an increased risk for rash and mucositis as compared to T-allele carriers ($P < 0.05$). No association with sunitinib treatment outcome was observed for rs776746 in *CYP3A5*.²⁹

In order to verify the value of earlier reported associations, Diekstra *et al.*³⁰ pooled patients from exploratory studies in The Netherlands, Spain, and the United States^{19,20,25,26,31}, resulting in a sample size of 333 mRCC patients. Data confirmed the association of *CYP3A5**1 with the need for dose reductions ($P<0.05$) as presented earlier by Garcia-Donas *et al.*^{20,30}

Pharmacogenetics to predict sunitinib efficacy

In a follow-up paper to van Erp *et al.*^{19,31}, the same group tested a comparable set of SNPs for associations with sunitinib efficacy. In a population of 136 mRCC patients, they observed that *CYP3A5**1, presence of the TCG copy in *ABCB1* (rs1045642, rs1128503, rs2032582), or absence of the CAT copy in *NR1I3* (rs2307424, rs2307418, rs4073054) were associated with an improved PFS. Carriers of the favorable genetic profile (at least an A-allele of *CYP3A5*, a TCG copy in *ABCB1*, or a missing CAT copy in *NR1I3*) showed an improved PFS and OS compared to the disadvantageous genetic profile carriers.³¹

With regard to SNPs in *VEGF* and *VEGF-R*, Garcia-Donas *et al.* reported SNPs rs307826 and rs307821 in *VEGF-R3* to be associated with a decrease in PFS in a study of 89 mRCC patients.²⁰ Kim *et al.*²⁶ showed that carriers of the combination of CC and GG genotypes of rs3025039 in *VEGF-A* and rs2305948 in *VEGF-R2*, respectively, showed a worse OS compared to noncarriers.²⁶ In another study of *VEGF* and *VEGF-R* SNPs in a group of 84 mRCC patients, Scartozzi *et al.*³² report that carriers of the favorable SNP genotypes of rs833061, rs6877011, rs2010963, and rs699947 showed a better overall response rate compared to the unfavorable genotype carriers.³² By using a similar approach with a slightly different SNP set, Motzer *et al.*³³ report conflicting results. They could not identify any significant associations of SNPs in *VEGF-A* or *VEGFR3* with efficacy outcomes in sunitinib treatment (PFS, OS, objective response, or time-to-tumor progression).³³

Beuselinck *et al.* assessed 16 SNPs in 10 genes for their relation with efficacy endpoints in 88 mRCC patients treated with sunitinib.³⁴ SNP rs1128503 in *ABCB1*, rs4073054 in *NR1I3*, and rs307821 in *VEGF-R3* were associated with PFS and OS. For PFS only, rs2981582 in *FGF-R2* and rs2276707 in *NR1I2* showed an association. For OS only, rs2307424 in *NR1I3* and rs307826 in *VEGF-R3* were associated ($P<0.05$).³⁴ In a separate study on 91 subjects, the CC genotypes of rs9582036 in *VEGF-R1* had a lower response rate and a decreased survival (PFS and OS) compared to A-allele carriers.³⁵ Furthermore, a decreased PFS was observed for the AA genotype of rs9554320 in *VEGF-R1* compared to C-allele carriers.³⁵ Teo *et al.*²⁹ reported that the CC genotype of rs1045642 in *ABCB1* was associated with progression of the disease as compared to T-allele carriers ($P<0.05$)²⁹ in 25 Chinese patients.

In the aforementioned study of Diekstra *et al.*, presence of CGT in the *ABCB1* haplotype (rs1128503, rs2032582, rs1045642) was confirmed to be associated with an improved PFS ($P<0.001$) as previously observed by van der Veldt *et al.*^{30,31}




Figure 2 Main results on associations of SNPs involved in pharmacokinetics or pharmacodynamics of TKIs with clinical treatment outcome in mRCC.

An increased risk for common sunitinib toxicities was seen for SNPs in *CYP1A1*, *NR1I3*, *ABCB1* and *ABCG2*. Furthermore, *CYP3A5*1* was associated with dose reductions due to toxicity and an increased time-to-dose-reduction (TTDR) was seen for *ABCB1* SNPs. Associations with the clearance of sunitinib and/or SU12662 were observed for SNPs in *CYP3A5*, *ABCB1* and *CYP3A4*. A higher exposure to sunitinib and progressive disease was reported for rs1045642 in *ABCB1* in Chinese patients. For efficacy, *CYP3A5*1* and SNPs in *ABCB1*, *NR1I2* and *NR1I3* were associated with PFS. Associations of *CYP3A5*1* with the need for dose reductions and of the *ABCB1* haplotype with PFS were confirmed in a larger study.³⁰ Regarding pharmacodynamics, SNPs in *VEGFR2*, *FLT3*, *VEGF-A* and *eNOS* were associated with sunitinib toxicities and with hypertension in particular. Associations with either response rate, PFS and/or OS were observed for SNPs in *VEGF-A*, *VEGF-R1*, *VEGF-R3* and *FGF-R2*.¹⁹⁻⁴⁴ For pazopanib, only articles from the group of Xu *et al.* were included in our search. The main findings were that *UGT1A1*28*, *UGT1A1*60* and *CYP1A2* (rs762551) were associated with bilirubin levels and SNPs in *HFE* were associated with increased ALT levels. Additionally, SNPs in *IL8*, *HIF1A*, *NR1I2* and *VEGFA* were associated with either PFS or response rate. For patients using either sunitinib or pazopanib, *UGT1A1* SNP genotypes *28, *37 and *6 showed an increased risk for hyperbilirubinemia and rs1126647 in *IL8* was associated with OS. Only two pharmacogenetic studies on sorafenib were found in which genetic polymorphisms in *ABCC2* and *HLA-A* were associated with high-grade skin rash and rs2071559 in *VEGF-R2* was associated with PFS and OS. For axitinib, no associations were reported.⁴⁰⁻⁴⁴

Pharmacogenetics to predict sunitinib PK

To investigate the effect of genetic variants on the PK of sunitinib, Mizuno *et al.* performed a population PK study in a set of 19 patients which identified the AA genotype of rs2231142 in *ABCG2* to be a predictor for increased clearance of sunitinib.³⁶ In the study of Teo *et al.*²⁹, including 25 patients of Chinese origin, the variant CC genotype of rs1045642 in *ABCB1* was associated with a higher exposure to sunitinib as compared to T-allele carriers ($P<0.05$).²⁹ Diekstra *et al.*³⁷ tested a selection of the earlier associated SNPs related to the PK of sunitinib for possible association with clearance of sunitinib and/or its active metabolite (SU12662). Fourteen SNPs were tested in a sample of 114 patients with RCC (N=69), neuro-endocrine tumors (N=14), GIST (N=8), or another solid tumor type (N=23). A 28% increase in clearance of SU12662 was observed for the AG genotype of rs776746 in *CYP3A5*. Furthermore, the TT variant genotype of rs2032582 in *ABCB1* showed an 18% increase in clearance of sunitinib ($P<0.05$). For T-allele carriers of rs35599367 in *CYP3A4* (*CYP3A4**22), a 22.5% decreased clearance was observed ($P<0.01$).³⁷

Pharmacogenetics to predict pazopanib toxicity

Fewer studies are available on pharmacogenetics of pazopanib, since pazopanib was approved 3 years after sunitinib. Pazopanib and sunitinib share most common toxicities, although for pazopanib hepatotoxicity with elevations of alanine transaminase levels in particular is a common adverse effect.³⁸ In 2010, Xu *et al.*³⁸ investigated 28 SNPs for their relation with levels of alanine aminotransferase (ALT) and bilirubin in pazopanib-treated mRCC patients from an exploratory cohort (N=116) and a replication cohort (N=130). In the exploratory cohort, SNPs rs8175347 (*UGT1A1**28), rs4148323 (*UGT1A1**60), and rs762551 in *CYP1A2* were associated with bilirubin levels, and none of the SNPs showed an association with ALT levels. In the replication cohort, only the TA-repeat polymorphism of *UGT1A1**28 was associated with hyperbilirubinemia. The TA7 homozygote patients developed more hyperbilirubinemia compared to other genotypes, probably due to a reduced expression of UGT1A1. It was suggested that this is not likely to affect pazopanib exposure nor toxicities, since pazopanib is not glucuronized by UGT1A1.³⁸ In roughly the same patient cohorts (N=115 explorative and N=128 for replication), Xu *et al.* studied 9308 SNPs in 282 candidate genes. For rs2858996 and rs707889 in *HFE*, which are in strong linkage disequilibrium, the TT genotypes were associated with increased ALT levels.³⁹

Pharmacogenetics to predict pazopanib efficacy

In a subsequent study, Xu *et al.*⁴⁰ investigated a slightly different set of 27 SNPs in candidate genes (SNPs in *IL-8* and *HIF1A*) to test whether these are related to pazopanib efficacy. In a cohort of 397 RCC patients, a decreased PFS was observed for TT variant

carriers of rs1126647 in *IL8* compared to wildtype AA. A worse PFS and response rate were observed for AG carriers of rs11549467 in *HIF1A* compared to wild-type GG. A reduced response rate was also observed for TT carriers of rs3814055 in *NR1I2* compared to wild-type CC and for wildtype CC of rs833061 in *VEGFA* compared to TT genotypes.⁴⁰

Pharmacogenetics to predict clinical outcome of sunitinib and pazopanib

In some of the included studies, a combined genetic association analyses was performed on patients who are either treated with sunitinib or pazopanib. Motzer *et al.*⁴¹ investigated the *UGT1A1* polymorphisms as reported by Xu *et al.*³⁸ in 719 patients from the COMPARZ trial: a phase III, randomized, clinical trial comparing pazopanib versus sunitinib in mRCC treatment.^{10,41} Subjects with an expected reduction in function of *UGT1A1* based on the SNP genotypes of rs8175347 and rs4148323 (*28, *37, and *6) showed increased levels of bilirubin at baseline and had a higher risk for developing hyperbilirubinemia on treatment with either sunitinib or pazopanib. None of the *UGT1A1* SNPs showed an association with elevated ALT levels.⁴¹ More recently, Xu *et al.*⁴² examined three separate patient cohorts of 186 patients on pazopanib in the first cohort, 690 patients using either pazopanib (N=353) or sunitinib (N=337) in the second cohort, and the third cohort comprised 88 patients on sunitinib. The variant T allele of SNP rs1126647 in *IL8* was associated with an inferior OS in every separate cohort ($P<0.05$). In a meta-analysis of the association results in the combined data set of 964 patients, this SNP was also associated with a worse OS.⁴²

Pharmacogenetics to predict clinical outcome of sorafenib or axitinib

Few studies are available on pharmacogenetics of sorafenib in mRCC and axitinib. In a study of 55 Japanese mRCC patients, Tsuchiya *et al.*⁴³ reported that the CC genotype of rs717620 in *ABCC2* compared to the CT genotype and the *HLA-A*24* allele were both associated with an increased risk for sorafenib-induced high-grade skin rash. In 2015, Escudier *et al.*⁴⁴ reported a set of 305 mRCC patients (N=159 axitinib and N=146 sorafenib), in which 15 SNPs were investigated in *VEGF-A*, *VEGF-R1*, *VEGF-R2*, and *HIF-1A* for the association with blood pressure or efficacy outcomes. It was observed that no significant association was reported for any of the SNPs with treatment outcome on axitinib. For sorafenib treatment, it was found that SNP rs2071559 in *VEGF-R2* was associated with PFS and OS ($P<0.01$), but the sensitivity/specificity analysis on this SNP showed a result of <80%. Therefore, it cannot be considered as a predictive SNP for survival outcome on sorafenib in mRCC.⁴⁴

Other associated biomarkers with TKI treatment outcome in mRCC patients

In addition to SNPs as potential biomarkers, many other biomarkers including histological subtypes, molecular subtypes, protein expression, immuno-expression, and microRNA were evaluated for potential association with TKI treatment outcome. Results are presented in **Supplementary document 3**.⁴⁵⁻⁷²

Histological and molecular subtypes in sunitinib treatment

In a case report by Choueiri *et al.*⁴⁵, a 49-year-old woman initially diagnosed with clear cell mRCC was later confirmed to have an Xp11.2 translocation RCC histological subtype which generally involves the *TFE3* gene. This patient developed severe HFS on sorafenib and, therefore, switched to sunitinib on which she showed a partial response followed by a durable response for more than 2 years.⁴⁵ This could indicate that the Xp11.2 translocation histological subtype is a biomarker that predicts a more favorable efficacy and toxicity profile on sunitinib compared to sorafenib. In a recent communication, Beuselinck *et al.*⁴⁶ showed that molecular subtypes of clear cell mRCC can predict the response to sunitinib with regard to response rate, PFS, and OS. These tumor subtypes are characterized based on chromosome copy-number aberrations, methylation status, and gene mutations in von Hippel-Lindau and the polybromo-1 protein (PBRM1).⁴⁶

Protein expression and immunoexpression in sunitinib treatment

With regard to protein expression, Paule *et al.*⁴⁷ investigated molecular biomarkers in primary tumors of 23 mRCC patients. Herein, levels of VEGF soluble isoforms VEGF121 and VEGF165 were associated with sunitinib response, and the ratio of VEGF121/VEGF165 was associated with prognosis. Gruenwald *et al.*⁴⁸ investigated another type of soluble markers and reported circulating endothelial cells (CEC) to be elevated within 28 days after start of sunitinib treatment in patients with a PFS above the median.⁴⁸ Later, in 2012, expression levels of the chemokine receptor CXCR4 were associated with objective response rate and survival on sunitinib as communicated by D'Alterio *et al.*⁴⁹ In a report of Garcia-Donas *et al.*⁵⁰, protein expression was investigated in 67 sunitinib-treated mRCC patients. A high expression of HIF2A and PDGFRB proteins showed an improved response to sunitinib, and VEGFR3 expression was associated with PFS. Furthermore, expression of HIF2A and VEGF-A or EGLN3 mRNA content were associated with OS.⁵⁰ Minardi *et al.*⁵¹ showed that VEGF expression was correlated with distant metastasis-free survival (DMFS) and OS in mRCC patients, but showed no association with regard to sunitinib treatment.⁵¹ In a set of 50 Japanese RCC patients, Sato *et al.*⁵² investigated the Extracellular Matrix MetalloProteinase Inducer (EMMPRIN), a cell-surface glycoprotein that belongs to the immunoglobulin superfamily encoded by a gene localized to 19p13.3. High expression

of EMMPRIN was seen for sunitinib-treated patients and sunitinib-resistant cells.⁵² In the above-mentioned study of Motzer *et al.*³³, baseline levels of angiopoietin-2 and matrix metalloproteinase were associated with tumor response, and HIF-1 α expression was associated with PFS.³³ The expression of phosphorylated VEGF-R2 was associated with PFS and OS in mRCC patients by Del Puerto-Nevado *et al.*⁵³ Gambini *et al.*⁵⁴ reported a case of a 60-year-old man with complete remission after 1 year on sunitinib. For this patient, a significant expression of neuron-specific enolase was observed together with the presence of pancreatic metastases. The hypothesis is that these neuroendocrine markers could predict the response to TKIs.⁵⁴

Considering immunoexpression, Mikami *et al.*⁵⁵ studied a subset of 25 Japanese mRCC patients on sunitinib. Herein, a high expression of the cancer stem cell marker CD44 was associated with a poor treatment outcome and responsible for developing sunitinib resistance. Also, the combination of high expression of TNF- α and CD44 was associated with a decreased PFS.⁵⁵ Dornbusch *et al.*⁵⁶ evaluated the immunoexpression of tumor protein markers in 42 patients with mRCC. They concluded that HIF-1 α , CA9, Ki67, CD31, pVEGFR1, VEGFR1 and -2, pPDGFR- α and - β were associated with the response to sunitinib ($P < 0.05$).⁵⁶

Micro RNA in sunitinib treatment

Some studies focused on microRNA (miRNA) as a possible predictor for sunitinib treatment outcome. miRNAs are noncoding RNAs that regulate gene expression.⁵⁷ Gámez-Pozo *et al.*⁵⁷ associated the expression of miRNA combinations (e.g. hsa-miR-141, hsa-miR-31, hsa-miR-125a-5p) in peripheral blood of RCC patients with poor or prolonged response to sunitinib.⁵⁷ Also, Berkers *et al.*⁵⁸ showed that miRNA was related to the response to sunitinib as tested in 20 mRCC patients: a significantly lower expression of miRNA-141 was observed for poor responders.⁵⁸ On miRNA expression of miRNA, Prior *et al.*⁵⁹ identified miRNA-942, miR-628-5p, miR-133a, and miR-484 as predictors for sunitinib efficacy (i.e. time to progression, OS, and resistance). It was postulated that high levels of miRNA-942 will stimulate the secretion of matrix metalloproteinase-9 and VEGF and thereby effectuate resistance to sunitinib.⁵⁹

Pazopanib

Other than SNPs, our literature search identified no associations of biomarkers with pazopanib treatment outcome in mRCC. Choueiri *et al.*⁶⁰ studied biomarkers along the VHL/HIF-1 α /HIF-2 α axis but reported no significant associations with pazopanib activity in patients with advanced clear-cell RCC.⁶⁰

Sorafenib

Mean VEGF plasma levels showed a significant increase in response to sorafenib, while plasma proteins VEGF-R2 and tissue inhibitor of metalloproteinase 1 (TIMP-1) were decreased in response to sorafenib as was observed by Peña *et al.*⁶¹ Feng *et al.*⁶² showed that high levels of circulating cell-free DNA (cfDNA) were associated with a poor response to sorafenib in 18 Chinese mRCC patients. Aziz *et al.*⁶³ reported that a high microvessel area was associated with a better response on sorafenib; smaller primary tumors were observed ($P=0.005$).⁶³ Another study showed that miRNA-30a is predictive for sorafenib efficacy.⁶⁴ Zheng *et al.*⁶⁴ explain that miRNA-30a inhibits its targeted gene Beclin-1, and consequently, the cytotoxic potential of sorafenib will increase.⁶⁴

Potential biomarkers from analyses including patients using different therapies

Some of the included articles evaluate more types of TKIs in the same analysis or in combination with other therapies in mRCC. In a cohort of 282 mRCC patients, Doberstein *et al.* showed that the expression of the L1 cell adhesion molecule was predictive both for OS and for therapy resistance in mRCC treatment with either sunitinib, rapamycin, or cisplatin.⁶⁵ Furuya *et al.*⁶⁶ studied 66 Japanese patients (26 received IFN-alpha and 21 patients received concomitant sorafenib as a second-line treatment) in which significantly higher serum levels of the IFN-alpha receptor messenger RNA were seen for patients with a better response and longer OS on IFN-alpha treatment with or without the addition of sorafenib.⁶⁶ Sharpe *et al.*⁶⁷ observed a significant reduction in vessel density (CD31), phospho-S6K expression, PDL-1 expression, and FOXP3 expression, but also a significant increase in cytoplasmic FGF-2, MET receptor expression in vessels, Fuhrman tumor grade, and Ki-67 in 85 patients who use either sunitinib or pazopanib. In addition, progression occurred later in patients with high levels of Ki-67 and CD31.⁶⁷

Yamada *et al.*⁶⁸ studied 27 cytokines as possible predictive biomarkers for TKI treatment in 13 mRCC patients treated with sunitinib (N=8), axitinib (N=4), or sorafenib (N=1). The plasma granulocyte macrophage colony-stimulating factor level was higher in patients who showed a partial response compared to patients with stable disease or progressive disease ($P<0.05$). Although not significant, higher IL-6 levels were observed in patients with progressive disease ($P=0.141$).⁶⁸ Interestingly, Yamada *et al.* refer to a paper of Tran *et al.*⁶⁹ who describe that for patients receiving pazopanib, high IL-6 was associated with an improved PFS compared to placebo ($P=0.009$).⁶⁹

Peters *et al.*⁷⁰ examined epigenetic biomarkers, namely the methylation status in primary tumor tissues in a cohort of 18 patients treated with either sunitinib (N=12), sorafenib (N=4), axitinib (N=1), or bevacizumab (N=1), as a first-line treatment. A decreased PFS and OS were observed for the hyper methylated status of CST6 and LAD1.

For the prediction of therapy failure on first-line therapy, sensitivity and specificity were determined and resulted in a specificity of 1.0 and a sensitivity of 0.73 for LAD1 methylation.⁷⁰ Dubrowskaja *et al.*⁷¹ of the same research group studied the same patient cohort and showed that a higher methylation status of the Neurofilament Heavy polypeptid gene was related to a decrease in OS.⁷¹

Finally, Choudhury *et al.*⁷² investigated 48 patients on molecular subtypes of clear-cell RCC for their association with response to either sunitinib or pazopanib as a first-, second-, or third-line treatment. Using a multigene assay (8 genes: CXCL5, EFNA5, EMCN, LAMB3, PLG, PRAME, RARRES1, SLC6A19), clinical prognostic subtypes were identified, which were shown to be predictive for response and survival on TKI treatment.⁷²

DISCUSSION

This systematic review outlines the available literature on associations of SNPs and other biomarkers with clinical treatment outcome of TKIs in mRCC. The most promising biomarkers on predicting TKI outcome are related to PK (genes encoding CYP450 enzymes, CYP450 regulators, or drug efflux transporters) and PD (genes encoding VEGF or VEGF receptors and interleukins).¹⁹⁻⁴⁴ For sunitinib, SNPs in *CYP1A1*, *CYP3A5*, *CYP3A4*, *NR1I2*, *NR1I3*, *ABCB1*, *ABCG2*, *VEGF-A*, *VEGFR1*, *VEGFR2*, *VEGF-R3*, *FGF-R2*, *FLT3*, *eNOS*, *UGT1A1*, and *IL8* were related to toxicity, efficacy, clearance, or drug exposure. Moreover, the potential of SNPs in *CYP3A5* and *ABCB1* to predict sunitinib related outcomes was confirmed in a large set of patients. With regard to pazopanib, SNPs in *UGT1A1*, *CYP1A2*, and *HFE* were associated with either bilirubin levels or ALT levels, and SNPs in *IL8*, *HIF1A*, *NR1I2*, and *VEGFA* were associated with efficacy outcomes on pazopanib. The association of rs1126647 in *IL8* with OS was confirmed in two independent data sets including patients treated with either sunitinib or pazopanib.^{40,42} Sorafenib toxicity and efficacy were related to genetic polymorphisms in *ABCC2*, *HLA-A*, and the only large study (N>300) available on sorafenib treatment showed that rs2071559 in *VEGF-R2* was related to efficacy. For axitinib, no SNP associations were reported.¹⁹⁻⁴⁴ Other reported biomarkers include proteins (e.g. VEGF), epigenetic markers (miRNA or methylation), and histological or molecular subtypes.⁴⁵⁻⁷² Clinical implementation of these biomarkers remains a difficult issue, since most studies present nonvalidated findings from retrospective cohort studies, effect sizes are rather small, heterogeneity between studies is large, sample sizes are limited, and above all, no consensus has been reached on the interpretation of these findings in terms of clinical utility on guiding individual dosing regimens of TKIs.^{17,18}

Our systematic approach minimizes the risk that we have missed relevant articles. Moreover, we only focused on TKIs (excluding other mRCC treatments) to ensure

that similar mechanisms and drug targets are compared. A meta-analysis is generally preferred to easily summarize the findings of all studies and subsequently value the clinical utility of associated biomarkers. However, in this case, a meta-analysis would be misjudging because of the large heterogeneity in study design, sample sizes, ethnicities, study endpoints, tested SNPs, evaluated treatments, and statistical approach. Sample sizes range from 1 to 451, in which the number of tested SNPs varied between 1 and 6852 SNPs.^{22,39} Furthermore, publication bias has very likely occurred and obscures a clear conclusion from a meta-analysis. In this review, we compared study results on the drugs of interest, sample sizes, ethnicity differences, and effect sizes with confidence intervals and *P*-values. One of the limitations in the studies in this field is that biomarkers are typically selected with a significance threshold value of $P < 0.05$. However, often a large number of hypotheses are tested in the same data set, increasing the chance for false-positive associations.⁷³ Another important issue is the difference in allele frequencies of SNPs among ethnicities. For example, SNP allele frequencies in *CYP3A5* and *ABCB1* differ between Caucasian and Asian patients. Furthermore, co-medication during TKI treatment and total exposure to the TKI was not taken into account in most studies, while drug-drug interactions may be especially important with regard to the effects on PK and PD. These matters have an important impact on the clinical validity and should be taken into account before assessing the possibilities on clinical implementation.

The SNPs tested in most of the included studies were selected according to the candidate gene approach. On PK, the results show that SNPs in genes coding for CYP enzymes, CYP regulators, and drug efflux transporters (*ABCB1*, *ABCC2*, and *ABCG2*) seem to play an important role in sunitinib treatment outcome, especially *CYP3A5* and *ABCB1* SNPs.^{19-21,29,31,34,37} In some of the articles on sunitinib, the intermediate endpoints clearance and exposure were investigated and confirmed the hypotheses that SNPs in *CYP3A4*, *CYP3A5*, and *ABCB1* influence drug exposure (**Figure 2**).^{22,29,36,37} Sunitinib and pazopanib are metabolized by CYP3A4 and probably CYP3A5 but are not known to be metabolized by UGT1A1.^{38,41,74} Pazopanib inhibits UGT1A1 in vitro. In pazopanib-treated patients, SNPs in *UGT1A1* may cause a reduced expression of UGT1A1 and, together with the inhibitory effect of pazopanib, would result in hyperbilirubinemia. For sunitinib, however, this enhanced effect on inhibition does not apply.^{38,41} Since sunitinib and pazopanib are both metabolized by the same CYP enzymes and are substrates for *ABCB1* and *ABCG2*, one could expect to find the same SNPs in these PK-related genes to be related to pazopanib outcomes. Only for a SNP in *NR1I2*, a relation with response rate on pazopanib was observed.⁴⁰ The articles in this review have not tested pazopanib toxicity outcomes for these SNPs related to PK. However, the effects on total drug exposure resulting from changes in CYP enzyme activity may be very different for sunitinib and pazopanib. Including both sunitinib- and pazopanib-treated patients in this type of

analyses is not very reasonable^{41,42}, since sunitinib is converted to an active metabolite with similar activity and potency and a longer half-life, while pazopanib only has less active metabolites. In addition, the extent of absorption for pazopanib and sunitinib may be very different (due to substrate specificity, for example), and its role in the total PK of the TKI is unclear. Changes in genes encoding efflux transporters can, therefore, be more important for sunitinib than for pazopanib. Regarding pharmacodynamics of sunitinib, SNPs in *eNOS*, *IL8*, *VEGF-A*, and *VEGF-R1,2,3* present promising biomarkers for toxicity or survival on sunitinib.^{19,20,25,26,32,34,35} However, results on *VEGF-A* and *VEGF-R1,2* or 3 genes are contradictory for both toxicity and efficacy outcomes since also negative associations were reported.^{30,33} In a recent analysis of our group, we associated SNPs in *IL8* and *IL13* with sunitinib toxicities (hypertension, leukopenia, and any toxicity >grade 2) and referred to the findings of Xu *et al.*^{40,42,75} Here, it was carefully hypothesized that IL8 and VEGF can be related in their effect on sunitinib-induced toxicities.⁷⁵ For sorafenib, only one study with a significant sample size is available that reported a SNP in *VEGF-R2* to be a potential predictor for efficacy (PFS and OS).⁴⁴ This effect was explained by a change in the promoter, subsequently altering transcription factor-binding and thereby the expression of VEGF-R2, but no clear downstream effect was presented.⁴⁴ In addition to SNPs, other types of biomarkers are potentially important. Results on epigenetics (miRNA and DNA methylation status) may influence gene expression and SNP effects, and results on protein markers can provide useful insight into mechanisms involved in SNP effects.⁴⁵⁻⁷² The associated SNPs and other biomarkers are different between TKIs, and therefore, we cannot assume a class effect pharmacogenetics in mRCC. While many targets, metabolizing enzymes, and drug transporters of the TKIs are comparable, significant differences in mechanisms of action can still exist.

A randomized clinical trial (RCT) is considered the hallmark of evidence to show utility for a biomarker in a clinical setting. However, an RCT is not critical for clinical application. It has already been proven that genetic substudies of RCTs in cancer are sufficient evidence to implement pharmacogenetic testing as is illustrated by testing *EGFR* for erlotinib and (*K*)*RAS* for cetuximab and panitumumab.⁷⁶ Funakoshi *et al.*¹⁴ assessed the level of evidence (LOE) of associated biomarkers by arranging the articles into different categories.^{77,78} Categories A and B represent the highest LOE with prospective RCT addressing biomarker questions and prospective studies not primarily designed to address biomarker questions. Categories C and D represent the lowest LOE with prospective and retrospective observational studies.^{14,77,78} In this field, these categories would be a false premise since no randomized studies and few prospective studies (category A or B) are available on pharmacogenetic biomarker studies in mRCC, which means only category C and D studies with a low LOE are left.

Gillis *et al.*¹⁸ mention that both clinical validity and clinical utility should be considered in translating the findings into a practical method to improve clinical practice. There is a lack of consensus on the definition of clinical utility. However, Gillis *et al.* are confident that validation of the associated biomarker in an independent replication cohort is the best evidence of clinical utility.¹⁸ Janssens *et al.* consider a pharmacogenetic test useful when it is 'sufficiently predictive for clinically important phenotypes, contributes to improved health outcomes, and is economically feasible to implement in community-based practice settings'.¹⁷ Clinical application is very likely for the reported SNP associations of *CYP3A5* and *ABCB1*, which reflect a considerably high predictive value since these SNPs have been associated with sunitinib PK³⁷, associations on clinical outcome are confirmed^{20,30,31}, and these findings are further supported by the exposure-effect relationship on sunitinib treatment.⁷ However, no genetic biomarkers for clinical outcome on TKIs in mRCC have been implemented yet in any of the innovative guidelines such as the Clinical Pharmacogenetics Implementation Consortium (CPIC) and the Dutch Pharmacogenetics Working Group (DPWG).^{79,80}

In conclusion, a decade of pharmacogenomics research on TKIs in mRCC patients has revealed *CYP3A5*1* and *ABCB1* as the most promising biomarkers for sunitinib treatment outcome. Implementation of pre-emptive pharmacogenetic testing will become likely when confirmed in independent cohorts preferably in prospective studies.

In the near future, results from genome-wide association studies from discovery and validation sets are expected. Many more SNPs in different genes will be investigated in the genome-wide association studies as compared to the candidate-gene approach that is used in most of the currently available studies. This could mean that earlier discovered SNPs will be affirmed, but these results may also identify additional genetic variants as biomarkers for outcome of TKI treatment in renal cell cancer. Additionally, we think that it is essential to further study both immunological and angiogenic factors (both protein expression and genetic polymorphisms) on prognosis and anti-VEGF treatment outcome in mRCC. Besides the effect on VEGF inhibition, sunitinib and other TKIs may influence the immune system, but we need a better understanding on the mechanisms involved.^{42,69,75} Anti-angiogenic therapy may result in a further induction of the immunosuppressive environment of the RCC tumor.⁸¹ Future findings may indicate that we need a combination therapy to treat both the immunogenic and pro-angiogenic character of RCC. Utilizing the pharmacogenetic profile of the patient may optimize TKI selection. Currently, the drug arsenal for mRCC is growing with new available TKIs or TKIs in development, and new promising immune-based therapies for RCC treatment are on their way, such as anti-programmed death-ligand 1 (PD-L1).⁸² Recent findings showed that in some patients, the efficacy of anti-angiogenic therapy might improve by PD-L1 blockade. However, as Liu *et al.* mention: 'The top priority and challenge is the selection of patients who are likely to respond to anti-PD-L1 therapy'.⁸¹

Within 5 years from now, we expect that some of the presented biomarkers here will be validated in replication cohorts and prospective validation studies with a longitudinal observation in time and a systematic data collection. These can be designed as noninterventional because blood samples (DNA) can be collected during routine monitoring visits of the patients. In retrospect, it will be investigated whether the determined genotypes of the patients have been a prediction for the outcome of TKI treatment. Two large prospective, nonrandomized clinical trials on precision medicine in oncology have already been set up as presented at the 2015 American Society of Clinical Oncology (ASCO) annual meeting: ASCO's Targeted Agent and Profiling Utilization Registry (TAPUR) study and the National Cancer Institute (NCI) Molecular Analysis for Therapy Choice (NCI-MATCH).^{83,84} In these trials, more than 20 different targeted therapies will be investigated including sunitinib, and these drugs will be made freely available. For the NCI-MATCH study, a number of 3000 patients is anticipated. Patients are treated with a drug based on the molecular profile of the tumor, and important clinical outcomes (e.g. overall response rate and PFS) will be recorded to see how patients can benefit from these drugs.^{83,84}

REFERENCES

Papers of special note have been highlighted as:

• of interest

•• of considerable interest

- Kashyap MK, Kumar A, Emelianenko N, et al. Biochemical and molecular markers in renal cell carcinoma: an update and future prospects. *Biomarkers*. 2005;10(4):258-294.
- Rini BI, Escudier B, Tomczak P, et al. Comparative effectiveness of axitinib versus sorafenib in advanced renal cell carcinoma (AXIS): a randomised phase 3 trial. *Lancet*. 2011;378(9807):1931-1939.
- Motzer RJ, Hutson TE, Tomczak P, et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2009;27(22):3584-3590.
- Escudier B, Eisen T, Stadler WM, et al. Sorafenib in advanced clear cell renal-cell carcinoma. *N Engl J Med*. 2007;356(2):125-134.
- Houk BE, Bello CL, Kang D, et al. A population pharmacokinetic meta-analysis of sunitinib malate (SU11248) and its primary metabolite (SU12662) in healthy volunteers and oncology patients. *Clin Cancer Res*. 2009;15(7):2497-2506.
- Suttle AB, Ball HA, Molimard M, et al. Relationships between pazopanib exposure and clinical safety and efficacy in patients with advanced renal cell carcinoma. *Br J Cancer*. 2014;111(10):1909-1916.
- Bhargava P, Robinson MO. Development of second-generation VEGFR tyrosine kinase inhibitors: current status. *Curr Oncol Rep*. 2011;13(2):103-111.
- Choueiri TK, Pal SK, McDermott DF, et al. A phase I study of cabozantinib (XL184) in patients with renal cell cancer. *Ann Oncol*. 2014;25(8):1603-1608.
- Escudier B, Porta C, Bono P, et al. Randomized, controlled, doubleblind, cross-over trial assessing treatment preference for pazopanib versus sunitinib in patients with metastatic renal cell carcinoma: PISCES study. *J Clin Oncol*. 2014;32(14):1412-1418.
- Motzer RJ, Hutson TE, Cella D, et al. Pazopanib versus sunitinib in metastatic renal-cell carcinoma. *N Engl J Med*. 2013;369(8):722-731.
- García-Donas J, Rodríguez-Moreno JF, Romero-Laorden N, et al. Renal carcinoma pharmacogenomics and predictors of response: steps toward treatment individualization. *Urol Oncol*. 2015;33(4):179-186.
- Rodríguez-Antona C, Taron M. Pharmacogenomic biomarkers for personalized cancer treatment. *J Intern Med*. 2015;277(2):201-217.
- Erdem L, Giovannetti E, Leon LG, et al. Polymorphisms to predict outcome to the tyrosine kinase inhibitors gefitinib, erlotinib, sorafenib and sunitinib. *Curr Top Med Chem*. 2012;12(15):1649-1659.
- Funakoshi T, Lee CH, Hsieh JJ. A systematic review of predictive and prognostic biomarkers for VEGF-targeted therapy in renal cell carcinoma. *Cancer Treat Rev*. 2014;40(4):533-547.
- Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*. 2015;349:g7647.

16. International prospective register of systematic reviews [Internet]. [cited 2015 Nov 16]. URL: www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42015016509#VR5WIE39laQ.
17. Janssens A, Deverka P. Useless until proven effective: the clinical utility of preemptive pharmacogenetic testing. *Clin Pharmacol Ther.* 2014;96(6):652-654.
18. Gillis N, Innocenti F. Evidence required to demonstrate clinical utility of pharmacogenetic testing: the debate continues. *Clin Pharmacol Ther.* 2014;96(6):655-657.
19. Van Erp N, Eechoute K, van der Veldt AA, et al. Pharmacogenetic pathway analysis for determination of sunitinib-induced toxicity. *J Clin Oncol.* 2009;27(26):4406-4412.
 - This is the first pharmacogenetic study on sunitinib-induced toxicities.
20. Garcia-Donas J, Esteban E, Leandro-García L, et al. Single nucleotide polymorphism associations with response and toxic effects in patients with advanced renal-cell carcinoma treated with first-line sunitinib: a multicentre, observational, prospective study. *Lancet Oncol.* 2011;12(12):1143-1150.
21. Beuselinck B, Lambrechts D, Van Brussel T, et al. Efflux pump ABCB1 single nucleotide polymorphisms and dose reductions in patients with metastatic renal cell carcinoma treated with sunitinib. *Acta Oncol.* 2014;53(10):1413-1422.
 - This report underlines the importance of genetic polymorphisms in ABCB1 in sunitinib treatment.
22. Mizuno T, Terada T, Kamba T, et al. ABCG2 421C>A polymorphism and high exposure of sunitinib in a patient with renal cell carcinoma. *Ann Oncol.* 2010;21(6):1382-1383.
23. Mizuno T, Fukudo M, Terada T, et al. Impact of genetic variation in breast cancer resistance protein (BCRP/ABCG2) on sunitinib pharmacokinetics. *Drug Metab Pharmacokinet.* 2012;27(6):631-639.
24. Miura Y, Imamura C, Fukunaga K, et al. Sunitinib-induced severe toxicities in a Japanese patient with the ABCG2 421 AA genotype. *BMC Cancer.* 2014;14:964.
25. Eechoute K, van der Veldt AA, Oosting S, et al. Polymorphisms in endothelial nitric oxide synthase (eNOS) and vascular endothelial growth factor (VEGF) predict sunitinib-induced hypertension. *Clin Pharmacol Ther.* 2012;92(4):503-510.
26. Kim JJ, Vaziri SA, Rini BI et al. Association of VEGF and VEGFR2 single nucleotide polymorphisms with hypertension and clinical outcome in metastatic clear cell renal cell carcinoma patients treated with sunitinib. *Cancer.* 2012;118(7):1946-1954.
27. Kim H, Park H, Kwon W, et al. Pharmacogenetic determinants associated with sunitinib-induced toxicity and ethnic difference in Korean metastatic renal cell carcinoma patients. *Cancer Chemother Pharmacol.* 2013;72(4):825-835.
28. Takayoshi K, Sagara K, Uchino K, et al. A case of metastatic renal cell carcinoma and bile duct carcinoma treated with a combination of sunitinib and gemcitabine. *BMC Cancer.* 2015;15(1):426.
29. Teo Y, Wee H, Chue X, et al. Effect of the CYP3A5 and ABCB1 genotype on exposure, clinical response and manifestation of toxicities from sunitinib in Asian patients. *Pharmacogenomics J.* 2016;16(1):47-53.

30. Diekstra MH, Swen JJ, Boven E, et al. CYP3A5 and ABCB1 polymorphisms as predictors for sunitinib outcome in metastatic renal cell carcinoma. *Eur Urol.* 2015;68(4):621-629.
••This is the largest confirmatory study of pharmacogenetics of sunitinib in mRCC patients.
31. Van der Veldt AA, Eechoute K, Gelderblom H, et al. Genetic polymorphisms associated with a prolonged progression-free survival in patients with metastatic renal cell cancer treated with sunitinib. *Clin Cancer Res.* 2011;17(3):620-629.
32. Scartozzi M, Bianconi M, Faloppi L, et al. VEGF and VEGFR polymorphisms affect clinical outcome in advanced renal cell carcinoma patients receiving first-line sunitinib. *Br J Cancer.* 2013;108(5):1126-1132.
33. Motzer R, Hutson T, Hudes G, et al. Investigation of novel circulating proteins, germ line single-nucleotide polymorphisms, and molecular tumor markers as potential efficacy biomarkers of first-line sunitinib therapy for advanced renal cell carcinoma. *Cancer Chemother Pharmacol.* 2014;74(4):739-750.
•This is a large published study that reports nonsignificant relations between SNPs in VEGF-A and VEGF-R3 on sunitinib efficacy outcomes.
34. Beuselinck B, Karadimou A, Lambrechts D, et al. Single-nucleotide polymorphisms associated with outcome in metastatic renal cell carcinoma treated with sunitinib. *Br J Cancer.* 2013;108(4):887-900.
35. Beuselinck B, Karadimou A, Lambrechts D, et al. VEGFR1 single nucleotide polymorphisms associated with outcome in patients with metastatic renal cell carcinoma treated with sunitinib - a multicentric retrospective analysis. *Acta Oncol.* 2014;53(1):103-112.
36. Mizuno T, Fukudo M, Fukuda T, et al. The effect of ABCG2 genotype on the population pharmacokinetics of sunitinib in patients with renal cell carcinoma. *Ther Drug Monit.* 2014;36(3):310-316.
37. Diekstra MH, Klümpen HJ, Lolkema MP, et al. Association analysis of genetic polymorphisms in genes related to sunitinib pharmacokinetics, specifically clearance of sunitinib and SU12662. *Clin Pharmacol Ther.* 2014;96(1):81-89.
38. Xu CF, Reck B, Xue Z, et al. Pazopanib-induced hyperbilirubinemia is associated with Gilbert's syndrome UGT1A1 polymorphism. *Br J Cancer.* 2010;102(9):1371-1377.
39. Xu CF, Reck B, Goodman V, et al. Association of the hemochromatosis gene with pazopanib-induced transaminase elevation in renal cell carcinoma. *J Hepatol.* 2011;54(6):1237-1243.
40. Xu CF, Bing N, Ball H, et al. Pazopanib efficacy in renal cell carcinoma: evidence for predictive genetic markers in angiogenesis-related and exposure-related genes. *J Clin Oncol.* 2011;29(18):2557-2564.
41. Motzer R, Johnson T, Choueiri T, et al. Hyperbilirubinemia in pazopanib- or sunitinib-treated patients in COMPARZ is associated with UGT1A1 polymorphisms. *Ann Oncol.* 2013;24(11):2927-2928.
42. Xu CF, Johnson T, Garcia-Donas J, et al. IL8 polymorphisms and overall survival in pazopanib- or sunitinib-treated patients with renal cell carcinoma. *Br J Cancer.* 2015;112(7):1190-1198.
••This article reports on a SNP in IL8 that can be relevant for both sunitinib and pazopanib treatment in mRCC.

43. Tsuchiya N, Narita S, Inoue T, et al. Risk factors for sorafenib-induced high-grade skin rash in Japanese patients with advanced renal cell carcinoma. *Anticancer Drugs*. 2013;24(3):310-314.
44. Escudier B, Rini B, Motzer R, et al. Genotype correlations with blood pressure and efficacy from a randomized phase III trial of second-line axitinib versus sorafenib in metastatic renal cell carcinoma. *Clin Genitourin Cancer*. 2015;13(4):328-337.
• This is the largest and only study available on SNPs related to efficacy in sorafenib-treated mRCC patients.
45. Choueiri TK, Mosquera JM, Hirsch MS. A case of adult metastatic Xp11 translocation renal cell carcinoma treated successfully with sunitinib. *Clin Genitourin Cancer*. 2009;7(3):E93-E94.
46. Beuselinck B, Job S, Becht E, et al. Molecular subtypes of clear cell renal cell carcinoma are associated with sunitinib response in the metastatic setting. *Clin Cancer Res*. 2015;21(6):1329-1339.
47. Paule B, Bastien L, Deslandes E, et al. Soluble isoforms of vascular endothelial growth factor are predictors of response to sunitinib in metastatic renal cell carcinomas. *Plos One*. 2010;5(5):e10715.
48. Gruenewald V, Beutel G, Schuch-Jantsch S, et al. Circulating endothelial cells are an early predictor in renal cell carcinoma for tumor response to sunitinib. *BMC Cancer*. 2010;10:695.
49. D'Alterio C, Portella L, Ottaiano A, et al. High CXCR4 expression correlates with sunitinib poor response in metastatic renal cancer. *Curr Cancer Drug Targets*. 2012;12(6):693-702.
50. Garcia-Donas J, Leandro-García L, González Del Alba A, et al. Prospective study assessing hypoxia-related proteins as markers for the outcome of treatment with sunitinib in advanced clear-cell renal cell carcinoma. *Ann Oncol*. 2013;24(9):2409-2414.
51. Minardi D, Lucarini G, Santoni M, et al. VEGF expression and response to sunitinib in patients with metastatic clear cell renal cell carcinoma. *Anticancer Res*. 2013;33(11):5017-5022.
52. Sato M, Nakai Y, Nakata W, et al. EMMPRIN promotes angiogenesis, proliferation, invasion and resistance to sunitinib in renal cell carcinoma, and its level predicts patient outcome. *Plos One*. 2013;8(9):e74313.
53. Del Puerto-Nevado L, Rojo F, Zazo S, et al. Active angiogenesis in metastatic renal cell carcinoma predicts clinical benefit to sunitinib-based therapy. *Br J Cancer*. 2014;110(11):2700-2707.
54. Gambini D, Locatelli E, Gianelli U. Sunitinib-induced complete response in metastatic renal cancer expressing neuroendocrine markers: a new predictive factor? *Anticancer Res*. 2014;34(12):7361-7365.
55. Mikami S, Mizuno R, Kosaka T, et al. Expression of TNF- α and CD44 is implicated in poor prognosis, cancer cell invasion, metastasis and resistance to the sunitinib treatment in clear cell renal cell carcinomas. *Int J Cancer*. 2015;136(7):1504-1514.
56. Dornbusch J, Zacharis A, Meinhardt M, et al. Analyses of potential predictive markers and survival data for a response to sunitinib in patients with metastatic renal cell carcinoma. *Plos One*. 2013;8(9):e76386.

57. Gámez-Pozo A, Antón-Aparicio L, Bayona C, et al. MicroRNA expression profiling of peripheral blood samples predicts resistance to first-line sunitinib in advanced renal cell carcinoma patients. *Neoplasia*. 2012;14(12):1144-1152.
58. Berkers J, Govaere O, Wolter P, et al. A possible role for microRNA-141 down-regulation in sunitinib resistant metastatic clear cell renal cell carcinoma through induction of epithelial-to-mesenchymal transition and hypoxia resistance. *J Urol*. 2013;189(5):1930-1938.
59. Prior C, Perez-Gracia J, Garcia-Donas J, et al. Identification of tissue microRNAs predictive of sunitinib activity in patients with metastatic renal cell carcinoma. *Plos One*. 2014;9(1):e86263.
60. Choueiri T, Fay A, Gagnon R, et al. The role of aberrant VHL/HIF pathway elements in predicting clinical outcome to pazopanib therapy in patients with metastatic clear-cell renal cell carcinoma. *Clin Cancer Res*. 2013;19(18):5218-5226.
61. Peña C, Lathia C, Shan M, et al. Biomarkers predicting outcome in patients with advanced renal cell carcinoma: results from sorafenib phase III treatment approaches in renal cancer global evaluation trial. *Clin Cancer Res*. 2010;16(19):4853-4863.
62. Feng G, Ye X, Fang F, et al. Quantification of plasma cell-free DNA in predicting therapeutic efficacy of sorafenib on metastatic clear cell renal cell carcinoma. *Dis Markers*. 2013;34(2):105-111.
63. Aziz S, Sznoł J, Albiges L, et al. Microvessel area as a predictor of sorafenib response in metastatic renal cell carcinoma. *Cancer Cell Int*. 2014;14(1):4.
64. Zheng B, Zhu H, Gu D, et al. MiRNA-30a-mediated autophagy inhibition sensitizes renal cell carcinoma cells to sorafenib. *Biochem Biophys Res Commun*. 2015;459(2):234-239.
65. Doberstein K, Wieland A, Lee S, et al. L1-CAM expression in ccRCC correlates with shorter patients survival times and confers chemoresistance in renal cell carcinoma cells. *Carcinogenesis*. 2011;32(3):262-270.
66. Furuya N, Kamai T, Shirataki H, et al. Serum interferon alpha receptor 2 mRNA may predict efficacy of interferon alpha with/without low-dose sorafenib for metastatic clear cell renal cell carcinoma. *Cancer Immunol Immunother*. 2011;60(6):793-808.
67. Sharpe K, Stewart G, Mackay A, et al. The effect of VEGF-targeted therapy on biomarker expression in sequential tissue from patients with metastatic clear cell renal cancer. *Clin Cancer Res*. 2013;19(24):6924-6934.
68. Yamada D, Matsushita H, Azuma T, et al. Granulocyte macrophage colony-stimulating factor as a predictor of the response of metastatic renal cell carcinoma to tyrosine kinase inhibitor therapy. *Mol Clin Oncol*. 2014;2(6):1023-1027.
69. Tran HT, Liu Y, Zurita AJ, et al. Prognostic or predictive plasma cytokines and angiogenic factors for patients treated with pazopanib for metastatic renal-cell cancer: a retrospective analysis of phase 2 and phase 3 trials. *Lancet Oncol*. 2012;13(8):827-837.
70. Peters I, Dubrowskaja N, Abbas M, et al. DNA methylation biomarkers predict progression-free and overall survival of metastatic renal cell cancer (mRCC) treated with antiangiogenic therapies. *Plos One*. 2014;9(3):e91440.

71. Dubrowskaja N, Gebauer K, Peters I, et al. Neurofilament Heavy polypeptide CpG island methylation associates with prognosis of renal cell carcinoma and prediction of antivasular endothelial growth factor therapy response. *Cancer Med.* 2014;3(2):300-309.
72. Choudhury Y, Wei X, Chu Y, et al. A multigene assay identifying distinct prognostic subtypes of clear cell renal cell carcinoma with differential response to tyrosine kinase inhibition. *Eur Urol.* 2015;67(1):17-20.
73. Maitland M, Ratain M, Cox N. Interpreting P values in pharmacogenetic studies: a call for process and perspective. *J Clin Oncol.* 2007;25(29):4513-4515.
74. Van Erp NP, Gelderblom H, Guchelaar HJ. Clinical pharmacokinetics of tyrosine kinase inhibitors. *Cancer Treat Rev.* 2009;35(8):692-706.
75. Diekstra MH, Liu X, Swen JJ, et al. Association of single nucleotide polymorphisms in IL8 and IL13 with sunitinib-induced toxicity in patients with metastatic renal cell carcinoma. *Eur J Clin Pharmacol.* 2015;71(12):1477-1484.
76. Voora D. Building the evidentiary framework for pharmacogenetic testing: is it time to move beyond randomized controlled trials? *Personalized Medicine.* 2013;10(1):1-3.
77. Hayes D, Bast R, Desch C, et al. Tumor marker utility grading system: a framework to evaluate clinical utility of tumor markers. *J Natl Cancer Inst.* 1996;88(20):1456-1466.
78. Simon R, Paik S, Hayes D. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst.* 2009;101(21):1446-1452.
79. Caudle KE, Klein TE, Hoffman JM, et al. Incorporation of pharmacogenomics into routine clinical practice: the Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline development process. *Curr Drug Metab.* 2014;15(2):209-217.
80. The Dutch Pharmacogenetics Working Group [Internet]. [cited 2015 Nov 23]. URL: <https://www.pharmgkb.org/page/dpwg>.
81. Liu X, Hoang A, Zhou L, et al. Resistance to anti-angiogenic therapy is associated with an immunosuppressive tumor microenvironment in metastatic renal cell carcinoma. *Cancer Immunol Res.* 2015;3(9):1017-1029.
82. Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Eng J Med.* 2015;373(19):1803-1813.
83. The Targeted Agent and Profiling Utilization Registry (TAPUR) study [Internet]. [cited 2015 Nov 23]. URL: <http://www.asco.org/practice-research/targeted-agent-and-profiling-utilization-registrystudy>.
84. NCI trial on Molecular Analysis for Therapy Choice (NCI-MATCH) [Internet]. [cited 2015 Nov 23]. URL: <http://www.cancer.gov/about-cancer/treatment/clinical-trials/nci-supported/ncimatch>.

SUPPLEMENTARY MATERIAL

Supplementary document 1: Systematic review protocol

1. TITLE

Systematic review: a decade of pharmacogenetic research on TKIs in metastatic renal cell carcinoma

2. REGISTRATION

PROSPERO registration number: CRD42015016509

3. AUTHORS

Corresponding Author: Henk-Jan Guchelaar, PharmD PhD, (H.J.Guchelaar@lumc.nl)

Author Affiliations: Leiden University Medical Center, Dept. of Clinical Pharmacy and Toxicology, Albinusdreef 2, 2333ZA, Leiden, The Netherlands. Postal address: P.O. Box 9600, 2300 RC Leiden, Post zone LO-P, Phone: +31 (0)71 52 64 018, Fax: +31(0)71 52 66 866

Email: Meta H.M. Diekstra M.H.M.Diekstra@lumc.nl, Jesse J. Swen J.J.Swen@lumc.nl, Hans Gelderblom A.J.Gelderblom@lumc.nl, Henk-Jan Guchelaar H.J.Guchelaar@lumc.nl

Contributions: MD and JS drafted the manuscript. Jan Schoones developed the search strategy. All authors read, provided feedback and approved the final manuscript

4. AMENDMENTS

In the event of protocol amendments, the date of each amendment will be accompanied by a description of the change and the rationale

5. SUPPORT

MD is supported by the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 259939.

6. RATIONALE

INTRODUCTION

In the last ten years a number of targeted agents for treatment of solid tumors have become available. Sunitinib, a tyrosine kinase inhibitor (TKI) was the first targeted drug for mRCC and is on the market since 2006. While pazopanib, axitinib and sorafenib have since then become available, sunitinib is still the first-choice treatment for metastatic renal cell carcinoma (mRCC).

Patients respond very differently to sunitinib with sometimes severe toxicities that need dose reductions and/or a poor response with regard to tumor shrinkage. Therefore, we are not yet able to predict the individual treatment outcome on sunitinib.

Previous studies identified Single Nucleotide Polymorphisms (SNPs) in genes encoding enzymes or transporters related to pharmacokinetics and pharmacodynamics of TKIs which have been associated with toxicity and efficacy of sunitinib. These genetic biomarkers are potential predictors for TKI treatment outcome in mRCC patients. Ideally, pharmacogenetic testing of genetic biomarkers prior to start of therapy, could result in giving the individual patient the most effective TKI with the least possible side effects. However, findings on these genetic biomarkers have not been validated yet. This means clinical implementation of pharmacogenetic testing to guide sunitinib dosing or other TKIs is not yet possible.¹⁻¹⁴

We perform a new systematic review up to January 2015 in order to have a more complete overview of all pharmacogenetic studies on sunitinib and other TKIs in mRCC to date and we only focus on genetic predictive biomarkers.

7. OBJECTIVES

The objective of this review is to systematically review the research findings on pharmacogenetics of sunitinib in mRCC to date and to evaluate the use of genetic biomarkers for sunitinib treatment outcome. Therefore, we try to answer the following questions:

What are the germline genetic variants that associate with sunitinib and other targeted agents (pazopanib, axitinib, sorafenib) and their efficacy and toxicity in mRCC patients?

What is the value and quality of the included studies in our review?

Based on the available research findings, what has been the result with regard to clinical implementation of predictive biomarkers for sunitinib treatment outcome up till now?

What should be the focus of future research in this field?

Supplementary document 1: Continued

8. ELIGIBILITY CRITERIA

Studies will be selected according to the criteria outlined below:

We will include studies with patients diagnosed with (metastatic) renal cell carcinoma (mRCC) receiving a targeted therapy (sunitinib, axitinib, sorafenib, pazopanib or other). In addition, we included studies that test the association of SNPs, any other genetic variants or gene/protein expression with the treatment outcome of these targeted therapies. Therefore, information that has to be available in the studies are DNA samples, genotyping data together with outcome data which are toxicity and efficacy data.

References were eligible when genetic biomarkers (i.e. SNPs, genetic variants, gene/protein expression) were investigated in relation to TKI treatment and mRCC. Excluded from our selection were review articles, preclinical studies (animal models and in vitro studies), phase I or II clinical trials and papers discussing pathology, prognosis, tumor biology or somatic mutations (e.g. Von Hippel-Lindau mutations). Accordingly, articles were excluded from our review if non-treatment related biomarkers were studied or if no TKI response was assessed, because these generally consider prognostic biomarkers.

Our literature search was done for articles that appeared between January 2004 and January 2015. Only English language papers will be included and conference abstracts are excluded from our search.

Excluded from our selection were review articles, preclinical studies (animal models and in vitro studies), phase I or II clinical trials and papers discussing pathology, prognosis or tumor biology, somatic mutations (e.g. VHL).

We have made sure that all authors of the review team agreed with the eligibility criteria prior to the selection process.

9. INFORMATION SOURCES

Search for references in the following databases:

PubMed

Embase (OVID version)

Web of Science

COCHRANE

CENTRAL

10. SEARCH STRATEGY

Together with a trained librarian, we will perform a search in PubMed, Embase (OVID version), Web of Science, COCHRANE Library and CENTRAL. The search consists of the combination of three subjects:

Renal cell carcinoma

Tyrosine kinase inhibitors (e.g. sunitinib, sorafenib)

Pharmacogenetics

The query will be applied in all databases taking into account the terminological and technical differences between these databases. Various synonyms and related terms for all subjects were used. Results will be limited to articles in the English language, and from the year 2004 onwards. Meeting abstracts are being excluded. See **Supplementary document 2: search strategy**

11. STUDY RECORDS**11a. DATA MANAGEMENT**

Results from our literature search will be uploaded to EndNote X6 and references will be de-duplicated with every new insertion of search results from one of the databases. The selection of articles will be made within EndNote in order to easily check which articles are excluded. An overview of our selection including the data collection will be processed in an excel file and eventually presented in tables in the manuscript.

11b. SELECTION PROCESS

First of all, papers will be selected by title. Titles will be judged on the topic of renal cell carcinoma, tyrosine kinase inhibitors and pharmacogenetics. The first two authors of the review (MD and JS) will screen the titles from the first original search independently from each other. Disagreement on the selection by title will be resolved through discussion with the last author on the review (HJG).

Secondly, a selection will be made by abstract with the included articles from the first selection by title. Abstracts will be judged on the topic of renal cell carcinoma, tyrosine kinase inhibitors and pharmacogenetics. The first two authors of the review (MD and JS) will screen the abstracts from papers included in the first selection step independently from each other. Disagreement on the selection by title will be resolved through discussion with the last author on the review (HJG).

Thirdly, the final aim is to get a total of roughly 80 references. Full-text will be judged on the topic of renal cell carcinoma, tyrosine kinase inhibitors and pharmacogenetics. The first two authors of the review (MD and JS) will screen the full texts papers included in the second selection step independently from each other. Disagreement on the selection by title will be resolved through discussion with the last author on the review (HJG).

For every selection step, the reason for exclusion of the paper will be recorded. An overview of our selection criteria and selection process will be saved in an excel file.

Supplementary document 1: Continued**11c. DATA COLLECTION PROCESS**

During the selection process, we will use excel-based forms to make a selection of articles. With the final selection of articles we will again use the excel file to extract relevant data to include in our review results.

12. DATA ITEMS

For the selection by title, abstract and full text of articles, we will make a selection based on the eligibility criteria mentioned in item 8. In addition, the excel file contains the reasons for exclusion.

The last separate worksheet is used for data extraction from the final selection of articles. Extracted data were: first author, year of publication, demographics (median age, number of patients, male percentage, ethnicity), sample material and origin (i.e. germline DNA from blood, serum or other), methods, tumor type, drug information (i.e. drug name, treatment line, dose), evaluated biomarkers, tested endpoints and statistical approach. The exact extraction data can be found in the excel file in the last worksheet.

13. OUTCOMES AND PRIORITISATION

End points to be tested for association with genetic variants will be toxicity and efficacy endpoints:

Progression-free Survival (PFS)

Overall Survival (OS)

Any type of toxicity (e.g. thrombocytopenia, leukopenia and many more) with an explanation on clinically relevant toxicities on severity grades.

Response to treatment (RECIST criteria; PD, SD, PR, CR)

Dose reductions

Any other endpoint related to toxicity or efficacy of the TKI treatment

14. RISK OF BIAS INDIVIDUAL STUDIES

Clinical validity (the robustness of the statistical association between a pharmacogenetic variant and outcome of drug therapy) and clinical utility (a test's health-care value) of the included articles was evaluated in a descriptive manner.^{17,18}

15. DATA SYNTHESIS**15a. CRITERIA FOR SYNTHESIS STUDY DATA**

Because of the diversity in study populations, ethnicities, outcomes, number of SNPs tested and differences in statistical approach in the studies, we will not perform a meta-analysis.

15b. PLANNED SUMMARY MEASURES

N/A

15c. ADDITIONAL ANALYSES

N/A

15d. PLANNED TYPE OF SUMMARY

We will visualize the study quality by using the NOS star system as described in item 14 and present the difference between studies in a more descriptive/qualitative way.

16. META BIAS(ES)

Meta bias was evaluated in a descriptive manner.

17. CONFIDENCE IN CUMULATIVE EVIDENCE

Clinical validity (the robustness of the statistical association between a pharmacogenetic variant and outcome of drug therapy) and clinical utility (a test's health-care value) of the included articles was evaluated in a descriptive manner.^{17,18}

Supplementary document 2: Search strategy

Research question:

pharmacogenetics of sunitinib in metastatic renal cell carcinoma

Search strategy

Together with a trained librarian, we performed a search in PubMed, Embase (OVID version), Web of Science, COCHRANE Library and CENTRAL. The search consisted of the combination of three subjects:

Renal cell carcinoma

Tyrosine kinase inhibitors (e.g. sunitinib, sorafenib)

Pharmacogenetics

Search results

This search resulted in a total of 1511 references. After removing duplicates automatically, a total of 973 references were obtained d.d. 14-01-2015, originating from:

PubMed: 680 [RefID 1-680]

Embase: 662, of which 256 unique [RefID 681-936]

Web of Science: 129, of which 35 unique [RefID 937-971]

COCHRANE: 20, of which 2 unique [RefID 972-973]

CENTRAL: 20, of which 0 unique

A final amount of **968 unique references** were obtained after removing the following 5 extra duplicates manually:

Bielecka, 2014

Dabora, 2011

Joosten, 2015

Pecuchet, 2013

Wu, 2012

Supplementary document 2: Continued

Database	Search Strategy	Number of references	Number of unique references
PubMed	("Carcinoma, Renal Cell"[Mesh] OR "renal cell carcinoma"[all fields] OR "renal tumor"[all fields] OR "renal tumour"[all fields] OR "renal tumors"[all fields] OR "renal tumours"[all fields] OR "RCC"[all fields] OR "mRCC"[all fields] OR "kidney cell carcinoma"[all fields] OR "kidney tumor"[all fields] OR "kidney tumour"[all fields] OR "kidney tumors"[all fields] OR "kidney tumours"[all fields] OR "renal cell adenocarcinoma"[all fields] OR "kidney cell adenocarcinoma"[all fields] OR "Grawitz"[all fields]) AND ("Genetic Variation"[Mesh] OR "genetic"[all fields] OR "genetics"[all fields] OR "genomics"[all fields] OR "genomic"[all fields] OR "genome"[all fields] OR "mutation"[all fields] OR "mutations"[all fields] OR "allelic"[all fields] OR "Allele"[all fields] OR "alleles"[all fields] OR "DNA"[all fields] OR "gene"[all fields] OR "genes"[all fields] OR "chromosome"[all fields] OR "chromosomes"[all fields] OR "germ-line"[all fields] OR "germline"[all fields] OR "INDEL"[all fields] OR "insertion"[all fields] OR "deletion"[all fields] OR "inversion"[all fields] OR "polymorphism"[all fields] OR "polymorphisms"[all fields] OR "polymorfism"[all fields] OR "polymorfisms"[all fields] OR "Genomic Structural Variation"[all fields] OR "Genomic Structural Variations"[all fields] OR "DNA Copy Number Variations"[all fields] OR "DNA Copy Number Variation"[all fields] OR "SNP"[all fields] OR "SNPs"[all fields] OR "nucleotide"[all fields] OR "nucleotides"[all fields] OR "Genetics"[Mesh] OR "pharmacogenetic"[all fields] OR "pharmacogenetics"[all fields] OR "hapmap"[all fields] OR "Pharmacogenetics"[Mesh] OR toxicogenetic*[all fields] OR "predictive marker"[all fields] OR "predictive markers"[all fields] OR "predictive biomarkers"[all fields] OR "predictive biomarker"[all fields] OR "Genetic Markers"[Mesh] OR "Genetic Phenomena"[Mesh] OR "GWAS"[all fields] OR "Genome-Wide Association Study"[Mesh] OR "Genome-Wide Association"[all fields] OR "Genetic Association Studies"[Mesh]) AND ("sunitinib"[Supplementary Concept] OR sunitinib*[all fields] OR "Sutent"[all fields] OR "SU 11248"[all fields] OR "SU11248"[all fields] OR "SU-11248"[all fields] OR "Protein-Tyrosine Kinases/antagonists and inhibitors"[Mesh] OR "tyrosine kinase inhibitor"[all fields] OR "Tyrosine kinase inhibitors"[all fields] OR "TKI"[all fields] OR "SU011248"[all fields] OR "SU-011248"[all fields] OR "SU 011248"[all fields] OR "Receptors, Vascular Endothelial Growth Factor"[Mesh] OR "vascular endothelial growth factor receptor"[all fields] OR "VEGF"[all fields] OR "VEGFR"[all fields] OR "VEGF-R1"[all fields] OR "VEGF-R2"[all fields] OR "VEGF-R3"[all fields] OR "VEGFR1"[all fields] OR "VEGFR2"[all fields] OR "VEGFR3"[all fields] OR "PDGF"[all fields] OR "PDGFR"[all fields] OR "sorafenib"[Supplementary Concept] OR "sorafenib"[All fields] OR "nexavar"[All fields] OR "BAY 545-9085"[All fields] OR "BAY-545-9085"[All fields] OR "BAY 43-9006"[All fields] OR "BAY-43-9006"[All fields] OR "BAY 439006"[All fields] OR "BAY43-9006"[All fields] OR "BAY439006"[All fields] OR "pazopanib"[Supplementary Concept] OR "votrient"[All fields] OR "GW 786034B"[All fields] OR "GW786034B"[All fields] OR "GW-786034B"[All fields] OR "GW 786034"[All fields] OR "GW786034"[All fields] OR "GW-786034"[All fields] OR "axitinib"[Supplementary Concept] OR "inlyta"[All fields] OR "AG 013736"[All fields] OR "AG013736"[All fields] OR "AG-013736"[All fields] OR "Angiogenesis Inhibitors"[Mesh] OR "angiogenesis inhibitors"[All fields] OR "angiogenesis inhibitor"[All fields] OR "anti angiogenic"[All fields] OR "anti-angiogenic"[All fields] OR "antiangiogenic"[All fields] OR "anti angiogenesis"[All fields] OR "anti-angiogenesis"[All fields] OR "antiangiogenesis"[All fields]) AND english[la] AND ("2004/01/01"[PDAT] : "3000/12/31"[PDAT])	680	680

Supplementary document 2: Continued

Database	Search Strategy	Number of references	Number of unique references
Embase (OVID-version)	("Kidney Carcinoma"/ OR "renal cell carcinoma".ti,ab OR "renal tumor".ti,ab OR "renal tumour".ti,ab OR "renal tumors".ti,ab OR "renal tumours".ti,ab OR "RCC".ti,ab OR "mRCC".ti,ab OR "kidney cell carcinoma".ti,ab OR "kidney tumor".ti,ab OR "kidney tumour".ti,ab OR "kidney tumors".ti,ab OR "kidney tumours".ti,ab OR "renal cell adenocarcinoma".ti,ab OR "kidney cell adenocarcinoma".ti,ab OR "Grawitz".ti,ab) AND (*"Genetic Variability"/ OR "genetic".ti,ab OR "genetics".ti,ab OR "genomics".ti,ab OR "genomic".ti,ab OR "genome".ti,ab OR "mutation".ti,ab OR "mutations".ti,ab OR "allelic".ti,ab OR "Allele".ti,ab OR "alleles".ti,ab OR "DNA".ti,ab OR "gene".ti,ab OR "genes".ti,ab OR "chromosome".ti,ab OR "chromosomes".ti,ab OR "germ-line".ti,ab OR "germline".ti,ab OR "INDEL".ti,ab OR "insertion".ti,ab OR "deletion".ti,ab OR "inversion".ti,ab OR exp *genetic polymorphism/ OR "polymorphism".ti,ab OR "polymorphisms".ti,ab OR "polymorfism".ti,ab OR "polymorfisms".ti,ab OR "Genomic Structural Variation".ti,ab OR "Genomic Structural Variations".ti,ab OR *"Copy Number Variation"/ OR "DNA Copy Number Variations".ti,ab OR "DNA Copy Number Variation".ti,ab OR "SNP".ti,ab OR "SNPs".ti,ab OR "nucleotide".ti,ab OR "nucleotides".ti,ab OR exp *"Genetics"/ OR "pharmacogenetic".ti,ab OR "pharmacogenetics".ti,ab OR "hapmap".ti,ab OR exp *"Pharmacogenetics"/ OR toxicogenetic*.ti,ab OR "predictive marker".ti,ab OR "predictive markers".ti,ab OR "predictive biomarkers".ti,ab OR "predictive biomarker".ti,ab OR exp *"Genetic Marker"/ OR "GWAS".ti,ab OR "Genome-Wide Association".ti,ab OR exp *"Genetic Association"/) AND (*"sunitinib"/ OR sunitinib*.ti,ab OR "Sutent".ti,ab OR "SU 11248".ti,ab OR "SU11248".ti,ab OR "SU-11248".ti,ab OR exp *"protein tyrosine kinase inhibitor"/ OR "tyrosine kinase inhibitor".ti,ab OR "Tyrosine kinase inhibitors".ti,ab OR "TKI".ti,ab OR "SU011248".ti,ab OR "SU-011248".ti,ab OR "SU 011248".ti,ab OR *vasculotropin receptor/ OR "vascular endothelial growth factor receptor".ti,ab OR "VEGF".ti,ab OR "VEGFR".ti,ab OR "VEGF-R1".ti,ab OR "VEGF-R2".ti,ab OR "VEGF-R3".ti,ab OR "VEGFR1".ti,ab OR "VEGFR2".ti,ab OR "VEGFR3".ti,ab OR "PDGF".ti,ab OR "PDGFR".ti,ab OR *"sorafenib"/ OR "sorafenib".ti,ab OR "nexavar".ti,ab OR "BAY 545-9085".ti,ab OR "BAY-545-9085".ti,ab OR "BAY 43-9006".ti,ab OR "BAY-43-9006".ti,ab OR "BAY 439006".ti,ab OR "BAY43-9006".ti,ab OR "BAY439006".ti,ab OR *"pazopanib"/ OR "votrient".ti,ab OR "GW 786034B".ti,ab OR "GW786034B".ti,ab OR "GW-786034B".ti,ab OR "GW 786034".ti,ab OR "GW786034".ti,ab OR "GW-786034".ti,ab OR *"axitinib"/ OR "inlyta".ti,ab OR "AG 013736".ti,ab OR "AG013736".ti,ab OR "AG-013736".ti,ab OR exp *"Angiogenesis Inhibitor"/ OR "angiogenesis inhibitors".ti,ab OR "angiogenesis inhibitor".ti,ab OR "anti angiogenic".ti,ab OR "anti-angiogenic".ti,ab OR "antiangiogenic".ti,ab OR "anti angiogenesis".ti,ab OR "anti-angiogenesis".ti,ab OR "antiangiogenesis".ti,ab) NOT conference abstract.pt AND english.la AND (2004 OR 2005 OR 2006 OR 2007 OR 2008 OR 2009 OR 201*).yr	662	256

Supplementary document 2: Continued

Database	Search Strategy	Number of references	Number of unique references
Web of Science	(TI=(“Kidney Carcinoma” OR “renal cell carcinoma” OR “renal tumor” OR “renal tumour” OR “renal tumors” OR “renal tumours” OR “RCC” OR “mRCC” OR “kidney cell carcinoma” OR “kidney tumor” OR “kidney tumour” OR “kidney tumors” OR “kidney tumours” OR “renal cell adenocarcinoma” OR “kidney cell adenocarcinoma” OR “Grawitz”) AND TS=(“Genetic Variability” OR “genetic” OR “genetics” OR “genomics” OR “genomic” OR “genome” OR “mutation” OR “mutations” OR “allelic” OR “Allele” OR “alleles” OR “DNA” OR “gene” OR “genes” OR “chromosome” OR “chromosomes” OR “germ-line” OR “germline” OR “INDEL” OR “insertion” OR “deletion” OR “inversion” OR genetic polymorphism OR “polymorphism” OR “polymorphisms” OR “polymorfism” OR “polymorfisms” OR “Genomic Structural Variation” OR “Genomic Structural Variations” OR “Copy Number Variation” OR “DNA Copy Number Variations” OR “DNA Copy Number Variation” OR “SNP” OR “SNPs” OR “nucleotide” OR “nucleotides” OR “Genetics” OR “pharmacogenetic” OR “pharmacogenetics” OR “hapmap” OR “Pharmacogenetics” OR toxicogenetic* OR “predictive marker” OR “predictive markers” OR “predictive biomarkers” OR “predictive biomarker” OR “Genetic Marker” OR “GWAS” OR “Genome-Wide Association” OR “Genetic Association”) AND TI=(“sunitinib” OR sunitinib* OR “Sutent” OR “SU 11248” OR “SU11248” OR “SU-11248” OR “protein tyrosine kinase inhibitor” OR “tyrosine kinase inhibitor” OR “Tyrosine kinase inhibitors” OR “TKI” OR “SU011248” OR “SU-011248” OR “SU 011248” OR vasculotropin receptor OR “vascular endothelial growth factor receptor” OR “VEGF” OR “VEGFR” OR “VEGF-R1” OR “VEGF-R2” OR “VEGF-R3” OR “VEGFR1” OR “VEGFR2” OR “VEGFR3” OR “PDGF” OR “PDGFR” OR “sorafenib” OR “sorafenib” OR “nexavar” OR “BAY 545-9085” OR “BAY-545-9085” OR “BAY 43-9006” OR “BAY-43-9006” OR “BAY 439006” OR “BAY43-9006” OR “BAY439006” OR “pazopanib” OR “votrient” OR “GW 786034B” OR “GW786034B” OR “GW-786034B” OR “GW 786034” OR “GW-786034” OR “GW-786034” OR “axitinib” OR “inlyta” OR “AG 013736” OR “AG013736” OR “AG-013736” OR “Angiogenesis Inhibitor” OR “angiogenesis inhibitors” OR “angiogenesis inhibitor” OR “anti angiogenic” OR “anti-angiogenic” OR “antiangiogenic” OR “anti angiogenesis” OR “anti-angiogenesis” OR “antiangiogenesis”)) AND la=english AND py=(2004 OR 2005 OR 2006 OR 2007 OR 2008 OR 2009 OR 2010 OR 2011 OR 2012 OR 2013 OR 2014 OR 2015) NOT meeting abstracts	129	35
COCHRANE Library	((“Kidney Carcinoma” OR “renal cell carcinoma” OR “renal tumor” OR “renal tumour” OR “renal tumors” OR “renal tumours” OR “RCC” OR “mRCC” OR “kidney cell carcinoma” OR “kidney tumor” OR “kidney tumour” OR “kidney tumors” OR “kidney tumours” OR “renal cell adenocarcinoma” OR “kidney cell adenocarcinoma” OR “Grawitz”) AND (“Genetic Variability” OR “genetic” OR “genetics” OR “genomics” OR “genomic” OR “genome” OR “mutation” OR “mutations” OR “allelic” OR “Allele” OR “alleles” OR “DNA” OR “gene” OR “genes” OR “chromosome” OR “chromosomes” OR “germ-line” OR “germline” OR “INDEL” OR “insertion” OR “deletion” OR “inversion” OR genetic polymorphism OR “polymorphism” OR “polymorphisms” OR “polymorfism” OR “polymorfisms” OR “Genomic Structural Variation” OR “Genomic Structural Variations” OR “Copy Number Variation” OR “DNA Copy Number Variations” OR “DNA Copy Number Variation” OR “SNP” OR “SNPs” OR “nucleotide” OR “nucleotides” OR “Genetics” OR “pharmacogenetic” OR “pharmacogenetics” OR “hapmap” OR “Pharmacogenetics” OR toxicogenetic* OR “predictive marker” OR “predictive markers” OR “predictive biomarkers” OR “predictive biomarker” OR “Genetic Marker” OR “GWAS” OR “Genome-Wide Association” OR “Genetic Association”) AND (“sunitinib” OR sunitinib* OR “Sutent” OR “SU 11248” OR “SU11248” OR “SU-11248” OR “protein tyrosine kinase inhibitor” OR “tyrosine kinase inhibitor” OR “Tyrosine kinase inhibitors” OR “TKI” OR “SU011248” OR “SU-011248” OR “SU 011248” OR vasculotropin receptor OR “vascular endothelial growth factor receptor” OR “VEGF” OR “VEGFR” OR “VEGF-R1” OR “VEGF-R2” OR “VEGF-R3” OR “VEGFR1” OR “VEGFR2” OR “VEGFR3” OR “PDGF” OR “PDGFR” OR “sorafenib” OR “sorafenib” OR “nexavar” OR “BAY 545-9085” OR “BAY-545-9085” OR “BAY 43-9006” OR “BAY-43-9006” OR “BAY 439006” OR “BAY43-9006” OR “BAY439006” OR “pazopanib” OR “votrient” OR “GW 786034B” OR “GW786034B” OR “GW-786034B” OR “GW 786034” OR “GW-786034” OR “GW-786034” OR “axitinib” OR “inlyta” OR “AG 013736” OR “AG013736” OR “AG-013736” OR “Angiogenesis Inhibitor” OR “angiogenesis inhibitors” OR “angiogenesis inhibitor” OR “anti angiogenic” OR “anti-angiogenic” OR “antiangiogenic” OR “anti angiogenesis” OR “anti-angiogenesis” OR “antiangiogenesis”))	20	2

Supplementary document 2: Continued

Database	Search Strategy	Number of references	Number of unique references
CENTRAL	(("Kidney Carcinoma" OR "renal cell carcinoma" OR "renal tumor" OR "renal tumour" OR "renal tumors" OR "renal tumours" OR "RCC" OR "mRCC" OR "kidney cell carcinoma" OR "kidney tumor" OR "kidney tumour" OR "kidney tumors" OR "kidney tumours" OR "renal cell adenocarcinoma" OR "kidney cell adenocarcinoma" OR "Grawitz") AND ("Genetic Variability" OR "genetic" OR "genetics" OR "genomics" OR "genomic" OR "genome" OR "mutation" OR "mutations" OR "allelic" OR "Allele" OR "alleles" OR "DNA" OR "gene" OR "genes" OR "chromosome" OR "chromosomes" OR "germ-line" OR "germline" OR "INDEL" OR "insertion" OR "deletion" OR "inversion" OR genetic polymorphism OR "polymorphism" OR "polymorphisms" OR "polymorfism" OR "polymorfisms" OR "Genomic Structural Variation" OR "Genomic Structural Variations" OR "Copy Number Variation" OR "DNA Copy Number Variations" OR "DNA Copy Number Variation" OR "SNP" OR "SNPs" OR "nucleotide" OR "nucleotides" OR "Genetics" OR "pharmacogenetic" OR "pharmacogenetics" OR "hapmap" OR "Pharmacogenetics" OR toxicogenetic* OR "predictive marker" OR "predictive markers" OR "predictive biomarkers" OR "predictive biomarker" OR "Genetic Marker" OR "GWAS" OR "Genome-Wide Association" OR "Genetic Association") AND ("sunitinib" OR sunitinib* OR "Sutent" OR "SU 11248" OR "SU11248" OR "SU-11248" OR "protein tyrosine kinase inhibitor" OR "tyrosine kinase inhibitor" OR "Tyrosine kinase inhibitors" OR "TKI" OR "SU011248" OR "SU-011248" OR "SU 011248" OR vasculotropin receptor OR "vascular endothelial growth factor receptor" OR "VEGF" OR "VEGFR" OR "VEGF-R1" OR "VEGF-R2" OR "VEGF-R3" OR "VEGFR1" OR "VEGFR2" OR "VEGFR3" OR "PDGF" OR "PDGFR" OR "sorafenib" OR "sorafenib" OR "nexavar" OR "BAY 545-9085" OR "BAY-545-9085" OR "BAY 43-9006" OR "BAY-43-9006" OR "BAY 439006" OR "BAY43-9006" OR "BAY439006" OR "pazopanib" OR "votrient" OR "GW 786034B" OR "GW786034B" OR "GW-786034B" OR "GW 786034" OR "GW786034" OR "GW-786034" OR "axitinib" OR "inlyta" OR "AG 013736" OR "AG013736" OR "AG-013736" OR "Angiogenesis Inhibitor" OR "angiogenesis inhibitors" OR "angiogenesis inhibitor" OR "anti angiogenic" OR "anti-angiogenic" OR "antiangiogenic" OR "anti angiogenesis" OR "anti-angiogenesis" OR "antiangiogenesis"))	20	0
Total			973

Supplementary document 3: Reported non-SNP biomarker associations ($P \leq 0.05$) with TKI treatment outcome in mRCC from 2009 to 2015.

First Author, year	TKI	No. of patients	ethnicity	Tumour type	Associated biomarkers	Study endpoint	Effect size	95%CI	p-value
Choueiri <i>et al.</i> , 2009 [45]	SUN	1	probably CAU (US)	CC on first diagnosis, then Xp11 translocation RCC	Xp11.2 translocation histological subtype (<i>TFE3</i> gene)	Response	N/A		
Beuselinck <i>et al.</i> , 2015 [46]	SUN	121	probably CAU (Belgium)	CC	cdRCC tumor molecular subtypes <i>crrcc1/crrcc4</i> vs. <i>crrcc2/crrcc3</i>	Response			$P=0.005$
						PFS	HR:0.42	0.21-0.86	$P=0.017$
						OS	HR:0.38	0.19-0.76	$P=0.0064$
Paule <i>et al.</i> , 2010 [47]	SUN	23	probably CAU (France)	CC	VEGF (isoforms 121, 165, and 189), VEGF121/VEGF165	Response	39%		$P=0.04$
						OS	HR:5.8	1.4-24.5	$P=0.02$
Gruenewald <i>et al.</i> , 2010 [48]	SUN	26	probably CAU (Germany)	80.8% CC	Level of circulating endothelial cells (CEC)	PFS	Increase from baseline 40 ± 41 CEC/ml to 111 ± 61 at day 28		$P=0.0109$
D'Alterio <i>et al.</i> , 2012 [49]	SUN	62	probably CAU (Italy)	Mostly CC	CXCR4 expression	Objective response	-		$P=0.0262$
					CXCR4 expression + Fuhrman grading				$P=0.0005$
					CXCR4 expression	PFS	HR:2.04	1.08-3.84	$P=0.0271$
García-Donas <i>et al.</i> , 2013 [50]	SUN	67	mostly European	CC	HIF1A protein expression	Objective response	HR:0.11	0.017-0.75	$P=0.024$
					PDGFRB protein expression		HR:0.04	0.002-0.68	$P=0.026$
					VEGFR3 protein expression	PFS	HR:0.403	0.20-0.82	$P=0.012$
					VEGFA protein expression	OS	HR:4.29	1.43-12.8	$P=0.0092$
					HIF2A protein expression		HR:0.39	0.15-0.99	$P=0.048$
					EGALN3 expression		HR:2.89	1.04-8.05	$P=0.042$

Supplementary document 3: Continued

First Author, year	TKI	No. of patients	ethnicity	Tumour type	Associated biomarkers	Study endpoint	Effect size	95%CI	p-value
Minardi <i>et al.</i> , 2013 [51]	SUN	41	probably European	CC	Endothelial cell VEGF expression VEGF staining (H-score)	distant metastasis free survival (DMFS)	-		$P=0.031$
					tumour cell VEGF expression	OS			$P=0.042$
					VEGF staining (H-score)				$P=0.0145$
					Endothelial cell VEGF expression				$P=0.004$
					EMMPRIN expression	Pathological T stage Clinical M stage			$P=0.016$
Sato <i>et al.</i> , 2013 [52]	SUN	50	JAP	82% CC		AJCC stage			$P<0.0001$
						Fuhrman grade			$P=0.0001$
						OS			$P=0.0345$
						PFS			$P=0.0066$
						MVA	$R^2=0.4801$		
Motzer <i>et al.</i> , 2014 [33]	SUN	292	88% CAU	mRCC	HIF-1 α	PFS	HR:1.55	1.03-2.34	$P=0.0341$
					baseline angiopoietin-2 baseline matrix baseline metalloproteinase-2	best overall response (CR/ PR vs. SD/PD) at baseline	-		$P=0.0197$
del Puerto-Nevado <i>et al.</i> , 2014 [53]	SUN	23	probably CAU (Spain)	CC	protein expression: phosphorylated KDR (pKDR-Y1775)	PFS	HR:5.35	1.49-19.13	$P=0.01$
Gambini <i>et al.</i> , 2014 [54]	SUN	1	probably CAU (Italy)	CC	neuroendocrine markers + pancreatic metastases	OS	HR:5.13	1.25-21.05	$P=0.02$
						response	N/A		
Mikami <i>et al.</i> , 2014 [55]	SUN	25	probably JAP	CC	CD44 expression	Time to treatment failure	-		$P=0.008$
						OS			$P=0.007$
					TNF- α /CD44 expression	PFS			$P=0.029$

Supplementary document 3: Continued

First Author, year	TKI	No. of patients	ethnicity	Tumour type	Associated biomarkers	Study endpoint	Effect size	95%CI	p-value
Pena <i>et al.</i> , 2010 [61]	SOR (N=451, placebo (N=452))	903	probably CAU (USA)	Mostly CC	VEGF plasma protein levels week 3 and week 12 sVEGFR2 plasma protein levels week 3 and week 12 TIMP-1 plasma protein levels week 3 TIMP-1 plasma protein levels week 12	Sorafenib use	-		P<0.0001 P<0.0001 P=0.002 P=0.006
Feng <i>et al.</i> , 2013 [62]	SOR	28 (18 mRCC vs 10 healthy)	probably CHI	CC	Plasma cell free DNA levels	Response week 8 Response week 12 Response week 16 Response week 24 Survival week 12 Survival week 16 Survival week 24	-		P=0.019 P=0.007 P=0.004 P<0.001 P=0.047 P=0.018 P=0.018
Aziz <i>et al.</i> , 2014 [63]	SOR	96	probably CAU (USA)	RCC (subtype not specified)	Microvessel area (MVA) in tumor samples	response	-	-	P=0.005 (after Bonferroni correction)
Zheng <i>et al.</i> , 2015 [64]	SOR	10	probably CHI	RCC (subtype not specified)	Micro RNA-30a (miR-30a)	Sorafenib resistance	-	-	P<0.05
Doberstein <i>et al.</i> , 2011 [65]	rapamycin, SUN and cisplatin	282	probably CAU (Germany)	CC=81.6% (n=230)	expression of L1 cell adhesion molecule (L1-CAM)	survival sunitinib resistance	RR:1.8	-	P=0.014 P<0.01

Supplementary document 3: Continued

First Author, year	TKI	No. of patients	ethnicity	Tumour type	Associated biomarkers	Study endpoint	Effect size	95%CI	p-value
Furuya <i>et al.</i> , 2011 [66]	SOR ± IFNa	66	JAP	CC	IFMAR2 mRNA levels	Response OS	-		P<0.0001
Sharpe <i>et al.</i> , 2013 [67]	SUN or PAZ	85	N/A, but probably all CAU (UIK or Belgium)	CC	vessel density (CD31)	Use of TKI			P<0.05
					phospho-S6K expression			P<0.05	
					PDL-1 expression			P<0.05	
					FOXP3 expression			P<0.05	
					increase in cytoplasmic FGF-2			P<0.05	
					MET receptor expression in vessels			P<0.05	
					Fuhrman tumor grade			P<0.05	
					Ki-67			P<0.05	
					levels of Ki-67		PD		P<0.05
levels of CD31		P<0.05							
Yamada <i>et al.</i> , 2014 [68]	SUN, SOR, AXI	13	probably JAP	CC: N=11, papillary; N=2	Plasma granulocyte macrophage colony-stimulating factor (GM-CSF) level	Response (PR vs SD or PD)	-	-	P=0.012
Tran <i>et al.</i> , 2012 [69]	PAZ, placebo	344		mRCC (subtype not specified)	IL-6	PFS	HR:0.55	0.28-0.71	P=0.009
Peters <i>et al.</i> , 2014 [70]	SUN, SOR, AXI, BEV	18	probably CAU (Hannover)	CC=88.9%	CST6 methylation status	PFS	HR:4.1	1.3-12.6	P=0.009
							HR:4.1	13.0-13.4	P=0.011
					LAD1 methylation status	OS	HR:6.4	1.6-26.0	P=0.004
Dubrowsinskaja <i>et al.</i> , 2014 [71]	SUN, SOR, BEV, AXI	18	-	CC: N=16, papillary; N=1, chromophobe; N=1	NEFH methylation	OS	HR:2.9	1.0-8.6	P=0.043
							-	-	P=0.028
Choudhury <i>et al.</i> , 2015 [72]	SUN, PAZ	48	Asian	CC	Prognostic classification based on multigene assay: CXCL5, EFNA5, EMCN, LAMB3, PLG, PRAME, RABRST1, SLC6A19	Radiologic response	-	-	P=5.96 x10 ⁻⁴
						OS	HR:3.58	1.162-11.03	P=0.0186

Abbreviations: ABCB1=ATP binding cassette member B1; ABCG2=ATP binding cassette member G2; sunitinib=SUN; pazopanib=PAZ; sorafenib=SOR; axitinib=AXI; bevacizumab=BEV; foretinib=FOR; valatinib=VAL; CAU=Caucasian; CHI=Chinese; JAP=Japanese; CC=clear cell renal cell carcinoma; GIST=gastrointestinal stromal tumor; PFS=progression-free survival; OS=overall survival; AST= aspartate aminotransferase; ALT= alanine aminotransferase