



Universiteit
Leiden
The Netherlands

Innovation and stasis : gymnosperms from the early Permian Jambi flora

Booi, M.

Citation

Booi, M. (2017, November 15). *Innovation and stasis : gymnosperms from the early Permian Jambi flora*. Retrieved from <https://hdl.handle.net/1887/57351>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/57351>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/57351> holds various files of this Leiden University dissertation

Author: Booij, Menno

Title: Innovation and stasis : gymnosperms from the early Permian Jambi flora

Date: 2017-11-15

References

- Anfodillo, A., Petit, G., Crivellaro, A., 2013. Axial conduit widening in woody species: a still neglected anatomical pattern. IAWA J. 34, 24–36.
- Arens, N.C., 1997. Responses of leaf anatomy to light environment in the tree fern *Cyathea caracasana* (*Cyatheaceae*) and its application to some ancient seed ferns. Palaios 12, 84-94.
- Asama, K. 1959. A systematic study of so-called *Gigantopteris*. Science reports of the Tohoku University. 2nd series, Geology 31, 1-72.
- Asama, K., 1960. Evolution of the leaf-forms through the ages explained by the successive retardation and neotony. Sci. Rep. of the Tohoku University, Sendai, Japan, 2nd Ser. (Geology) 31 (1), 1-72.
- Asama, K., 1962. Evolution of Shansi Flora and Origin of Simple Leaf. Science Reports of the Tohoku University, Sendai, Japan, 2nd Ser. (Geology), Special Volume 5, 274-273.
- Asama, K., 1966. Permian plants from Phetchabun, Thailand and problems of floral migration from Gondwanaland. Bull. Nat. Sci. Mus. 9(2), 171-211.
- Asama, K. 1975. Summary of the Carboniferous and Permian plants from Thailand, Malaysia and adjacent areas. Geology and Palaeontology of Southeast Asia 15, 77-101.
- Asama, K. 1984. *Gigantopteris* flora in China and Southeast Asia. Geology and Palaeontology of Southeast Asia 25, 311-323.
- Bailey I.W. & Faull A.F. 1934. The cambium and its derivative tissues. No. IX. Structural variability in the redwood *Sequoia sempervirens*, and its significance in the identification of fossil woods. J. Arnold Arbor. 15, 233–254.
- Bamford, M.K. & Philippe, M. 2001. Jurassic–Early Cretaceous Gondwanan homoxylous woods: a nomenclatural revision of the genera with taxonomic notes. Review Palaeobot. Palynol.113: 287--297.
- Barbacka, M. & Van Konijnenburg-Van Cittert, J.H.A. 1998. Sun and shade leaves in two Jurassic species of Pteridosperms. Rev. Palaeobot. Palynol.103(3-4), 209-221.
- Barber, A. J. & Crow, M.J. 2003. Evaluation of plate tectonic models for the development of Sumatra. Gondwana Research, v. 20, 1-28.
- Beauchamp, B., 1994. Permian climatic cooling in the Canadian Arctic. In: Klein, G.D. (Ed.), Pangea: Paleoclimate, Tectonics and Sedimentation During Accretion, Zenith and Breakup of a Supercontinent. Geological Society America of Special Paper 288, 229-246.
- Barber, A.J., Crow, M.J. & De Smet, M.E.M. 2005. Tectonic Evolution. In: Barber, A.J., Crow, M.J., Milsom, J.S. (Eds.) Sumatra-Geology, resources and tectonic evolution, Geological Society Memoir 31, 300pp.
- Barthel, M. 1968. “*Pecopteris*” *feminaeformis* (Schlotheim) Sterzel und “*Araucarites*” *spiciformis* Andrae in Germar – Coenopterideen des Stephans und Unteren Perms. Paläontologische Abhandlungen, Abt. B, 2, 727-742.

- Bashforth, A.R. & Zodrow, E.L. 2007. Partial reconstruction and palaeoecology of *Sphenophyllum costae* (Middle Pennsylvanian, Nova Scotia, Canada). *Bulletin of Geosciences* 82: 365-382.
- Beck, A.L. & Labandeira C.C. 1998. Early Permian folivory on a gigantopterid-dominated riparian flora from north-central Texas. *Palaeogeography, Palaeoclimatology, Palaeoecology* 142, 139-173.
- Beerling, D.J., Osborne, C.P. & Chaloner W.G. 2001. Evolution of leaf-form in land plants linked to atmospheric CO₂ decline in the Late Palaeozoic era. *Nature* 410, 352-354.
- Berner, R.A. & Kothavala, Z. 2001. Geocarb III: A revised model of atmospheric CO₂ over Phanerozoic time. *American Journal of Science* 301, 182-204.
- Berthelin, M., Broutin, J., Kerp, H., Crasquin-Soleau, S., Platel, J.-P. & Roger, J. 2003. The Oman Gharif mixed paleoflora: a useful tool for testing Permian Pangea reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 196, 85-98.
- Booi, M., Van Waveren, I.M., Van Konijnenburg-Van Cittert, J.H.A. & De Boer, P.L. 2008. New material of *Macrlethopteris* from the Early Permian Jambi flora (Middle Sumatra, Indonesia) and its palaeoecological implications. *Review of Palaeobotany and Palynology* 152 (3-4), 101-112.
- Booi, M., van Waveren, I.M. & van Konijnenburg-van Cittert, J.H.A. 2009a. *Comia* and *Rhachiphyllum* from the Early Permian of Sumatra, Indonesia. *Review of Palaeobotany and Palynology*, 156, 418-435.
- Booi, M., van Waveren, I.M. & van Konijnenburg-van Cittert, J.H.A. 2009b. The Jambi gigantopterids and their place in gigantopterid classification. *Bot. J. Lin. Soc.* 161(3), 302-328.
- Booi, M., Van Waveren, I.M. & Van Konijnenburg-Van Cittert, J.H.A. 2014. Gymnosperm permineralized wood from the Early Permian Jambi flora, Sumatra, Indonesia. *Permian Araucarioids. ISAWA Journal*, 35, 307-331.
- Boureau E. & Doubinger J. 1975. Gigantopterideae et formes apparentées. In: Boureau E, ed. *Traité de Paléobotanique*. 4 (2), *Pteridophylla* (Première partie), Paris: Masson et Cie.
- Boyce, C.K. 2005. Patterns of segregation and convergence in the evolution of fern and seed plant leaf morphologies. *Paleobiology* 31: 117-140.
- Boyce, C.K. 2007. Mechanisms of laminar growth in morphologically convergent leaves and flower petals. *International Journal of Plant Science* 168: 1151-1156.
- Boyce, C.K. 2008. The fossil record of plant physiology and development - what leaves can tell us. In: Kelley PH, Bambach RK, eds. *From evolution to geobiology: research questions driving paleontology at the start of a new century*, Paleontological Society Short Course, October 4, 2008. *Paleontological Society Papers* 14: 133-146.
- Boyce, C.K. & Knoll, A.H. 2002. Evolution of developmental potential and the multiple independent origins of leaves in Paleozoic vascular plants. *Paleobiology* 28: 70-100.
- Brea, M., Artabe, A.E. & Spallotti, L.A. 2009. Darwin forest at Agua de la Zorra: the first in situ forest discovered in South America by Darwin in 1835. *Revista de la Asociación Geológica Argentina* 64 (1): 21 – 31.

- Broutin, J. 1977. Nouvelles données sur la flore des bassins Autuno-Stephaniens des environ de Guadalcanal (Province de Seville-Espagne). Cuadernos Geología Ibérica 4: 91-98.
- Calder, J.H., Gibling, M.R., Scott, A.C., Davies, S.J. & Hebert, B.L. 2006. A fossil lycopsid forest succession in the classic Joggins section of Nova Scotia: paleoecology of a disturbance-prone Pennsylvanian wetland. In Wetlands through time. Edited by S. Greb and W.A. DiMichele. Geological Society of America Special Paper 399: 169–195.
- Carlquist S. 1975. Ecological strategies of xylem evolution. University of California Press, Berkeley.
- Carlquist S. 1977. Ecological factors in wood evolution: A floristic approach. Amer. J. Bot. 64: 887–896.
- Carpentier, A. 1920. Notes paléophytiques sur le Westphalien du Nord de la France. Ann. Soc. Géol. Nord 44: 137–150.
- Cas, R.A.F. & Wright, J.V. 1987. Volcanic Successions. Modern and Ancient, London, Boston, Sydney, Wellington: Allen & Unw 528 pp.
- Chaloner, W.G. & Meyen, S.V. 1973. Carboniferous and Permian floras of the Northern continents. In: Hallam, A. (ed.) Atlas of Palaeobiogeography, Elsevier, Amsterdam, 169–186.
- Champagne, C. & Sinha, N. 2004. Compound leaves: equal to the sum of their parts? Development 131(18): 4401-4412.
- Chaney, D.S. & DiMichele, W.A. 2007. Paleobotany of the classic redbeds (Clear Fork Group-Early Permian) of north central Texas. In: Wong, Th. (Ed.) Proceedings of the XVth International Congress on Carboniferous and Permian Stratigraphy. Utrecht, The Netherlands, 10-16 August 2003. Royal Netherlands Academy of Arts and Sciences, The Netherlands
- Chaney, D.S., Mamay, S.H., DiMichele, W.A. & Kerp, H. 2009. *Auritifolia* gen. nov., probable seed plant foliage with comioid affinities from the Early Permian of Texas, U.S.A. International Journal of Plant Science 170(2): 247-266.
- Cleal, C.J. 1993. Gymnospermophyta. In: Benton MJ, ed. The fossil record 2. London: Chapman and Hall, pp.795-808.
- Cleal, C.J. & Thomas B.A. 2005. Palaeozoic tropical rainforests and their effect on global climates: is the past the key to the present? Geobiology 3(1): 13-31.
- Cleal, C.J., Uhl, D., Cascales-Minana, B., Thomas, B.A., Bashforth, A.R., King, S.C. & Zodrow, E.L. 2012. Plant biodiversity changes in Carboniferous tropical wetlands. Earth-Science Reviews 114: 124–155.
- Cridland, A.A. 1964. *Amyelon* in American coal balls. Palaeontology 7: 189-209.
- Cridland, A.A. & Morris, J.E. 1960. *Spermopteris*, a new genus of Pteridosperms from the Upper Pennsylvanian Series of Kansas. American Journal of Botany 47(10): 855-859.
- Crippa, G., Angiolini, L., Van Waveren, I., Crow, M., Hasibuan, F., Stephenson, M. & Ueno, K. 2014. Brachiopods, fusulines and palynomorphs from the Mengkarang Formation (early Permian, Sumatra) and their geographical significance. Journal of Asian Earth Sciences 79: 206–223.

- Crow M.J. & Van Waveren I.M. 2010. A preliminary account of the Karing Volcanic Complex in the Permian West Sumatra Volcanic Arc. Symposium of the International Geological Correlation Programme Project 516 (IGCP 516), Kuala Lumpur, Malaysia, 36.
- Darwin, C. 1846. Geological observations on South America. Being the third part of the Geology of the voyage of the Beagle, during the years 1832 to 1836. Smith, Elder and Co. 280 p., London.
- Dawson, J.W. 1871. The fossil plants of the Devonian and Upper Silurian formations of Canada. Geological Survey of Canada, part 1, 1-92.
- De Laet J., Clinckemaillie D., Jansen S. & Smets E. 1995. Floral ontogeny in the Plumbaginaceae. *Journal of Plant Research* 108: 289-304.
- DiMichele, W.A., Chaney, D.S., Dixon, W.H., Nelson, W.J. & Hook, R.W. 2000. An Early Permian Coastal Flora from the Central Basin Platform of Gaines County, West Texas. *Palaios* 15: 524-534.
- DiMichele, W.A., Chaney, D.S., Nelson, W.J., Lucas, S.G., Looy, C.V., Quick, K. & Wang, J. 2007. A low diversity, seasonal tropical landscape dominated by conifers and peltasperms: Early Permian Abo Formation, New Mexico. *Review of Palaeobotany and Palynology* 145: 249-273.
- DiMichele, W.A., Gastaldo, R.A. & Pfefferkorn, H.W. 2005. Plant Biodiversity Partitioning in the Late Carboniferous and Early Permian and Its Implications for Ecosystem Assembly. *Proceedings of the California Academy of Sciences* 56, suppl. I, no.4, 32-49.
- DiMichele, W.A., Kerp, H., Krings, M., Chaney, D.S. 2005. The Permian peltasperm radiation: evidence from the Southwestern United States. In: Lucas, S.G., Zeigler, K.E. (Eds.), *The Nonmarine Permian*. New Mexico Museum of Natural History and Science Bulletin 30: 67-79.
- DiMichele, W.A., Kerp, H., Tabor, N.J. & Looy, C.V. 2008. Revisiting the so-called “Paleophytic-Mesophytic” transition in equatorial Pangea: vegetational integrity and climatic tracking. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268(3-4): 152-163.
- DiMichele, W.A., Mamay, S.H., Chaney, D.S., Hook, R.W. & Nelson, W.J. 2001a. An Early Permian Flora with Late Permian and Mesozoic Affinities from North-Central Texas. *Journal of Paleontology* 75(2): 449-460.
- DiMichele, W.A., Pfefferkorn, H.W. & Gastaldo, R.A. 2001b. Response of Late Carboniferous and Early Permian Plant Communities to Climate Change. *Annual Review of Plant Sciences* 29: 461-487.
- DiMichele, W.A. & Phillips, T.L. 1994. Paleobotanical and paleoecological constraints on models of peat formation in the Late Carboniferous of Euramerica. *Palaeogeography, Palaeoclimatology, Palaeoecology* 106: 39-90.
- DiMichele, W.A. & Phillips, T.L. 1996. Climate change, plant extinctions and vegetational recovery during the Middle-Late Pennsylvanian Transition: the Case of tropical peat-forming environments in North America. In: Hart, M.B. (ed.) *Biotic Recovery from Mass Extinction Events*, Geological Society Publication 102: 201-221.
- DiMichele, W.A. & Phillips, T.L. 2002. The ecology of Paleozoic ferns. *Review of Palaeobotany and Palynology* 119: 143-159.

- DiMichele, W.A., Phillips, T.L. & Pfefferkorn, H.W. 2006a. Paleoecology of Late Paleozoic pteridosperms from tropical Euramerica. *Journal of the Torrey Botanical Society* 133: 83-118.
- DiMichele, W.A., Tabor, N.J., Chaney, D.S. & Nelson, W.J. 2006b. From wetlands to wet spots: Environmental tracking and the fate of Carboniferous elements in Early Permian tropical floras. In: Greb, S.F., and DiMichele, W.A., *Wetlands through time: Geological Society of America Special Paper* 399: 223–248.
- Donovan, S., Van Waveren, I.M. & Portell, R. 2013. Island slopes and jumbled shell beds. *J. Geol. Soc.* 170: 527–534.
- Endress P.K. 1995. Origins of flower morphology. *Journal of Experimental Zoology (Molecular and Developmental Evolution)* 291: 105-115.
- Endlicher, S.L. 1847. *Synopsis coniferarum*. Scheitlin und Zollikofer, Sangalli (Sankt Gallen).
- Esau, K. 1965. Plant anatomy. New York: John Wiley and Sons.
- Evans, W.C., Bergfeld, D., McGeehin, J.P., King, J.C. & Haesler, H. 2010. Tree-ring ^{14}C links seismic swarm to CO₂ spike at Yellowstone, USA. *Geology* 38: 1075-1078.
- Falcon-Lang, H. J. 2005. Intra-tree variability in wood anatomy and its implications for fossil wood systematics and palaeoclimatic studies. *Palaeontology* 48(1): 171-183.
- Falcon-Lang, H.J. & Bashforth, A.R. 2004. Pennsylvanian uplands were forested by giant cordaitalean trees. *Geology* 32(5): 417-420.
- Falcon-Lang, H.J. & Bashforth, A.R. 2005. Morphology, anatomy, and upland ecology of large cordaitalean trees from the Middle Pennsylvanian of Newfoundland. *Review of Palaeobotany and Palynology* 135: 223-243.
- Farr, E. R. & G. Zijlstra, eds. *Index Nominum Genericorum (Plantarum)*. 1996+. <<http://botany.si.edu/ing/>> 03/01/2010
- Fefilova, L.A. 1973. The Permian ferns of the North of the Cis-Ural depression. Nauka, Leningrad. (in Russian)
- Ferrari, O.M., Hochard, C. & Stampfli, G.M. 2008. An alternative plate tectonic model for the Palaeozoic-early Mesozoic palaeo-Tethyan evolution of southeast Asia (northern Thailand-Burma). *Tectonophysics* 451: 346-365.
- Fontaine, H. & Gafoer, S. 1989. The pre-Tertiary fossils of Sumatra and their environments: CCOP Technical Paper 19: 1-356.
- Frentzen K. 1931. Studien über die fossilen Hölzer der Sammelgattung *Dadoxylon* Endlicher. Abhandlungen der Heidelberger Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse 16, 1-93 and 19: 1-51.
- Friedman, W.E., Moore, R.C. & Purugganan, M.D. 2004. The evolution of plant development. *American Journal of Botany* 91: 1726-1741.

- Fritz, W.J. 1980. Stumps transported and deposited upright by Mount St. Helens mud flows. *Geology* 8: 568-588.
- Galtier, J. 1996. Coal-ball floras of the Namurian-Westphalian of Europe. *Review of Palaeobotany and Palynology* 95: 51-72.
- Galtier, J. & Broutin, J. 2008. Floras from red beds of the Permian Basin of Lodève (Southern France). *Journal of Iberian Geology* 34(1): 57-72.
- Gälweiler L., Guan C.-H., Müller A., Wisman E., Mendgen K., Yephremov A. & Palme K., 1998. Regulation of polar auxin transport by AtPIN1 in *Arabidopsis* vascular tissue. *Science* 282: 2226-2230.
- Gand, G., Kerp, H., Parsons, C. & Martínez-García, E. 1997. Palaeoenvironmental and stratigraphic aspects of animal traces and plant remains in Spanish Permian red beds (Peña Sagra, Cantabrian mountains, Spain). *Géobios* 30(2): 295.
- Gastaldo, R.A. & Demko, T.M. 2011. Long term hydrology controls the plant fossil record. In: Allison, P.A., and Bottjer, D.J., eds., *Taphonomy, Second Edition: Processes and Bias Through Time: Topics in Geobiology*, 32: 249-285.
- Gastaldo, R. A., DiMichele, W. A. & Pfefferkorn, H. W. 1996. Out of the icehouse into the greenhouse: A Late Paleozoic Analog for modern global vegetational change. *GSA Today* 6(10): 1-7.
- Gifford, A.S. & Gifford, E.M. 1959. Comparative morphology of vascular plants. San Francisco and London: W.H. Freeman and company. First edition.
- Giraud, B. 1991. Les espèces du genre *Dadoxylon* depuis 1962: leur répartition et leur évolution du Permien à la fin du Mésozoïque. *Review Palaeobot. Palynol.* 67: 13-39.
- Givnish, T.J. 1976. Leaf form in relation to environment: A theoretical study. PhD thesis, Princeton University.
- Givnish, T.J. 1979. On the adaptive significance of leaf form. In: Solbrig, O. T., Jain, S., Johnson, G. B. & Raven, P. H. (Eds.) *Topics in Plant Population Biology*, New York, Columbia University Press: 375-407.
- Givnish, T.J. 1988. Adaptation to sun and shade: A whole plant perspective. *Australian Journal of Plant Physiology* 15: 63-92.
- Glasspool, I.J., Hilton, J., Collinson, M.E. & Wang, S.-J. 2004a. Defining the gigantopterid concept: A reinvestigation of *Gigantopteris (Megalopteris) nicotianaefolia* Schenck and its taxonomical implications. *Palaeontology* 47: 1339-1361.
- Glasspool, I.J., Hilton, J., Collinson, M.E., Wang, S.-J., Sen, L.-C. 2004b. Foliar physiognomy in Cathaysian gigantopterids and the potential to track Palaeozoic climates using an extinct plant group. *Palaeogeography, Palaeoclimatology, Palaeoecology* 205: 69-110.
- Gothan W. 1905. Zur Anatomie lebender und fossiler Gymnospermenholzer. *Abhandlungen der Königlich Preussischen Geologischen Landesanstalt* 44: 1-108.
- Gothan, W., Weyland, H. 1954. Lehrbuch der Paläobotanik. Akademie-Verlag, Berlin.

- Götze J., Plötze M., Habermann D. 2001. Origin, spectral characteristics and practical applications of the cathodoluminescence (CL) of quartz – a review. *J. Mineralogy and Petrology* 71: 225-250.
- Grand'Eury, F.C. 1877. Mémoire sur la flore carbonifère du département de la Loire et du Centre de la France, étudiée aux trois points de vue botanique, stratigraphique et géognostique. Mémoires présentés par divers savants à l'Académie des Sciences de l'Institut de France 24, 624pp.
- Greb, S.F., DiMichele, W.A. & Gastaldo, R.A. 2006. Evolution and importance of wetlands in earth history. In: Greb, S.F. & DiMichele, W.A., eds., *Wetlands through time*. Geological Society of America Special Paper 399: 1-40.
- Greuter, W., McNeill, J., Barrie, F.R., Burdet, H.M., Demoulin, V., Filgueiras T.S., Nicolson, D.H., Silva, P.C., Skog, J.E., Trehane, P., Turland, N.J. & Hawksworth, D.L. 2000. International Code of Botanical Nomenclature (St. Louis Code). *Regnum Vegetabile* 138. Königstein: Koeltz Scientific Books.
- 'Gu and Zhi' (=Writing Group of The Institute of Geology and Palaeontology, The Institute of Botany, Academia Sinica). 1974. Palaeozoic plants from China. *Fossil Plants of China*, vol.1. Science Press, Beijing, 226pp. (in Chinese)
- Guo, Y.-T., Tian, B.-L. & Chang, J.-L. 1993. The epidermal and anatomical study of gigantopterids leaves. *Chinese Journal of Botany* 5: 92-96.
- Hagemann, W. & Gleissberg, S. 1996. Organogenetic capacity of leaves: The significance of marginal blastozones in angiosperms. *Plant Systematics and Evolution* 199: 121-152.
- Halle, T.G. 1927. Palaeozoic plants from Central Shansi. *Palaeontologica Sinica*, Ser. A, II (I), 1-316.
- Halle ,T.G. 1929. On the habit of *Gigantopteris*. *Geologiska Föreningens i Stockholm Förhandlingar* 51: 236-242.
- Hammer, Ø., Harper, D.A.T. & Ryan, P.D. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1): 1-9.
- Hammer, Ø. & Harper, D.A.T. 2006. Paleontological data analysis. Blackwell Publishing, Oxford, 351pp.
- Hasibuan, F., Andi-Mangga S. & Suyoko. 2000. *Stereochia semi-reticulatus* (Martin) dari Formasi Mengkarang, Jambi, Sumatera. *Palaeontological series GRDC* 10: 59-69.
- Hickey, L.J. 1973. Classification of the architecture of dicotyledonous leaves. *American Journal of Botany* 60: 17-33.
- Hill, C.R. & Camus J.R. 1986. Evolutionary cladistics of Marattialean ferns. *Bulletin of the British Museum of Natural History. Botany* 14: 219-300.
- Hill, C.R., Wagner, R.H. & El-Khayal, A.A. 1985. *Qasimia* gen. nov., an early Marattia-like fern from the Permian of Saudi Arabia. *Scripta Geologica* 79, 1-50.
- Hill, M.O., Gauch, H.G., 1980. Detrended correspondence analysis: An improved ordination technique. *Vegetatio* 42: 47-58.
- Hilton, J. & Cleal, C.J. 2007. The relationship between Euramerican and Cathaysian tropical floras in the Late Palaeozoic: Palaeobiogeographical and palaeogeographical implications. *Earth-Science Reviews* 85: 85-116.

- Hopping, C.H. & Wagner, R.H. 1962. Plant fossils. In: Visser, W.A., Hermes, J.J.(Eds.). Geological results of the exploration for oil in Netherlands New Guinea. Koninklijk Nederlands Geologisch Mijnbouwkundig Genootschap. Geologische Serie, Speciaal nummer 20, Enclosure 17: 1-11.
- Huang, B.-H. 1977. Permian flora from the southeastern part of the Xiao Xingan Lin, NE China. Geol. Press, Beijing (in Chinese with English abstract), 79pp.
- Huang, B.-H. 1980. The plant kingdom. In: Shenyang Institute of Geology and Mineral Resource, ed. Palaeontological Atlas of Northeast China.1. Palaeozoic. Beijing: Geological Publishing House, 526-573 (in Chinese).
- Hutchison, C.S. 2007. Geological Evolution of South-East Asia, 2nd ed. Kuala Lumpur, Geological Society of Malaysia, Kuala Lumpur, 433 pp.
- IAWA Committee. 2004. Richter H.G., Grosser D., Heinz I. & Gasson P.E. (eds.), IAWA list of microscopic features for softwood identification. IAWA J. 25: 1-70.
- Jeffrey, E.C. 1913. The history, comparative anatomy and evolution of the *Araucarioxylon* type. Proc. American Academy of Arts and Sciences 48: 531-571.
- Jongmans, W.J. 1940, Beiträge zur Kenntnis der Karbonflora von Niederländisch Neu-Guinea. Mededelingen van de Geologische Stichting (behorende bij het jaarverslag over 1938 en 1939): 263-273.
- Jongmans, W.J. 1941. Elementen der *Glossopteris*-flora in het Karboon van Nieuw Guinea. Jaarverslag 1940 van de Geologische Stichting, Bijlage III: 33-35.
- Jongmans, W.J. & Gothan, W. 1925. Beiträge zur Kenntnis der Flora des Oberkarbons von Sumatra. Verhandelingen Geologisch Mijnbouwkundig Genootschap voor Nederland en Koloniën (Geologie series) 8:279-304.
- Jongmans, W.J. & Gothan, W. 1935. Die Ergebnisse der paläobotanischen Djambi-Expedition 1925, 2. Die paläobotanische Ergebnisse. Jaarboek van het Mijnwezen in Nederlandsch-Indië 59 (3): 71-121.
- Kawasaki, S., 1934. The flora of the Heian System, Part II. Bull. of the Geological Survey of Chosen (Korea) 6 (4), 47-311.
- Kerp, J.H.F. 1988. Aspects of Permian palaeobotany and palynology. X. The West and Central European species of the genus *Autunia* Krasser emend. Kerp (*Peltaspermaceae*) and the form-genus *Rhachiphyllum* Kerp (callipterid foliage). Review of Palaeobotany and Palynology 54(3-4): 249-360.
- Kerp, H. 1990. The study of fossil gymnosperms by means of cuticular analysis. Palaios 5: 548-569.
- Kerp, H. 1996. Post-Variscan late Palaeozoic Northern Hemisphere gymnosperms: the onset to the Mesozoic. Rev. Palaeobot. Palynol. 90: 263-285.
- Kerp, H. 2000. The modernization of landscapes during the Late Paleozoic-Early Mesozoic. In: Gastaldo, R.A., DiMichele, W.A.(eds.) Phanerozoic Terrestrial Ecosystems, Paleontological Society Papers 6: 80-113.

- Kerp, J.H.F. & Haubold, H. 1988. Aspects of Permian palaeobotany and palynology. VIII. On the reclassification of the West- and Central European species of the form-genus *Callipteris* Brongniart 1849. Review of Palaeobotany and Palynology 54: 135-150.
- Kerp, H. & Krings, M. 1998. Climbing and scrambling growth habits: Common life strategies among Late Carboniferous seed ferns. C.R. Acad. Sci. Paris, Sér. IIa 326: 583-588.
- Kerp, H. & Krings, M. 1999. Light microscopy of fossil cuticles. In: Jones, T.P. & Rowe, N.P. (Eds.), Fossil Plants and Spores: Modern Techniques. Geological Society, London, 52-56.
- Koidzumi, G. 1934. Notes on *Gigantopteris* flora. Acta Phytotaxonomica et Geobotanica 3: 112-114. (in Japanese)
- Koidzumi, G. 1936. On the *Gigantopteris* flora. Acta Phytotaxonomica et Geobotanica 5: 130-139. (in Japanese with English summary)
- Kon'no, E. 1963. Some Permian plants from Thailand. Jap. Jour. Geol. Geogr. 34(2-4): 139-159.
- Kramer, K. 1974a. Die tertiären Hölzer Südost-Asiens (unter Ausschluss der Dipterocarpaceae) 1. Teil. Palaeontographica, Abt.B, 144: 45-181.
- Kramer, K. 1974b. Die tertiären Hölzer Südost-Asiens (unter Ausschluss der Dipterocarpaceae) 2. Teil. Palaeontographica, Abt.B 145: 1-150.
- Krasser, F. 1919. Studien über die fertile Region der Cycadophyten aus den Lunzer Schichten: Makrosporophylle. Denkschr. Kaiserl. Akad. Wiss., Mathem.-naturw. Kl. 97: 1-32.
- Kraus, G. 1870. In: Schimper, W.Ph. Traité Pal. végét. 2 (1870-72)(1), 363-385.
- Kraus, G. 1882. (II.) Zur Diagnostik des Coniferenholzes. Abhandlungen naturforsch. Ges. Halle, 16-33.
- Krings, M. & Kerp, H. 1997. An improved method for obtaining large pteridosperm cuticles. Review of Palaeobotany and Palaeontology 96: 453-456.
- Krings, M., Grewing, A. & Taylor, T.N. 2003a. Mariopterid pteridosperms, scrambling and climbing vines of the Late Carboniferous swamp forest: a case study from the Namurian B of Hagen-Vorhalle (Germany). Botanische Jahrbücher, 124(4): 427-448
- Krings, M., Kerp, H., Taylor, T.N. & Taylor, E.L. 2003b. How Palaeozoic vines and lianas got off the ground: On scrambling and climbing Carboniferous-Early Permian pteridosperms. The Botanical Review 69(2): 204-224.
- Krings, M., Klavins, S.D., DiMichele, W.A., Axsmith, B.J., Kerp, H. & Taylor, T.N. 2005. Epidermal anatomy of *Glenopteris splendens* Sellards nov. emend., an enigmatic seed plant from the Lower Permian of Kansas (U.S.A.). Review of Palaeobotany and Palynology 136: 159-180.
- Kristofovich, A.N. 1957. Paleobotanika. Gostoptekhizdat naučno-techničeskoe izd-vo neftjanoj igorno-toplivnoj literatury. Leningrad.
- Kustatscher, E., Wachtler, M. & Van Konijnenburg-van Cittert, J.H.A. 2007. Horsetails and seed ferns from the Middle Triassic (Anisian) locality Kühwiesenkopf (Monte Prá della Vacca) in the Dolomites (Northern Italy). Palaeontology 50(5): 1277-1298.

- Laveine, J.-P., Coquel, R., Loboziak, S. 1977. Phylogénie générale des Calliptéridiacées (Pteridospermopsida). *Geobios* 10 (6): 757-847.
- Li, H.-Q., Taylor, D.W. 1998. *Aculovinea yunguiensis* gen et sp. nov. (Gigantopteridales), a new taxon of gigantopterid stem from the Upper Permian of Guizhou Province, China. International Journal of Plant Science 159: 1023-1033.
- Li, H.-Q. & Taylor, D.W. 1999. Vessel-bearing stems of *Vasovinea tianii* gen. et sp. nov. (Gigantopteridales) from the Upper Permian of Guizhou Province, China. American Journal of Botany 86: 1563-1575.
- Li, H.-Q. & Tian, B.-L. 1990. Anatomic study of foliage leaf of *Gigantonoclea guizhouensis* Gu et Zhi. *Acta Palaeontologica Sinica* 29: 216-227 (in Chinese with English summary).
- Li, H.-Q., Tian, B.-L., Taylor, E.L. & Taylor, T.N. 1994. Foliar anatomy of *Gigantonoclea guizhouensis* (Gigantopteridales) from the Upper Permian of Guizhou Province, China. American Journal of Botany 81: 678-689.
- Li, X.-X. & Shen, G.-L. 1996. A brief review of the Permian macrofloras in southeast Asia and their phytogeographical delimitation. *Journal of Asian Earth Sciences* 13(3-5): 161-170.
- Li, X.-X. & Yao, Z.-Q. 1980. An outline of recent researches on the Cathaysia flora in Asia. Institute Geol. & Paleont., Acad. Sinica, Nanjing, China: 1-15.
- Li, X.-X. & Yao, Z.-Q. 1982. A review of recent research on the Cathaysia flora in Asia. American Journal of Botany 69(3): 479-486.
- Li, X.-X. & Yao, Z.-Q. 1983a. Current studies of gigantopterids. *Palaeontologia Cathayana* 1: 319-326.
- Li, X.-X. & Yao, Z.-Q. 1983b. Fructifications of gigantopterids from South China. *Palaeontographica, Abt.B* 185 (Lfg.1-3): 11-26.
- Li, X.-X., Yao, Z.-Q. & Deng, L.-H. 1982. An early Late Permian flora from Toba, Qamdo District, Eastern Xizang. *Palaeont. Xizang* V: 17-44.
- Li, Z.-X. & Powell, C.McA. 2001. An outline of the palaeogeographic evolution of the Australasian region since the beginning of the Neoproterozoic. *Earth-Science Reviews* 53: 237-277.
- Liu, L.-J., Wang, X.-F. & Yao, Z.-Q. 1996. *Emplectopteris* and allied plant fossils from the Kaiping Basin, China. *Palaeontographica Abt. B* 239: 109-136.
- Liu, L.-J. & Yao Z.-Q. 1992. On venation of *Emplectopteridium alatum* Kawasaki. *Acta Palaeontologica Sinica* 31: 646-656. (In Chinese and English.)
- Liu, L.-J. & Yao, Z.-Q. 2002. Comparison in leaf architecture between Chinese and American species of *Gigantopteridium*. *Acta Palaeontologica Sinica* 41: 322-333.
- Looy, C.V., Kerp, H., Duijnstee, I.A.P. & DiMichele, W.A. 2014. The late Paleozoic ecological-evolutionary laboratory, a land-plant fossil record perspective. *The Sedimentary Record* 12(4): 4-10.

- Lyell, C. & Dawson, J.W. 1853. On the remains of a reptile (*Dendrerpeton acadianum*) Wyman and Owen), and of a land shell discovered in the interior of an erect fossil tree in the coal measures of Nova Scotia. Quarterly Journal of the Geological Society [London] 9: 58–63.
- McLoughlin, S. 2001. The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. Aust. J. Bot. 49: 271–300.
- Mamay, S.H. 1986. New species of Gigantopteridaceae from the Lower Permian of Texas. Phytologia 61: 311–315.
- Mamay, S.H. 1989. *Evolsonia*, a new genus of Gigantopteridaceae from the Lower Permian Vale Formation, North-Central Texas. American Journal of Botany 76: 1299–1311.
- Mamay, S.H. & Breed, W.J. 1970. Early Permian plants from the Cutler Formation in Monument Valley, Utah. U.S. Geological Survey, Prof. Paper 700-B: B109–B117.
- Mamay, S.H., Chaney, D.S. & DiMichele, W.A. 2009. *Comia*, a seed plant possibly of peltaspermous affinity: A brief review of the genus and description of two new species from the Early Permian (Artinskian) of Texas, *C. greggii* sp. nov. and *C. craddockii* sp. nov. International Journal of Plant Sciences 170(2): 267–282.
- Mamay, S.H., Miller, J.M., Rohr, D.M. & Stein Jr., W.E. 1984. *Delnortea*, a new genus of Permian plants from West Texas. Phytologia 60: 345–346.
- Mamay S.H., Miller, J.M., Rohr, D.M. & Stein Jr., W.E. 1988. Foliar morphology and anatomy of the gigantopterid plant *Delnortea abbottiae*, from the Lower Permian of West Texas. American Journal of Botany 75: 1409–1433.
- Matysová, P., Rößler, R., Gotze, J., Leichmann, J., Forbes, G., Taylor, E.L., Sakala, J. & Grygar, T. 2010. Alluvial and volcanic pathways to silicified plant stems (Upper Carboniferous-Triassic) and their taphonomic and paleoenvironmental meaning. Paleogeography Paleoceanography Paleoclimatology 292: 127–143.
- Matysová, P., Booij, M., Crow, M.C., Hasibuan, F., Perdoni, A.P., van Waveren, I.M. & Donovan, S.K. in press. Burial and preservation of a fossil forest on an Early Permian (Asselian) volcano (Merangin River, Sumatra, Indonesia). Geological Journal.
- McNeill, J., Barrie, F.R., Burdet, H.M., Demoulin, V., Hawksworth, D.L., Marhold, K., Nicolson, D.D.H., Prado, J., da Silva, P.C., Skog, J.E., Wiersema, J.H. & Turland, N.J. 2006. International Code of Botanical Nomenclature (Vienna Code). Regnum Vegetabile 146. Vienna: A.R.G. Gantner Verlag KG.
- Metcalfe, I. 2006. Palaeozoic and Mesozoic tectonic evolution and palaeogeography of East Asian crustal fragments: The Korean Peninsula in context. Gondwana Research 9: 24–46.
- Metcalfe, I. 2009. Comment on “An alternative plate tectonic model for the Palaeozoic-Early Mesozoic Palaeotethyan evolution of Southeast Asia (Northern Thailand-Burma)” by O.M. Ferrari, C. Hochard and Stampfli, G.M. Tectonophysics 451: 346–365.

- Metcalfe, I. 2011. Tectonic framework and Phanerozoic evolution of Sundaland. *Gondwana Research* 19: 3-21.
- Metcalfe, I. & Sone, M. 2008. Biostratigraphy and palaeobiogeography of Lower Permian (lower Kungurian) conodonts from the Tak Fa Formation (Saraburi Limestone), Thailand. *Palaeogeography, Palaeoclimatology, Palaeoecology* 257: 139-151.
- Meyen, S.V. 1982. The Carboniferous and Permian floras of Angaraland (a synthesis). *Biol. Mem.* 7: 1-109.
- Meyen, S.V. 1984. Basic features of gymnosperm systematics and phylogeny as evidenced by the fossil record. *The Botanical Review* 50: 1-111.
- Meyen, S.V. 1987. Fundamentals of palaeobotany. London: Chapman and Hall.
- Montañez, I.P. & Poulsen, C.J. 2013. The Late Paleozoic Ice Age: An Evolving Paradigm. *Annu. Rev. Earth Planet. Sci.* 41: 629-56.
- Montanez, I.P., Tabor, N.J., Niemeier, D., DiMichele, W.A., Frank, T.D., Fielding, C.R., Isbell, J.L., Birgenheier, L.P. & Rygel, M.C. 2007. CO₂-forced climate and vegetation instability during Late Palaeozoic deglaciation. *Science* 315: 87-91.
- Naugolnykh, S.V. 1999. A new species of *Compsopteris Zalessky* from the Upper Permian of the Kama River Basin (Perm Region). *Paleontological Journal* 33(6): 686-697.
- Naugolnykh, S.V. 2000. Mixed Permian floras of Eurasia: a new concept with significance for paleophytogeographic reconstructions. *Paleontological Journal* 34, Suppl.1: S99-S105.
- Opluštík, S., Pšenička, J., Libertín, M., Bashforth, A.R., Šimůnek, Z., Drábková, J. & Daškova, J. 2009. A Middle Pennsylvanian (Bolsovian) peat-forming forest preserved *in situ* in volcanic ash of the Whetstone Horizon in the Radnice Basin, Czech Republic. *Review of Palaeobotany and Palynology* 155: 234-274.
- Parkhurst, D.F. & Loucks, O.L. 1972. Optimal Leaf Size in Relation to Environment. *The Journal of Ecology* 60 (2): 505-537.
- Patzkowsky, M.E., Smith, L.H., Markwick, P.J., Engberts, C.J. & Gyllenhaal, E.D. 1991. Application of the Fujita-Ziegler paleoclimate model: Early Permian and Late Cretaceous examples. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 86: 67-85.
- Pearson, T. & Scott, A.C. 1999. Large palynomorphs and debris. In: *Fossil Plants and Spores: Modern Techniques*, Jones, T.P., Rowe, N.P. (eds). Geological Society: London; 245-250.
- Phillips, T.L., Peppers, R.A. & DiMichele, W.A. 1985. Stratigraphic and interregional changes in Pennsylvanian coal-swamp vegetation: Environmental inferences. *International Journal of Coal Geology* 5: 43-109.
- Phillips, T.L. & DiMichele, W.A. 1992. Comparative ecology and life-history biology of arborescent lycopsids in Late Carboniferous swamps of Euramerica. *Annals of the Missouri Botanical Garden* 79: 560-588.

- Philippe, M. 1993. Nomenclature générique des trachéidoxyles fossiles mésozoïques à champs araucarioïdes. *Taxon* 42: 74-80.
- Philippe, M. 2011. How many species of *Araucarioxylon*? *Paleovol* 10: 201-208.
- Playford, G. & Rigby, J.F. 2008. Permian palynoflora of the A nim and Aiduna formations, West Papua. *Revista española de micropaleontología* 40(1-2): 1-57.
- Poli, D., Jacobs, M. & Cooke, T.J. 2003. Auxin regulation of axial growth in bryophyte sporophytes: Its potential significance for the evolution of early land plants. *American Journal of Botany* 90: 1405-1415.
- Poole, I. 1994. "Twig"- wood anatomical characters as palaeoecological indicators. *Review of Palaeobotany and Palynology* 81: 33-52.
- Posthumus, O. 1927. Some remarks concerning the Paleozoic Flora of Djambi. *Proceed. Kon. Akad. Amsterdam* 30: 628-634.
- Potonié, H. 1902. Über die fossilen Filicales im Allgemeinen und die Reste derselben zweifelhafter Verwandtschaft. In: Engler A, Pranti K, eds. *Die natürlichen Pflanzenfamilien*, 1. Leipzig, 473-515.
- Ramalho, R.S., Quartau, R., Trenhaile, A.S., Mitchell, N.C., Woodroffe, C.D. & Avila, S.P. 2013. Coastal evolution on volcanic oceanic islands: a complex interplay between volcanism, erosion, sedimentation, sea-level change and biogenic production. *Earth-Science Reviews* 127: 140-170.
- Raskassova, E.S. 1963. Novyi Predstavitel' Roda Comia iz Noril'skogo Raiona. *Uchenya Zapiski Paleontologiya i Biostratigrafiya* 3: 63-73.
- Raymond, A., Lambert, L., Costanza, S., Slone, E.J. & Cutlip, P.C. 2010. Cordaitans in paleotropical wetlands: An ecological re-evaluation. *International Journal of Coal Geology*: 248-265.
- Raymond, A. & Phillips, T.L. 1983. Evidence for an Upper Carboniferous mangrove community. In: Teas, H.J., ed., *Tasks for vegetation science*. Dr. W. Junk Publishers, The Hague, 19-30.
- Read, C.B. & Mamay, S.H. 1964. Upper Palaeozoic floral zones and floral provinces of the United States. *U.S. Geological Survey Professional Paper* 454-K: K1-K32.
- Rees, P.McA. & Cleal, C.J. 1993. Marked polymorphism in *Archangelskya furcata*, a pteridospermous frond from the Jurassic of Antarctica. *Special Papers in Palaeontology* 49: 85-100.
- Remy, W. 1953. Untersuchungen über einige Fruktifikationen von Farnen und Pteridospermen aus dem mitteleuropäischen Karbon und Perm. *Abhandlungen der Deutschen Akademie der Wissenschaften zu Berlin, Klasse für Mathematik und allgemeine Naturwissenschaften*, Jahrgang 1952 (2), 1-38.
- Renault, B. 1879. Structure comparée de quelques tiges de la flore Carbonifère. *Nouv. Arch. Mus.* 2: 213-348
- Ricardo-Branco, F. 2008. Venezuelan paleoflora of the Pennsylvanian-Early Permian: Paleobiogeographical relationships to central and western equatorial Pangea. *Gondwana Research* 14: 297-305.

- Rigby, J.F. 1996. The significance of a Permian flora from Irian Jaya (West New Guinea) containing elements related to coeval floras of Gondwanaland and Cathaysia land. *The Palaeobotanist* 45: 295-302.
- Rigby, J.F. 2001. A review of the Early Permian flora from Papua (West New Guinea). In: Metcalfe, I., Smith, J.M.B., Morwood, M. & Davidson, I. (Eds.). *Faunal and floral migrations and evolution in SE Asia-Australasia*. A.A. Balkema Publishers, Lisse, 85-95.
- Rößler, R., Philippe, M., Van Konijnenburg-Van Cittert, J.H.A., McLoughlin, S., Sakala, J., Zijlstra, G. et al. 2014. Which name(s) should be used for *Araucaria*-like fossil wood? – Results of a poll. *Taxon* 63: 177-184.
- Rößler, R., Zierold, T., Feng, Z., Kretzschmar, R., Merbitz, M., Annacker, V. & Schneider, J.W. 2012. A snapshot of an Early Permian ecosystem preserved by explosive volcanism: new results from the Petrified Forest of Chemnitz, Germany. *Palaios* 27: 814-34.
- Roth-Nebelsick, A., Uhl, D., Mosbrugger, V. & Kerp, H. 2001. Evolution and function of leaf venation architecture: a review. *Annals of Botany* 87: 553-566.
- Rowley, D.B., Raymond, A., Parrish, J.T., Lottes, A.L., Scotese, C.R. & Ziegler, A.M. 1985. Carboniferous paleogeographic, phytogeographic, and paleoclimatic reconstructions. In: Phillips, T.L. & Cecil, C.B. (Eds.), *Paleoclimatic Controls on Coal Resources of the Pennsylvanian System of North America*. International Journal of Coal Geology 5: 7-42.
- Royer, D.L., Wilf, P., Janesko, D.A., Kowalski, E.A. & Dilcher, D.L. 2005. Correlations of climate and plant ecology to leaf size and shape: potential proxies for the fossil record. *American Journal of Botany* 92(7): 1141-1151.
- Scarpella, E., Marcos, D., Friml, J. & Berleth, Th. 2006. Control of leaf vascular patterning by polar auxin transport. *Genes & Development* 20: 1015-1027.
- Schenk, A. 1883. Pflanzen aus der Steinkohlen-Formation. In: Richthofen, F. von, Beiträge zur Paläontologie von China, 4. Berlin: Verlag von Dietrich Reimer, 211-269.
- Schimper, W.Ph. 1870-1872. *Traité de paléontologie végétale*. Tome second. J.B. Baillière et Fils, Paris.
- Schmid, R. 1981. Descriptive nomenclature and classification of pyroclastic deposits and fragments: recommendations of the IUGS Subcommission on the Systematics of Igneous Rocks. *Geology* 9: 41-43.
- Shannon, C.E. 1948. A mathematical theory of communication. *The Bell System Technical Journal* 27: 379-423, 623-656.
- Shen G.-L. 1995. Permian Floras. In: Li Xingxue (ed.) *Fossil Floras of China through the geological ages*, Guangdong Science and Technology Press, Guangzhou, China, 127-223.
- Shi, G.R. 2006. The marine Permian of East and Northeast Asia: an overview of biostratigraphy, palaeobiogeography and palaeogeographical implications. *Journal of Asian Earth Sciences* 26: 175-206.

- Schultze-Motel, J. 1962. Literatur über die Gattung *Dadoxylon* Endlicher (*Araucarioxylon* Kraus). Geologie 11: 716-731.
- Schultze-Motel, J., 1966. Gymnospermen-Hölzer aus den oberkretazischen Umzamba-Schichten von Ost-Pondoland (S-Afrika). Senckenbergiana lethaea 47(3), 279-337.
- Schweitzer, H-J. 1958. Die fossilen Diptercarpaceen-Hölzer. Palaeontographica Abt. B 105: 1-66.
- Šimůnek, Z. 1996. Carboniferous Alethopterids of the Karviná and Žacléř Formations (Czech Republic). Sborník geologických věd, Paleontologie 33: 5-37.
- Šimůnek, Z. 2000. Cuticles of Cordaites from the Westphalian, Stephanian and Autunian of the Bohemian Massif. Acta Palaeobotanica 40: 25-34.
- Šimůnek, Z. 2007. New classification of the genus Cordaites from the Carboniferous and Permian of the Bohemian Massif based on cuticle morphology. Acta Musei Nationalis Pragae, Ser. B, Historia Naturalis 62: 97-210.
- Sisó, S., Camarero, J. & Gil-Pelegín, E. 2001, Relationship between hydraulic resistance and leaf morphology in broadleaf *Quercus* species: a new interpretation of leaf lobation. Trees 15: 341-345.
- Stamberg, S. 1997. New discoveries of palaeoniscoid fishes and other fauna and flora from the northern region of Boskovice Furrow, Czech Republic. Journal of the Czech Geological Society 42(1-2): 111-120.
- Stephens, M.A. 1974. EDF statistics for goodness of fit and some comparisons. J. Amer. Statist. Assoc. 69 (347): 730-737.
- Stephenson, N.L. & Demetry A. 1995. Estimating ages of giant sequoias. Canadian Journal of Forestry Research 25: 223-233.
- Sterzel, J.T. 1918. Die organischen Reste des Kulms und Rotliegenden der Gegend von Chemnitz. Abhandlungen der Mathematisch-Physischen Klasse der Königlichen Sächsischen Gesellschaft der Wissenschaften, XXXV (V): 205-315.
- Stidd, B. M. 1981. The current status of medullosan seed ferns. Review of Palaeobotany and Palynology 32: 63-101.
- Stockmans, F. & Mathieu, F.F. 1939. La flore paléozoïque du Bassin houiller de Kaiping (Chine). Patrimonie Mus. Roy. Hist. Nat. Belgique, Bruxelles, 49-165.
- Sun Ke-Qin. 2006. The Cathaysia flora and the mixed Late Permian Cathaysian-Angaran floras in East Asia. Journal of Integrative Plant Biology 48(4): 381-389.
- Sun, Y., Jaochimski, M.M., Wignall, P.B., Yan, C., Chen, Y., Jiang, H., Wang, L. & Lai, X. 2012. Lethally hot temperatures during the Early Triassic greenhouse. Science 388: 366-370.
- Suwarna, N. 2006. Permian Mengkarang coal facies and environment, based on organic petrology studies. Jurnal Geologi Indonesia 1: 1-8.
- Suwarna, N., Suharsono, Amiruddin & Hermanto, B. 1998. Geological map of the Bangko Quadrangle, Sumatera Scale 1, 100 000: Geological Research and Development Centre, Bandung.

- Suwarna, N., Suharsono, Gafoer, S., Amin, T.C., Kusnama & Hermanto, B. 1994. Geology of the Sarolangun, Sumatra. Geological Research and Development Center, Bandung, West Java, 100 pp.
- Sze, H.C. 1954a. A preliminary study of the fossil flora of the Pingtingshan Coal Field in central Honan. *Acta Palaeobotanica Sinica* 2 (3): 289-306.
- Sze, H.C. 1954b. On the occurrence of a new species of *Callipteris* of the Kusnezk type in the Cathaysia Flora in Northwestern Shansi. *Acta Scientia Sinica* III(1): 97-103.
- Takahashi, T. & Tsujimoto, H. 2000. A mechanical model for the Merapi-type pyroclastic flow. *Journal of Volcanology and geothermal Research* 98: 91-115.
- Taylor T.N. & Taylor E.L. 1993. The biology and evolution of fossil plants. Englewood Cliffs: Prentice Hall.
- Thompson, M.L. 1936. Lower Permian fusulinids from Sumatra. *Journal of Paleontology* 10 (7): 587-592.
- Thouret, J.C. 1999. Volcanic geomorphology-an overview: *Earth-Science Reviews* 47: 95-131.
- Tobler, A. 1917. Über Deckenbau im Gebiet von Djambi. *Verhdl. d. Naturf. Ges in Basel*, XXVIII: 2: 123-147.
- Tobler, A. 1922. Djambi-verslag. Uitkomsten van het geologisch en mijnbouwkundig onderzoek in de Residentie Djambi, 1906-1912. *Jaarboek van het Mijnwezen van Nederlandsch Oost-Indië*, Verhandelingen 1919, 48: 1-585.
- Tobler, A. 1923. Unsere paläontologische Kenntnis von Sumatra. *Ecologae Geologicae Helvetiae* 18: 313-342.
- Tobler, A. 1924. Fundorte fossiler Pflanzen im südlichen Sumatra. In: R. Kräusel, Fossile Hölzer aus dem Tertiär von Süd-Sumatra. *Verhdl. Geol. Mijnbk. Genootschap v. Nederland en Kol. Geologische Serie V*: 236-237.
- Tomescu, A.M.F. 2008. Megaphylls, microphylls and the evolution of leaf development. *Trends in Plant Science* 14: 5-12.
- Traiser, C., Klotz, S., Uhl, D. & Mosbrugger, V. 2005. Environmental signals from leaves – a physiognomic analysis of European vegetation. *New Phytologist* 166: 465-484.
- Trivett, M.L. & Pigg, K.B. 1996. A survey of reticulate venation among fossil and living land plants. In: Taylor DW, Hickey LJ, eds. *Flowering plant origin, evolution and phylogeny*. New York: Chapman and Hall, 8-31.
- Ueno K., Nishikawa S., Van Waveren I.M., Booi, M., Hasibuan F., Suyoko, Iskandar E.P.A., De Boer P.L., King C.I., De Leeuw J.H.V.M. & Chaney D.S. 2007. Early Permian fusuline faunas from Jambi, Sumatra, Indonesia: Faunal characteristics and paleobiogeographic implications. Abstract of 16th International Congress on the Carboniferous and Permian. *Journal of Stratigraphy* 31(supplement I): 138-139.

- Uhl, D. & Mosbrugger, V. 2002. Some data on the potential use of leaf venation density as a palaeoclimatic proxy in leaves from the Permocarboniferous. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 223 (3): 405-432.
- Uhl, D., Walther, H. & Mosbrugger, V. 2002. Leaf venation density in angiosperm leaves as related to climatic and environmental conditions – problems and potential for palaeoclimatology. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 224 (1): 49-95.
- Utami, W., Herdianita, N.R. & Atmaja, R.W. 2014. The Effect of Temperature and pH on the Formation of Silica Scaling of Dieng Geothermal Field, Central Java, Indonesia. Proceedings, Thirty-Ninth Workshop on Geothermal Reservoir Engineering Stanford University, Stanford, California, February 24-26: 202.
- Vachard, D. 1989. A rich algal Microflora from the Lower Permian of Jambi Province. In: Fontaine, H., Gafoer, S. (Eds.) *The Pre-Tertiary Fossils of Sumatra and their Environments*. CCOP Technical Secretariat, 59-69.
- Van de Laar, J.G.M. & Van der Zwan, C.J., 1996. Palynostratigraphy and palynofacies reconstruction of the Carboniferous of borehole “De Lutte-06” (East Twente, the Netherlands). *Mededelingen Rijks Geologische Dienst* 55: 61-82.
- Van Waveren, I. M. 1993. Planktonic organic matter in surficial sediments of the Banda Sea (Indonesia) -a palynological approach. *Geologica Ultraiectina, Mededelingen van de Faculteit Aardwetenschappen Universiteit Utrecht* 104. 237pp.
- Van Waveren, I.M., Booij, M. & Van Konijnenburg-van Cittert, J.H.A. 2006. Basinal and extrabasinal fossil plant associations from the Early Permian of Sumatra (Jambi province), Indonesia. (Abstract) *Abstracts of the 7th European Palaeobotany-Palynology Conference (7EPPC)*: 157.
- Van Waveren, I.M., Hasibuan, F., Suyoko, Makmur, de Boer, P.L., Chaney, D., Ueno, K., Booij, M., Iskandar, E.A.P., King, C., De Leeuw, J.M. & Van Konijnenburg-van Cittert, J.H.A. 2005. Taphonomy, Palaeobotany and Sedimentology of the Mengkarang Formation (Early Permian, Jambi, Sumatra, Indonesia). In: Lucas, E. (Ed.), *The Nonmarine Permian*. *Bulletin of the New Mexico Museum of Natural History and Science* 30: 333-341.
- Van Waveren, I.M., Iskandar, E.A.P., Booij, M. & Van Konijnenburg-van Cittert, J.H.A. 2007. Composition and palaeogeographic position of the Lower Permian Jambi flora from Sumatra. *Scripta Geologica* 135: 1-28.
- Van Waveren, I. M. & Visscher, H. 1994. Palynological analysis of the composition and selective preservation of particulate organic matter in surface sediments from a high productivity deep sea area (eastern Banda Sea, Indonesia). *Paleogeography Paleoclimatology Paleoecology* 112: 85-111.
- Van Waveren, I.M., Crow, M.J., Hasibuan, F., Schmitz, M.D., Hettling, H., Booij, M., Zwaan, H., Braumuller, Y. In preparation. Climatic and paleoecologic conditions for the early appearances and the late presence of paleofloral taxa on a far eastern Permian volcano.
- Vogellehner, D. 1964. Zur Nomenklatur der fossilen Holzgattung *Dadoxylon* Endlicher 1874. *Taxon* 13(7): 233-237.

- Vozenin-Serra, C. 1985. Bois homoxylés du Permien inférieur de Sumatra. Implications paléogéographiques. 110e Congrès national des Sociétés savantes, Montpellier, 1985, fasc. V: 55-63.
- Vozenin-Serra, C. 1989. Two gymnospermous woods from the Lower Permian of Jambi, Sumatra. In: Fontaine, H., Gafoer, S. (eds.) The Pre-Tertiary fossils of Sumatra and their environments The Pre-Tertiary Fossils of Sumatra and their Environments. CCOP Technical Secretariat, 168-171.
- Wagner, R. H. 1961. Some *Alethopteridae* from the South-Limburg coal field. Mededelingen van de Geologische Stichting, N. S. 14: 5-13.
- Wagner, R. H. 1968. Upper Westphalian and Stephanian species of *Alethopteris* from Europe, Asia Minor and North America. Mededelingen van de Rijks Geologische Dienst, Serie C, III-1, no. 6, 318 p., 55 fig., 64 pl. in a separate atlas, Serie C, III-1, no.5.
- Wagner, R.H. & Mayoral, E.J. 2007. The Early Permian of Valdevar in Sevilla province, SW Spain, basin history and climatic/palaeogeographic implications: Journal of Iberian Geology 33: 93-124
- Wang, Z.-Q. 1996. Past global floristic changes: the Permian great Eurasian floral interchange. Palaeontology 39(1): 189-217.
- Wang, Z.-Q. 1997. Permian *Supaia* fronds and an associated *Autunia* fructification from Shanxi, China. Palaeontology 40(1): 245-277.
- Wang Z.-Q. 1999. *Gigantonoclea*: An enigmatic Permian plant from North China. Palaeontology 42: 329-373.
- Weber, R. 1997. How old is the Triassic flora of Sonora and Tamaulipas and news on Leonardian floras in Puebla and Hidalgo, Mexico. Revista Mexicana de Ciencias Geológicas 14(2): 225-243.
- Wheeler, E. A. & Baas, P. 1998. Wood identification: review. International Association of Wood Anatomists' Bulletin 19: 241–264.
- Whitaker, M.F. 1984. The usage of palynology in definition of Troll Field geology: 6th Offshore Northern Seas Conference and Exhibition, Stavanger 1984, Paper G6.
- White, D., 1893. A new Taeniopteroid fern and its allies. Bull. of the Geological Society of America 4, 119-132.
- White, D. 1912. The characters of the fossil plant *Gigantopteris* Schenk and its occurrence in North America. Proceedings of the United States National Museum 41: 493-516.
- White, D. 1929. Flora of the Hermit Shale, Grand Canyon, Arizona. Publication, vol. 405. Carnegie Institution of Washington, 1-221
- Wolfe, J.A. 1979. Temperature parameters of humid to mesic forests of eastern Asia and relation to forests of other regions of the northern hemisphere and Australasia. U.S. Geological Survey Professional Paper 1106: 1-37.
- Wu S.-Z. 1985. New Late Permian *Comia* species from Kuqa, Xinjiang. Geological Review 31: 213-216.
- Yao, Z.-Q. 1978. On the age of “*Gigantopteris* Coal Series” and *Gigantopteris*-flora in South China. Acta Palaeontologica Sinica 17: 81-89.

- Yao, Z.-Q. & Crane, P.R. 1986. Gigantopterid leaves with cuticles from the Lower Permian of China. American Journal of Botany 73: 715-716.
- Yao, Z.-Q. & Liu L.-J. 2002. Discovery of dichotomously forked gigantopterid fronds from China, with special comparison between Asian and American gigantopterids. Acta Palaeontologica Sinica 41: 314-321.
- Yao, Z.-Q. & Liu L.-J. 2004. A new gigantopterid plant with cuticles from the Permian of South China. Review of Palaeobotany and Palynology 131: 29-48.
- Yang G.-X. 1987. The evolution of the Permian gigantopterids in Yuxian County, Western Henan and its geological significance. Geoscience 1: 173-195. (In Chinese with English summary)
- Zalessky, M.D. 1927. Flore Permienne des limites ouraliennes de l'Angaride (Atlas). Mémoires du Comité Géologique, Nouvelle série, Livraison 176, pl.II, fig.1&2, pl.VII, fig.1-3, pl.VIII, fig.1, pl.X, fig.4.
- Zalessky, M.D. 1934. Observations sur les végétaux permiens du bassin de la Petchora. I. Bulletin de l'Académie des Sciences de l'URSS 2-3: 241-290.
- Zeiller, R. 1907. Note sur quelques empreintes végétales des gites de charbon du Yunnan méridional. Annales des Mines, 10e série, 10: 5-27.
- Zhao, X.-H., Mo, Z.-G., Zhang, S.-Z. & Yao, Z.-Q. 1980. Late Permian flora in Western Guizhou and Eastern Yunnan. In: Nanking Institute of Geology and Palaeontology, Academia Sinica, eds. Stratigraphy and palaeontology of Upper Permian coalbearing formation in Western Guizhou and Eastern Yunnan, China. Beijing: Science Press, 70-122. (in Chinese with English summary)
- Zhu, J.-N. & Zhang, X.-S. 1995. *Trinerviopteris cardiophylla* (Zhu et Geng) Zhu gen. et comb. nov. and the classification of gigantopterids. Acta Botanica Sinica 37: 314-320 (in Chinese with English summary).
- Ziegler, A.M. 1990. Phytogeographic patterns and continental configurations during the Permian Period. In: McKerrow, W.S. & Scotese, C.R. (eds.), 1990, Palaeozoic Palaeogeography and Biogeography, Geological Society Memoir 12: 363-379.
- Ziegler, A.M., Eshel, G., McAllister Rees, P., Rothfus, T.A., Rowley, D.B. & Sunderlin, D. 2003. Tracing the tropics across land and sea: Permian to present. Lethaia 36: 227-254.
- Ziegler, A.M., Hulver, M.L. & Rowley, D.B. 1997. Permian world topography and climate. In: Martini, I.P. (Ed.), Late glacial and post-glacial environmental changes – Quaternary, Carboniferous-Permian and Proterozoic. Oxford University Press, New York, 111-146.
- Zodrow, E.L. 2004. Note on the different kinds of attachments of the trigonocarpalean (Medullosales) ovules from the Pennsylvanian Sydney Coalfield, Canada. Atlantic Geology 40: 197-206.
- Zodrow, E.L. & Cleal, C.J. 1998. Revision of the Pteridosperm foliage Alethopteris and Lonchopteridium (Upper Carboniferous), Sydney Coalfield, Nova Scotia, Canada. Palaeographica Abt. B 247 (3-6): 65-122.
- Zurakowski, K.A. & Gifford, E.M. 1988. Quantitative studies of pinnules development in the ferns *Adiantum raddianum* and *Cheilanthes viridis*. American Journal of Botany 75: 1559-1570.

Zwieniecki, M.A., Boyce, C.K. & Holbrook, N.M. 2004. Hydraulic limitations imposed by crown placement determine final size and shape of *Quercus rubra* L. leaves. Plant, Cell and Environment 27: 357-365.

Zwierzycki, J. 1935. Die Ergebnisse der paläobotanischen Djambi-expedition 1925, 1. Die geologischen Ergebnisse. Jaarboek Mijnwezen Nederlandsch-Indië 59: 1-70.