



Universiteit
Leiden
The Netherlands

Transport coefficients and low energy excitations of a strongly interacting holographic fluid

Poovuttikul, N.

Citation

Poovuttikul, N. (2017, November 16). *Transport coefficients and low energy excitations of a strongly interacting holographic fluid*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/57561>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/57561>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/57561> holds various files of this Leiden University dissertation

Author: Poovuttikul, N.

Title: Transport coefficients and low energy excitations of a strongly interacting holographic fluid

Date: 2017-11-16

Casimir PHD Series 2017-35
ISBN 978-90-8593-319-9

TRANSPORT COEFFICIENTS AND LOW ENERGY EXCITATIONS
OF A STRONGLY INTERACTING HOLOGRAPHIC FLUID

N. POOVUTTİKUL

TRANSPORT COEFFICIENTS AND LOW ENERGY EXCITATIONS OF A STRONGLY INTERACTING HOLOGRAPHIC FLUID

N. Poovuttikul

$$\tau_{\parallel} = \frac{2m^2}{N_c^2} \left(\frac{v}{2\pi\sqrt{w}} \right),$$

$$\mathcal{J}_{5,CS}^{\mu} = 2\kappa A_t B_5^{\mu} + 2\gamma V_t B^{\mu} + \left(\kappa A_t^2 + \lambda \frac{g(r^2, f')^2}{2f} \right) \omega^{\mu}$$

$$\tau_{\perp} = \frac{2\pi^2}{N_c^2} \left(\frac{\sqrt{w}}{2\pi r_h} \right) \left(\frac{b_{\perp}(\frac{1}{2})}{b_{\perp}(0)} \right)^2,$$

$$S = \frac{N_c^2}{8\pi^2} \left[\int d^5 X \sqrt{-G} \left(R + 12 - \frac{1}{3e_H^2} H_{abc} H^{abc} \right) + \int_{\partial M} d^4 x \sqrt{-g} \left(2 \text{Tr} K - 6 + \frac{1}{e_H^2} \mathcal{H}_{\mu\nu} \mathcal{H}^{\mu\nu} \ln C \right) \right],$$

$$\lim_{\omega \rightarrow 0} \omega^{-1} \text{Im} G_{T^{\mu\nu}}^R(\omega, 0) < s/4\pi$$

$$\frac{\omega^2}{k^2} = \left(\frac{\mu\rho \cos^2 \theta}{\varepsilon + p} \right) - i \left[\left(\frac{\mu r_{\perp}}{\rho} + \frac{\eta_{\parallel}}{\varepsilon + p} \right) \cos^2 \theta + \left(\frac{\mu r_{\parallel}}{\rho} + \frac{\eta_{\perp}}{\varepsilon + p} \right) \sin^2 \theta \right] \omega + \frac{\mu}{2\rho(\varepsilon + p)} (r_{\perp} \cos^2 \theta + 2r_{\parallel} \sin^2 \theta) (\eta_{\perp} \sin^2 \theta + \eta_{\parallel} \cos^2 \theta) k^2.$$

$$\sigma_{J_5\omega} = \kappa\mu_5^2 + \gamma\mu^2 + 2\lambda(2\pi T)^2$$