



Universiteit
Leiden
The Netherlands

Bone marrow transplantation in mice as a tool to study M2 macrophage activation in atherogenesis

Ren, B.

Citation

Ren, B. (2017, December 14). *Bone marrow transplantation in mice as a tool to study M2 macrophage activation in atherogenesis*. Retrieved from <https://hdl.handle.net/1887/57798>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/57798>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/57798> holds various files of this Leiden University dissertation

Author: Ren, Baoyan

Title: Bone marrow transplantation in mice as a tool to study M2 macrophage activation in atherogenesis

Date: 2017-12-14

1.

General introduction

Table of contents

Chapter 1 General introduction	9
1 Atherosclerosis	11
1.1 Lipoproteins: metabolism and association with atherosclerosis	11
1.2 General pathogenesis underlying atherosclerotic lesion development.....	14
2 The role of monocytes and macrophages in the pathology of atherosclerosis.....	15
2.1 Monocyte heterogeneity	15
2.2 M1/M2 macrophages and atherosclerosis	16
2.3 Macrophage phenotype and foam cell susceptibility	17
2.4 Macrophage phenotype switch	17
3 Experimental mouse models and strategies for studying atherosclerosis	17
3.1 Mice and atherosclerosis susceptibility	17
3.2 ApoE KO mice and LDLr KO mice.....	18
3.3 Bone marrow transplantation.....	18
4 Signaling pathways and atherosclerosis	19
4.1 Nitric oxide/L-arginine pathway in atherosclerosis.....	20
4.2 Protein kinase B (Akt) in atherosclerosis	22
4.3 MAPKs/MKPs pathway in atherosclerosis.....	23
4.4 Usf transcription factors and atherosclerosis	25
5 Outline of this Thesis.....	26
6 References.....	26

1 Atherosclerosis

Atherosclerosis is a blood vessel narrowing and hardening disease characterized by the deposition of cholesterol locally in the arterial wall, leading to a low-grade chronic inflammation.¹ Atherosclerotic lesions take decades to become large enough as to have a significant effect and cause cardiovascular complications in humans.² Atherosclerosis can eventually lead to serious problems such as chest pain (angina), heart failure, heart attack, stroke, ischemic attack, aneurysms, or even death.³ Nowadays, atherosclerotic cardiovascular disease is the leading cause of mortality worldwide, accounting for 31% of global deaths in 2015.⁴ Although it is closely related to contemporary lifestyles, atherosclerosis is not only found in modern human beings.⁵ In contrast, the disease has been found in 4000 years old mummies and the earliest literature that described the pathological changes of atherosclerosis can be traced back to 442 years ago. In the year 1575, the Italian anatomist Gabriel Fallopius had already described calcification of the arterial wall, a pathological phenomenon of atherosclerotic plaques.⁶ Later on, several researchers and medical doctors had observed atherosclerosis in large arteries,⁷⁻⁹ and proposed a connection of atherosclerosis with angina and ischemic heart disease.¹⁰ The first description of plaque rupture was reported in 1844.¹⁰ In 1829, Jean Lobstein for the first time introduced the word “arteriosclerosis” in his unfinished book “*Traité d’ Anatomie Pathologique*” which was ultimately published in the year 1933,^{11,12} almost 100 years after his death. However, in general the German pathologist Felix Marchand (1904) is recognized as the first using the term “atherosclerosis”, stemming from the Greek words of porridge “athero” and hardening “sclerosis”, to describe the fat-rich materials that accumulated inside a hardened artery.^{13,14}

Since its discovery, researchers have been trying to uncover the etiopathogenesis of atherosclerosis. At the end of 18th century, theories underlying atherosclerosis development were proposed by Carl von Rokitansky and Rudolf Virchow¹⁵ who both recognized the presence of inflammation. However, it was unclear whether inflammation played a causative role. Carl von Rokitansky considered atherosclerosis to be the result of the buildup of fibrin or other blood elements, which subsequently was modified to a lipid-rich plaque in the arterial wall.¹⁴ In contrast, Rudolf Virchow suggested that the inflammatory response to lipid insudation or intimal injury is the cause of atherosclerosis.¹⁵ The exact pathogenesis of atherosclerosis remained unclear until the “response to injury hypothesis”, initially proposed by Rudolf Virchow and revived by Russell Ross in 1999, became a widely accepted theory.¹⁶

1.1 Lipoproteins: metabolism and association with atherosclerosis

Hyperlipidemia plays a leading role in triggering and promoting atherosclerosis development.¹⁷ Lipoproteins, being the main carrier of cholesterol in the circulation, were firstly associated to cardiovascular disease in 1949 by John Gofman and his colleagues. They found that increased levels of low-density lipoprotein (LDL) cholesterol are associated with an increased cardiovascular risk, and that patients with familial hypercholesterolemia are predisposed to the development of premature atherosclerosis.¹⁸ These patients have an overall cholesterol elevation in their plasma, which can mainly be attributed to an increase in LDL and intermediate-density lipoprotein (IDL).¹⁴ The hypothesis that high plasma lipid levels are associated with increased cardiovascular risk was further supported by a cooperative study performed in the 1950s and 1960s, which confirmed the connection of cardiovascular risk to plasma cholesterol levels using patient cohorts from seven different countries.¹⁹ Circulating lipids are transported by lipoproteins, particles composed of a shell of a monolayer of phospholipids with free cholesterol and apolipoproteins (Apo) and a lipid-

rich core constituted of esterified cholesterol and triglycerides.²⁰ Based on the proportion of each component as well as particle density, lipoproteins are classed into 5 groups: high-density lipoprotein (HDL), low-density lipoprotein (LDL), intermediate-density lipoprotein (IDL), very-low-density lipoprotein (VLDL), and chylomicrons. Except for HDL, all these subclasses contain apolipoprotein B (ApoB) as their major apolipoprotein and are considered athero-promoting lipoproteins. In contrast, HDL has apolipoprotein A as its primary apolipoprotein and is considered to act as an athero-protective lipoprotein.

1.1.1 ApoB-containing lipoproteins and atherosclerosis

Chylomicrons, ApoB48-containing lipoproteins, are the major carriers for lipids absorbed from the diet in the intestine, and represent a major source of triglycerides (TG) for various tissues. In addition to TG, chylomicrons transport cholesterol, but only a limited amount.²¹ Therefore, for quite a long time chylomicrons were believed not to contribute to atherogenesis.^{22,23} However, later on ApoB48 was found in atherosclerotic plaques^{24,25} and an ApoB 48-specific receptor was detected in human and murine macrophages,^{26,27} thereby highlighting the contribution of chylomicrons to atherosclerosis development. In general, increased chylomicron levels seem to generate a pro-atherogenic profile, however, there is still not much evidence showing a direct link between high chylomicron levels and atherogenesis.²⁸

In humans, and specifically in women with elevated TG levels, increased risk for cardiovascular events was shown in a 11.4 years follow-up study in America.²⁹ Interestingly, both fasting and non-fasting TG levels are associated with cardiovascular disease, with postprandial TG levels showing the strongest association with future cardiovascular risk.²⁹⁻³¹ Interestingly, the strong linear correlation between plasma TG and atherosclerosis is most likely gender-dependent, of which TG levels in females have shown to be the best predictor for cardiovascular risk in both human and mice.^{29,32}

Fasting TG levels are determined by the amount of TG transported by VLDL particles. VLDL is synthesized by hepatocytes,³³ and serves as the precursor for IDL and LDL. Similar to chylomicrons, VLDL is also a TG-rich lipoprotein. In contrast to chylomicrons that carry exogenous (dietary) lipids, VLDL transports endogenous lipid products to peripheral tissues. High VLDL levels are considered a risk factor for coronary artery disease. Likely the lipid composition of the VLDL particle determines cardiovascular risk. For example, VLDL is a strong predictor for cardiovascular risk in females with a significantly higher VLDL cholesterol/TG ratio than males.³²

IDL, as the remnant of VLDL and precursor of LDL, is also considered a causal factor for the development of atherosclerosis.³⁴ The IDL concentration has been associated with the incidence of coronary artery disease, especially in patients with normal cholesterol levels.³⁵⁻³⁸ However, as an intermediate form between VLDL and LDL, the exact role of IDL in atherosclerosis is less-well defined. The circulating IDL particles are quickly taken up by the liver, or converted to LDL by undergoing triglyceride hydrolysis in peripheral tissues.

In humans, LDL is the primary carrier of cholesterol, accounting for 70-80% of the total cholesterol concentration in the circulation. Importantly, each LDL particle contains a single copy of ApoB100.³⁹ Hence, by analyzing the ApoB100 concentration, LDL particle numbers in the circulation can be calculated. Plasma LDL cholesterol is highly associated with atherosclerosis development. Noteworthy, not only the concentration of LDL cholesterol, but also the

heterogeneity of the LDL particles, plays an important role in atherogenesis. LDL is comprised of multiple subclasses that differ in size and density and each contribute distinctly to the susceptibility for cardiovascular disease.^{40,41} The size and density of LDL varies with its lipid content.⁴² Small and dense LDL particles are more atherogenic, due to their higher penetration capabilities of the endothelial barrier^{43,44} and greater oxidation potential compared to the larger, less dense LDL particles.⁴⁵ Epidemiological studies showed that the small and dense LDL particles are associated with a cluster of cardiovascular disease risk factors, including elevated levels of plasma TG and ApoB, reduced concentrations of HDL cholesterol, and impaired insulin sensitivity.^{46,47} Thus, atherogenesis is not only affected by the amount of cholesterol transported by LDL, but also by the characteristics and heterogeneity of the LDL particles.⁴⁸

1.1.2 High-density lipoproteins and atherosclerosis

In contrast to LDL that promotes atherogenesis, high-density lipoprotein (HDL) is considered to protect against atherosclerosis. HDL removes cholesterol out of lipid-rich tissues and transports it to the liver, a process commonly referred to as reverse cholesterol transport (RCT).^{49,50} RCT is a complicated process involving various steps. First, cholesterol efflux from peripheral cells, including foam cells in the arterial wall, is facilitated by the ATP-binding cassette (ABC) transporters ABCA1 or ABCG1, which mediate the efflux of intracellular lipids to lipid-poor ApoA1 (nascent HDL) and mature HDL in the circulation, respectively.^{51,52} Upon uptake by HDL, the effluxed cholesterol is esterified, via lecithin cholesterol acyltransferase (LCAT), and transferred to the core of HDL, resulting in the remodeling and maturation of the HDL particle. Next, the cholesterol in HDL is transferred to the liver, either via selective uptake of HDL-cholesterol by scavenger receptor BI (SR-BI), holoparticle uptake via the LDLr, or indirectly, via the transfer of cholesterol to other, TG-rich, lipoproteins, through cholesteryl ester transfer protein (CETP) which are subsequently also fluxed back to the liver. Here, the cholesterol taken up by the liver is excreted into bile and feces or used as substrate for *de novo* cholesterol synthesis. Importantly, mice and rats naturally lack CETP activity, and hence CETP-induced cholesterol transfer does not occur in these animals.⁵³

Since HDL can mediate cholesterol efflux from lipid-rich macrophages, HDL has been identified as an important anti-atherogenic particle.⁵⁴ The anti-atherogenic properties of HDL, however, extend beyond the removal of excess lipid from the vascular wall.^{54,55} For example, HDL can also exert antioxidant effects, as under oxidizing conditions, the presence of HDL can significantly decrease lipid peroxide concentrations within the LDL particle.^{56,57} In addition, HDL inhibits monocyte adhesion to the vessel wall, by suppressing the expression of endothelial adhesion and migration molecules.⁵⁸ Finally, HDL also protects against damage inflicted by inflammatory mediators to the endothelium, and prevents thrombosis by upregulating nitric oxide (NO) production in endothelial cells.⁵⁹⁻⁶¹

Considering the wide array of atheroprotective functions of HDL, high HDL concentrations were long thought to be associated with a reduction in coronary artery disease (CAD) risk. Indeed, epidemiological studies have indicated that low HDL cholesterol is associated with an increased risk. However, pharmacological induction of plasma HDL cholesterol levels did not reduce CAD risk.⁶² In line, increased macrophage cholesterol efflux capability of human serum is also independent of the HDL cholesterol level.^{63,64} HDL cholesterol efflux capacity, however, did strongly correlate with the concentration of lipid-poor ApoA1 (pre- β HDL).^{65,66} To explore the development of a novel HDL based-atheroprotective therapy against it is thus important to focus

on modulating nascent HDL (with high cholesterol efflux capacity) rather than HDL cholesterol levels.⁶⁷

1.2 General pathogenesis underlying atherosclerotic lesion development

The development of atherosclerosis is the consequence of a chronic inflammatory reaction of the vascular wall, in response to dyslipidemia and endothelial distress, involving the inflammatory recruitment of leukocytes and the activation of resident vascular cells.⁶⁸ According to the response-to-injury hypothesis, the development of atherosclerosis is initiated by dysfunction of the arterial endothelium.⁶⁹ Cardiovascular risk factors, such as smoking, hypertension, inflammation, age, and lipids (in particular LDL), are known to aggravate endothelium dysfunction and activation. This causes the activated endothelial cells to start expressing surface factors that stimulate the infiltration of monocytes from the blood stream into the intima and subintimal space, where they differentiate into macrophages.⁷⁰ LDL retained in the arterial wall, mostly after extensive oxidative modification⁷¹, is phagocytized by the monocyte-derived macrophages, leading to the formation of lipid-rich foam cells as well the start of a chronic inflammatory process. Monocytes, T cells and mast cells all migrate to the site of action in response to inflammatory signals produced at the site of the early atherosclerotic lesion. These cells in turn will contribute to the immune reaction, creating a progressive inflammatory environment in the developing plaque which further accelerates atherosclerosis development. When the plaque macrophages are unable to sufficiently efflux their excess cholesterol, they become heavily lipid-laden foam cells. Ultimately, these foam cells grow in size and die, thereby releasing all cellular cholesterol into the plaque, causing intraplaque cytotoxicity and further aggravation of the inflammatory response. At this stage, the atherosclerotic plaque consists of a lipid core which contains cholesterol, cellular debris and infiltrated immune cells, covered by a fibrous cap. This collagen-rich fibrous cap overlying the lipid-core of the plaque is the consequence of vascular smooth muscle cells (VSMC) proliferation and provides stability to the plaque. However, thinning of this fibrous cap by mediators secreted by inflammatory cells can ultimately result in plaque rupture.⁷¹ In this advanced stage of lesion development, plaque stability and correlated susceptibility to plaque rupture, is determined by the balance between VSMCs that protect the plaque, and cytotoxic factors released by immune cells/endothelial cells that damage the fibrous cap.⁷² A stable plaque is usually rich in extracellular matrix and smooth muscle cells and in most cases does not cause acute clinical symptoms. Rupture of the atherosclerotic plaque, or erosion of the endothelial layer lead to the formation of a thrombus on top of the atherosclerotic lesion, the culprit for the development of acute cardiovascular events (**Figure 1**).




Figure 1 Progression of atherosclerosis development.

This schematic diagram shows the changes occurring in the arterial wall from a normal healthy artery to an artery with advanced atherosclerosis, and the development of atherothrombosis. a) A healthy artery, composed of three layers, the intima, media and adventitia. The inner layer of the artery, the intima, is formed by a monolayer of endothelial cells. The middle layer of the artery, the media, is principally made up of smooth muscle cells (SMCs). The outer layer, the adventitia is composed of elastic collagen fibers and fibroblasts. b) Upon initiation of atherosclerosis, blood leukocytes, including monocytes are recruited to the activated endothelium. Monocytes, as the largest population during recruitment, differentiate into macrophages, take up oxLDL, and become foam cells. c) During lesion progression, vascular SMCs migrate from the media to the intima and start producing collagen, elastin and proteoglycans that form a fibrous cap overlying the lesion. In advanced lesions, apoptosis of cells in the lipid core contribute to the formation of a necrotic core. d) Once atherosclerotic lesions rupture, a thrombus rapidly develops on top of the lesions. Thrombosis impedes blood flow to a varying extent, but could totally block the blood flow and cause an acute infarction or cerebral stroke. **Painting by Mengyin An, adapted from Libby *et al.***⁷³

2 The role of monocytes and macrophages in the pathology of atherosclerosis

2.1 Monocyte heterogeneity

The bone marrow is a hematopoietic organ responsible for generating new blood cell components, including monocytes. Monocytes are the precursors of macrophages which represent the primary cell type in atherosclerosis.⁷⁴ Monocytes originate from bone marrow myeloid progenitor cells, which are also the progenitor cells of neutrophils.⁷⁵ Notably, circulating monocytes show

morphological heterogeneity. In humans, monocytes are identified by the expression of surface receptors CD14 and CD16. Based on the expression levels of these two markers, monocytes are divided into two main subtypes. The first subtype is a so-called 'classical' monocyte, expressing high levels of CD14 and very low levels of CD16 (CD14^{hi}CD16^{lo}). This classical monocyte population also has a high producing capacity for pro-inflammatory cytokines. The second subtype is the 'non-classical' monocyte, which is characterized by low expression levels of CD14 and high expression levels of CD16 (CD14^{lo}CD16^{hi}). These monocytes are also called patrolling monocytes, as they constantly patrol the vasculature, and are involved in the early responses to pathogens and tissue repair.^{76,77}

In mice, monocytes are classified into two subtypes based on the expression of Ly6C and CCR2, with Ly6C^{hi}CCR2⁺ monocytes representing the equivalent of human CD14^{hi}CD16^{lo} monocytes, and Ly6C^{lo}CCR2⁻ being equivalent to human CD14^{lo}CD16^{hi} monocytes.⁷⁸⁻⁸⁰ After infiltration into the arterial wall, the monocytes are exposed to specific environmental factors, triggering their differentiation into different types of macrophages that differentially contribute to the progression of atherosclerosis.⁸¹

2.2 M1/M2 macrophages and atherosclerosis

Similar as their monocyte precursors, multiple types of macrophages can be distinguished.⁸² Since different macrophage activation forms were proposed by Mackness⁸³ in 1962 and Gordon⁸⁴ in 1992, for M1 and M2 macrophages, respectively, more and more researchers have started investigating the relationship between macrophage phenotypes and atherosclerosis. Both M1 and M2 macrophages have been implicated in atherosclerosis. In vitro, the M1 macrophage phenotype can be induced by incubation with LPS or IFN- γ .⁸⁵ Upon LPS activation, macrophages secrete high levels of pro-inflammatory cytokines, including interleukin 1 β (IL-1 β), IL-6, IL-12, tumor necrosis factor α (TNF- α), but low levels of anti-inflammatory cytokines such as IL-10. As such, M1 macrophages actively contribute to the persistent inflammatory environment in the atherosclerotic plaque, and thereby accelerate atherosclerosis development. In contrast, M2 macrophages are known to protect against atherosclerosis. M2 macrophages can be induced by IL-4, IL-10 and IL-13 and, upon activation, produce high levels of anti-inflammatory cytokines, including IL-10, and low levels of pro-inflammatory cytokines, such as IL-12. Besides producing anti-inflammatory cytokines, M2 macrophages also enhance the production of pro-fibrotic factors, including collagen, and thereby promote tissue repair and remodeling.⁸⁶ Hence, M2 macrophages protect against atherosclerosis not only by decreasing the local inflammatory status of the plaque but also by increasing plaque stability. In mice, M1 macrophages express high levels of inducible nitric oxide synthase (iNOS), which renders iNOS a murine M1 marker gene. In addition to iNOS, high expression levels of the pro-inflammatory cytokines TNF α , IL-1 β , and IL-12 are also considered as M1 markers. In contrast, murine M2 macrophages are known to express high levels of arginase 1 (Arg1). Additionally, YM1 and FIZZ1 and scavenger receptors (CD204,⁸⁷ CD163⁸⁸) are also considered M2 macrophage markers.⁸¹

2.3 Macrophage phenotype and foam cell susceptibility

Besides playing a role in the immune response, another important function of macrophages is to engulf foreign agents, including oxidized LDL (oxLDL).⁸⁹ Macrophages take up oxLDL and become foam cells, a process which is considered to be one of the hallmarks of atherosclerosis. SR-A⁹⁰ and CD36 (scavenger receptor class B member 3) are the main receptors involved in foam cell formation, being responsible for up to 90% of the oxLDL uptake by macrophages *in vitro*.⁹¹ Interestingly, both SR-A⁹² and CD36⁹³ are upregulated during M2 macrophage differentiation, suggesting an increased susceptibility of M2 macrophages to become foam cells. Indeed, van Tits *et al.* found that compared to M1 macrophages, M2 macrophages are more prone to take up oxLDL and become foam cells.⁹⁴ This suggests that, in contrast to their atheroprotective anti-inflammatory role, M2 macrophages are also likely to play a pro-atherogenic role by promoting macrophage foam cell formation.

2.4 Macrophage phenotype switch

The process of macrophage polarization is dynamic, as macrophages can rapidly switch from one phenotype to another in response to a changing microenvironment.⁹⁵⁻⁹⁷ In atherosclerosis, numerous factors affect the lesional microenvironment, including cholesterol oxidation, inflammation mediators, infiltrated immune cells, growth factors, dead cells and other substances.⁹⁸ Therefore, the lesional microenvironment changes with the different stages of atherosclerosis development, thereby further influencing macrophage polarization.⁹⁹ Da Silva and colleagues found that cholesterol loading of human macrophages limited their capability to be primed to M1 macrophages, but not to M2 macrophages, suggesting an anti-inflammatory property of foam cells.¹⁰⁰ Furthermore, in response to oxidized phospholipids, a product of lipid oxidation, macrophages are primed to a so-called Mox phenotype.¹⁰¹ Finally, in response to haem and haemoglobin exposure after intraplaque hemorrhage, macrophages can be polarized towards an Mhem phenotype.¹⁰² Noteworthy, both Mox and Mhem macrophages display a reduced capacity to engulf oxLDL, and are thus considered less prone to foam cell formation.^{101,103} All M1, M2, Mox, and Mhem macrophage phenotypes have been demonstrated in atherosclerotic lesions. However, M1 and M2 macrophages are suggested to act as the main precursors for macrophage foam cells.¹⁰⁴

3 Experimental mouse models and strategies for studying atherosclerosis

3.1 Mice and atherosclerosis susceptibility

Multiple animal species have been used as experimental models to study atherosclerotic lesion development, including pigs, rabbits, monkeys, non-human primates and mice.¹⁰⁵⁻¹¹⁰ Among these non-human models, mice now are considered the best choice for studying atherogenesis, due to their low cost, high reproduction rate and short time frame for disease development.¹⁰⁶ Although C57BL/6 (hereafter referred to as WT mice) is the mouse strain most sensitive to the development of atherosclerosis, as compared to other murine strains, C57BL/6 mice are still relatively resistant to diet-induced atherosclerosis.¹⁰⁶ Persistent hypercholesterolemia, reflecting plasma cholesterol levels exceeding 300 mg/dL, is needed to induce atherosclerosis development in mice.¹¹¹ The main reason that mice in general are resistant to atherosclerosis is their distinct plasma lipoprotein profile, as compared to humans.¹¹² The fact that mice lack the cholesterol ester transfer protein (CETP) and exert a low ability to absorb dietary cholesterol causes a cardiometabolic lipid profile,

reflected by consistently high plasma levels of HDL cholesterol and low plasma levels of LDL cholesterol.¹¹² Therefore, exposing mice to high concentrations of dietary cholesterol alone is not sufficient to induce atherosclerosis development. To enhance the atherosclerosis susceptibility of mice, genetic modification is required for induction of a sufficiently high pro-atherogenic lipoprotein profile. Since the 1990s, the technique of homologous recombination in embryonic stem cells made it possible to selectively knock out genes involved in the metabolism of pro-atherogenic lipoproteins.¹¹³ Currently, the most frequently used mouse models to study atherosclerosis are LDL receptor (LDLr) knockout (KO) and ApoE KO mice.¹¹²

3.2 ApoE KO mice and LDLr KO mice

ApoE is a constituent of non-LDL lipoproteins and serves as an essential ligand for the uptake of these lipoproteins by the liver.¹¹⁴ Therefore, mice lacking ApoE show impaired clearance of plasma cholesterol, resulting in severe hypercholesterolemia. ApoE KO mice fed a regular chow diet display plasma cholesterol levels of > 500 mg/dL, which can mainly be attributed to increased levels of chylomicrons and VLDL, whereas plasma HDL-cholesterol is decreased.¹¹⁵ ApoE KO mice develop extensive atherosclerotic lesions.^{116,117} Under normal chow conditions, spontaneous atherosclerotic lesion development is observed in the aortic sinus within 3-4 months of age. In the older mice, atherosclerotic lesions are visible throughout the aorta at locations of principal branches.^{118,119} Importantly, this process can be accelerated by feeding ApoE KO mice a high fat/high cholesterol diet.¹²⁰

The LDL-receptor regulates plasma cholesterol levels by removing IDL and LDL from plasma. Mice lacking the LDLr, as compared to WT mice, display a 2-fold higher plasma cholesterol level (~230 mg/dL) when fed a regular chow diet, as compared to WT mice. This increase can mainly be attributed to an increase in cholesterol within the IDL/LDL fraction.^{121,122} Furthermore, LDLr deficiency also leads to a small increase in VLDL-cholesterol levels. Moreover, HDL-cholesterol levels are increased.¹²² The mild hypercholesterolemia induced by LDLr deficiency however, is not sufficient to effectively induce atherosclerosis development in mice on chow. Interestingly, plasma cholesterol levels of LDLr KO mice are highly responsive to dietary interventions,^{123,124} and a high-fat/high-cholesterol (Western-type) diet is known to induce severe hypercholesterolemia and rapid atherosclerosis development.^{122,125,126} Similar to ApoE KO mice, atherosclerotic lesion development in LDLr KO mice on Western-type diet is initiated in the aortic root.¹²⁷ Intermediate aortic lesion development occurs within 3 months of Western-type diet feeding, and advanced lesions are present in the aorta after 5 months of dietary challenge.^{123,124,126} A commonly used approach to investigate the function of a specific gene in atherosclerosis, is crossbreeding of mice deficient for the gene of interest with the hypercholesterolemic LDLr KO or ApoE KO mice. Since generating double knockout mice is a time consuming and costly approach, there is an ongoing search for alternative methods to induce hypercholesterolemia in mice.

3.3 Bone marrow transplantation

Hematopoietic stem cell transplantation to an established mouse model of atherosclerosis, such as the apoE or LDLr KO mice, is an effective strategy to generate chimeric mice with target gene alterations in bone marrow-derived cells of an atherosclerosis-prone background. One of the strengths of this model is that in the recipients specifically the genotype of the bone marrow-derived cells, which represent the major players in atherosclerosis, is altered and thus allows analysis of the specific contribution of a gene of interest in blood cells. Hematopoietic cell

transplantation not only helps to ease the time and money needed for the generation of sophisticated cell type-specific knockout mouse models, but also allows a closer mechanistic insight into the cellular biology underlying atherosclerosis development.¹²⁸

Before birth, blood cells are derived from the fetal liver and spleen, however, after birth the bone marrow becomes the primary origin for the generation of blood cells. Therefore, bone marrow is normally the source of hematopoietic cells for transplantation, especially in murine models. In bone marrow transplantation studies (**Figure 2**), LDLr KO mice are normally chosen over ApoE KO mice as the atherosclerosis-prone recipients. This is because 1) the lipid profile of LDLr KO mice, characterized by a high IDL/LDL cholesterol fraction, resembles strongly the plasma lipid profiles of human dyslipidemic patients,¹²⁹ 2) the morphology of the atherosclerotic lesions in LDLr KO mice resemble human atherosclerotic plaques,^{130,131} and, most importantly, 3) the presence of the LDLr in the donor bone marrow does not affect the pro-atherogenic lipoprotein profile and atherosclerosis susceptibility of the LDLr KO recipients.¹³²⁻¹³⁴ In contrast, several studies have demonstrated that restoration of ApoE in bone marrow-derived cells normalizes serum cholesterol to WT levels and reduces atherosclerosis development in ApoE KO recipients.¹³⁵⁻¹³⁸




Figure 2 Schematic diagram of a bone marrow transplantation (BMT) procedure in mice for studying atherosclerosis.

1) Original bone marrow of atherosclerosis-prone recipient mice (often LDLr KO mice) is destroyed by a lethal dose of irradiation (9 Gy). 2) The recipients receive five-million donor bone marrow cells, lacking or overexpressing the gene of interest. 3) Recipients are allowed to recover for 8 weeks on a chow diet. One week before the irradiation, and throughout the complete recovery period, recipient mice receive antibiotics via drinking water. 4) After recovery, the recipients are challenged with an atherogenic Western-type diet to induce atherosclerosis development.

4 Signaling pathways and atherosclerosis

As mentioned before, macrophages are the main cell type in atherosclerotic lesions and their phenotype and activation status influence their exact role in the pathogenesis of the disease. Importantly, macrophage phenotype and activation are highly dependent on the lesional microenvironment and the intracellular signaling pathways that are activated within the macrophages.

Macrophage activation is a very complicated process. From receiving an initial stimulus to the point that eventually the macrophage's phenotypic functional protein production is altered, this process entails activation of a complex set of signaling pathways, and transcriptional and post-transcriptional regulatory networks.^{96,139} In a simplified summary, as shown in **figure 3** the following key steps can be distinguished: 1) macrophage surface receptors recognize environmental stimuli; 2) the signals are amplified and transmitted to the nucleus by protein

kinase transducers; and 3) the nuclear transcriptional/post-transcriptional factors regulate macrophage-specific gene expression and thus dictate macrophage polarization and functions.¹³⁹ This complicated activation process provides multiple possibilities to design novel strategies, addressing the different activation steps, to reprogram specific macrophage phenotypes for therapeutic benefit. In this thesis, four genes involved in macrophage activation at different levels, were investigated. Macrophage polarization to the M1 and M2 phenotype has long served as a paradigm for studying atherosclerosis. Inducing new effector activities by activated macrophages is considered as an attractive therapeutic approach for atherosclerosis treatment.¹³⁹⁻¹⁴³ To explore potential novel therapeutic targets, we first evaluated and discussed the role of the M2 macrophage signature gene Arg1, a key player in the nitric oxide/L-arginine pathway, in atherosclerosis development. Protein kinases play essential role in the transcriptional and epigenetic regulation of macrophage polarization,^{139, 144,145} and are the most intensively studied protein targets in pharmacology research.¹⁴⁶⁻¹⁵⁰ In this thesis the atherosclerotic role of Akt2 and MKP2, key members of the protein kinase B and mitogen-activated protein kinases family respectively, are addressed. Furthermore, transcription factors, critical regulators of gene expression, have long been proposed to execute essential regulatory functions in the pathogenesis of atherosclerosis.¹⁵¹ In addition, in this thesis, we focus on the upstream stimulatory factors (Usfs), recently identified lipid-related transcription factors,¹⁵² that are regulators of several important cellular processes¹⁵³ and hence are expected to influence atherosclerosis development. Below the background of 1) the Nitric oxide/L-arginine and Arg1, 2) protein kinases and their inhibitors Akt2 and MKP2, and 3) Usfs in atherosclerosis is described in more detail below.

4.1 Nitric oxide/L-arginine pathway in atherosclerosis

Nitric oxide (NO) is an important signaling molecule that influences many cellular processes.¹⁵⁴ The cardiometabolic related functions of this molecule include: 1) prevention of endothelial cell apoptosis,^{155,156} 2) reduction of oxidative stress, induced by reactive oxygen species (ROS),¹⁵⁷ 3) inhibition of smooth muscle cell proliferation, and 4) inhibition of vascular cell adhesion molecule-1 (VCAM-1) expression and, hence, inhibition of monocyte recruitment.¹⁵⁸ In line with these athero-protective functions of NO, several animal studies have confirmed that decreasing NO production induces atherosclerosis,¹⁵⁹ while increasing NO production attenuates atherosclerosis.^{160,161}

Important to note is that there are indications that the protective role of NO is both tissue-specific^{159,160,162-164} and dose-dependent.^{160,162,165} NO is a product of nitric oxide synthases (NOSs). The NOS family has three members, endothelial NOS (eNOS), neuronal NOS (nNOS), and inducible NOS (iNOS).¹⁵⁴ eNOS and nNOS are constitutively expressed.¹⁶⁶ As indicated by the name, eNOS is primarily produced by endothelial cells and, nNOS by neurons. The expression of iNOS can be induced by stimulators, especially inflammatory cytokines.¹⁶⁷ More importantly, iNOS is expressed by macrophages, and highly upregulated in response to lipopolysaccharide and inflammatory cytokines,¹⁶⁸ leading to an enhanced production of NO.




Figure 3 Regulation in macrophage activation and target factors Arg1, Akt2, MKP2 and Usf1 studied in more detail in this thesis.

1) Environmental stimuli activate cellular effector receptors. 2) The signals are amplified and transmitted to the nucleus by protein kinase transducers, including Akt2 or MKP2. 3) Transcription factors, such as Usf1, regulate the stimulus-specific gene expression, eventually leading to phenotypic functional protein production, such as Arg1.

Increasing evidence indicates that NO produced by nNOS and eNOS is atheroprotective, while NO produced by iNOS is more likely pro-atherogenic.¹⁶⁹ Diminished levels of bioavailable NO are associated with endothelial dysfunction,¹⁷⁰ the indicator of early atherosclerosis. Deletion of eNOS in ApoE deficient mice leads to increased atherosclerosis development,¹⁵⁹ and eNOS gene transfer improves atherosclerosis,^{160,161} further evidencing the athero-protective role of NO derived from eNOS. The beneficial effects of nNOS in atherosclerosis described by Kuhlencordt *et al.*,¹⁷¹ were surprising at first, since nNOS is primarily detected outside of cardiovascular system.¹⁷²⁻¹⁷⁴ However, in their study, nNOS expression was demonstrated in the atherosclerotic lesion, predominantly in smooth muscle cells, and to a lesser extent in macrophages, but not in endothelial cells.¹⁷⁵ nNOS-derived hydrogen peroxide induces endothelial dysfunction, but the mechanism underlying the protective effects of nNOS in atherosclerosis is not clear yet.^{169,176} NO produced by iNOS of lesional macrophages, is pro-atherogenic in humans.¹⁷⁷ In support, deletion of iNOS in mice reduces their susceptibility to the development of atherosclerosis.^{163,164,178} In addition to its cell type-specific effects on atherosclerosis, NO also influences atherogenesis in a dose-dependent manner: low concentrations of NO improves atherosclerosis,^{160,161} but massive NO production worsens atherosclerosis.¹⁶²

iNOS, a signature marker of M1 macrophages, uses the substrate L-arginine, to produce NO. Interestingly, this same substrate is also used by Arginase-1 (Arg1), an important M2 macrophage marker, which catalyzes the conversion of L-arginine to urea and ornithine, the latter being a precursor for collagen production. Deletion of iNOS is known to cause a decrease in NO production, resulting in an increased availability of its substrate L-arginine for Arg1 mediated-conversion into ornithine. Hence, iNOS deletion can indirectly lead to an enhanced production of collagen,¹⁷⁹ thereby improving atherosclerotic plaque stability. Conversely, deletion of Arg1 might lead to increased substrate availability for the production of NO. As mentioned above, NO in atherosclerosis is well-studied, however, the role of Arg1 in atherosclerotic plaque development is currently still unknown.

4.2 Protein kinase B (Akt) in atherosclerosis

Protein kinase B is a serine/threonine-specific protein kinase, of which three isoforms can be distinguished: Akt1, Akt2, and Akt3.¹⁸⁰ Akt plays an important role in many cellular processes, such as apoptosis,¹⁸¹ proliferation,¹⁸² migration,¹⁸³ transcription,¹⁸⁴ and insulin responsiveness.¹⁸⁵ Importantly, Akt signaling is also known to influence atherosclerotic lesion development.

Akt1 is widely expressed in all tissues, whereas Akt2 expression is limited to metabolic tissues, such as adipose tissue, liver, and skeletal muscle, and Akt3 is preferentially expressed in brain.^{180,186,187} Interestingly, macrophages express all three Akt isoforms.¹⁸⁸ Recent studies showed that Akt isoforms differentially contribute to macrophage polarization.^{189,190} For example, upon *Staphylococcus aureus* infection, Akt1 deficient macrophages showed upregulated expression of the M1 signature gene iNOS.¹⁸⁹ Conversely, upon LPS stimulation, Akt2 knockout macrophages display an M2-like phenotype, as evidenced by augmented expression of the M2 macrophage markers Arg1, YM-1, and FIZZ-1.^{190,191} So far, the role of Akt3 in macrophage polarization remains unknown. In addition to their roles in macrophage polarization, Akts also differentially influence macrophage foam cell formation. Previous studies have shown that Akt1 does not affect oxLDL-induced cholesterol accumulation in macrophages,¹⁹² whereas Akt2 promotes acLDL-induced foam cell formation,¹⁹¹ and Akt3 protects macrophages against acLDL-induced foam cell formation.¹⁹³

In line with the different roles of the Akt isoforms in macrophage polarization and foam cell formation, Akt isoforms also distinctly contribute to atherosclerosis development. Akt1 has been reported to have an atheroprotective role.¹⁹² However, this effect is likely due to Akt1 of vascular origin and not macrophage Akt1. Indeed, bone marrow-specific deletion of Akt1 did not influence atherosclerosis susceptibility.¹⁹² Moreover, Fernandez-Hernandez and colleagues showed that whole body Akt1 deletion inhibits the proliferation and migration of vascular smooth muscle cells (VSMCs), leading to the development of vulnerable atherosclerotic plaques with increased necrosis and a smaller collagen-rich fibrous cap.¹⁹⁴ Akt3 is barely detectable in the healthy vasculature.¹⁹⁵ However, in line with the importance of Akt3 to limit macrophage foam cell formation, an increased susceptibility to atherosclerosis development was observed in Akt3 total body and bone marrow-specific knockout mice.¹⁹³

The role of Akt2 in atherosclerosis is more complex. Rensing and colleagues found that total body deletion of Akt2 induces smaller but unstable atherosclerotic lesions, with a major causative role for VSMC derived Akt2 in the decreased lesional collagen content and increased necrotic core formation.¹⁹⁶ The unstable phenotype induced by Akt2 loss is likely due to disturbances in VSMCs

migration, proliferation, and metalloproteinase production.¹⁹⁶ However, a study from another group found that total body Akt2 deletion does not affect atherosclerosis development.¹⁹¹ To exclude the contribution of smooth muscle cell Akt2, Babaev *et al.* used fetal liver cell transplantation and Rotllan *et al.* used bone marrow transplantation to specifically delete Akt2 in hematopoietic cells of LDLr KO mice.^{188,191} Both studies indicated that hematopoietic Akt2 deficiency protects LDLr KO mice against diet-induced atherosclerosis. Akt2 deletion in macrophages induced M2 macrophage polarization, decreased macrophage migration and inhibited macrophage foam cell formation; processes that are likely responsible for the observed decrease in atherosclerosis susceptibility of the LDLr KO recipients.^{191,197} These findings indicate that both VSMC Akt2 and hematopoietic Akt2 play a role in atherogenesis. Importantly, the protective effect of hematopoietic Akt2 deficiency was independent of Akt1 and Akt3, because bone marrow-specific deletion of Akt1 and Akt3 led to unchanged or increased atherosclerosis development, while the phenotype of the reduced susceptibility to atherosclerosis persisted upon combined Akt2/Akt1 or Akt2/Akt3 deletion in bone marrow.^{192,193,198-200}

4.3 MAPKs/MKPs pathway in atherosclerosis

4.3.1 MAPKs and atherosclerosis

MAPKs are a family of protein kinases that specifically phosphorylate serine/threonine residues. Three major subfamilies can be distinguished: extra cellular regulated protein kinase (ERK), c-Jun N-terminal kinase (JNK) and p38. All these subfamilies have been reported to participate in atherosclerosis development.

4.3.1.1 ERK and atherosclerosis

In vivo, ERK expression is increased in atherosclerotic lesions of cholesterol-fed rabbits.²⁰¹ *In vitro*, ERK is rapidly activated upon oxLDL stimulation in macrophages.²⁰² These findings indicate a role for ERK in foam cell formation. Indeed, inhibition of ERK1/2, by their upstream MEK1/2 inhibitor U0126, significantly decreased foam cell formation both *in vivo* and *in vitro*, which is most likely the consequence of upregulated cholesterol efflux transporters ABCA1 and ABCG1.^{203,204} In agreement with the decreased susceptibility to foam cell formation, atherosclerotic lesion development was reduced in ApoE knockout mice treated with the inhibitor U0126. These findings suggested an anti-atherosclerotic role for ERK1/2 inhibition.

ERK also influences macrophage polarization. Inhibition of ERK by the inhibitors U0126, or PD0325901 led to an M2-like macrophage phenotype, reflected by increased M2 marker gene expression.^{205,206} Furthermore, the ERK inhibitor-dependent increase of M2 macrophage marker gene expression is likely independent of the pre-existing polarization state of the macrophage,²⁰⁶ as re-priming of LPS-polarized M1 macrophages by IL-4/IL-13 still induced a shift towards the M2 phenotype. These findings suggest that skewing of macrophages towards an M2 phenotype might also contribute to the observed athero-protective effect of ERK inhibition.

4.3.1.2 P38 and atherosclerosis

P38, also called mitogen-activated protein kinase 11 (MAPK11), has 4 isoforms: p38 α , p38 β , p38 γ , and p38 δ .²⁰⁷ p38 α , the most well-studied isoform of p38, is rapidly phosphorylated in macrophages in response to LPS and is responsible for the subsequent induction in the production of pro-inflammatory cytokines.²⁰⁸⁻²¹¹ Genetic deletion of p38 α in macrophages results in an

impaired TLR4-mediated LPS-induced innate immune response, reflected by a decreased production of the pro-inflammatory cytokines TNF- α and IL-12.²¹² Furthermore, p38 activation is enhanced in IL-4-induced alternatively activated macrophages, and inactivation of p38 led to decreased IL-4-induced M2 marker expression.²¹³ Collectively, these findings suggest that p38 activation is likely needed for both M1 and M2 macrophage polarization and function.

The role of p38 in foam cell formation is not clear yet. Inactivation of pan p38 by a pharmaceutical inhibitor prevents foam cell formation *in vitro*.^{214,215} However, genetic deficiency of p38 α in macrophages does not affect foam cell susceptibility,²¹⁶ albeit it does enhance macrophage apoptosis.²¹⁷ Several studies have been performed to address the role of p38 in atherosclerosis. Systemic p38 inhibition, either via pharmaceutical inhibition or genetic deletion of the p38 substrate MK2, protects ApoE KO mice against atherosclerosis,^{218,219} indicating a pro-atherogenic role of p38. Interestingly, Seimon *et al.* showed that macrophage-specific deletion of p38 α promotes atherosclerosis development.²¹⁷ In contrast, Kardakis *et al.*, using the same experimental set up, found that p38 α deficiency had no effect on atherosclerosis development.²¹⁶ Hence, the role of macrophage p38 in atherosclerosis remains controversial.

4.3.1.3 JNK and atherosclerosis

c-Jun N-terminal kinase (JNK), also named mitogen-activated protein kinase 8 (MAPK8), is present in three isoforms: JNK1, JNK2, and JNK3.²²⁰ Of these three isoforms, JNK1 and JNK2 are ubiquitously expressed, whereas JNK3 is mainly expressed in brain, and to a lesser extent in the heart and testes.²²⁰ JNK1 and JNK2 are both expressed in macrophages,²²¹ and their activation is known to regulate various macrophage functions, including polarization, foam cell formation, and programmed cell death.

Inhibition of JNK activation leads to impaired macrophage development, proliferation, and survival,²²² suggesting a broad function of JNK in macrophages. *In vitro*, JNK1 was shown to be responsible for cytokine and NO production by LPS-stimulated M1 macrophages.^{223,224} Furthermore, *in vivo*, deletion of JNK1 reduced macrophage migration and infiltration in a murine arthritis model.²²⁵ On the other hand, JNK2 was shown to stimulate oxLDL-induced foam cell formation.²⁰² Conversely, oxLDL triggers JNK2 activation in macrophages and facilitates scavenger-mediated foam cell formation in a CD36-JNK-SR-A loop manner.²²⁶ In accordance, enhanced activation of JNK2 is observed in macrophage-rich atherosclerotic lesions.^{227,228}

In line with the distinct effects of JNK isoforms on macrophage function, JNK isoforms also contribute to atherosclerosis differently. Genetic deletion of JNK2 protects ApoE KO mice against high-fat diet induced atherosclerosis, and this is likely attributed to the lack of JNK2-induced foam cell formation.²²⁷ Interestingly, hematopoietic JNK1 deficiency promotes atherosclerosis development in LDLr KO mice, likely caused by the lack of JNK1-mediated regulation of macrophage survival.²²¹

4.3.1.4 Interactions between MAPK members

MAPK cascades can be activated by either intra- or extracellular stimulators and signaling molecules. Depending on the cell type, stimulus signal strength and dynamics, MAPK can be activated differently and serve distinct functions. Importantly, each member of MAPK family closely interacts with the other members, and in most cases is subject to negative feedback regulation.²²⁹

4.3.2 MAP kinase phosphatases and atherosclerosis

The activity of MAPKs is tightly regulated by MAP kinase phosphatases (MKPs) that belong to the group of dual-specificity phosphatases (DUSPs), which includes at least 10 members.²³⁰ MKPs are normally referred to as the typical DUSPs.²³¹ Atypical DUSPs lack the MAPK-binding motif or kinase-interacting motif (MKB/KIM), which determines the dephosphorylation activity of DUSPs.^{232,233} Based on their subcellular localization, MKPs are divided into three groups. The first group consists of MKP1 (DUSP1), PAC1 (DUSP2), MKP2 (DUSP4), and DUSP5, which are all inducible MKPs that are located in the nucleus. The second group is represented by MKP3 (DUSP6), MKP-X (DUSP7) and MKP4 (DUSP9), which are all located in the cytoplasm. Finally, the third group is comprised of DUSP8, MKP5 (DUSP10) and MKP-7 (DUSP16), which are located in both the nucleus and the cytoplasm of a cell.^{230,234,235} MKPs inactivate MAPKs by dephosphorylating their phosphoserine/threonine and phosphotyrosine residues.^{236,237} Noteworthy, the expression of group 1 members, being inducible MKPs,^{230,234,235} is induced in response to stimulation with LPS,²³⁸ a potent activator of MAPKs. Therefore, this group of MKPs is likely to act as MAPKs feedback loop regulator.^{230,239}

All three members of the MAPK family are involved in macrophage polarization and foam cell formation, indicating that MKPs are most likely also involved in the regulation of these processes and thereby influence atherosclerosis development. Indeed, several studies have demonstrated that various macrophage functions are affected by depletion of MKPs, especially the inducible MKPs. For instance, deletion of MKP1 led to the upregulation of several M1 signature genes, including TNF- α , IL-6, IL-1 β , and CCL2.^{240,241} In addition, PAC1 deficient macrophages displayed an increased production of pro-inflammatory cytokines in response to LPS stimulation.²⁴² These data suggest an important role for MKP1 and PAC1 in regulating LPS-induced M1 polarization. Unlike the clear roles of MKP1 and PAC1 in the regulation of M1 polarization, the current findings on the role of MKP2 in macrophage polarization remain contradictory. Al-Mutairi *et al.* found that upon LPS stimulation, genetic loss of MKP2 in bone marrow-derived macrophages (BMDM) led to enhanced production of pro-inflammatory cytokines, while the production of anti-inflammatory cytokines was decreased.²⁴³ In contrast, Cornell and colleagues found that upon LPS stimulation, MKP2 deletion in BMDMs resulted in decreased TNF- α and IL-10 production.²⁴⁴ DUSP5 deficient macrophages showed no altered cytokine or chemokine production in response to LPS stimulation.²⁴⁵ However, despite some indications for a role of MKPs in macrophage polarization and function, so far only limited research has been done addressing the role of MKPs in atherosclerosis.²⁴⁶⁻²⁴⁸

4.4 Usf transcription factors and atherosclerosis

Upstream stimulatory factors (Usfs) are DNA-binding proteins, featured as a helix-loop-helix motif and leucine repeat, that serve as transcription factors.²⁴⁹ By binding to target DNA as Usf homo- and heterodimers, Usfs regulate target gene expression.^{139,250} Accumulating evidence showed that disturbed Usf signaling normally leads to metabolic disorders, especially in the case of Usf1.²⁵¹⁻²⁵⁴ Usf1 is ubiquitously expressed, and has a broad range of target genes, of which the genes involved in lipid and glucose metabolism are most widely studied.²⁵⁵ Usf1 is also known to regulate gene expression in response to stressors, such as ultraviolet (UV) irradiation,²⁵⁶ insulin,²⁵⁷ and growth factor.^{258,259} Moreover, in response to PI3 kinase/Akt signaling, Usf1 regulates the transcription of genes implicated in cellular apoptosis and cell cycle arrest.²⁶⁰

Usf1 is a direct target of p38 and AMPK, which both have been shown to be involved in macrophage polarization.^{254,256,261-263} Furthermore, Usf1 is phosphorylated by protein kinase CK2.²⁶⁴ Protein kinases are important for regulation of various cellular functions. However, the exact role of Usf1 in cell function regulation is still unclear. Importantly, mutations in Usf1 are strongly associated with familial combined hyperlipidemia, a common hereditary dyslipidemia with a prevalence of 20% in CVD patients.²⁶⁵ In support of this association, a strong pro-atherogenic role for Usf1 is found in LDLr KO mice lacking Usf-1, which display a remarkable decreased in their susceptibility to atherosclerotic lesion development.¹⁵² The mechanism behind this effect, however, is poorly defined.

5 Outline of this Thesis

Currently, therapeutic strategies to prevent atherosclerosis are primarily based on the use of cholesterol lowering drugs, e.g. statins. However, the morbidity and mortality of cardiovascular disease caused by atherosclerosis remains high. Therefore, identification of novel pharmaceutical targets suitable for the development of novel drugs is strongly needed. The aim of this thesis is to unravel the role of various candidate genes in macrophage activation and their subsequent role in atherosclerosis.

In **Chapter 2**, we determined the role of hematopoietic Arg1, a classic marker of M2 macrophage activation, in atherosclerosis. We show that although Arg1 deficiency promotes macrophage foam cell formation, it does not impact on atherosclerosis development.

Akt2 is a key player in the PI3K/Akt transduction pathway that regulates M2 macrophage polarization.¹⁹⁰ In **Chapter 3**, we addressed the effects of bone marrow Akt2-reconstitution, in Akt2/LDLr dKO mice. We show that hematopoietic Akt2 promotes foam cell formation, but does not alter atherosclerosis development in Akt2/LDLr dKO mice.

Next, we determined the effects of the upstream regulators MKP2, of the MAPK transduction pathway, in atherosclerosis. MKP2 was found to play an important role in regulating macrophage function, and in **Chapter 4** we show that MKP2 deficiency skews macrophages to an M2 phenotype associated with an enhanced susceptibility to foam cell formation. In line, deletion of MKP2 in bone marrow-derived cells leads to increased atherosclerosis development.

Additionally, we focused on the bone marrow-specific effects of Usf1, an upstream transcription stress sensor in atherosclerosis. In **Chapter 5**, we show that Usf1 in bone marrow-derived cells protects against atherosclerosis.

Finally, the overall conclusions and future perspectives of this thesis are discussed in **Chapter 6**.

6 References

1. Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. 2002;105:1135-1143
2. Davies MJ, Woolf N. Atherosclerosis: What is it and why does it occur? *Br Heart J*. 1993;69:S3-11
3. Ross R. The pathogenesis of atherosclerosis: A perspective for the 1990s. *Nature*. 1993;362:801-809
4. Townsend N, Nichols M, Scarborough P, Rayner M. Cardiovascular disease in europe--epidemiological update 2015. *Eur Heart J*. 2015;36:2696-2705

5. Thompson RC, Allam AH, Lombardi GP, Wann LS, Sutherland ML, Sutherland JD, Soliman MA, Frohlich B, Mininberg DT, Monge JM, Vallodolid CM, Cox SL, Abd el-Maksoud G, Badr I, Miyamoto MI, el-Halim Nur el-Din A, Narula J, Finch CE, Thomas GS. Atherosclerosis across 4000 years of human history: The horus study of four ancient populations. *Lancet*. 2013;381:1211-1222
6. Coiter V. De avium sceletis et praecipius musculis. 1575
7. Khan IA, Mehta NJ. Initial historical descriptions of the angina pectoris. *J Emerg Med*. 2002;22:295-298
8. Parry CH. *An inquiry into the symptoms and causes of the syncope anginosa commonly called angina pectoris; illustrated by dissections: By caleb hillier parry, md member of the college of physicians of london, and of the royal medical society of edinburgh; and one of the physicians of the bath general hospital*. R. Cruttwell; and sold by Cadell and Davies, Strand, London; 1799.
9. LaRosa JC, Pedersen TR. Foreword. *European Heart Journal Supplements*. 2004;6:C1-C4
10. Slijkhuis W, Mali W, Appelman Y. A historical perspective towards a non-invasive treatment for patients with atherosclerosis. *Neth Heart J*. 2009;17:140-144
11. Ascher E, Hollier LH, Strandness DE, Towne JB, Haimovici H, Calligaro K, Kent KC, Moneta GL, Pearce WH, Ricotta JJ. *Haimovici's vascular surgery*. John Wiley & Sons; 2008.
12. Tedgui A, Mallat Z. Cytokines in atherosclerosis: Pathogenic and regulatory pathways. *Physiol Rev*. 2006;86:515-581
13. Mehta NJ, Khan IA. Cardiology's 10 greatest discoveries of the 20th century. *Tex Heart Inst J*. 2002;29:164-171
14. Gotto AM, Jr. Evolving concepts of dyslipidemia, atherosclerosis, and cardiovascular disease: The louis f. Bishop lecture. *J Am Coll Cardiol*. 2005;46:1219-1224
15. Mayerl C, Lukasser M, Sedivy R, Niederegger H, Seiler R, Wick G. Atherosclerosis research from past to present--on the track of two pathologists with opposing views, carl von rokitansky and rudolf virchow. *Virchows Arch*. 2006;449:96-103
16. Ross R. Atherosclerosis--an inflammatory disease. *N Engl J Med*. 1999;340:115-126
17. Ross R, Harker L. Hyperlipidemia and atherosclerosis. *Science*. 1976;193:1094-1100
18. Gofman JW, Lindgren FT, Elliott H. Ultracentrifugal studies of lipoproteins of human serum. *J Biol Chem*. 1949;179:973-979
19. Keys A. Coronary heart disease in seven countries. *Circulation*. 1970;41:186-195
20. Prassl R, Laggner P. *Lipoprotein structure and dynamics: Low density lipoprotein viewed as a highly dynamic and flexible nanoparticle*. INTECH Open Access Publisher; 2012.
21. Nordestgaard BG, Tybjaerg-Hansen A. Idl, vldl, chylomicrons and atherosclerosis. *Eur J Epidemiol*. 1992;8 Suppl 1:92-98
22. Nikkila E. Familial lipoprotein lipase deficiency and related disorders of chylomicron metabolism. *Metabolic basis of inherited disease/[edited by] John B. Stanbury...[et al.]*. 1983
23. Nordestgaard BG, Zilversmit DB. Large lipoproteins are excluded from the arterial wall in diabetic cholesterol-fed rabbits. *J Lipid Res*. 1988;29:1491-1500
24. Proctor SD, Mamo JC. Intimal retention of cholesterol derived from apolipoprotein b100--and apolipoprotein b48--containing lipoproteins in carotid arteries of watanabe heritable hyperlipidemic rabbits. *Arteriosclerosis, thrombosis, and vascular biology*. 2003;23:1595-1600
25. Pal S, Semorine K, Watts GF, Mamo J. Identification of lipoproteins of intestinal origin in human atherosclerotic plaque. *Clin Chem Lab Med*. 2003;41:792-795
26. Gianturco SH, Ramprasad MP, Lin AH, Song R, Bradley WA. Cellular binding site and membrane binding proteins for triglyceride-rich lipoproteins in human monocyte-macrophages and thp-1 monocytic cells. *J Lipid Res*. 1994;35:1674-1687
27. Gianturco SH, Brown SA, Via DP, Bradley WA. The beta-vldl receptor pathway of murine p388d1 macrophages. *J Lipid Res*. 1986;27:412-420
28. Tomkin GH, Owens D. The chylomicron: Relationship to atherosclerosis. *Int J Vasc Med*. 2012;2012:784536
29. Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM. Fasting compared with nonfasting triglycerides and risk of cardiovascular events in women. *JAMA*. 2007;298:309-316

30. Al-Aubaidy HA, Jelinek HF. Oxidative stress and triglycerides as predictors of subclinical atherosclerosis in prediabetes. *Redox Report*. 2014;19:87-91
31. Ridker PM. Fasting versus nonfasting triglycerides and the prediction of cardiovascular risk: Do we need to revisit the oral triglyceride tolerance test? *Clinical chemistry*. 2008;54:11-13
32. VanderLaan PA, Reardon CA, Thisted RA, Getz GS. VLDL best predicts aortic root atherosclerosis in *Idl* receptor deficient mice. *J Lipid Res*. 2009;50:376-385
33. Gibbons GF, Wiggins D, Brown AM, Hebbachi AM. Synthesis and function of hepatic very-low-density lipoprotein. *Biochem Soc Trans*. 2004;32:59-64
34. Kugiyama K, Doi H, Motoyama T, Soejima H, Misumi K, Kawano H, Nakagawa O, Yoshimura M, Ogawa H, Matsumura T, Sugiyama S, Nakano T, Nakajima K, Yasue H. Association of remnant lipoprotein levels with impairment of endothelium-dependent vasomotor function in human coronary arteries. *Circulation*. 1998;97:2519-2526
35. Krauss RM, Lindgren FT, Williams PT, Kelsey SF, Brensike J, Vranizan K, Detre KM, Levy RI. Intermediate-density lipoproteins and progression of coronary artery disease in hypercholesterolaemic men. *Lancet*. 1987;2:62-66
36. Kugiyama K, Doi H, Takazoe K, Kawano H, Soejima H, Mizuno Y, Tsunoda R, Sakamoto T, Nakano T, Nakajima K, Ogawa H, Sugiyama S, Yoshimura M, Yasue H. Remnant lipoprotein levels in fasting serum predict coronary events in patients with coronary artery disease. *Circulation*. 1999;99:2858-2860
37. Fukushima H, Kugiyama K, Sugiyama S, Honda O, Koide S, Nakamura S, Kawano H, Soejima H, Miyamoto S, Yoshimura M, Sakamoto T, Ogawa H. Comparison of remnant-like lipoprotein particles in postmenopausal women with and without coronary artery disease and in men with coronary artery disease. *American Journal of Cardiology*. 2001;88:1370-1373
38. Masuoka H, Kamei S, Wagayama H, Ozaki M, Kawasaki A, Tanaka T, Kitamura M, Katoh S, Shintani U, Misaki M, Sugawa M, Ito M, Nakano T. Association of remnant-like particle cholesterol with coronary artery disease in patients with normal total cholesterol levels. *American Heart Journal*. 2000;139:305-310
39. Wasan KM, Brocks DR, Lee SD, Sachs-Barrable K, Thornton SJ. Impact of lipoproteins on the biological activity and disposition of hydrophobic drugs: Implications for drug discovery. *Nat Rev Drug Discov*. 2008;7:84-99
40. Krauss RM, Burke DJ. Identification of multiple subclasses of plasma low density lipoproteins in normal humans. *J Lipid Res*. 1982;23:97-104
41. Coresh J, Kwiterovich PO, Jr. Small, dense low-density lipoprotein particles and coronary heart disease risk: A clear association with uncertain implications. *JAMA*. 1996;276:914-915
42. Rajman I, Eacho PI, Chowienczyk PJ, Ritter JM. *Ldl* particle size: An important drug target? *Br J Clin Pharmacol*. 1999;48:125-133
43. de Graaf J, Hak-Lemmers HL, Hectors MP, Demacker PN, Hendriks JC, Stalenhoef AF. Enhanced susceptibility to in vitro oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arterioscler Thromb*. 1991;11:298-306
44. Hurt-Camejo E, Olsson U, Wiklund O, Bondjers G, Camejo G. Cellular consequences of the association of apob lipoproteins with proteoglycans. Potential contribution to atherogenesis. *Arterioscler Thromb Vasc Biol*. 1997;17:1011-1017
45. Vakkilainen J, Makimattila S, Seppala-Lindroos A, Vehkavaara S, Lahdenpera S, Groop PH, Taskinen MR, Yki-Jarvinen H. Endothelial dysfunction in men with small *Idl* particles. *Circulation*. 2000;102:716-721
46. Austin MA, King MC, Vranizan KM, Krauss RM. Atherogenic lipoprotein phenotype. A proposed genetic marker for coronary heart disease risk. *Circulation*. 1990;82:495-506
47. Berneis K, Rizzo M. *Ldl* size: Does it matter. *Swiss Med Wkly*. 2004;134:720-724
48. Rajman I, Eacho PI, Chowienczyk P, Ritter J. *Ldl* particle size: An important drug target? *British journal of clinical pharmacology*. 1999;48:125-133
49. Spady DK. Reverse cholesterol transport and atherosclerosis regression. *Circulation*. 1999;100:576-578

50. Favari E, Chroni A, Tietge UJ, Zanotti I, Escola-Gil JC, Bernini F. Cholesterol efflux and reverse cholesterol transport. *Handb Exp Pharmacol*. 2015;224:181-206
51. Rosenson RS, Brewer HB, Jr., Davidson WS, Fayad ZA, Fuster V, Goldstein J, Hellerstein M, Jiang XC, Phillips MC, Rader DJ, Remaley AT, Rothblat GH, Tall AR, Yvan-Charvet L. Cholesterol efflux and atheroprotection: Advancing the concept of reverse cholesterol transport. *Circulation*. 2012;125:1905-1919
52. Duffy D, Rader DJ. Update on strategies to increase hdl quantity and function. *Nat Rev Cardiol*. 2009;6:455-463
53. Hogarth CA, Roy A, Ebert DL. Genomic evidence for the absence of a functional cholesteryl ester transfer protein gene in mice and rats. *Comp Biochem Physiol B Biochem Mol Biol*. 2003;135:219-229
54. Barter P. The role of hdl-cholesterol in preventing atherosclerotic disease. *European heart journal Supplements*. 2005;7:F4-F8
55. Nofer JR, Kehrel B, Fobker M, Levkau B, Assmann G, von Eckardstein A. Hdl and arteriosclerosis: Beyond reverse cholesterol transport. *Atherosclerosis*. 2002;161:1-16
56. Mackness MI, Arrol S, Durrington PN. Paraonase prevents accumulation of lipoperoxides in low-density lipoprotein. *FEBS Lett*. 1991;286:152-154
57. Mackness MI, Arrol S, Abbott C, Durrington PN. Protection of low-density lipoprotein against oxidative modification by high-density lipoprotein associated paraonase. *Atherosclerosis*. 1993;104:129-135
58. Clay MA, Pyle DH, Rye KA, Vadas MA, Gamble JR, Barter PJ. Time sequence of the inhibition of endothelial adhesion molecule expression by reconstituted high density lipoproteins. *Atherosclerosis*. 2001;157:23-29
59. Yuhanna IS, Zhu Y, Cox BE, Hahner LD, Osborne-Lawrence S, Lu P, Marcel YL, Anderson RG, Mendelsohn ME, Hobbs HH, Shaul PW. High-density lipoprotein binding to scavenger receptor-bi activates endothelial nitric oxide synthase. *Nat Med*. 2001;7:853-857
60. Mineo C, Yuhanna IS, Quon MJ, Shaul PW. High density lipoprotein-induced endothelial nitric-oxide synthase activation is mediated by akt and map kinases. *J Biol Chem*. 2003;278:9142-9149
61. Riwanto M, Landmesser U. High density lipoproteins and endothelial functions: Mechanistic insights and alterations in cardiovascular disease. *J Lipid Res*. 2013;54:3227-3243
62. Keene D, Price C, Shun-Shin MJ, Francis DP. Effect on cardiovascular risk of high density lipoprotein targeted drug treatments niacin, fibrates, and cetp inhibitors: Meta-analysis of randomised controlled trials including 117 411 patients. *Bmj-Brit Med J*. 2014;349
63. Khera AV, Cuchel M, de la Llera-Moya M, Rodrigues A, Burke MF, Jafri K, French BC, Phillips JA, Mucksavage ML, Wilensky RL, Mohler ER, Rothblat GH, Rader DJ. Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. *New England Journal of Medicine*. 2011;364:127-135
64. Rohatgi A, Khera A, Berry JD, Givens EG, Ayers CR, Wedin KE, Neeland IJ, Yuhanna IS, Rader DR, de Lemos JA, Shaul PW. Hdl cholesterol efflux capacity and incident cardiovascular events. *N Engl J Med*. 2014;371:2383-2393
65. Annema W, Dijkers A, de Boer JF, van Greevenbroek MMJ, van der Kallen CJH, Schalkwijk CG, Stehouwer CDA, Dullaart RPF, Tietge UJF. Impaired hdl cholesterol efflux in metabolic syndrome is unrelated to glucose tolerance status: The codam study. *Sci Rep-Uk*. 2016;6
66. Saleheen D, Scott R, Javad S, Zhao W, Rodrigues A, Picataggi A, Lukmanova D, Mucksavage ML, Luben R, Billheimer J, Kastelein JJP, Boekholdt SM, Khaw KT, Wareham N, Rader DJ. Association of hdl cholesterol efflux capacity with incident coronary heart disease events: A prospective case-control study. *Lancet Diabetes Endo*. 2015;3:507-513
67. Siddiqi HK, Kiss D, Rader D. Hdl-cholesterol and cardiovascular disease: Rethinking our approach. *Curr Opin Cardiol*. 2015;30:536-542
68. Weber C, Schober A, Zernecke A. Micrnas in arterial remodelling, inflammation and atherosclerosis. *Current Drug Targets*. 2010;11:950-956

69. Hansson GK. Mechanisms of disease - inflammation, atherosclerosis, and coronary artery disease. *New England Journal of Medicine*. 2005;352:1685-1695
70. Ley K, Laudanna C, Cybulsky MI, Nourshargh S. Getting to the site of inflammation: The leukocyte adhesion cascade updated. *Nature Reviews Immunology*. 2007;7:678-689
71. Singh RB, Mengi SA, Xu YJ, Arneja AS, Dhalla NS. Pathogenesis of atherosclerosis: A multifactorial process. *Exp Clin Cardiol*. 2002;7:40-53
72. Fishbein MC. The vulnerable and unstable atherosclerotic plaque. *Cardiovasc Pathol*. 2010;19:6-11
73. Libby P, Ridker PM, Hansson GK. Progress and challenges in translating the biology of atherosclerosis. *Nature*. 2011;473:317-325
74. Rubin R, Strayer DS, Rubin E. *Rubin's pathology: Clinicopathologic foundations of medicine*. Lippincott Williams & Wilkins; 2008.
75. Volkman A, Gowans JL. The origin of macrophages from bone marrow in the rat. *Br J Exp Pathol*. 1965;46:62-70
76. Passlick B, Flieger D, Ziegler-Heitbrock HW. Identification and characterization of a novel monocyte subpopulation in human peripheral blood. *Blood*. 1989;74:2527-2534
77. Ziegler-Heitbrock HW, Fingerle G, Strobel M, Schraut W, Stelter F, Schutt C, Passlick B, Pforte A. The novel subset of cd14+/cd16+ blood monocytes exhibits features of tissue macrophages. *Eur J Immunol*. 1993;23:2053-2058
78. Geissmann F, Jung S, Littman DR. Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity*. 2003;19:71-82
79. Palframan RT, Jung S, Cheng CY, Weninger W, Luo Y, Dorf M, Littman DR, Rollins BJ, Zweerink H, Rot A, von Andrian UH. Inflammatory chemokine transport and presentation in heV: A remote control mechanism for monocyte recruitment to lymph nodes in inflamed tissues. *Journal of Experimental Medicine*. 2001;194:1361-1373
80. Yang J, Zhang L, Yu C, Yang X-F, Wang H. Monocyte and macrophage differentiation: Circulation inflammatory monocyte as biomarker for inflammatory diseases. *Biomarker research*. 2014;2:1
81. Mantovani A, Garlanda C, Locati M. Macrophage diversity and polarization in atherosclerosis: A question of balance. *Arterioscler Thromb Vasc Biol*. 2009;29:1419-1423
82. Colin S, Chinetti-Gbaguidi G, Staels B. Macrophage phenotypes in atherosclerosis. *Immunol Rev*. 2014;262:153-166
83. Mackaness GB. Cellular resistance to infection. *Journal of Experimental Medicine*. 1962;116:381-&
84. Stein M, Keshav S, Harris N, Gordon S. Interleukin 4 potently enhances murine macrophage mannose receptor activity: A marker of alternative immunologic macrophage activation. *J Exp Med*. 1992;176:287-292
85. Martinez FO, Sica A, Mantovani A, Locati M. Macrophage activation and polarization. *Front Biosci-Landmrk*. 2008;13:453-461
86. Chistiakov DA, Bobryshev YV, Nikiforov NG, Elizova NV, Sobenin IA, Orekhov AN. Macrophage phenotypic plasticity in atherosclerosis: The associated features and the peculiarities of the expression of inflammatory genes. *Int J Cardiol*. 2015;184:436-445
87. Dewals BG, Marillier RG, Hoving JC, Leeto M, Schwegmann A, Brombacher F. Il-4alpha-independent expression of mannose receptor and ym1 by macrophages depends on their il-10 responsiveness. *PLoS Negl Trop Dis*. 2010;4:e689
88. Roszer T. Understanding the mysterious m2 macrophage through activation markers and effector mechanisms. *Mediat Inflamm*. 2015
89. Aderem A, Underhill DM. Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol*. 1999;17:593-623
90. de Winther MPJ, van Dijk KW, Havekes LM, Hofker MH. Macrophage scavenger receptor class a - a multifunctional receptor in atherosclerosis. *Arterioscl Throm Vas*. 2000;20:290-297
91. Kunjathoor VV, Febbraio M, Podrez EA, Moore KJ, Andersson L, Koehn S, Rhee JS, Silverstein R, Hoff HF, Freeman MW. Scavenger receptors class ai/ii and cd36 are the principal receptors responsible for the uptake of modified low density lipoprotein leading to lipid loading in macrophages. *Journal of Biological Chemistry*. 2002;277:49982-49988

92. Canton J, Neculai D, Grinstein S. Scavenger receptors in homeostasis and immunity. *Nature Reviews Immunology*. 2013;13:621-634
93. Huang SCC, Everts B, Ivanova Y, O'Sullivan D, Nascimento M, Smith AM, Beatty W, Love-Gregory L, Lam WY, O'Neil CM, Yan C, Du H, Abumrad NA, Urban JF, Artyomov MN, Pearce EL, Pearce EJ. Cell-intrinsic lysosomal lipolysis is essential for alternative activation of macrophages. *Nature Immunology*. 2014;15:846-855
94. van Tits LJ, Stienstra R, van Lent PL, Netea MG, Joosten LA, Stalenhoef AF. Oxidized ldl enhances pro-inflammatory responses of alternatively activated m2 macrophages: A crucial role for kruppel-like factor 2. *Atherosclerosis*. 2011;214:345-349
95. Zheng XF, Hong YX, Feng GJ, Zhang GF, Rogers H, Lewis MAO, Williams DW, Xia ZF, Song B, Wei XQ. Lipopolysaccharide-induced m2 to m1 macrophage transformation for il-12p70 production is blocked by candida albicans mediated up-regulation of ebi3 expression. *Plos One*. 2013;8
96. Wang N, Liang HW, Zen K. Molecular mechanisms that influence the macrophage m1-m2 polarization balance. *Front Immunol*. 2014;5
97. Shirey KA, Pletneva LM, Puche AC, Keegan AD, Prince GA, Blanco JCG, Vogel SN. Control of rsv-induced lung injury by alternatively activated macrophages is il-4r alpha-, tlr4-, and ifn-beta-dependent. *Mucosal Immunol*. 2010;3:291-300
98. Yurdagul A, Finney AC, Woolard MD, Orr AW. The arterial microenvironment: The where and why of atherosclerosis. *Biochemical Journal*. 2016;473:1281-1295
99. Tabas I, Bornfeldt KE. Macrophage phenotype and function in different stages of atherosclerosis. *Circulation Research*. 2016;118:653-667
100. da Silva RF, Lappalainen J, Lee-Rueckert M, Kovanen PT. Conversion of human m-csf macrophages into foam cells reduces their proinflammatory responses to classical m1-polarizing activation. *Atherosclerosis*. 2016;248:170-178
101. Kadl A, Meher AK, Sharma PR, Lee MY, Doran AC, Johnstone SR, Elliott MR, Gruber F, Han J, Chen WS, Kensler T, Ravichandran KS, Isakson BE, Wamhoff BR, Leitinger N. Identification of a novel macrophage phenotype that develops in response to atherogenic phospholipids via nrf2. *Circulation Research*. 2010;107:737-U155
102. Boyle JJ, Harrington HA, Piper E, Elderfield K, Stark J, Landis RC, Haskard DO. Coronary intraplaque hemorrhage evokes a novel atheroprotective macrophage phenotype. *American Journal of Pathology*. 2009;174:1097-1108
103. Finn AV, Nakano M, Polavarapu R, Karmali V, Saeed O, Zhao XQ, Yazdani S, Otsuka F, Davis T, Habib A, Narula J, Kolodgie FD, Virmani R. Hemoglobin directs macrophage differentiation and prevents foam cell formation in human atherosclerotic plaques. *Journal of the American College of Cardiology*. 2012;59:166-177
104. Leitinger N, Schulman IG. Phenotypic polarization of macrophages in atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2013;33:1120-1126
105. Freiman PC, Mitchell GG, Heistad DD, Armstrong ML, Harrison DG. Atherosclerosis impairs endothelium-dependent vascular relaxation to acetylcholine and thrombin in primates. *Circulation Research*. 1986;58:783-789
106. Getz GS, Reardon CA. Animal models of atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2012;32:1104-1115
107. Luginbühl H, Pauli B, Ratcliffe HL. Atherosclerosis in swine and swine as a model for the study of atherosclerosis. *Comparative Pathology of the Heart*. 119-126
108. Russell J. Of mice and men, rats and atherosclerosis. *Cardiovascular Research*. 2003;59:810-811
109. Shore B, Shore V. Rabbits as a model for the study of hyperlipoproteinemia and atherosclerosis. *Advances in Experimental Medicine and Biology*. 1976:123-141
110. Vesselinovitch D, Getz GS, Hughes RH, Wissler RW. Atherosclerosis in the rhesus monkey fed three food fats. *Atherosclerosis*. 1974;20:303-321
111. Getz GS, Reardon CA. Diet and murine atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2006;26:242-249

112. Kapourchali FR, Surendiran G, Chen L, Uitz E, Bahadori B, Moghadasian MH. Animal models of atherosclerosis. *World J Clin Cases*. 2014;2:126-132
113. Whitman SC. A practical approach to using mice in atherosclerosis research. *Clinical Biochemist Reviews*. 2004;25:81-93
114. Kuipers F, van Ree JM, Hofker MH, Wolters H, In't Veld G, Havinga R, Vonk RJ, Princen HM, Havekes LM. Altered lipid metabolism in apolipoprotein e-deficient mice does not affect cholesterol balance across the liver. *Hepatology*. 1996;24:241-247
115. Plump AS, Forte TM, Eisenberg S, Breslow JL. Atherogenic beta-vldl in the apo e-deficient mouse - composition, origin, and fate. *Circulation*. 1993;88:2-2
116. Meir KS, Leitersdorf E. Atherosclerosis in the apolipoprotein-e-deficient mouse: A decade of progress. *Arterioscler Thromb Vasc Biol*. 2004;24:1006-1014
117. Breslow JL. Mouse models of atherosclerosis. *Science*. 1996;272:685-688
118. Reddick RL, Zhang SH, Maeda N. Atherosclerosis in mice lacking apo-e - evaluation of lesional development and progression. *Arteriosclerosis and Thrombosis*. 1994;14:141-147
119. Nakashima Y, Plump AS, Raines EW, Breslow JL, Ross R. Apoe-deficient mice develop lesions of all phases of atherosclerosis throughout the arterial tree. *Arteriosclerosis and Thrombosis*. 1994;14:133-140
120. Ishibashi S, Brown MS, Goldstein JL, Gerard RD, Hammer RE, Herz J. Hypercholesterolemia in low density lipoprotein receptor knockout mice and its reversal by adenovirus-mediated gene delivery. *J Clin Invest*. 1993;92:883-893
121. Zadelaar S, Kleemann R, Verschuren L, de Vries-Van der Weij J, van der Hoorn J, Princen HM, Kooistra T. Mouse models for atherosclerosis and pharmaceutical modifiers. *Arterioscler Thromb Vasc Biol*. 2007;27:1706-1721
122. Ishibashi S, Goldstein JL, Brown MS, Herz J, Burns DK. Massive xanthomatosis and atherosclerosis in cholesterol-fed low density lipoprotein receptor-negative mice. *J Clin Invest*. 1994;93:1885-1893
123. Kowala MC, Recce R, Beyer S, Gu C, Valentine M. Characterization of atherosclerosis in ldl receptor knockout mice: Macrophage accumulation correlates with rapid and sustained expression of aortic mcp-1/jc. *Atherosclerosis*. 2000;149:323-330
124. Ma Y, Wang W, Zhang J, Lu Y, Wu W, Yan H, Wang Y. Hyperlipidemia and atherosclerotic lesion development in ldlr-deficient mice on a long-term high-fat diet. *PLoS One*. 2012;7:e35835
125. Tangirala RK, Rubin EM, Palinski W. Quantitation of atherosclerosis in murine models: Correlation between lesions in the aortic origin and in the entire aorta, and differences in the extent of lesions between sexes in ldl receptor-deficient and apolipoprotein e-deficient mice. *J Lipid Res*. 1995;36:2320-2328
126. Lichtman AH, Clinton SK, Iiyama K, Connelly PW, Libby P, Cybulsky MI. Hyperlipidemia and atherosclerotic lesion development in ldl receptor-deficient mice fed defined semipurified diets with and without cholate. *Arterioscl Thromb Vas*. 1999;19:1938-1944
127. Ishibashi S, Goldstein JL, Brown MS, Herz J, Burns DK. Massive xanthomatosis and atherosclerosis in cholesterol-fed low-density-lipoprotein receptor-negative mice. *Journal of Clinical Investigation*. 1994;93:1885-1893
128. Linton MF, Fazio S. Macrophages, lipoprotein metabolism, and atherosclerosis: Insights from murine bone marrow transplantation studies. *Current Opinion in Lipidology*. 1999;10:97-105
129. Chorro FJ, Such-Belenguer L, Lopez-Merino V. Animal models of cardiovascular disease. *Rev Esp Cardiol*. 2009;62:69-84
130. Ma YL, Wang WY, Zhang J, Lu YL, Wu WY, Yan H, Wang YP. Hyperlipidemia and atherosclerotic lesion development in ldlr-deficient mice on a long-term high-fat diet. *Plos One*. 2012;7
131. Rosenfeld ME, Polinsky P, Virmani R, Kauser K, Rubanyi G, Schwartz SM. Advanced atherosclerotic lesions in the innominate artery of the apoe knockout mouse. *Arterioscl Thromb Vas*. 2000;20:2587-2592
132. Herijgers N, VanEck M, Groot PHE, Hoogerbrugge PM, VanBerkel TJC. Effect of bone marrow transplantation on lipoprotein metabolism and atherosclerosis in ldl receptor knockout mice. *Arterioscl Thromb Vas*. 1997;17:1995-2003

133. Boisvert WA, Spangenberg J, Curtiss LK. Role of leukocyte-specific ldl receptors on plasma lipoprotein cholesterol and atherosclerosis in mice. *Arterioscl Throm Vas*. 1997;17:340-347
134. Fazio S, Hasty AH, Carter KJ, Murray AB, Price JO, Linton MF. Leukocyte low density lipoprotein receptor (ldl-r) does not contribute to ldl clearance in vivo: Bone marrow transplantation studies in the mouse. *J Lipid Res*. 1997;38:391-400
135. Linton MF, Atkinson JB, Fazio S. Prevention of atherosclerosis in apolipoprotein e-deficient mice by bone-marrow transplantation. *Science*. 1995;267:1034-1037
136. Fazio S, Linton MF. Murine bone marrow transplantation as a novel approach to studying the role of macrophages in lipoprotein metabolism and atherogenesis. *Trends Cardiovas Med*. 1996;6:58-65
137. Boisvert WA, Spangenberg J, Curtiss LK. Treatment of severe hypercholesterolemia in apolipoprotein e-deficient mice by bone-marrow transplantation. *Journal of Clinical Investigation*. 1995;96:1118-1124
138. VanEck M, Herijgers N, Yates J, Pearce NJ, Hoogerbrugge PM, Groot PHE, VanBerkel TJC. Bone marrow transplantation in apolipoprotein e deficient mice - effect of apoe gene dosage on serum lipid concentrations, (beta)vldl catabolism, and atherosclerosis. *Arterioscl Throm Vas*. 1997;17:3117-3126
139. Lawrence T, Natoli G. Transcriptional regulation of macrophage polarization: Enabling diversity with identity. *Nat Rev Immunol*. 2011;11:750-761
140. Bekkering S, Joosten LA, van der Meer JW, Netea MG, Riksen NP. The epigenetic memory of monocytes and macrophages as a novel drug target in atherosclerosis. *Clin Ther*. 2015;37:914-923
141. Sica A, Mantovani A. Macrophage plasticity and polarization: In vivo veritas. *J Clin Invest*. 2012;122:787-795
142. Lee CH, Chawla A, Urbiztondo N, Liao D, Boisvert WA, Evans RM, Curtiss LK. Transcriptional repression of atherogenic inflammation: Modulation by ppardelta. *Science*. 2003;302:453-457
143. Moore KJ, Sheedy FJ, Fisher EA. Macrophages in atherosclerosis: A dynamic balance. *Nat Rev Immunol*. 2013;13:709-721
144. Wang X, Cao Q, Yu L, Shi H, Xue B, Shi H. Epigenetic regulation of macrophage polarization and inflammation by DNA methylation in obesity. *JCI Insight*. 2016;1:e87748
145. Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajani-Refah A, Matarese F, Cheng SC, Ratter J, Berentsen K, van der Ent MA, Sharifi N, Janssen-Megens EM, Ter Huurne M, Mandoli A, van Schaik T, Ng A, Burden F, Downes K, Frontini M, Kumar V, Giamarellos-Bourboulis EJ, Ouweland WH, van der Meer JW, Joosten LA, Wijmenga C, Martens JH, Xavier RJ, Logie C, Netea MG, Stunnenberg HG. Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science*. 2014;345:1251086
146. Rask-Andersen M, Zhang J, Fabbro D, Schioth HB. Advances in kinase targeting: Current clinical use and clinical trials. *Trends in Pharmacological Sciences*. 2014;35:60-76
147. Liu PX, Cheng HL, Roberts TM, Zhao JJ. Targeting the phosphoinositide 3-kinase pathway in cancer. *Nature Reviews Drug Discovery*. 2009;8:627-644
148. Hommes DW, Peppelenbosch MP, van Deventer SJ. Mitogen activated protein (map) kinase signal transduction pathways and novel anti-inflammatory targets. *Gut*. 2003;52:144-151
149. English JM, Cobb MH. Pharmacological inhibitors of mapk pathways. *Trends Pharmacol Sci*. 2002;23:40-45
150. Kaminska B. Mapk signalling pathways as molecular targets for anti-inflammatory therapy - from molecular mechanisms to therapeutic benefits. *Bba-Proteins Proteom*. 2005;1754:253-262
151. Tugal D, Liao X, Jain MK. Transcriptional control of macrophage polarization. *Arterioscler Thromb Vasc Biol*. 2013;33:1135-1144
152. Laurila PP, Soronen J, Kooijman S, Forsstrom S, Boon MR, Surakka I, Kaiharju E, Coomans CP, Van Den Berg SA, Autio A, Sarin AP, Kettunen J, Tikkanen E, Manninen T, Metso J, Silvennoinen R, Merikanto K, Ruuth M, Perttinen J, Makela A, Isomi A, Tuomainen AM, Tikka A, Ramadan UA, Seppala I, Lehtimäki T, Eriksson J, Havulinna A, Jula A, Karhunen PJ, Salomaa V, Perola M, Ehnholm C, Lee-Rueckert M, Van Eck M, Roivainen A, Taskinen MR, Peltonen L, Mervaala E, Jalanko A, Hohtola E,

- Olkkonen VM, Ripatti S, Kovanen PT, Rensen PC, Suomalainen A, Jauhiainen M. Usp1 deficiency activates brown adipose tissue and improves cardiometabolic health. *Sci Transl Med*. 2016;8:323ra313
153. Horbach T, Gotz C, Kietzmann T, Dimova EY. Protein kinases as switches for the function of upstream stimulatory factors: Implications for tissue injury and cancer. *Front Pharmacol*. 2015;6
 154. Hou YC, Janczuk A, Wang PG. Current trends in the development of nitric oxide donors. *Curr Pharm Des*. 1999;5:417-441
 155. Forstermann U, Sessa WC. Nitric oxide synthases: Regulation and function. *Eur Heart J*. 2012;33:829-837, 837a-837d
 156. Xu F, Sun YN, Chen YG, Sun Y, Li RJ, Liu CX, Zhang C, Wang R, Zhang Y. Endothelial cell apoptosis is responsible for the formation of coronary thrombotic atherosclerotic plaques. *Tohoku J Exp Med*. 2009;218:25-33
 157. Pierini D, Bryan NS. Nitric oxide availability as a marker of oxidative stress. *Advanced Protocols in Oxidative Stress III*. 2015:63-71
 158. Cybulsky MI, Iiyama K, Li H, Zhu S, Chen M, Iiyama M, Davis V, Gutierrez-Ramos JC, Connelly PW, Milstone DS. A major role for vcam-1, but not icam-1, in early atherosclerosis. *J Clin Invest*. 2001;107:1255-1262
 159. Kuhlencordt PJ, Gyrko R, Han F, Scherrer-Crosbie M, Aretz TH, Hajjar R, Picard MH, Huang PL. Accelerated atherosclerosis, aortic aneurysm formation, and ischemic heart disease in apolipoprotein e/endothelial nitric oxide synthase double-knockout mice. *Circulation*. 2001;104:448-454
 160. Hayashi T, Sumi D, Juliet PAR, Matsui-Hirai H, Asai-Tanaka Y, Kano H, Fukatsu A, Tsunekawa T, Miyazaki A, Iguchi A, Ignarro LJ. Gene transfer of endothelial no synthase, but not enos plus inducible nos, regressed atherosclerosis in rabbits. *Cardiovascular Research*. 2004;61:339-351
 161. Mujynya-Ludunge K, Viswambharan H, Driscoll R, Ming XF, von Segesser LK, Kappenberger L, Yang ZH, Vassalli G. Endothelial nitric oxide synthase gene transfer restores endothelium-dependent relaxations and attenuates lesion formation in carotid arteries in apolipoprotein e-deficient mice. *Basic Research in Cardiology*. 2005;100:102-111
 162. Ozaki M, Kawashima S, Yamashita T, Hirase T, Namiki M, Inoue N, Hirata K, Yasui H, Sakurai H, Yoshida Y, Masada M, Yokoyama M. Overexpression of endothelial nitric oxide synthase accelerates atherosclerotic lesion formation in apoe-deficient mice. *Journal of Clinical Investigation*. 2002;110:331-340
 163. Detmers PA, Hernandez M, Mudgett J, Hassing H, Burton C, Mundt S, Chun S, Fletcher D, Card DJ, Lisnock J, Weikel R, Bergstrom JD, Shevell DE, Hermanowski-Vosatka A, Sparrow CP, Chao YS, Rader DJ, Wright SD, Pure E. Deficiency in inducible nitric oxide synthase results in reduced atherosclerosis in apolipoprotein e-deficient mice. *Journal of Immunology*. 2000;165:3430-3435
 164. Kuhlencordt PJ, Chen JQ, Han F, Astern J, Huang PL. Genetic deficiency of inducible nitric oxide synthase reduces atherosclerosis and lowers plasma lipid peroxides in apolipoprotein e-knockout mice. *Circulation*. 2001;103:3099-3104
 165. Mujynya-Ludunge K, Viswambharan H, Driscoll R, Ming XF, von Segesser LK, Kappenberger L, Yang Z, Vassalli G. Endothelial nitric oxide synthase gene transfer restores endothelium-dependent relaxations and attenuates lesion formation in carotid arteries in apolipoprotein e-deficient mice. *Basic Res Cardiol*. 2005;100:102-111
 166. Fish JE, Marsden PA. Endothelial nitric oxide synthase: Insight into cell-specific gene regulation in the vascular endothelium. *Cell Mol Life Sci*. 2006;63:144-162
 167. Green SJ, Scheller LF, Marletta MA, Seguin MC, Klotz FW, Slayter M, Nelson BJ, Nacy CA. Nitric-oxide - cytokine-regulation of nitric-oxide in host-resistance to intracellular pathogens. *Immunology Letters*. 1994;43:87-94
 168. Forstermann U, Sessa WC. Nitric oxide synthases: Regulation and function. *European Heart Journal*. 2012;33:829-+
 169. Lowenstein CJ. Beneficial effects of neuronal nitric oxide synthase in atherosclerosis. *Arterioscl Throm Vas*. 2006;26:1417-1418

170. Verbeuren TJ, Coene MC, Jordaens FH, Vanhove CE, Zonnekeyn LL, Herman AG. Effect of hypercholesterolemia on vascular reactivity in the rabbit .2. Influence of treatment with dipyridamole on endothelium-dependent and endothelium-independent responses in isolated aortas of control and hypercholesterolemic rabbits. *Circulation Research*. 1986;59:496-504
171. Kuhlencordt PJ, Hotten S, Schodel J, Rutzel S, Hu K, Widder J, Marx A, Huang PL, Ertl G. Atheroprotective effects of neuronal nitric oxide synthase in apolipoprotein e knockout mice. *Arterioscl Throm Vas*. 2006;26:1539-1544
172. Bredt DS, Glatt CE, Hwang PM, Fotuhi M, Dawson TM, Snyder SH. Nitric oxide synthase protein and mrna are discretely localized in neuronal populations of the mammalian cns together with nadph diaphorase. *Neuron*. 1991;7:615-624
173. Mungrue IN, Bredt DS. Nnos at a glance: Implications for brain and brawn. *J Cell Sci*. 2004;117:2627-2629
174. Papapetropoulos A, Rudic RD, Sessa WC. Molecular control of nitric oxide synthases in the cardiovascular system. *Cardiovascular Research*. 1999;43:509-520
175. Schodel J, Padmapriya P, Marx A, Huang PL, Ertl G, Kuhlencordt PJ. Expression of neuronal nitric oxide synthase splice variants in atherosclerotic plaques of apoe knockout mice. *Atherosclerosis*. 2009;206:383-389
176. Capettini LSA, Cortes SF, Silva JF, Alvarez-Leite JJ, Lemos VS. Decreased production of neuronal nos-derived hydrogen peroxide contributes to endothelial dysfunction in atherosclerosis. *Brit J Pharmacol*. 2011;164:1738-1748
177. BATTERY LD, Springall DR, Chester AH, Evans TJ, Standfield EN, Parums DV, Yacoub MH, Polak JM. Inducible nitric oxide synthase is present within human atherosclerotic lesions and promotes the formation and activity of peroxynitrite. *Lab Invest*. 1996;75:77-85
178. Miyoshi T, Li YH, Shih DM, Wang XP, Laubach VE, Matsumoto AH, Helm GA, Lusis AJ, Shi WB. Deficiency of inducible no synthase reduces advanced but not early atherosclerosis in apolipoprotein e-deficient mice. *Life Sciences*. 2006;79:525-531
179. Albina JE, Abate JA, Mastrofrancesco B. Role of ornithine as a proline precursor in healing wounds. *J Surg Res*. 1993;55:97-102
180. Gonzalez E, McGraw TE. The akt kinases isoform specificity in metabolism and cancer. *Cell Cycle*. 2009;8:2502-2508
181. Franke TF, Hornik CP, Segev L, Shostak GA, Sugimoto C. Pi3k/akt and apoptosis: Size matters. *Oncogene*. 2003;22:8983-8998
182. Liang J, Slingerland JM. Multiple roles of the pi3k/pkb (akt) pathway in cell cycle progression. *Cell Cycle*. 2003;2:339-345
183. Morales-Ruiz M, Fulton D, Sowa G, Languino LR, Fujio Y, Walsh K, Sessa WC. Vascular endothelial growth factor-stimulated actin reorganization and migration of endothelial cells is regulated via the serine/threonine kinase akt. *Circulation Research*. 2000;86:892-896
184. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, Anderson MJ, Arden KC, Blenis J, Greenberg ME. Akt promotes cell survival by phosphorylating and inhibiting a forkhead transcription factor. *Cell*. 1999;96:857-868
185. Zeng GY, Nystrom FH, Ravichandran LV, Cong LN, Kirby M, Mostowski H, Quon MJ. Roles for insulin receptor, pi3-kinase, and akt in insulin-signaling pathways related to production of nitric oxide in human vascular endothelial cells. *Circulation*. 2000;101:1539-1545
186. Yu HX, Littlewood T, Bennett M. Akt isoforms in vascular disease. *Vasc Pharmacol*. 2015;71:57-64
187. Fayard E, Tintignac LA, Baudry A, Hemmings BA. Protein kinase b/akt at a glance. *J Cell Sci*. 2005;118:5675-5678
188. Babaev VR, Hebron KE, Wiese CB, Toth CL, Ding L, Zhang Y, May JM, Fazio S, Vickers KC, Linton MF. Macrophage deficiency of akt2 reduces atherosclerosis in ldlr null mice. *J Lipid Res*. 2014;55:2296-2308
189. Xu F, Kang Y, Zhang H, Piao Z, Yin H, Diao R, Xia J, Shi L. Akt1-mediated regulation of macrophage polarization in a murine model of staphylococcus aureus pulmonary infection. *J Infect Dis*. 2013;208:528-538

190. Arranz A, Doxaki C, Vergadi E, Martinez de la Torre Y, Vaporidi K, Lagoudaki ED, Ieronymaki E, Androulidaki A, Venihaki M, Margioris AN, Stathopoulos EN, Tsielis PN, Tsatsanis C. Akt1 and akt2 protein kinases differentially contribute to macrophage polarization. *Proc Natl Acad Sci U S A*. 2012;109:9517-9522
191. Rotllan N, Chamorro-Jorganes A, Araldi E, Wanschel AC, Aryal B, Aranda JF, Goedeke L, Salerno AG, Ramirez CM, Sessa WC, Suarez Y, Fernandez-Hernando C. Hematopoietic akt2 deficiency attenuates the progression of atherosclerosis. *Faseb Journal*. 2015;29:597-610
192. Fernandez-Hernando C, Ackah E, Yu J, Suarez Y, Murata T, Iwakiri Y, Prendergast J, Miao RQ, Birnbaum MJ, Sessa WC. Loss of akt1 leads to severe atherosclerosis and occlusive coronary artery disease. *Cell Metab*. 2007;6:446-457
193. Ding L, Biswas S, Morton RE, Smith JD, Hay N, Byzova TV, Febbraio M, Podrez EA. Akt3 deficiency in macrophages promotes foam cell formation and atherosclerosis in mice. *Cell Metab*. 2012;15:861-872
194. Fernandez-Hernando C, Jozsef L, Jenkins D, Di Lorenzo A, Sessa WC. Absence of akt1 reduces vascular smooth muscle cell migration and survival and induces features of plaque vulnerability and cardiac dysfunction during atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2009;29:2033-2040
195. Yu H, Littlewood T, Bennett M. Akt isoforms in vascular disease. *Vascul Pharmacol*. 2015;71:57-64
196. Rensing KL, de Jager SCA, Stroes ES, Vos M, Twickler MTB, Dallinga-Thie GM, de Vries CJM, Kuiper J, Bot I, von der Thusen JH. Akt2/Ildlr double knockout mice display impaired glucose tolerance and develop more complex atherosclerotic plaques than Ildlr knockout mice. *Cardiovascular Research*. 2014;101:277-287
197. Babaev VR, Hebron KE, Wiese CB, Toth CL, Ding L, Zhang YM, May JM, Fazio S, Vickers KC, Linton MF. Macrophage deficiency of akt2 reduces atherosclerosis in Ildlr null mice. *Journal of Lipid Research*. 2014;55:2296-2308
198. Pasterkamp G, van der Laan SW, Haitjema S, Foroughi Asl H, Siemelink MA, Bezemer T, van Setten J, Dichgans M, Malik R, Worrall BB, Schunkert H, Samani NJ, de Kleijn DP, Markus HS, Hoefer IE, Michoel T, de Jager SC, Bjorkegren JL, den Ruijter HM, Asselbergs FW. Human validation of genes associated with a murine atherosclerotic phenotype. *Arterioscler Thromb Vasc Biol*. 2016;36:1240-1246
199. Blackhart B, Yao Z, McCarthy B. An expression system for human apolipoprotein b100 in a rat hepatoma cell line. *Journal of Biological Chemistry*. 1990;265:8358-8360
200. Babaev VR, Ding L, Zhang Y, May JM, Linton MF. Loss of macrophage akt2/akt3 or akt1/akt2 decreases cell survival and suppresses early atherosclerosis in Ildlr null mice. 2015
201. Hu Y, Dietrich H, Metzler B, Wick G, Xu Q. Hyperexpression and activation of extracellular signal-regulated kinases (erk1/2) in atherosclerotic lesions of cholesterol-fed rabbits. *Arterioscler Thromb Vasc Biol*. 2000;20:18-26
202. Rahaman SO, Lennon DJ, Febbraio M, Podrez EA, Hazen SL, Silverstein RL. A cd36-dependent signaling cascade is necessary for macrophage foam cell formation. *Cell Metabolism*. 2006;4:211-221
203. Zhang L, Chen Y, Yang X, Yang J, Cao X, Li X, Li L, Miao QR, Hajjar DP, Duan Y, Han J. Mek1/2 inhibitors activate macrophage abcg1 expression and reverse cholesterol transport-an anti-atherogenic function of erk1/2 inhibition. *Biochim Biophys Acta*. 2016;1861:1180-1191
204. Chen Y, Duan Y, Yang X, Sun L, Liu M, Wang Q, Ma X, Zhang W, Li X, Hu W, Miao RQ, Xiang R, Hajjar DP, Han J. Inhibition of erk1/2 and activation of Ixr synergistically reduce atherosclerotic lesions in apoe-deficient mice. *Arterioscler Thromb Vasc Biol*. 2015;35:948-959
205. Park SJ, Lee KP, Kang S, Lee J, Sato K, Chung HY, Okajima F, Im DS. Sphingosine 1-phosphate induced anti-atherogenic and atheroprotective m2 macrophage polarization through il-4. *Cell Signal*. 2014;26:2249-2258
206. Long ME, Eddy WE, Gong KQ, Lovelace-Macon LL, McMahan RS, Charron J, Liles WC, Manicone AM. Mek1/2 inhibition promotes macrophage reparative properties. *J Immunol*. 2017;198:862-872

207. Yang Y, Kim SC, Yu T, Yi YS, Rhee MH, Sung GH, Yoo BC, Cho JY. Functional roles of p38 mitogen-activated protein kinase in macrophage-mediated inflammatory responses. *Mediators Inflamm.* 2014;2014:352371
208. Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, Landvatter SW, Strickler JE, McLaughlin MM, Siemens IR, Fisher SM, Livi GP, White JR, Adams JL, Young PR. A protein-kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature.* 1994;372:739-746
209. Kim SH, Kim J, Sharma RP. Inhibition of p38 and erk map kinases blocks endotoxin-induced nitric oxide production and differentially modulates cytokine expression. *Pharmacological Research.* 2004;49:433-439
210. Zhu W, Downey JS, Gu J, Di Padova F, Gram H, Han JH. Regulation of tnfr expression by multiple mitogen-activated protein kinase pathways. *Journal of Immunology.* 2000;164:6349-6358
211. Perregaux DG, Dean D, Cronan M, Connelly P, Gabel CA. Inhibition of interleukin-1 beta production by skf86002: Evidence of two sites of in vitro activity and of a time and system dependence. *Mol Pharmacol.* 1995;48:433-442
212. Kang YJ, Chen J, Otsuka M, Mols J, Ren S, Wang Y, Han J. Macrophage deletion of p38alpha partially impairs lipopolysaccharide-induced cellular activation. *J Immunol.* 2008;180:5075-5082
213. Jimenez-Garcia L, Herranz S, Luque A, Hortelano S. Critical role of p38 mapk in il-4-induced alternative activation of peritoneal macrophages. *European Journal of Immunology.* 2015;45:273-286
214. Zhao M, Liu YW, Wang XF, New LG, Han JH, Brunk UT. Activation of the p38 map kinase pathway is required for foam cell formation from macrophages exposed to oxidized ldl. *Apmis.* 2002;110:458-468
215. Yin R. Ppar phosphorylation mediated by jnk mapk: A potential role in macrophage-derived foam cell formation. *Acta pharmacologica sinica.* 2006;27:1146-1152
216. Kardakaris R, Gareus R, Xanthoulea S, Pasparakis M. Endothelial and macrophage-specific deficiency of p38alpha mapk does not affect the pathogenesis of atherosclerosis in apoe-/- mice. *PLoS One.* 2011;6:e21055
217. Seimon TA, Wang Y, Han S, Senokuchi T, Schrijvers DM, Kuriakose G, Tall AR, Tabas IA. Macrophage deficiency of p38alpha mapk promotes apoptosis and plaque necrosis in advanced atherosclerotic lesions in mice. *J Clin Invest.* 2009;119:886-898
218. Seeger FH, Sedding D, Langheinrich AC, Haendeler J, Zeiher AM, Dimmeler S. Inhibition of the p38 map kinase in vivo improves number and functional activity of vasculogenic cells and reduces atherosclerotic disease progression. *Basic Res Cardiol.* 2010;105:389-397
219. Jagavelu K, Tietge UJ, Gaestel M, Drexler H, Schieffer B, Bavendiek U. Systemic deficiency of the map kinase-activated protein kinase 2 reduces atherosclerosis in hypercholesterolemic mice. *Circ Res.* 2007;101:1104-1112
220. Waetzig V, Herdegen T. Context-specific inhibition of jnks: Overcoming the dilemma of protection and damage. *Trends Pharmacol Sci.* 2005;26:455-461
221. Babaev VR, Yeung M, Erbay E, Ding L, Zhang Y, May JM, Fazio S, Hotamisligil GS, Linton MF. Jnk1 deficiency in hematopoietic cells suppresses macrophage apoptosis and increases atherosclerosis in low-density lipoprotein receptor null mice. *Arterioscler Thromb Vasc Biol.* 2016;36:1122-1131
222. Himes SR, Sester DP, Ravasi T, Cronau SL, Sasmono T, Hume DA. The jnk are important for development and survival of macrophages. *J Immunol.* 2006;176:2219-2228
223. Sanchez-Tillo E, Comalada M, Xaus J, Farrera C, Villedor AF, Caelles C, Lloberas J, Celada A. Jnk1 is required for the induction of mkp1 expression in macrophages during proliferation and lipopolysaccharide-dependent activation. *Journal of Biological Chemistry.* 2007;282:12566-12573
224. Utsugi M, Dobashi K, Ono A, Ishizuka T, Hisada T, Koga Y, Shimizu Y, Kawata T, Matsuzaki S, Aoki H, Kamide Y, Mori M. Jnk1 and jnk2 differently regulate il-12 production in thp-1 macrophage cells. *Cytokine.* 2010;51:127-131
225. Guma M, Ronacher LM, Firestein GS, Karin M, Corr M. Jnk-1 deficiency limits macrophage-mediated antigen-induced arthritis. *Arthritis Rheum.* 2011;63:1603-1612

226. Muslin AJ. Mapk signalling in cardiovascular health and disease: Molecular mechanisms and therapeutic targets. *Clin Sci*. 2008;115:203-218
227. Ricci R, Sumara G, Sumara I, Rozenberg I, Kurrer M, Akhmedov A, Hersberger M, Eriksson U, Eberli FR, Becher B, Boren J, Chen M, Cybulsky MI, Moore KJ, Freeman MW, Wagner EF, Matter CM, Luscher TF. Requirement of jnk2 for scavenger receptor a-mediated foam cell formation in atherogenesis. *Science*. 2004;306:1558-1561
228. Metzler B, Hu YH, Dietrich H, Xu QB. Increased expression and activation of stress-activated protein kinases/c-jun nh(2)-terminal protein kinases in atherosclerotic lesions coincide with p53. *American Journal of Pathology*. 2000;156:1875-1886
229. Fey D, Croucher DR, Kolch W, Kholodenko BN. Crosstalk and signaling switches in mitogen-activated protein kinase cascades. *Front Physiol*. 2012;3
230. Caunt CJ, Keyse SM. Dual-specificity map kinase phosphatases (mkps): Shaping the outcome of map kinase signalling. *FEBS J*. 2013;280:489-504
231. Salojin K, Oravecz T. Regulation of innate immunity by mapk dual-specificity phosphatases: Knockout models reveal new tricks of old genes. *J Leukoc Biol*. 2007;81:860-869
232. Huang CY, Tan TH. Dusps, to map kinases and beyond. *Cell Biosci*. 2012;2
233. Patterson KI, Brummer T, O'Brien PM, Daly RJ. Dual-specificity phosphatases: Critical regulators with diverse cellular targets. *Biochemical Journal*. 2009;418:475-489
234. Theodosiou A, Ashworth A. Map kinase phosphatases. *Genome Biol*. 2002;3:REVIEWS3009
235. Camps M, Nichols A, Arkinstall S. Dual specificity phosphatases: A gene family for control of map kinase function. *FASEB J*. 2000;14:6-16
236. Lang R, Hammer M, Mages J. Dusp meet immunology: Dual specificity mapk phosphatases in control of the inflammatory response. *Journal of Immunology*. 2006;177:7497-7504
237. Sim J, Yi K, Kim H, Ahn H, Chung Y, Rehman A, Jang SM, Lee KH, Jang K, Paik SS. Immunohistochemical expression of dual-specificity protein phosphatase 4 in patients with colorectal adenocarcinoma. *Gastroenterol Res Pract*. 2015;2015:283764
238. Ham JE, Oh EK, Kim DH, Choi SH. Differential expression profiles and roles of inducible dusps and erk1/2-specific constitutive dusp6 and dusp7 in microglia. *Biochem Biophys Res Commun*. 2015;467:254-260
239. Kondoh K, Nishida E. Regulation of map kinases by map kinase phosphatases. *Biochim Biophys Acta*. 2007;1773:1227-1237
240. Abraham SM, Clark AR. Dual-specificity phosphatase 1: A critical regulator of innate immune responses. *Biochem Soc T*. 2006;34:1018-1023
241. Smallie T, Ross EA, Ammit AJ, Cunliffe HE, Tang T, Rosner DR, Ridley ML, Buckley CD, Saklatvala J, Dean JL, Clark AR. Dual-specificity phosphatase 1 and tristetraprolin cooperate to regulate macrophage responses to lipopolysaccharide. *J Immunol*. 2015;195:277-288
242. Jeffrey KL, Brummer T, Rolph MS, Liu SM, Callejas NA, Grumont RJ, Gillieron C, Mackay F, Grey S, Camps M, Rommel C, Gerondakis SD, Mackay CR. Positive regulation of immune cell function and inflammatory responses by phosphatase pac-1. *Nature Immunology*. 2006;7:274-283
243. Al-Mutairi MS, Cadalbert LC, McGachy HA, Shweash M, Schroeder J, Kurnik M, Sloss CM, Bryant CE, Alexander J, Plevin R. Map kinase phosphatase-2 plays a critical role in response to infection by leishmania mexicana. *Plos Pathog*. 2010;6
244. Cornell TT, Rodenhouse P, Cai Q, Sun L, Shanley TP. Mitogen-activated protein kinase phosphatase 2 regulates the inflammatory response in sepsis. *Infection and Immunity*. 2010;78:2868-2876
245. Rushworth LK, Kidger AM, Delavaine L, Stewart G, van Schelven S, Davidson J, Bryant CJ, Caddy E, East P, Caunt CJ, Keyse SM. Dual-specificity phosphatase 5 regulates nuclear erk activity and suppresses skin cancer by inhibiting mutant harvey-ras (hrasq61l)-driven serpinb2 expression. *Proc Natl Acad Sci U S A*. 2014;111:18267-18272
246. Luo LJ, Liu F, Wang XY, Dai TY, Dai YL, Dong C, Ge BX. An essential function for mkp5 in the formation of oxidized low density lipid-induced foam cells. *Cell Signal*. 2012;24:1889-1898

247. Shen J, Chandrasekharan UM, Ashraf MZ, Long E, Morton RE, Liu Y, Smith JD, DiCorleto PE. Lack of mitogen-activated protein kinase phosphatase-1 protects apoe-null mice against atherosclerosis. *Circ Res*. 2010;106:902-910
248. Imaizumi S, Grijalva V, Priceman S, Wu L, Su F, Farias-Eisner R, Hama S, Navab M, Fogelman AM, Reddy ST. Mitogen-activated protein kinase phosphatase-1 deficiency decreases atherosclerosis in apolipoprotein e null mice by reducing monocyte chemoattractant protein-1 levels. *Mol Genet Metab*. 2010;101:66-75
249. Gregor PD, Sawadogo M, Roeder RG. The adenovirus major late transcription factor usf is a member of the helix-loop-helix group of regulatory proteins and binds to DNA as a dimer. *Genes & Development*. 1990;4:1730-1740
250. Sirito M, Lin Q, Maity T, Sawadogo M. Ubiquitous expression of the 43- and 44-kda forms of transcription factor usf in mammalian cells. *Nucleic Acids Res*. 1994;22:427-433
251. Pajukanta P, Lilja HE, Sinsheimer JS, Cantor RM, Lusi AJ, Gentile M, Duan XJ, Soro-Paavonen A, Naukkarinen J, Saarela J, Laakso M, Ehnholm C, Taskinen MR, Peltonen L. Familial combined hyperlipidemia is associated with upstream transcription factor 1 (usf1). *Nat Genet*. 2004;36:371-376
252. Ng MC, Miyake K, So WY, Poon EW, Lam VK, Li JK, Cox NJ, Bell GI, Chan JC. The linkage and association of the gene encoding upstream stimulatory factor 1 with type 2 diabetes and metabolic syndrome in the chinese population. *Diabetologia*. 2005;48:2018-2024
253. Coon H, Xin Y, Hopkins PN, Cawthon RM, Hasstedt SJ, Hunt SC. Upstream stimulatory factor 1 associated with familial combined hyperlipidemia, ldl cholesterol, and triglycerides. *Hum Genet*. 2005;117:444-451
254. Sanchez AP, Zhao JH, You Y, Decleves AE, Diamond-Stanic M, Sharma K. Role of the usf1 transcription factor in diabetic kidney disease. *Am J Physiol-Renal*. 2011;301:F271-F279
255. Naukkarinen J, Gentile M, Soro-Paavonen A, Saarela J, Koistinen HA, Pajukanta P, Taskinen MR, Peltonen L. Usf1 and dyslipidemias: Converging evidence for a functional intronic variant. *Human Molecular Genetics*. 2005;14:2595-2605
256. Galibert MD, Carreira S, Goding CR. The usf-1 transcription factor is a novel target for the stress-responsive p38 kinase and mediates uv-induced tyrosinase expression. *Embo Journal*. 2001;20:5022-5031
257. Nowak M, Helleboid-Chapman A, Jakel H, Martin G, Duran-Sandoval D, Staels B, Rubin EM, Pennacchio LA, Taskinen MR, Fruchart-Najib J, Fruchart JC. Insulin-mediated down-regulation of apolipoprotein a5 gene expression through the phosphatidylinositol 3-kinase pathway: Role of upstream stimulatory factor. *Mol Cell Biol*. 2005;25:1537-1548
258. Imagawa S, Fujii S, Dong J, Furumoto T, Kaneko T, Zaman T, Satoh Y, Tsutsui H, Sobel BE. Hepatocyte growth factor regulates e box-dependent plasminogen activator inhibitor type 1 gene expression in hepg2 liver cells. *Arterioscl Thromb Vas*. 2006;26:2407-2413
259. Chanda D, Li TG, Song KH, Kim YH, Sim JG, Lee CH, Chiang JYL, Choi HS. Hepatocyte growth factor family negatively regulates hepatic gluconeogenesis via induction of orphan nuclear receptor small heterodimer partner in primary hepatocytes. *Journal of Biological Chemistry*. 2009;284:28510-28521
260. Terragni J, Nayak G, Banerjee S, Medrano JL, Graham JR, Brennan JF, Sepulveda S, Cooper GM. The e-box binding factors max/mnt, mitf, and usf1 act coordinately with foxo to regulate expression of proapoptotic and cell cycle control genes by phosphatidylinositol 3-kinase/akt/glycogen synthase kinase 3 signaling. *J Biol Chem*. 2011;286:36215-36227
261. Corre S, Primot A, Baron Y, Le Seyec J, Goding C, Galibert MD. Target gene specificity of usf-1 is directed via p38-mediated phosphorylation-dependent acetylation. *Journal of Biological Chemistry*. 2009;284:18851-18862
262. Corre S, Primot A, Sviderskaya E, Bennett DC, Vaulont S, Goding CR, Galibert MD. Uv-induced expression of key component of the tanning process, the pomc and mc1r genes, is dependent on the p-38-activated upstream stimulating factor-1 (usf-1). *Journal of Biological Chemistry*. 2004;279:51226-51233

263. Sag D, Carling D, Stout RD, Suttles J. Adenosine 5'-monophosphate-activated protein kinase promotes macrophage polarization to an anti-inflammatory functional phenotype. *J Immunol.* 2008;181:8633-8641
264. Lupp S, Gotz C, Khadouma S, Horbach T, Dimova EY, Bohrer AM, Kietzmann T, Montenarh M. The upstream stimulatory factor usf1 is regulated by protein kinase ck2 phosphorylation. *Cell Signal.* 2014;26:2809-2817
265. Naukkarinen J, Ehnholm C, Peltonen L. Genetics of familial combined hyperlipidemia. *Current Opinion in Lipidology.* 2006;17:285-290