



Universiteit
Leiden
The Netherlands

Lab-on-a-tissue : optimization of on-tissue chemistry for improved mass spectrometry imaging

Heijs, B.P.A.M.

Citation

Heijs, B. P. A. M. (2018, February 1). *Lab-on-a-tissue : optimization of on-tissue chemistry for improved mass spectrometry imaging*. Retrieved from <https://hdl.handle.net/1887/60212>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/60212>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/60212> holds various files of this Leiden University dissertation.

Author: Heijs, B.P.A.M.

Title: Lab-on-a-tissue : optimization of on-tissue chemistry for improved mass spectrometry imaging

Issue Date: 2018-02-01

Bibliography

- Abdelmoula, W.M., Carreira, R.J., Shyti, R., Balluff, B., van Zeijl, R.J.J., Tolner, E.A., Lelieveldt, B.F., van den Maagdenberg, A.M.J.M., McDonnell, L.A., and Dijkstra, J. (2014). Automatic registration of mass spectrometry imaging data sets to the Allen brain atlas. *Anal Chem* 86, 3947–3954.
- Adamczyk, B., Tharmalingam, T., and Rudd, P.M. (2012). Glycans as cancer biomarkers. *Biochim Biophys Acta* 1820, 1347–1353.
- Addie, R.D., Balluff, B., Bovée, J.V., Morreau, H., and McDonnell, L.A. (2015). Current state and future challenges of mass spectrometry imaging for clinical research. *Anal Chem* 87, 6426–6433.
- Aerni, H.R., Cornett, D.S., and Caprioli, R.M. (2006). Automated acoustic matrix deposition for MALDI sample preparation. *Anal Chem* 78, 827–834.
- Aichler, M., and Walch, A. (2015). MALDI Imaging mass spectrometry: current frontiers and perspectives in pathology research and practice. *Lab Invest* 95, 422–431.
- Aichler, M., Elsner, M., Ludyga, N., Feuchtinger, A., Zangen, V., Maier, S., Balluff, B., Schöne, C., Hierber, L., Braselmann, H., et al. (2013). Clinical response to chemotherapy in oesophageal adenocarcinoma patients is linked to defects in mitochondria. *J Pathol* 230, 410–419.
- Altelaar, M., van Minnen, J., Jiménez, C., Heeren, R., and Piersma, S. (2005). Direct molecular imaging of *Lymnaea stagnalis* nervous tissue at subcellular spatial resolution by mass spectrometry. *Anal Chem* 77, 735–741.
- Altelaar, M., Munoz, J., and Heck, A. (2012). Next-generation proteomics: towards an integrative view of proteome dynamics. *Nat Rev Genet* 14, 35–48.
- Amini, A., Dormady, S.J., Riggs, L., and Regnier, F.E. (2000). The impact of buffers and surfactants from micellar electrokinetic chromatography on matrix-assisted laser desorption ionization (MALDI) mass spectrometry of peptides. Effect of buffer type and concentration on mass determination by MALDI-time-of-flight mass spectrometry. *J Chromatogr A* 894, 345–355.
- Amstalden van Hove, E.R., Blackwell, T.R., Klinkert, I., Eijkel, G.B., Heeren, R.M., and Glunde, K. (2010). Multimodal mass spectrometric imaging of small molecules reveals distinct spatio-molecular signatures in differentially metastatic breast tumor models. *Cancer Res* 70, 9012–9021.
- Angel, P.M., Spraggins, J.M., Baldwin, H.S., and Caprioli, R.M. (2012). Enhanced sensitivity for high spatial resolution lipid analysis by negative ion mode matrix assisted laser desorption ionization imaging mass spectrometry. *Anal Chem* 84, 1557–1564.
- Antonescu, C.R., Tschernevsky, S.J., Decuseara, R., Leung, D.H., Woodruff, J.M., Brennan, M.F., Bridge, J.A., Neff, J.R., Goldblum, J.R., and Ladanyi, M. (2001). Prognostic impact of P53 status, TLS-CHOP fusion transcript structure, and histological grade in myxoid liposarcoma: a molecular and clinicopathologic study of 82 cases. *Clin Cancer Res* 7, 3977–3987.
- Anugrahan, M., Jacob, F., Nixdorf, S., Everest-Dass, A.V., Heinzelmann-Schwarz, V., and Packer, N.H. (2014). Specific glycosylation of membrane proteins in epithelial ovarian cancer cell lines: glycan structures reflect gene expression and DNA methylation status. *Mol Cell Proteomics* 13, 2213–2232.
- Atkinson, S.J., Loadman, P.M., Sutton, C., Patterson, L.H., and Clench, M.R. (2007). Examination of the distribution of the bioreductive drug AQ4N and its active metabolite AQ4 in solid tumours by imaging matrix-assisted laser desorption/ionisation mass spectrometry. *Rapid Commun Mass Spectrom* 21, 1271–1276.
- Awad, H., Khamis, M.M., and El-Aneed, A. (2015). Mass spectrometry, review of the basics: ionization. *Appl Spectrosc Rev* 50, 158–175.
- Balluff, B., Elsner, M., Kowarsch, A., Rauser, S., Meding, S., Schuhmacher, C., Feith, M., Herrmann, K., Röcken, C., Schmid, R., et al. (2010). Classification of HER2/neu status in gastric cancer using a breast-cancer derived proteome classifier. *J Proteome Res* 9, 6317–6322.
- Balluff, B., Rauser, S., Meding, S., Elsner, M., Schöne, C., Feuchtinger, A., Schuhmacher, C., Novotny, A., Jütting, U., Maccarrone, G., et al. (2011). MALDI imaging identifies prognostic seven-protein signature of novel tissue markers in intestinal-type gastric cancer. *Am J Pathol* 179, 2720–2729.
- Balluff, B., Frese, C.K., Maier, S.K., Schöne, C., Kuster, B., Schmitt, M., Aubele, M., Höfler, H., Deelder, A.M.M., Heck, A., et al. (2015). De novo discovery of phenotypic intratumour heterogeneity using imaging mass spectrometry. *J Pathol* 235, 3–13.
- Barry, J., Robichaud, G., and Muddiman, D. (2013). Mass recalibration of FT-ICR mass spectrometry imaging data using the average frequency shift of ambient ions. *J Am Soc Mass Spectrom* 24, 1137–1145.
- Beaudry, F., Le Blanc, Y., Coutu, Ramier, I., Moreau, J.P., and Brown, N.K. (1999). Metabolite profiling study of propranolol in rat using LC/MS/MS analysis. *Biomed Chromatogr* 13, 363–369.
- Bhatia, V.N., Perlman, D.H., Costello, C.E., and McComb, M.E. (2009). Software tool for researching annotations of proteins: open-source protein annotation software with data visualization. *Anal Chem* 81, 9819–9823.
- Bovée, J.V.M.G., and Hogendoorn, P.C. (2010). Molecular pathology of sarcomas: concepts and clinical implications. *Virchows Arch* 456, 193–199.
- Briggs, M.T., Ho, Y.Y., Kaur, G., Oehler, M.K., Everest-Dass, A.V., Packer, N.H., and Hoffmann, P. (2017). N-glycan matrix-assisted laser desorption/ionization mass spectrometry imaging protocol for formalin-fixed paraffin-embedded tissues. *Rapid Commun Mass Spectrom* epub ahead of print.
- Brown, R.S., and Lennon, J.J. (1995). Mass resolution improvement by incorporation of pulsed ion extraction in a matrix-assisted laser desorption/ionization linear time-of-flight mass spectrometer. *Anal Chem* 67, 1998–2003.
- Brownridge, P., and Beynon, R.J. (2011). The importance of the digest: proteolysis and absolute quantification in proteomics. *Methods* 54, 351–360.

- Buck, A., Ly, A., Balluff, B., Sun, N., Gorzolka, K., Feuchtinger, A., Janssen, K.-P.P., Kuppen, P.J., van de Velde, C.J., Weirich, G., et al. (2015). High-resolution MALDI-FT-ICR MS imaging for the analysis of metabolites from formalin-fixed, paraffin-embedded clinical tissue samples. *J Pathol* 237, 123–132.
- Budnik, B., Haselmann, K., and Zubarev, R. (2001). Electron detachment dissociation of peptide di-anions: an electron–hole recombination phenomenon. *Chem Phys Lett* 342, 299–302.
- Buse, J., Purves, R.W., Verrall, R.E., Badea, I., Zhang, H., Mulligan, C.C., Bailey, J., Headley, J.V., and El-Aneed, A. (2014). The development and assessment of high-throughput mass spectrometry-based methods for the quantification of a nanoparticle drug delivery agent in cellular lysate. *J Mass Spectrom* 49, 1171–1180.
- Caldwell, R.L., Holt, G.E., and Caprioli, R.M. (2005). Tissue Profiling by MALDI Mass Spectrometry Distinguishes Clinical Grades of Soft Tissue Sarcomas. *Cancer Genom Proteom* 2, 333–345.
- Caldwell, R.L., Gonzalez, A., Oppenheimer, S.R., Schwartz, H.S., and Caprioli, R.M. (2006). Molecular Assessment of the Tumor Protein Microenvironment Using Imaging Mass Spectrometry. *Cancer Genom Proteom* 3, 279–287.
- Calvano, C.D., Carulli, S., and Palmisano, F. (2010). 1H-Pteridine-2,4-dione (lumazine): a new MALDI matrix for complex (phospho) lipid mixtures analysis. *Anal Bioanal Chem* 398, 499–507.
- Caprioli, R.M., Farmer, T.B., and Gile, J. (1997). Molecular imaging of biological samples: localization of peptides and proteins using MALDI-TOF MS. *Anal Chem* 69, 4751–4760.
- Carlson, B., Bernstein, D., Annis, D., Misenheimer, T., Hannah, B., Mosher, D., and Keck, J. (2005). Structure of the calcium-rich signature domain of human thrombospondin-2. *Nat Struct Mol Biology* 12, 910–914.
- Carreira, R.J., Shyti, R., Balluff, B., Abdelmoula, W.M., van Heiningen, S.H., van Zeijl, R.J., Dijkstra, J., Ferrari, M.D., Tolner, E.A., McDonnell, L.A., et al. (2015). Large-scale mass spectrometry imaging investigation of consequences of cortical spreading depression in a transgenic mouse model of migraine. *J Am Soc Mass Spectrom* 26, 853–861.
- Casadonte, R., and Caprioli, R.M. (2011). Proteomic analysis of formalin-fixed paraffin-embedded tissue by MALDI imaging mass spectrometry. *Nat Protoc* 6, 1695–1709.
- Casadonte, R., Kriegsmann, M., Zweynert, F., Friedrich, K., Baretton, G., Bretton, G., Otto, M., Deininger, S.-O., Paape, R., Belau, E., et al. (2014). Imaging mass spectrometry to discriminate breast from pancreatic cancer metastasis in formalin-fixed paraffin-embedded tissues. *Proteomics* 14, 956–964.
- Cazares, L.H., Troyer, D.A., Mendrinos, S., Lance, R.A., Nyalwidhe, J.O., Beydoun, H.A., Clements, M.A., Drake, R.R., and Semmes, O.J. (2009). Imaging mass spectrometry of a specific fragment of mitogen-activated protein kinase/extracellular signal-regulated kinase kinase kinase 2 discriminates cancer from uninvolved prostate tissue. *Clin Cancer Res* 15, 5541–5551.
- Cazares, L.H., Troyer, D.A., Wang, B., Drake, R.R., and Semmes, O.J. (2011). MALDI tissue imaging: from biomarker discovery to clinical applications. *Anal Bioanal Chem* 401, 17–27.
- Cazares, L.H., Van Tongeren, S.A., Costantino, J., Kenny, T., Garza, N.L., Donnelly, G., Lane, D., Panchal, R.G., and Bavari, S. (2015). Heat fixation inactivates viral and bacterial pathogens and is compatible with downstream MALDI mass spectrometry tissue imaging. *BMC Microbiol* 15, 1–11.
- Chacon, A., Zagol-Ikapitte, I., Amarnath, V., Reyzer, M.L., Oates, J.A., Caprioli, R.M., and Boutaud, O. (2011). On-tissue chemical derivatization of 3-methoxysalicylamine for MALDI-imaging mass spectrometry. *J Mass Spectrom* 46, 840–846.
- Chaurand, P., Schwartz, S.A., Billheimer, D., Xu, B.J., Crecelius, A., and Caprioli, R.M. (2004). Integrating histology and imaging mass spectrometry. *Anal Chem* 76, 1145–1155.
- Chaurand, P., Cornett, D.S., Angel, P.M., and Caprioli, R.M. (2011). From whole-body sections down to cellular level, multiscale imaging of phospholipids by MALDI mass spectrometry. *Mol Cell Proteomics* 10, O110.004259.
- Chaurand, P., Stoeckli, M., Caprioli, R.M. (1999). Direct profiling of proteins in biological tissue sections by MALDI mass spectrometry. *Anal Chem* 71, 5263–5270.
- Chen, R., Hui, L., Sturm, R.M., and Li, L. (2009a). Three dimensional mapping of neuropeptides and lipids in crustacean brain by mass spectral imaging. *J Am Soc Mass Spectrom* 20, 1068–1077.
- Chen, R., Jiang, X., Sun, D., Han, G., Wang, F., Ye, M., Wang, L., and Zou, H. (2009b). Glycoproteomics analysis of human liver tissue by combination of multiple enzyme digestion and hydrazide chemistry. *J Proteome Res* 8, 651–661.
- Choudhary, G., Wu, S.-L., Shieh, P., and Hancock, W.S. (2003). Multiple enzymatic digestion for enhanced sequence coverage of proteins in complex proteomic mixtures using capillary LC with ion trap MS/MS. *J Proteome Res* 2, 59–67.
- Chughtai, K., Jiang, L., Greenwood, T.R., Klinkert, I., Amstalden van Hove, E.R., Heeren, R.M.A., and Glunde, K. (2012). Fiducial markers for combined 3-dimensional mass spectrometric and optical tissue imaging. *Anal Chem* 84, 1817–1823.
- Clemis, E.J., Smith, D.S., Camenzind, A., Danell, R.M., Parker, C.E., and Borchers, C.H. (2012). Quantitation of spatially-localized proteins in tissue samples using MALDI-MRM imaging. *Anal Chem* 84, 3514–3522.
- Cole, L.M., Djidja, M.-C., Bluff, J., Claude, E., Carolan, V.A., Paley, M., Tozer, G.M., and Clench, M.R. (2011). Investigation of protein induction in tumour vascular targeted strategies by MALDI MSI. *Methods* 54, 442–453.
- Comisarow, M.B., and Marshall, A.G. (1974). Fourier transform ion cyclotron resonance spectroscopy. *Chem Phys Lett* 25, 282–283.
- Coon, J.J., Shabanowitz, J., Hunt, D.F., and Syka, J.E.P. (2005). Electron transfer dissociation of peptide anions. *J Am Soc Mass Spectrom* 16, 880–882.
- Cornett, D.S., Mobley, J.A., Dias, E.C., Andersson, M., Arteaga, C.L., Sanders, M.E., and Caprioli, R.M. (2006). A novel histology-directed strategy for MALDI-MS tissue profiling that improves throughput and cellular specificity in human breast cancer. *Mol Cell Proteomics* 5, 1975–1983.

- Cotter, R.J. (1992). Time-of-flight mass spectrometry for the structural analysis of biological molecules. *Anal Chem* 64, 1027A–1039A.
- Cristofanilli, M., Budd, G.T., Ellis, M.J., Stopeck, A., Matera, J., Miller, M.C., Reuben, J.M., Doyle, G.V., Allard, J., Terstappen, L.W.M.M., et al. (2004). Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 351, 781–791.
- Dalerba, P., Kalisky, T., Sahoo, D., Rajendran, P.S., Rothenberg, M.E., Leyrat, A.A., Sim, S., Okamoto, J., Johnston, D.M., Qian, D., et al. (2011). Single-cell dissection of transcriptional heterogeneity in human colon tumors. *Nat Biotechnol* 29, 1120–1127.
- D'Arrigo, A., Belluco, C., Ambrosi, A., Digito, M., Esposito, G., Bertola, A., Fabris, M., Nofrate, V., Mammano, E., Leon, A., et al. (2005). Metastatic transcriptional pattern revealed by gene expression profiling in primary colorectal carcinoma. *Int J Cancer* 115, 256–262.
- Dawson, J.H.J., and Guilhaus, M. (1989). Orthogonal acceleration time of flight mass spectrometer. *Rapid Commun Mass Spectrom* 3, 155–159.
- de Graaf, E.L., Altelaar, A.F.M., van Breukelen, B., Mohammed, S., and Heck, A.J.R. (2011). Improving SRM assay development: a global comparison between triple quadrupole, ion trap, and higher energy CID peptide fragmentation spectra. *J Proteome Res* 10, 4334–4341.
- de Graaff, M.A., Yu, J.S., Beird, H.C., Ingram, D.R., Nguyen, T., Juehui Liu, J., Bolshakov, S., Szuhai, K., Åman, P., Torres, K., et al. (2016). Establishment and characterization of a new human myxoid liposarcoma cell line (DL-221) with the FUS-DDIT3 translocation. *Lab Invest* 96, 885–894.
- de Graaff, M., Malu, S., Guardiola, I., Kruisselbrink, A., de Jong, Y., Corver, W., Gelderblom, H., Hwu, P., Nielsen, T., Lazar, J., et al. (2017). High-throughput screening of myxoid liposarcoma cell lines reveals survivin as a potential novel druggable target. Leiden University.
- de Hoffmann, E. (1996). Tandem mass spectrometry: a primer. *J Mass Spectrom* 31, 129–137.
- de Leoz, M.L., Young, L.J., An, H.J., Kronewitter, S.R., Kim, J., Miyamoto, S., Borowsky, A.D., Chew, H.K., and Lebrilla, C.B. (2011). High-mannose glycans are elevated during breast cancer progression. *Mol Cell Proteomics* 10, M110.002717.
- de Vreeze, R., de Jong, D., Nederlof, P., Ruijter, H.J., Boerrigter, L., Haas, R., van Coevorden, F. (2010). Multifocal myxoid liposarcoma - Metastasis or second primary tumor? A molecular biological analysis. *J Mol Diagn* 12, 238–243.
- Deininger, S.-O., Ebert, M.P., Fütterer, A., Gerhard, M., and Röcken, C. (2008). MALDI imaging combined with hierarchical clustering as a new tool for the interpretation of complex human cancers. *J Proteome Res* 7, 5230–5236.
- Dekker, T.J., Balluff, B.D., Jones, E.A., Schöne, C.D., Schmitt, M., Aubele, M., Kroep, J.R., Smit, V.T., Tollenaar, R.A., Mesker, W.E., et al. (2014). Multicenter matrix-assisted laser desorption/ionization mass spectrometry imaging (MALDI MSI) identifies proteomic differences in breast-cancer-associated stroma. *J Proteome Res* 13, 4730–4738.
- Delcorte, A., and Garrison, B.J. (2000). High yield events of molecular emission induced by kiloelectronvolt particle bombardment. *J Phys Chem* 104, 6785–6800.
- Demetriou, M., Nabi, I., Coppolino, M., Dedhar, S., and Dennis, J. (1995). Reduced contact-inhibition and substratum adhesion in epithelial cells expressing GlcNAc-transferase V. *J Cell Biol* 130, 383–392.
- Demicco, E., Torres, K., Ghadimi, M., Colombo, C., Bolshakov, S., Hoffman, A., Peng, T., Bovée, J., Wang, W.-L., Lev, D., et al. (2011). Involvement of the PI3K/Akt pathway in myxoid/round cell liposarcoma. *Modern Pathology* 25, 212–221.
- Dennis, J.W., Laferte, S., Waghorne, C., Breitman, M.L., and Kerbel, R.S. (1987). Beta 1-6 branching of Asn-linked oligosaccharides is directly associated with metastasis. *Science* 236, 582–585.
- Dennis, J.W., Granovsky, M., and Warren, C.E. (1999). Glycoprotein glycosylation and cancer progression. *Biochim Biophys Acta* 1473, 21–34.
- Dice, L.R. (1945). Measures of the amount of ecologic association between species. *Ecology* 26, 297–302.
- Diehl, H.C., Beine, B., Elm, J., Trede, D., Ahrens, M., Eisenacher, M., Marcus, K., Meyer, H.E., and Henkel, C. (2015). The challenge of on-tissue digestion for MALDI MSI— a comparison of different protocols to improve imaging experiments. *Anal Bioanal Chem* 407, 2223–2243.
- Dilillo, M., Ait-Belkacem, R., Esteve, E., Pellegrini, D., Nicolardi, S., Costa, M., Vannini, E., de Graaf, E.L., Caleo, M., McDonnell, L.A. (2017). Ultra-high mass resolution MALDI imaging mass spectrometry of proteins and metabolites in a mouse model of glioblastoma. *Sci Rep* 7, 603.
- Djidja, M.-C., Francesc, S., Loadman, P.M., Sutton, C.W., Scriven, P., Claude, E., Snel, M.F., Franck, J., Salzet, M., and Clench, M.R. (2009a). Detergent addition to tryptic digests and ion mobility separation prior to MS/MS improves peptide yield and protein identification for *in situ* proteomic investigation of frozen and formalin-fixed paraffin-embedded adenocarcinoma tissue sections. *Proteomics* 9, 2750–2763.
- Djidja, M.-C., Claude, E., Snel, M.F., Scriven, P., Francesc, S., Carolan, V., and Clench, M.R. (2009b). MALDI-ion mobility separation-mass spectrometry imaging of glucose-regulated protein 78 kDa (Grp78) in human formalin-fixed, paraffin-embedded pancreatic adenocarcinoma tissue sections. *J Proteome Res* 8, 4876–4884.
- Djidja, M.-C., Claude, E., Snel, M.F., Francesc, S., Scriven, P., Carolan, V., and Clench, M.R. (2010). Novel molecular tumour classification using MALDI-mass spectrometry imaging of tissue micro-array. *Anal Bioanal Chem* 397, 587–601.
- Domon, B., and Aebersold, R. (2006). Mass spectrometry and protein analysis. *Science* 312, 212–217.
- Dong, X., Cheng, J., Li, J., and Wang, Y. (2010). Graphene as a novel matrix for the analysis of small molecules by MALDI-TOF MS. *Anal Chem* 82, 6208–6214.

- Dong, Y., Li, B., Malitsky, S., Rogachev, I., Aharoni, A., Kaftan, F., Svatoš, A., and Franceschi, P. (2016). Sample preparation for mass spectrometry imaging of plant tissues: a review. *Front Plant Sci* 7, 60.
- Drake, P.M., Cho, W., Li, B., Prakobphol, A., Johansen, E., Anderson, N.L., Regnier, F.E., Gibson, B.W., and Fisher, S.J. (2010). Sweetening the pot: adding glycosylation to the biomarker discovery equation. *Clin Chem* 56, 223–236.
- Dreisewerd, K., Schürenberg, M., and Karas, M., Hillenkamp, F. (1995). Influence of the laser intensity and spot size on the desorption of molecules and ions in matrix-assisted laser desorption/ionization with a uniform beam profile. *Int J Mass Spectrom Ion Process* 141, 127–148.
- Dreisewerd, K. (2003). The desorption process in MALDI. *Chem Rev* 103, 395–426.
- Dreisewerd, K. (2014). Recent methodological advances in MALDI mass spectrometry. *Anal Bioanal Chem* 406, 2261–2278.
- Easterling, M.L., Mize, T.H., and Amster, J. (1999). Routine part-per-million mass accuracy for high-mass ions: space-charge effects in MALDI FT-ICR. *Anal Chem* 71, 624–632.
- Eberlin, L.S., Liu, X., Ferreira, C.R., Santagata, S., Agar, N.Y.R., and Cooks, R.G. (2011). Desorption electrospray ionization then MALDI mass spectrometry imaging of lipid and protein distributions in single tissue sections. *Anal Chem* 83, 8366–8371.
- Eikel, D., Vavrek, M., Smith, S., Bason, C., Yeh, S., Korfmacher, W.A., and Henion, J.D. (2011). Liquid extraction surface analysis mass spectrometry (LESA-MS) as a novel profiling tool for drug distribution and metabolism analysis: the terfenadine example. *Rapid Commun Mass Spectrom* 25, 3587–3596.
- El Ayed, M., Bonnel, D., Longuespée, R., Castelier, C., Franck, J., Vergara, D., Desmons, A., Tasiemski, A., Kenani, A., Vinatier, D., et al. (2010). MALDI imaging mass spectrometry in ovarian cancer for tracking, identifying, and validating biomarkers. *Med Sci Monit* 16, BR233–45.
- Ellis, S.R., Cappell, J., Potočnik, N.O., Balluff, B., Hamaide, J., der Linden, A., and Heeren, R.M. (2016). More from less: high-throughput dual polarity lipid imaging of biological tissues. *Analyst* 141, 3832–3841.
- Elsner, M., Rauser, S., Maier, S., Schöne, C., Balluff, B., Meding, S., Jung, G., Nipp, M., Sarıoglu, H., Maccarrone, G., et al. (2012). MALDI imaging mass spectrometry reveals COX7A2, TAGLN2 and S100-A10 as novel prognostic markers in Barrett's adenocarcinoma. *J Proteomics* 75, 4693–4704.
- Endo, M., Graaff, M., Ingram, D., Lim, S., Lev, D., Bruijn, I., Somaiah, N., Bovée, J., Lazar, A., and Nielsen, T. (2015). NY-ESO-1 (CTAG1B) expression in mesenchymal tumors. *Modern Pathology* 28, 587–595.
- Enthaler, B., Bussmann, T., Pruns, J.K., Rapp, C., Fischer, M., and Vietzke, J.-P. (2013a). Influence of various on-tissue washing procedures on the entire protein quantity and the quality of matrix-assisted laser desorption/ionization spectra. *Rapid Commun Mass Spectrom* 27, 878–884.
- Enthaler, B., Trusch, M., Fischer, M., Rapp, C., Pruns, J.K., and Vietzke, J.-P. (2013b). MALDI imaging in human skin tissue sections: focus on various matrices and enzymes. *Anal Bioanal Chem* 405, 1159–1170.
- Esteve, C., Marina, M.L., and García, M.C. (2015). Novel strategy for the revalorization of olive (*Olea europaea*) residues based on the extraction of bioactive peptides. *Food Chem* 167, 272–280.
- Esteve, C., Tolner, E.A., Shyti, R., van den Maagdenberg, A.M.J.M., and McDonnell, L.A. (2016). Mass spectrometry imaging of amino neurotransmitters: a comparison of derivatization methods and application in mouse brain tissue. *Metabolomics* 12, 30.
- Fenn, J.B., Mann, M., Meng, C.K., Wong, S.F., and Whitehouse, C.M. (1989). Electrospray ionization for mass spectrometry of large biomolecules. *Science* 246, 64–71.
- Fenn, J.B., Mann, M., Meng, C.K., Wong, S.F., and Whitehouse, C.M. (1990). Electrospray ionization—principles and practice. *Mass Spectrom Rev* 9, 37–70.
- Fernandes, B., Sagman, U., Auger, M., Demetrio, M., and Dennis, J.W. (1991). β 1–6 branched oligosaccharides as a marker of tumor progression in human breast and colon neoplasia. *Cancer Res* 51, 718–723.
- Fischer, W.H., Rivier, J.E., and Craig, A.G. (1993). In situ reduction suitable for matrix-assisted laser desorption/ionization and liquid secondary ionization using tris(2-carboxyethyl)phosphine. *Rapid Commun Mass Spectrom* 7, 225–228.
- Fletcher, C.D.M., Bridge, J.A., Hogendoorn, P.C.W., and Mertens, F. (2013). World Health Organization classification of tumours. Pathology and genetics of tumours of soft tissue and bone. 4th ed. Lyon: IARC Press.
- Fletcher, C.D.M. (2014). The evolving classification of soft tissue tumours - an update based on the new 2013 WHO classification. *Histopathology* 64, 2–11.
- Franck, J., El Ayed, M., Wisztorski, M., Salzet, M., and Fournier, I. (2009). On-tissue N-terminal peptide derivatizations for enhancing protein identification in MALDI mass spectrometric imaging strategies. *Anal Chem* 81, 8305–8317.
- Frandsen, T.J., Sherman, M.G., Hunter, R.L., Locke, M.J., Bowers, W.D., and McIver, R.T. (1983). Experimental determination of the effects of space charge on ion cyclotron resonance frequencies. *Int J Mass Spectrom Ion Process* 54, 189–199.
- Frankevich, V.E., Zhang, J., Friess, S.D., Dashtiev, M., and Zenobi, R. (2003). Role of electrons in laser desorption/ionization mass spectrometry. *Anal Chem* 75, 6063–6067.
- Freeze, H.H. (2006). Genetic defects in the human glycome. *Nat Rev Genet* 7, 537–551.
- Frese, C.K., Altelaar, A.F.M., Hennrich, M.L., Nolting, D., Zeller, M., Griep-Raming, J., Heck, A.J.R., and Mohammed, S. (2011). Improved peptide identification by targeted fragmentation using CID, HCD and ETD on an LTQ-Orbitrap Velos. *J Proteome Res* 10, 2377–2388.
- Frese, C.K., Mikhaylova, M., Stucchi, R., Gautier, V., Liu, Q., Mohammed, S., Heck, A.J.R., Altelaar, A.F.M., and Hoogenraad, C.C. (2017). Quantitative map of proteome dynamics during neuronal differentiation. *Cell Rep* 18, 1527–1542.

- Fritchie, K.J., Goldblum, J.R., Tubbs, R.R., Sun, Y., Carver, P., Billings, S.D., and Rubin, B.P. (2012). The expanded histologic spectrum of myxoid liposarcoma with an emphasis on newly described patterns: implications for diagnosis on small biopsy specimens. *Am J Clin Pathol* 137, 229–239.
- Ganisl, B., Valovka, T., Hartl, M., Taucher, M., Bister, K., and Breuker, K. (2011). Electron detachment dissociation for top-down mass spectrometry of acidic proteins. *Chem Eur J* 17, 4460–4469.
- Gatlin, C.L., Eng, J.K., Cross, S.T., Detter, J.C., and Yates, J.R. (2000). Automated identification of amino acid sequence variations in proteins by HPLC/microspray tandem mass spectrometry. *Anal Chem* 72, 757–763.
- Gerlinger, M., Rowan, A., Horswell, S., Larkin, J., Endesfelder, D., Gronroos, E., Martinez, P., Matthews, N., Stewart, A., Tarpey, P., et al. (2012). Intratumor Heterogeneity and Branched Evolution Revealed by Multiregion Sequencing. *N Eng J Med* 366, 883–892.
- Glish, G.L., and Vachet, R.W. (2003). The basics of mass spectrometry in the twenty-first century. *Nat Rev Drug Discov* 2, 140–150.
- Goodwin, R.J. (2012). Sample preparation for mass spectrometry imaging: small mistakes can lead to big consequences. *J Proteomics* 75, 4893–4911.
- Graadt van Roggen, J.F., Hogendoorn, P.C., and Fletcher, C.D. (1999). Myxoid tumours of soft tissue. *Histopathology* 35, 291–312.
- Granovsky, M., Fata, J., Pawling, J., Muller, W.J., Khokha, R., and Dennis, J.W. (2000). Suppression of tumor growth and metastasis in Mgat5-deficient mice. *Nat Med* 6, 306–312.
- Grassl, J., Taylor, NL, and Millar, AH (2011). Matrix-assisted laser desorption/ionisation mass spectrometry imaging and its development for plant protein imaging. *Plant Methods* 7, 21.
- Greaves, and Maley (2012). Clonal evolution in cancer. *Nature* 481, 306–313.
- Green-Mitchell, S.M., Cazares, L.H., Semmes, O.J., Nadler, J.L., and Nyalwidhe, J.O. (2011). On-tissue identification of insulin: *in situ* reduction coupled with mass spectrometry imaging. *Proteomics Clin Appl* 5, 448–453.
- Grey, A.C., Chaurand, P., Caprioli, R.M., and Schey, K.L. (2009). MALDI imaging mass spectrometry of integral membrane proteins from ocular lens and retinal tissue. *J Proteome Res* 8, 3278–3283.
- Groeneveld, G., de Puit, M., Bleay, S., Bradshaw, R., and Francese, S. (2015). Detection and mapping of illicit drugs and their metabolites in fingermarks by MALDI MS and compatibility with forensic techniques. *Sci Rep* 5, 11716.
- Groseclose, M.R., and Castellino, S. (2013). A mimetic tissue model for the quantification of drug distributions by MALDI imaging mass spectrometry. *Anal Chem* 85, 10099–10106.
- Groseclose, M.R., Andersson, M., Hardesty, W.M., and Caprioli, R.M. (2007). Identification of proteins directly from tissue: *in situ* tryptic digestions coupled with imaging mass spectrometry. *J Mass Spectrom* 42, 254–262.
- Groseclose, M.R., Massion, P.P., Chaurand, P., and Caprioli, R.M. (2008). High-throughput proteomic analysis of formalin-fixed paraffin-embedded tissue microarrays using MALDI imaging mass spectrometry. *Proteomics* 8, 3715–3724.
- Guilhaus, M., Mlynářík, V., and Selby, D. (1997). Perfect timing: time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* 11, 951–962.
- Gustafsson, O.J.R., Oehler, M.K., Ruszkiewicz, A., McColl, S.R., and Hoffmann, P. (2011). MALDI Imaging Mass Spectrometry (MALDI-IMS) - application of spatial proteomics for ovarian cancer classification and diagnosis. *Int J Molecular Sci* 12, 773–794.
- Gustafsson, O.J.R., Eddes, J.S., Meding, S., McColl, S.R., Oehler, M.K., and Hoffmann, P. (2013). Matrix-assisted laser desorption/ionization imaging protocol for *in situ* characterization of tryptic peptide identity and distribution in formalin-fixed tissue. *Rapid Commun Mass Spectrom* 27, 655–670.
- Gustafsson, O.J.R., Briggs, M.T., Condina, M.R., Winderbaum, L.J., Pelzing, M., McColl, S.R., Everest-Dass, A.V., Packer, N.H., and Hoffmann, P. (2015). MALDI imaging mass spectrometry of N-linked glycans on formalin-fixed paraffin-embedded murine kidney. *Anal Bioanal Chem* 407, 2127–2139.
- Hamm, G., Bonnel, D., Legouffe, R., Pamlard, F., Delbos, J.-M., Bouzom, F., and Stauber, J. (2012). Quantitative mass spectrometry imaging of propranolol and olanzapine using tissue extinction calculation as normalization factor. *J Proteomics* 75, 4952–4961.
- Han, H., Pappin, D., Ross, P., and McLuckey, S. (2008). Electron transfer dissociation of iTRAQ labeled peptide ions. *J Proteome Res* 7, 3643–3648.
- Hankin, J.A., Barkley, R.M., and Murphy, R.C. (2007). Sublimation as a method of matrix application for mass spectrometric imaging. *J Am Soc Mass Spectrom* 18, 1646–1652.
- Hanton, S.D., Cornelio Clark, P.A., and Owens, K.G. (1999). Investigations of matrix-assisted laser desorption/ionization sample preparation by time-of-flight secondary ion mass spectrometry. *J Am Soc Mass Spectrom* 10, 104–111.
- Hattori, K., Kajimura, M., Hishiki, T., Nakanishi, T., Kubo, A., Nagahata, Y., Ohmura, M., Yachie-Kinoshita, A., Matsura, T., Morikawa, T., et al. (2010). Paradoxical ATP elevation in ischemic penumbra revealed by quantitative imaging mass spectrometry. *Antioxid Redox Signal* 13, 1157–1167.
- Heij, B., Carreira, R.J., Tolner, E.A., de Ru, A.H., Maagdenberg, A.M., van Veelen, P.A., and McDonnell, L.A. (2015). Comprehensive analysis of the mouse brain proteome sampled in mass spectrometry imaging. *Anal Chem* 87, 1867–1875.
- Heij, B., Holst, S., Briaire-de Bruijn, I.H., Pelt, G., de Ru, A.H., van Veelen, P.A., Drake, R.R., Mehta, A.S., Mesker, W.E., Tollenaar, R.A., et al. (2016). Multimodal mass spectrometry imaging of N-glycans and proteins from the same tissue section. *Anal Chem* 88, 7745–7753.
- Himmelsbach, M. (2012). 10 years of MS instrumental developments – impact on LC–MS/MS in clinical chemistry. *J Chromatogr B Analyt Technol Biomed Life Sci* 883–884, 3–17.

- Hoffman, A., Ghadimi, M.P.H., Demicco, E.G., Creighton, C.J., Torres, K., Colombo, C., Peng, T., Lusby, K., Ingram, D., Hornick, J.L., et al. (2013) Localized and metastatic myxoid/round cell liposarcoma: clinical and molecular observations. *Cancer* 119, 1868–1877.
- Hohmann, L., Sherwood, C., Eastham, A., Peterson, A., Eng, J.K., Eddes, J.S., Shteynberg, D., and Martin, D.B. (2009). Proteomic analyses using Grifola frondosa metalloendoprotease Lys-N. *J Proteome Res* 8, 1415–1422.
- Holle, A., Haase, A., Kayser, M., and Höhndorf, J. (2006). Optimizing UV laser focus profiles for improved MALDI performance. *J Mass Spectrom* 41, 705–716.
- Holst, S., Wührer, M., and Rombouts, Y. (2015). Advances in Cancer Research.
- Holst, S., Heijis, B., de Haan, N., van Zeijl, R.J.M., Braire-de Brujin, I.H., van Pelt, G.W., Mehta, A.S., Angel, P.M., Mesker, W.E., Tollenaar, R.A., et al. (2016). Linkage-specific *in situ* sialic acid derivatization for N-glycan mass spectrometry imaging of formalin-fixed paraffin-embedded tissues. *Anal Chem* 88, 5904–5913.
- Houel, S., Abernathy, R., Renganathan, K., Meyer-Arendt, K., Ahn, N.G., and Old, W.M. (2010). Quantifying the impact of chimera MS/MS spectra on peptide identification in large-scale proteomics studies. *J Proteome Res* 9, 4152–4160.
- Hurvitz, S.A., Hu, Y., O'Brien, N., and Finn, R.S. (2013). Current approaches and future directions in the treatment of HER2-positive breast cancer. *Cancer Treat Rev* 39, 219–229.
- Ibáñez, A.J., Muck, A., and Svatoš, A. (2007). Dissipation of charge on MALDI-TOF polymeric chips using an electron-acceptor: analysis of proteins. *J Mass Spectrom* 42, 634–640.
- Itkonen, H., and Mills, I. (2015). Receptor Tyrosine Kinases (Springer).
- Jansen, B.C., Reiding, K.R., Bondt, A., Ederveen, A.L., Palmblad, M., Falck, D., and Wührer, M. (2015). MassyTools: a high-throughput targeted data processing tool for relative quantitation and quality control developed for glycomic and glycoproteomic MALDI-MS. *J Proteome Res* 14, 5088–5098.
- Jennings, K.R. (1968). Collision-induced decompositions of aromatic molecular ions. *Int J Mass Spectrom Ion Phys* 1, 227–235.
- Jensen, L.J., Kuhn, M., Stark, M., Chaffron, S., Creevey, C., Muller, J., Doerks, T., Julien, P., Roth, A., Simonovic, M., et al. (2009). STRING 8 - a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res* 37, D412–D416.
- Johnson, R.S., Martin, S.A., Biemann, K., Stults, J.T., Watson, J.T. (1987). Novel fragmentation process of peptides by collision-induced decomposition in a tandem mass spectrometer: differentiation of leucine and isoleucine. *Anal Chem* 56, 2621–2625.
- Jones, E.A., van Remoortere, A., van Zeijl, R.J.M., Hogendoorn, P.C., Bovée, J.V., Deelder, A.M., and McDonnell, L.A. (2011). Multiple statistical analysis techniques corroborate intratumor heterogeneity in imaging mass spectrometry datasets of myxofibrosarcoma. *PLoS ONE* 6, e24913.
- Jones, E.A., Schmitz, N., Waaijer, C.J., Frese, C.K., van Remoortere, A., van Zeijl, R.J.J., Heck, A.J., Hogendoorn, P.C., Deelder, A.M.M., Altehaar, A.F., et al. (2013). Imaging mass spectrometry-based molecular histology differentiates microscopically identical and heterogeneous tumors. *J Proteome Res.* 12, 1847–1855.
- Kaftan, F., Vrkoslav, V., Kynast, P., Kulkarni, P., Böcker, S., Cvačka, J., Knaden, M., and Svatoš, A. (2014). Mass spectrometry imaging of surface lipids on intact *Drosophila melanogaster* flies. *J Mass Spectrom* 49, 223–232.
- Källback, P., Shariatgorji, M., Nilsson, A., and Andrén, P.E. (2012). Novel mass spectrometry imaging software assisting labeled normalization and quantitation of drugs and neuropeptides directly in tissue sections. *J Proteomics* 75, 4941–4951.
- Kaprio, T., Satomaa, T., Heiskanen, A., Hokke, C.H., Deelder, A.M., Mustonen, H., Hagström, J., Carpen, O., Saarinen, J., and Haglund, C. (2015). N-glycomics profiling as a tool to separate rectal adenomas from carcinomas. *Mol Cell Proteomics* 14, 277–288.
- Karas, M., and Hillenkamp, F. (1988). Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *Anal Chem* 60, 2299–2301.
- Karas, M., Bachmann, D., Bahr, U., and Hillenkamp, F. (1987). Matrix-assisted ultraviolet laser desorption of non-volatile compounds. *Int J Mass Spectrom Ion Proc* 78, 53–68.
- Kaspar, S., Peukert, M., Svatos, A., Matros, A., and Mock, H.P. (2011). MALDI-imaging mass spectrometry - an emerging technique in plant biology. *Proteomics* 11, 1840–1850.
- Kaufman, M., and Bard, J. (1999). The anatomical basis of mouse development. 1st ed. San Diego: Academic Press.
- Kawamoto, T. (2003). Use of a new adhesive film for the preparation of multi-purpose fresh-frozen sections from hard tissues, whole-animals, insects and plants. *Arch Histol Cytol* 66, 123–143.
- Kero, F.A., Pedder, R.E., and Yost, R.A. (2005). Quadrupole mass analyzers: theoretical and practical considerations. *Encyclopedia of Genetics, Genomics, Proteomics and Bioinformatics*. 3:3.1:8.
- Kertesz, V., and Berkel, G.J. (2010). Fully automated liquid extraction-based surface sampling and ionization using a chip-based robotic nanoelectrospray platform. *J Mass Spectrom* 45, 252–260.
- Keuschning, J., Henttinen, T., Auvinen, K., Karikoski, M., Salmi, M., and Jalkanen, S. (2009). The prototype endothelial marker PAL-E is a leukocyte trafficking molecule. *Blood* 114, 478–484.
- Khalil, S.M., Römpf, A., Pretzel, J., Becker, K., and Spengler, B. (2015). Phospholipid topography of whole-body sections of the *Anopheles stephensi* mosquito, characterized by high-resolution atmospheric-pressure scanning microprobe matrix-assisted laser desorption/ionization mass spectrometry imaging. *Anal Chem* 87, 11309–11316.
- Klein, S., Staring, M., Murphy, K., Viergever, M.A., and Pluim, J.P. (2010). Elastix: a toolbox for intensity-based medical image registration. *IEEE Trans Med Imaging* 29, 196–205.
- Klein, S., Staring, M., Andersson, P., and Pluim, J.P. (2011). Preconditioned stochastic gradient descent optimisation for monomodal image registration. *Med Image Comput Comput Assist Interv* 14, 549–556.

- Knochenmuss, R. (2004). Photoionization pathways and free electrons in UV-MALDI. *Anal Chem* 76, 3179–3184.
- Krause, E., Wenschuh, H., and Jungblut P.R. (1999). The dominance of arginine-containing peptides in MALDI-derived tryptic mass fingerprints of proteins. *Anal Chem* 71, 4160–4165.
- Kriegsmann, M., Casadonte, R., Kriegsmann, J., Dienemann, H., Schirmacher, P., Hendrik Kobarg, J., Schwamborn, K., Stenzinger, A., Warth, A., and Weichert, W. (2016). Reliable entity subtyping in non-small cell lung cancer by matrix-assisted laser desorption/ionization imaging mass spectrometry on formalin-fixed paraffin-embedded tissue specimens. *Mol Cell Proteomics* 15, 3081–3089.
- Krusemark, C.J., Frey, B.L., Belshaw, P.J., and Smith, L.M. (2009). Modifying the charge state distribution of proteins in electrospray ionization mass spectrometry by chemical derivatization. *J Am Soc Mass Spectrom* 20, 1617–1625.
- Kulak, N.A., Pichler, G., Paron, I., Nagaraj, N., and Mann, M. (2014). Minimal, encapsulated proteomic-sample processing applied to copy-number estimation in eukaryotic cells. *Nat Methods* 11, 319–324.
- Lanekoff, I., Burnum-Johnson K., Thomas, M., Short, J., Carson, J.P., Cha, J., Dey, S.K., Yang, P., Prieto Conaway, M.C., Laskin, J. (2013). High-speed tandem mass spectrometric *in situ* imaging by nanospray desorption electrospray ionization mass spectrometry. *Anal Chem* 85, 9596–9603.
- Ledford, E.B. Jr, Rempel, D.L., and Gross M.L. (1984). Space charge effects in Fourier transform mass spectrometry. Mass calibration. *Anal Chem* 56, 2744–2748.
- Lee, J.Y., Kim, J.Y., Park, G.W., Cheon, M.H., Kwon, K.-H., Ahn, Y.H., Moon, M.H., Lee, H.J., Paik, Y.K., and Yoo, J.S. (2011). Targeted mass spectrometric approach for biomarker discovery and validation with nonglycosylated tryptic peptides from N-linked glycoproteins in human plasma. *Mol Cell Proteomics* 10, M111.009290.
- Lemaire, R., Wisztorski, M., Desmons, A., Tabet, J.C., Day, R., Salzet, M., and Fournier, I. (2006). MALDI-MS direct tissue analysis of proteins: improving signal sensitivity using organic treatments. *Anal Chem* 78, 7145–7153.
- Lemaire, R., Desmons, A., Tabet, J.C., Day, R., Salzet, M., and Fournier, I. (2007). Direct analysis and MALDI imaging of formalin-fixed, paraffin-embedded tissue sections. *J Proteome Res* 6, 1295–1305.
- Lemeur, M., Mattei, J.-C., Souteyrand, P., Chagnaud, C., Curvale, G., and Rochwerger, A. (2015). Prognostic factors for the recurrence of myxoid liposarcoma: 20 cases with up to 8 years follow-up. *Orthop Traumatol Surg Res* 101, 103–107.
- Lin, T.-W., Chang, H.-T., Chen, C.-H., Chen, C.-H., Lin, S.-W., Hsu, T.-L., and Wong, C.-H. (2015). Galectin-3 binding protein and galectin-1 interaction in breast cancer cell aggregation and metastasis. *J Am Chem Soc* 137, 9685–9693.
- Liu, D.Q., and Hop, C.E. (2005). Strategies for characterization of drug metabolites using liquid chromatography-tandem mass spectrometry in conjunction with chemical derivatization and on-line H/D exchange approaches. *J Pharm Biomed Anal* 37, 1–18.
- Liu, X., Idé, J.L., Norton, I., Marchionni, M.A., Ebling, M.C., Wang, L.Y., Davis, E., Sauvageot, C.M., Kesari, S., Kellersberger, K.A., et al. (2013a). Molecular imaging of drug transit through the blood-brain barrier with MALDI mass spectrometry imaging. *Sci Rep* 3, 2859.
- Liu, X., Nie, H., Zhang, Y., Yao, Y., Maitikabili, A., Qu, Y., and Shi, S (2013b). Cell surface-specific N-glycan profiling in breast cancer. *PLOS One* 8, e72704.
- Lou, S., Balluff, B., de Graaff, M.A., Cleven, A.H., Briare-de Brujin, I., Bovée, J.V., and McDonnell, L.A. (2016a). High-grade sarcoma diagnosis and prognosis: biomarker discovery by mass spectrometry imaging. *Proteomics* 16, 1802–1813.
- Lou, S., Balluff, B., Cleven, A.H., Bovée, J.V., and McDonnell, L.A. (2016b). An experimental guideline for the analysis of histologically heterogeneous tumors by MALDI-TOF mass spectrometry imaging. *Biochim Biophys Acta* S1570-9639, 30205–30209.
- Lou, S., Balluff, B., Cleven, A.H., Bovée, J.V., and McDonnell, L.A. (2017). Prognostic Metabolite Biomarkers for Soft Tissue Sarcomas Discovered by Mass Spectrometry Imaging. *J. Am. Soc. Mass Spectrom.* 28, 376–383.
- Maes, F., Collignon, A., Vandermeulen, D., Marchal, G., and Suetens, P. (1997). Multimodality image registration by maximization of mutual information. *IEEE Trans Med Imaging* 16, 187–198.
- Maier, S.K., Hahne, H., Gholami, A.M., Balluff, B., Meding, S., Schoene, C., Walch, A.K., and Kuster, B. (2013). Comprehensive identification of proteins from MALDI imaging. *Mol Cell Proteomics* 12, 2901–2910.
- Makarov, A. (2000). Electrostatic axially harmonic orbital trapping: a high-performance technique of mass analysis. *Anal Chem* 72, 1156–1162.
- Maleki, H., Kondalaji, S.G., Khakinejad, M., and Valentine, S.J. (2016). Structural assignments of sulfur-containing compounds in crude oil using ion mobility spectrometry-mass spectrometry (IMS-MS). *Energy Fuels* 30, 9150–9161.
- Mamyrin, B.A., Karataev, V.I., Shmikk, D.V., and Zagulin, V.A. (1973). The mass-reflectron, a new nonmagnetic time-of-flight mass spectrometer with high resolution. *Zh Eksp Teor Fiz* 64, 82–89.
- Markesberry, W.R. (1997). Oxidative stress hypothesis in Alzheimer's disease. *Free Radic Biol Med* 23, 134–147.
- Marshall, A.G., Hendrickson, C.L., and Jackson, G.S. (1998). Fourier transform ion cyclotron resonance mass spectrometry: A primer. *Mass Spectrom Rev* 17, 1–35.
- Marshall, A.G. (1998). Fourier transform ion cyclotron resonance mass spectrometry. *AIP Conf Proc* 430, 3.
- Martin-Lorenzo, M., Alvarez-Llamas, G., McDonnell, L.A., and Vivanco, F. (2015). Molecular histology of arteries: mass spectrometry imaging as a novel ex vivo tool to investigate atherosclerosis. *Expert Rev Proteomics* 13, 69–81.
- McDonnell, L.A., and Heeren, R.M. (2007). Imaging mass spectrometry. *Mass Spectrom Rev* 26, 606–643.
- McDonnell, L.A., Walch, A.K., Stoeckli, M., and Corthals, G.L. (2014). MSiMass list: a public database of identifications for protein MALDI MS imaging. *J Proteome Res* 13, 1138–1142.

- McLafferty, F.W., Bente, P.F., Kornfeld, R., Tsai, S.-C., and Howe, I. (1973). Metastable ion characteristics. XXII. Collisional activation spectra of organic ions. *J Am Chem Soc* 95, 2120–2129.
- McLean, J.A., Ridenour, W.B., and Caprioli, R.M. (2007). Profiling and imaging of tissues by imaging ion mobility-mass spectrometry. *J Mass Spectrom* 42, 1099–1105.
- Meding, S., Nitsche, U., Balluff, B., Elsner, M., Rauser, S., Schöne, C., Nipp, M., Maak, M., Feith, M., Ebert, M., et al. (2012). Tumor classification of six common cancer types based on proteomic profiling by MALDI imaging. *J Proteome Res* 11, 1996–2003.
- Meding, S., Martin, K., Gustafsson, O.J.R., Eddes, J.S., Hack, S., Oehler, M.K., and Hoffmann, P. (2013). Tryptic peptide reference data sets for MALDI imaging mass spectrometry on formalin-fixed ovarian cancer tissues. *J Proteome Res* 12, 308–315.
- Melikian, A.A., O'Connor, R., Prahalad, A.K., Hu, P., Li, H., Kagan, M., and Thompson, S. (1999). Determination of the urinary benzene metabolites S-phenylmercapturic acid and trans,trans-muconic acid by liquid chromatography-tandem mass spectrometry. *Carcinogenesis* 20, 719–726.
- Min, Y., Kristiansen, K., and Boggs, J.M., Husted, C., Zasadzinski, J.A., Israelachvili, J. (2009). Interaction forces and adhesion of supported myelin lipid bilayers modulated by myelin basic protein. *Proc Natl Acad Sci USA* 106, 3154–3159.
- Miyoshi, E., and Nakano, M. (2008). Fucosylated haptoglobin is a novel marker for pancreatic cancer: detailed analyses of oligosaccharide structures. *Proteomics* 8, 3257–3262.
- Morth, J.P., Poulsen, H., Toustrup-Jensen, M.S., Schack, V.R., Egebjerg, J., Andersen, J.P., Vilse, B., and Nissen, P. (2009). The structure of the Na⁺,K⁺-ATPase and mapping of isoform differences and disease-related mutations. *Philos Trans R Soc Lond B Biol Sci* 364, 217–227.
- Mótyán, J.A., Tóth, F., and Tőzsér, J. (2013). Research applications of proteolytic enzymes in molecular biology. *Biomolecules* 3, 923–942.
- Muinelo-Romay, L., Vázquez-Martín, C., Villar-Portela, S., Cuevas, E., Gil-Martín, E., and Fernández-Briera, A. (2008). Expression and enzyme activity of alpha(1,6)fucosyltransferase in human colorectal cancer. *Int J Cancer* 123, 641–646.
- Mulder, I.A., Esteve, C., Wermer, M.J., Hoehn, M., Tolner, E.A., van den Maagdenberg, A.M.J.M., and McDonnell, L.A. (2016). Funnel-freezing versus heat-stabilization for the visualization of metabolites by mass spectrometry imaging in a mouse stroke model. *Proteomics* 16, 1652–1659.
- Muller, L., Kailas, A., Jackson, S.N., Roux, A., Barbacci, D.C., Schultz, J.A., Balaban, C.D., and Woods, A.S. (2015). Lipid imaging within the normal rat kidney using silver nanoparticles by matrix-assisted laser desorption/ionization mass spectrometry. *Kidney Int* 88, 186–192.
- Negri, T., Virdis, E., Brich, S., Bozzi, F., Tamborini, E., Tarantino, E., Jocolè, G., Cassinelli, G., Grossi, F., Sanfilippo, R., et al. (2010). Functional mapping of receptor tyrosine kinases in myxoid liposarcoma. *Clin Cancer Res* 16, 3581–3593.
- Nemes, P., and Vertes, A. (2007). Laser ablation electrospray ionization for atmospheric pressure, *in vivo*, and imaging mass spectrometry. *Anal Chem* 79, 8098–8106.
- Nezu, Y., Hagiwara, K., Yamamoto, Y., Fujiwara, T., Matsuo, K., Yoshida, A., Kawai, A., Saito, T., and Ochiya, T. (2016). miR-135b, a key regulator of malignancy, is linked to poor prognosis in human myxoid liposarcoma. *Oncogene* 35, 6177–6188.
- Nicklay, J.J., Harris, G.A., Schey, K.L., and Caprioli, R.M. (2013). MALDI imaging and *in situ* identification of integral membrane proteins from rat brain tissue sections. *Anal Chem* 85, 7191–7196.
- Nikolaev, E.N., Kostyukevich, Y.I., and Vladimirov, G.N. (2016). Fourier transform ion cyclotron resonance (FT ICR) mass spectrometry: theory and simulations. *Mass Spectrom Rev* 35, 219–258.
- Nishida, Y., Tsukushi, S., Nakashima, H., and Ishiguro, N. (2010). Clinicopathologic prognostic factors of pure myxoid liposarcoma of the extremities and trunk wall. *Clin Orthop Relat Res* 468, 3041–3046.
- Nonami, H., Fukui, S., and Erra-Balsells, R. (1997). β-carboline alkaloids as matrices for matrix-assisted ultraviolet laser desorption time-of-flight mass spectrometry of proteins and sulfated oligosaccharides: a comparative study using phenylcarbonyl compounds, carbazoles and classical matrices. *J Mass Spectrom* 32, 287–296.
- Nordström, A., Want, E., Northen, T., Lehtio, J., and Siuzdak, S. (2008). Multiple ionization mass spectrometry strategy used to reveal the complexity of metabolomics. *Anal Chem* 80, 421–429.
- Oviaño, M., Sparbier, K., Barba, M.J., Kostrzewa, M., and Bou, G. (2016). Universal protocol for the rapid automated detection of carbapenem-resistant Gram-negative bacilli directly from blood cultures by matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF/MS). *Int J of Antimicrob Agents* 48, 655–660.
- Park, S.-Y., Lee, S.-H., Kawasaki, N., Itoh, S., Kang, K., Hee Ryu, S., Hashii, N., Kim, J.-M., Kim, J.-Y., and Hoe Kim, J. (2011). α1-3/4 fucosylation at Asn 241 of β-haptoglobin is a novel marker for colon cancer: a combinatorial approach for development of glycan biomarkers. *Int J Cancer* 130, 2366–2376.
- Pecoraro, V., Roli, L., Plebani, M., and Trenti, T. (2016). Clinical utility of the (-2) proPSA and evaluation of the evidence: a systematic review. *Clin Chem Lab Med* 54, 1123–1132.
- Peukert, M., Matros, A., Lattanzio, G., Kaspar, S., Abadía, J., and Mock, H.P. (2012). Spatially resolved analysis of small molecules by matrix-assisted laser desorption/ionization mass spectrometric imaging (MALDI-MSI). *New Phytol* 193, 806–815.
- Pinho, S.S., and Reis, C.A. (2015). Glycosylation in cancer: mechanisms and clinical implications. *Nat Rev Cancer* 15, 540–555.
- Pluim, J.P., Maintz, J.B., and Viergever, M.A. (2003). Mutual-information-based registration of medical images: a survey. *IEEE Trans Med Imaging* 22, 986–1004.

- Porta, T., Lesur, A., Varesio, E., and Hopfgartner, G. (2015). Quantification in MALDI-MS imaging: what can we learn from MALDI-selected reaction monitoring and what can we expect for imaging? *Anal Bioanal Chem* 407, 2177–2187.
- Porter, A., Irwin, R., Miller, J., Horan, D.J., Robling, A.G., and McCabe, L.R. (2017). Quick and inexpensive paraffin-embedding method for dynamic bone formation analyses. *Sci Rep* 7, 42505.
- Poté, N., Alexandrov, T., Faouder, J., Laouirem, S., Léger, T., Mebarki, M., Belghiti, J., Camadro, J.-M., Bedossa, P., and Paradis, V. (2013). Imaging mass spectrometry reveals modified forms of histone H4 as new biomarkers of microvascular invasion in hepatocellular carcinomas. *Hepatology* 58, 983–994.
- Poulsen, H., Khandelia, H., Morth, J.P., Bublitz, M., Mouritsen, O.G., Egebjerg, J., and Nissen, P. (2010). Neurological disease mutations compromise a C-terminal ion pathway in the Na(+)/K(+) -ATPase. *Nature* 467, 99–102.
- Powers, M.P., Wang, W.-L.L., Hernandez, V.S., Patel, K.S., Lev, D.C., Lazar, A.J., and López-Terrada, D.H. (2010). Detection of myxoid liposarcoma-associated *FUS-DDIT3* rearrangement variants including a newly identified breakpoint using an optimized RT-PCR assay. *Mod Pathol* 23, 1307–1315.
- Powers, T.W., Jones, E.E., Betesh, L.R., Romano, P.R., Gao, P., Copland, J.A., Mehta, A.S., and Drake, R.R. (2013). Matrix assisted laser desorption ionization imaging mass spectrometry workflow for spatial profiling analysis of N-linked glycan expression in tissues. *Anal Chem* 85, 9799–9806.
- Powers, T.W., Neely, B.A., Shao, Y., Tang, H., Troyer, D.A., Mehta, A.S., Haab, B.B., and Drake, R.R. (2014). MALDI imaging mass spectrometry profiling of N-glycans in formalin-fixed paraffin embedded clinical tissue blocks and tissue microarrays. *PLoS ONE* 9, e106255.
- Powers, T.W., Holst, S., Wuhrer, M., Mehta, A.S., and Drake, R.R. (2015). Two-dimensional N-glycan distribution mapping of hepatocellular carcinoma tissues by MALDI-imaging mass spectrometry. *Biomolecules* 5, 2554–2572.
- Prideaux, B., and Stoeckli, M. (2012). Mass spectrometry imaging for drug distribution studies. *J Proteomics* 75, 4999–5013.
- Proskuryakov, S.Y., Konoplyannikov, A.G., and Gabai, V.L. (2003). Necrosis: a specific form of programmed cell death? *Exp Cell Res* 283, 1–16.
- Puolitaival, S.M., Burnum, K.E., Cornett, D.S., and Caprioli, R.M. (2008). Solvent-free matrix dry-coating for MALDI imaging of phospholipids. *J Am Soc Mass Spectrom* 19, 882–886.
- Qi, Y., and Volmer, D.A. (2017). Electron-based fragmentation methods in mass spectrometry: an overview. *Mass Spectrom Rev* 36, 4–15.
- Qiao, H., Spicer, V., and Ens, W. (2008). The effect of laser profile, fluence, and spot size on sensitivity in orthogonal-injection matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* 22, 2779–2790.
- Radionova, A., Filippov, I., and Derrick, P.J. (2016). In pursuit of resolution in time-of-flight mass spectrometry: a historical perspective. *Mass Spectrom Rev* 35, 738–757.
- Rao, W., Celiz, A.D., Scurr, D.J., Alexander, M.R., and Barrett, D.A. (2013). Ambient DESI and LESA-MS analysis of proteins adsorbed to a biomaterial surface using in-situ surface tryptic digestion. *J Am Soc Mass Spectrom* 24, 1927–1936.
- Reilly, J.P. (2009). Ultraviolet photofragmentation of biomolecular ions. *Mass Spectrom Rev* 28, 425–447.
- Reyzer, M.L., Hsieh, Y., Ng, K., Korfomacher, W.A., and Caprioli, R.M. (2003). Direct analysis of drug candidates in tissue by matrix-assisted laser desorption/ionization mass spectrometry. *J Mass Spectrom* 38, 1081–1092.
- Rietschel, B., Arrey, T.N., Meyer, B., Bornemann, S., Schuerken, M., Karas, M., and Poetsch, A. (2008). Elastase digests: new ammunition for shotgun membrane proteomics. *Mol Cell Proteomics* 8, 1029–1043.
- Riley, N.M., Rush, M.J., Rose, C.M., Richards, A.L., Kwiecien, N.W., Bailey, D.J., Hebert, A.S., Westphall, M.S., and Coon, J.J. (2015). The negative mode proteome with activated ion negative electron transfer dissociation (AI-NETD). *Mol Cell Proteom* 14, 2644–2660.
- Roepstorff, P., Fohlman, J. (1984). Proposal for a common nomenclature for sequence ions in mass spectra of peptides. *Biomed Mass Spectrom* 11, 601.
- Ronci, M., Bonanno, E., Colantoni, A., Pieroni, L., Ilio, C., Spagnoli, L.G., Federici, G., and Urbani, A. (2008). Protein unlocking procedures of formalin-fixed paraffin-embedded tissues: application to MALDI-TOF imaging MS investigations. *Proteomics* 8, 3702–3714.
- Schober, Y., Schramm, T., Spengler, B., and Römpf, A. (2011). Protein identification by accurate mass matrix-assisted laser desorption/ionization imaging of tryptic peptides. *Rapid Commun Mass Spectrom* 25, 2475–2483.
- Schober, Y., Guenther, S., Spengler, B., and Römpf, A. (2012). High-resolution matrix-assisted laser desorption/ionization imaging of tryptic peptides from tissue. *Rapid Commun Mass Spectrom* 26, 1141–1146.
- Schwartz, S.A., Reyzer, M.L., and Caprioli, R.M. (2003). Direct tissue analysis using matrix-assisted laser desorption/ionization mass spectrometry: practical aspects of sample preparation. *J Mass Spectrom* 38, 699–708.
- Scigelova, M., Hornshaw, M., Giannakopoulos, A., and Makarov, A. (2011). Fourier transform mass spectrometry. *Mol Cell Proteomics* 10, M111.009431.
- Selman, M.H., Hemayatkar, M., Deelder, A.M., and Wuhrer, M. (2011). Cotton HILIC SPE microtips for microscale purification and enrichment of glycans and glycopeptides. *Anal Chem* 83, 2492–2499.
- Shariatgorji, M., Källback, P., Gustavsson, L., Schintu, N., Svenssonsson, P., Goodwin, R.J., and Andren, P.E. (2012). Controlled-pH tissue cleanup protocol for signal enhancement of small molecule drugs analyzed by MALDI-MS imaging. *Anal Chem* 84, 4603–4607.
- Shariatgorji, M., Nilsson, A., Goodwin, R.J., Källback, P., Schintu, N., Zhang, X., Crossman, A.R., Bezard, E., Svenssonsson, P., Andren, P.E. (2014). Direct targeted quantitative molecular imaging of neurotransmitters in brain tissue sections. *Neuron* 84, 697–707.

- Shi, S.R., Taylor, C.R., Fowler, C.B., and Mason, J.T. (2013). Complete solubilization of formalin-fixed, paraffin-embedded tissue may improve proteomic studies. *Proteomics Clin Appl* 7, 264–272.
- Shi, S.R., Key, M.E., and Kalra, K.L. (1991). Antigen retrieval in formalin-fixed, paraffin-embedded tissues: an enhancement method for immunohistochemical staining based on microwave oven heating of tissue sections. *J Histochem Cytochem* 39, 741–748.
- Shimma, S., Furuta, M., Ichimura, K., Yoshida, Y., and Setou, M. (2006). Direct MS/MS analysis in mammalian tissue sections using MALDI-QIT-TOFMS and chemical inkjet technology. *Surf Interface Anal* 38, 1712–1714.
- Shimma, S., Sugiura, Y., Hayasaka, T., Zaima, N., Matsumoto, M., Setou, M. (2008). Mass imaging and identification of biomolecules with MALDI-QIT-TOF-based system. *Anal Chem* 80, 878–885.
- Shroff, R., and Svatoš, A. (2009). Proton sponge: a novel and versatile MALDI matrix for the analysis of metabolites using mass spectrometry. *Anal Chem* 81, 7954–7959.
- Siegel, G.J., Agranoff, B.W., Albers, R.W., Fisher, S.K., and Uhler, M.D. (1999). Basic neurochemistry. Philadelphia: Lippincott-Raven.
- Signor, L., and Boeri Erba, E. (2013). Matrix-assisted laser desorption/ionization time of flight (MALDI-TOF) mass spectrometric analysis of intact proteins larger than 100 kDa. *J Vis Exp* 79.
- Singh, R., Harshman, S., Ruppert, A.S., Mortazavi, A., Lucas, D.M., Thomas-Ahner, J.M., Clinton, S.K., Byrd, J.C., Freitas, M.A., and Parthun, M.R. (2015). Proteomic profiling identifies specific histone species associated with leukemic and cancer cells. *Clin Proteomics* 12, 22.
- De Sio, G., Smith, A.J., Galli, M., Garancini, M., Chinello, C., Bono, F., Pagni, F., and Magni, F. (2015). A MALDI-mass spectrometry imaging method applicable to different formalin-fixed paraffin-embedded human tissues. *Mol Biosyst* 11, 1507–1514.
- Slodzian, G., Daigne, B., Girard, F., Boust, F., and Hillion, F. (1992). Scanning secondary ion analytical microscopy with parallel detection. *Biol Cell* 74, 43–50.
- Smargiasso, N., Quinton, L., and De Pauw, E. (2012). 2-Aminobenzamide and 2-aminobenzoic acid as new MALDI matrices inducing radical mediated in-source decay of peptides and proteins. *J Am Soc Mass Spectrom* 23, 469–474.
- Smith, D.F., Klein, G.C., Yen, A.T., Squicciarini, M.P., Rodgers, R.P., and Marshall, A.G. (2008). Crude oil polar chemical composition derived from FT-ICR mass spectrometry accounts for asphaltene inhibitor specificity. *Energy Fuels* 22, 3112–3117.
- Spraggins, J.M., Rizzo, D.G., Moore, J.L., Rose, K.L., Hammer, N.D., Skaar, E.P., and Caprioli, R.M. (2015). MALDI FTICR IMS of intact proteins: using mass accuracy to link protein images with proteomics data. *J Am Soc Mass Spectrom* 26, 974–985.
- Stauber, J., MacAleese, L., Franck, J., Claude, E., Snel, M., Kaletas, B.K., Wiel, I.M., Wisztorski, M., Fournier, I., and Heeren, R.M. (2010). On-tissue protein identification and imaging by MALDI-ion mobility mass spectrometry. *J Am Soc Mass Spectrom* 21, 338–347.
- Steven, R.T., and Bunch, J. (2013). Repeat MALDI MS imaging of a single tissue section using multiple matrices and tissue washes. *Anal Bioanal Chem* 405, 4719–4728.
- Stoeckli, M., and Staab, D. (2015). Reproducible matrix deposition for MALDI MSI based on open-source software and hardware. *J Am Soc Mass Spectrom* 26, 911–914.
- Stoeckli, M., Farmer, T.B., Caprioli, R.M. (1999). Automated mass spectrometry imaging with a matrix-assisted laser desorption ionization time-of-flight instrument. *J Am Soc Mass Spectrom* 10, 67–71.
- Stoeckli, M., Chaurand, P., Hallahan, D.E., Caprioli, R.M. (2001) Imaging mass spectrometry: A new technology for the analysis of protein expression in mammalian tissues. *Nat Med* 7, 493–496.
- Stoeckli, M., Staab, D., and Schweitzer, A. (2007). Compound and metabolite distribution measured by MALDI mass spectrometric imaging in whole-body tissue sections. *Int J Mass Spectrom* 260, 195–202.
- Strohalm, M., Hassman, M., Kosata, B., and Kodíček, M. (2008). mMass data miner: an open source alternative for mass spectrometric data analysis. *Rapid Commun Mass Spectrom* 22, 905–908.
- Strohalm, M., Strohalm, J., Kaftan, F., Krásný, L., Volný, M., Novák, P., Ulbrich, K., and Havlíček, V. (2011). Poly[N-(2-hydroxypropyl)methacrylamide]-based tissue-embedding medium compatible with MALDI mass spectrometry imaging experiments. *Anal Chem* 83, 5458–5462.
- Sturm, R.M., Greer, T., Woodards, N., Gemperline, E., and Li, L. (2013). Mass spectrometric evaluation of neuropeptidomic profiles upon heat stabilization treatment of neuroendocrine tissues in crustaceans. *J Proteome Res* 12, 743–752.
- Sturtevant, D., Lee, Y.J., and Chapman, K.D. (2016). Matrix assisted laser desorption/ionization-mass spectrometry imaging (MALDI-MSI) for direct visualization of plant metabolites in situ. *Curr Opin Biotechnol* 37, 53–60.
- Sugiura, Y., Honda, K., Kajimura, M., and Suematsu, M. (2014). Visualization and quantification of cerebral metabolic fluxes of glucose in awake mice. *Proteomics* 14, 829–838.
- Svensson, M., Boren, M., Sköld, K., Fälth, M., Sjögren, B., Andersson, M., Svenssonsson, P., and Andren, P.E. (2009). Heat stabilization of the tissue proteome: a new technology for improved proteomics. *J Proteome Res* 8, 974–981.
- Takáts, Z., Wiseman, J.M., Gologan, B., and Cooks, R.G. (2004). Mass spectrometry sampling under ambient conditions with desorption electrospray ionization. *Science* 306, 471–473.
- Takáts, Z., Wiseman, J.M., and Cooks, R.G. (2005). Ambient mass spectrometry using desorption electrospray ionization (DESI): instrumentation, mechanisms and applications in forensics, chemistry, and biology. *J Mass Spectrom* 40, 1261–1275.
- Tanca, A., Pagnozzi, D., and Addis, M.F. (2012). Setting proteins free: progresses and achievements in proteomics of formalin-fixed, paraffin-embedded tissues. *Proteomics Clin Appl* 6, 7–21.

- Taylor, B.S., Barretina, J., Maki, R.G., Antonescu, C.R., Singer, S., and Ladanyi, M. (2011). Advances in sarcoma genomics and new therapeutic targets. *Nat Rev Cancer* 11, 541–557.
- te Brake, L., Dian, S., Ganiem, A.R., Ruesen, C., Burger, D., Donders, R., Ruslami, R., van Crevel, R., and Aarnoutse, R. (2015). Pharmacokinetic/pharmacodynamic analysis of an intensified regimen containing rifampicin and moxifloxacin for tuberculous meningitis. *Int J Antimicrob Agents* 45, 496–503.
- Thavarajah, R., Mudimbaimannar, V., Elizabeth, J., Rao, U.K., and Ranganathan, K. (2012). Chemical and physical basics of routine formaldehyde fixation. *J Oral Maxillofac Pathol* 16, 400–405.
- Thirunavukarasu, P., Sukumar, S., Sathaiah, M., Mahan, M., Pragatheeshwar, K.D., Pingpank, J.F., Zeh, H., Bartels, C.J., Lee, K.K., and Bartlett, D.L. (2011). C-stage in colon cancer: implications of carcinoembryonic antigen biomarker in staging, prognosis, and management. *J Nat Cancer Inst* 103, 689–697.
- Thomas, A., and Chaurand, P. (2014). Advances in tissue section preparation for MALDI imaging MS. *Bioanalysis* 6, 967–982.
- Thomas, A., Patterson, N.H., Laveaux Charbonneau, J., and Chaurand, P. (2013). Orthogonal organic and aqueous-based washes of tissue sections to enhance protein sensitivity by MALDI imaging mass spectrometry. *J Mass Spectrom* 48, 42–48.
- Thomson, J.J. (1912). Further experiments on positive rays. *Philos Mag (series 6)* 24, 209–253.
- Touboul, D., Halgand, F., Brunelle, A., Kersting, R., Tallarek, E., Hagenhoff, B., and Laprévote, O. (2004). Tissue molecular ion imaging by gold cluster ion bombardment. *Anal Chem*, 76, 1550–1559.
- Toue, S., Sugiura, Y., Kubo, A., Ohmura, M., Karakawa, S., Mizukoshi, T., Yoneda, J., Miyano, H., Noguchi, Y., Kobayashi, T., et al. (2014). Microscopic imaging mass spectrometry assisted by on-tissue chemical derivatization for visualizing multiple amino acids in human colon cancer xenografts. *Proteomics* 14, 810–819.
- Trim, P.J., Henson, C.M., Avery, J.L., McEwen, A., Snel, M.F., Claude, E., Marshall, P.S., West, A., Princivalle, A.P., and Clench, M.R. (2008). Matrix-assisted laser desorption/ionization-ion mobility separation-mass spectrometry imaging of vinblastine in whole body tissue sections. *Anal Chem* 80, 8628–8634.
- Unser, M., Aldroubi, A., and Gerfen, C. (1993). A multiresolution image registration procedure using spline pyramids. *Proc SPIE* 2034, 160–170.
- Valentine, S.J., Anderson, J.G., Ellington, A.D., and Clemmer, D.E. (1997). Disulfide-intact and -reduced lysozyme in the gas phase: conformations and pathways of folding and unfolding. *J Phys Chem B* 101, 3891–3900.
- Van Berkel, G.J., Sanchez, A.D., and Quirke, J.M. (2002). Thin-layer chromatography and electrospray mass spectrometry coupled using a surface sampling probe. *Anal Chem* 74, 6216–6223.
- Van Berkel, G.J., Kertesz, V., Koepflinger, K.A., Vavrek, M., and Kong, A.N. (2008). Liquid microjunction surface sampling probe electrospray mass spectrometry for detection of drugs and metabolites in thin tissue sections. *J Mass Spectrom* 43, 500–508.
- Vandermarkiere, E., Mueller, M., and Martens, L. (2013). Getting intimate with trypsin, the leading protease in proteomics. *Mass Spectrom Rev* 32, 453–465.
- Varki, A., Cummings, R.D., Aebi, M., Packer, N.H., Seeberger, P.H., Esko, J.D., Stanley, P., Hart, G., Darvill, A., Kinoshita, T., et al. (2015). Symbol nomenclature for graphical representations of glycans. *Glycobiology* 25, 1323–1324.
- Vasicek, L.A., Ledvina, A.R., Shaw, J., Griep-Raming, J., Westphall, M.S., Coon, J.J., and Brodbelt, J.S. (2011). Implementing photodissociation in an Orbitrap mass spectrometer. *J Am Soc Mass Spectrom* 22, 1105–1108.
- Végvári, Á., Fehniger, T.E., Gustavsson, L., Nilsson, A., Andrén, P.E., Kenne, K., Nilsson, J., Laurell, T., Marko-Varga, G. (2010). Essential tactics of tissue preparation and matrix nano-spotting for successful compound imaging mass spectrometry. *J Proteomics* 73, 1270–1278.
- Venter, A., Sojka, P.E., and Cooks, R.G. (2006). Droplet dynamics and ionization mechanisms in desorption electrospray ionization mass spectrometry. *Anal Chem* 78, 5849–8555.
- Vestal, M., and Juhasz, P. (1998). Resolution and mass accuracy in matrix-assisted laser desorption ionization-time-of-flight. *J Am Soc Mass Spectrom* 9, 892–911.
- Wa, C., Cerny, R., and Hage, D.S. (2006). Obtaining high sequence coverage in matrix-assisted laser desorption time-of-flight mass spectrometry for studies of protein modification: analysis of human serum albumin as a model. *Anal Biochem* 349, 229–241.
- Wang, Z., Udeshi, N.D., Slawson, C., Compton, P.D., Sakabe, K., Cheung, W.D., Shabanowitz, J., Hunt, D.F., and Hart, G.W. (2010). Extensive crosstalk between O-GlcNAcylation and phosphorylation regulates cytokinesis. *Sci Signal* 3, ra2.
- Wang, T., Goodman, M.A., McGough, R.L., Weiss, K.R., and Rao, U.N. (2014). Immunohistochemical analysis of expressions of RB1, CDK4, HSP90, cPLA2G4A, and CHMP2B is helpful in distinction between myxofibrosarcoma and myxoid liposarcoma. *Int J Surg Pathol* 22, 589–599.
- Wang, T.Y., Jia, Y.L., Zhang, X., Sun, Q.L., Li, Y.-C., Zhang, J.H., Zhao, C.P., Wang, X.Y., and Wang, L. (2015a). Treating colon cancer cells with FK228 reveals a link between histone lysine acetylation and extensive changes in the cellular proteome. *Sci Rep* 5, 18443.
- Wang, S., Xiao, Z., Xiao, C., Wang, H., Wang, B., Li, Y., Chen, X., and Guo, X. (2015b). (E)-Propyl α-Cyano-4-Hydroxy Cinnamylate: A High Sensitive and Salt Tolerant Matrix for Intact Protein Profiling by MALDI Mass Spectrometry. *J Am Soc Mass Spectr* 27, 709–718.
- Watson, G., Paxinos, G., and Puelles, L. (2012). The mouse nervous system. London: Academic.
- Wells, J.M., and McLuckey, S.A. (2005). Collision-induced dissociation (CID) of peptides and proteins. *Methods Enzymol* 402, 148–185.

- Wiley, W.C., and McLaren, I.H. (1955). Time-of-flight mass spectrometer with improved resolution. *Rev Sci Instrum* 26, 1150–1157.
- Willems, S.M., van Remoortere, A., van Zeijl, R.J.M., Deelder, A.M., McDonnell, L.A., and Hogendoorn, P.C.W. (2010a). Imaging mass spectrometry of myxoid sarcomas identifies proteins and lipids specific to tumour type and grade, and reveals biochemical intratumour heterogeneity. *J Pathol* 222, 400–409.
- Willems, S.M., Wiweger, M., Graadt van Roggen, J.F., Hogendoorn, P.C.W. (2010b). Running GAGs: myxoid matrix in tumor pathology revisited. *Virchows Arch* 456, 181–192.
- Wilm, M. (2011). Principles of electrospray ionization. *Mol Cell Proteomics* 10, M111.009407.
- Wolstenholme, R., Bradshaw, R., Clench, M.R., and Francese, S. (2009). Study of latent fingermarks by matrix-assisted laser desorption/ionisation mass spectrometry imaging of endogenous lipids. *Rapid Commun Mass Spectrom* 23, 3031–3039.
- Wu, J.M., Halushka, M.K., Argani, P. (2010) Intratumoral heterogeneity of HER-2 gene amplification and protein overexpression in breast cancer. *Hum Pathol* 41, 914–917.
- Wu, C., Dill, A., Eberlin, L.S., Cooks, R.G., and Ifa, D.R. (2013). Mass spectrometry imaging under ambient conditions. *Mass Spectrom Rev* 32, 218–243.
- Wysocki, V.H., Resing, K.A., Zhang, Q., and Cheng, G. (2005). Mass spectrometry of peptides and proteins. *Methods* 35, 211–222.
- Yamashita, M., and Fenn, J.B. (1984). Electrospray ion source. Another variation on the free-jet theme. *J Phys Chem* 88, 4451–4459.
- Yang, J., and Caprioli, R.M. (2011). Matrix sublimation/recrystallization for imaging proteins by mass spectrometry at high spatial resolution. *Anal Chem* 83, 5728–5734.
- Zagon, I.S., Higbee, R., Riederer, B.M., and Goodman, S.R. (1986). Spectrin subtypes in mammalian brain: an immunoelectron microscopic study. *J Neurosci* 6, 2977–2986.
- Zavalin, A., Todd, E.M., Rawhouser, P.D., Yang, J., Norris, J.L., and Caprioli, R.M. (2012). Direct imaging of single cells and tissue at sub-cellular spatial resolution using transmission geometry MALDI MS. *J Mass Spectrom* 47, i.
- Zenobi, R., and Knochenmuss, R. (1998). Ion formation in MALDI mass spectrometry. *Mass Spectrom Rev* 17, 337–366.
- Zhang, S.-Y., Lin, M., and Zhang, H.-B. (2015). Diagnostic value of carcinoembryonic antigen and carcinoma antigen 19-9 for colorectal carcinoma. *Int J Clin Exp Pathol* 8, 9404–9409.
- Zhao, Y., Takahashi, M., Gu, J., Miyoshi, E., Matsumoto, A., Kitazume, S., and Taniguchi, N. (2008). Functional roles of N-glycans in cell signalling and cell adhesion in cancer. *Cancer Science* 99, 1304–1310.
- Zlokovic, B.V. (2011). Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat Rev Neurosci* 12, 723–738.

APPENDIX

List of abbreviations

9-AA	9-aminoacridine		
ACN	Acetonitrile	LAESI	Laser ablation electrospray ionization
ADC	Adenocarcinoma		Liquid chromatography
APCI	Atmospheric pressure chemical ionization	LC LESA	Liquid extraction surface analysis
CE	Capillary electrophoresis		Lipid-linked oligosaccharide
CEA	Carcinoembryonic antigen	LLO	Liquid microjunction surface sampling
CHCA	α -cyano-4-hydroxycinnamic acid	LMJ-SSP	Leiomyosarcoma
CID	Collision-induced dissociation	LMS	Liposarcoma
CMC	Carboxymethyl cellulose	LPS	Matrix-assisted laser desorption/ionization
COX7A	Cytochrome C oxidase 7A2	MALDI	Microtubule associated protein 6
CRC	Colorectal carcinoma		Myelin basic protein
CSV	Comma-separated values	MAP6	Myxofibrosarcoma
Da	Dalton		Myxoid liposarcoma
DC	Direct current	MBP	Matrix metalloproteinase 2
DESI	Desorption electrospray ionization	MFS	Millimass units
DHB	2,5-dihydroxybenzoic acid	MMP2	Multi reaction monitoring
DSC	Dice similarity coefficient	mmu	Mass spectrometry
DTT	Dithiothreitol	MRM	Mass spectrometry imaging
ECM	Extracellular matrix	MS	Tandem mass spectrometry
EDD	Electron-detachment dissociation	MS/MS	Molecular weight cut off
EDTA	Ethylenediaminetetraacetic acid	MWCO	Myxoid MLS
ER	Endoplasmatic reticulum	MYX	mass-to-charge ratio
ESI	Electrospray ionization	<i>m/z</i>	Negative electron transfer
ETD	Electron-transfer dissociation	NETD	dissociation
FA	Formic acid	NSCLC	Non-small cell lung cancer
FDR	False discovery rate	OCT	Optimal cutting temperature
FFPE	Formalin-fixed paraffin embedded	OS	Osteosarcoma
FID	Free-induction-decay	PC	Phosphatidylcholines
FT	Fourier transform	PEG	Polyethylene glycol
FTICR	Fourier transform ion cyclotron resonance	PEV	Polyethylene vinyl
FTMS	Fourier transform mass spectrometry	pHPMA	Poly[N-(2-hydroxy-propyl)-methacryl-amide]
GnT-V	<i>N</i> -acetylglucosaminyl-transferase V	PIK3CA	Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform
HCD	Higher-energy collisional dissociation	PNGase F	Peptide N-glycosidase F
H&E	Haematoxylin & eosin staining	ppm	Parts-per-million
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid	PTEN	Phosphatase and tensin homolog
HG-MSI	Histology-guided mass spectrometry imaging	qTOF	Quadrupole-time-of-flight
HIAR	Heat-induced antigen retrieval	RC	Round cell MLS
HILIC	Hydrophilic interaction liquid chromatography	RF	Radio frequency
HRAM	High-resolution, accurate mass	ROI	Region of interest
ICR	Ion cyclotron resonance	SA	Sinapic acid
IGF1R	Insulin-like growth factor 1 receptor	SIMS	Secondary ion mass spectrometry
IHC	Immunohistochemistry	S/N	Signal-to-noise
ILRS	Isotopically-labeled reference standard	SPE	Solid phase extraction
IMS	Ion mobility separation	SPTN1	Spectrin-1
INT	Intermediate grade MLS	SqCC	Squamous cell carcinoma
IRMPD	Infrared multi-photon dissociation	STS	Soft tissue sarcoma
ITO	Indium-tin-oxide	TCEP	Tris(2-carboxyethyl)phosphine
		TFA	Trifluoroacetic acid
		TG	Triglyceride lipids
		TMA	Tissue microarray
		TOF	Time-of-flight
		TSP2	Thrombospondin-2
		UPS	Undifferentiated pleomorphic sarcoma