



Universiteit
Leiden
The Netherlands

Disconnected self: influence of dissociation on emotional distractibility in Borderline Personality Disorder: a neuroimaging approach

Krause, A.D.

Citation

Krause, A. D. (2017, November 16). *Disconnected self: influence of dissociation on emotional distractibility in Borderline Personality Disorder: a neuroimaging approach*. Retrieved from <https://hdl.handle.net/1887/60261>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/60261>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/60261> holds various files of this Leiden University dissertation.

Author: Krause, A.D.

Title: Disconnected self: influence of dissociation on emotional distractibility in Borderline Personality Disorder: a neuroimaging approach

Issue Date: 2017-11-16

Disconnected Self

**Influence of Dissociation on Emotional Distractibility
in Borderline Personality Disorder: A Neuroimaging Approach**

Annegret Krause-Utz

Annegret D. Krause-Utz

Disconnected Self:

Influence of Dissociation on Emotional Distractibility in
Borderline Personality Disorder: A Neuroimaging Approach

Cover and chapter illustrations by A. Krause-Utz

Layout and design: A. Krause-Utz

Print: Drukkerij Mostert & Van Onderen, Leiden

© 2017 A. D. Krause-Utz, Leiden, The Netherlands

All rights reserved.

Seltsam im Nebel zu wandern!
Einsam ist jeder Busch und Stein,
Kein Baum sieht den anderen,
Jeder ist allein.
Voll von Freunden war mir die Welt,
Als noch mein Leben licht war;
Nun, da der Nebel fällt,
Ist keiner mehr sichtbar.
Wahrlich, keiner ist weise,
Der nicht das Dunkel kennt,
Das unentrinnbar und leise
Von allem ihn trennt.

Vreemd, te wandelen in de mist!
Eenzaam is elke struik en steen,
Geen boom ziet de andere,
Ieder is alleen.
Vol vrienden was mijn wereld,
Toen mijn leven nog licht was;
Maar nu de mist daalt
Is niemand meer te zien.
Waarlijk, niemand is wijs
Die de duisternis niet kent
Die hem onstuitbaar en zacht
Van iedereen scheidt.

Hermann Hesse
Im Nebel
(vertaling: Jan Gielkens & Ton Naaijken, 1984)

Disconnected Self

**Influence of Dissociation on Emotional Distractibility
in Borderline Personality Disorder: A Neuroimaging Approach**

Proefschrift

ter verkrijging van

de graad van Doctor aan de Universiteit Leiden

op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker

volgens besluit van het College voor Promoties

ter verdediging op donderdag 16 november 2017

klokke 16.15 uur

door

Annegret Dorothea Krause-Utz

geboren te Mutlangen, Duitsland

in 1980

Committee

Promotors:

Prof. dr. B. M. Elzinga (Universiteit Leiden)

Prof. dr. Ph. Spinhoven (Universiteit Leiden)

Co-promotor: Prof. dr. Ch. Schmahl (Central Institute of Mental Health Mannheim)

Manuscriptcommissie:

Prof. dr. A. J. W. van der Does, voorzitter

Prof. dr. E. A. M. Crone

Prof. dr. K. Roelofs (Universiteit Nijmegen)

Prof. dr. D. Veltman (Vrije Universiteit Amsterdam)

Table of Contents

List of Contents	3
List of Tables	8
List of Figures	9
List of Abbreviations	13
Chapter 1: General Introduction	11
1.1. Borderline Personality Disorder	13
1.1.1. Epidemiology and course	13
1.1.2. Pathogenesis	14
1.1.3. Psychopathology	16
1.1.3.1. Emotion dysregulation	16
1.1.3.2. Cognitive disturbances and emotional distractibility	18
1.1.3.3. Dissociation	18
1.2. Functional magnetic resonance imaging	22
1.3. Brain networks relevant to the current thesis	23
1.3.1. Default mode network	23
1.3.2. Salience network	23
1.3.3. Amygdala and medial temporal lobe network	23
1.3.4. Networks implicated in emotional distractibility	24
1.3.5. Networks implicated in dissociation	24
1.4. Thesis outline: Research questions and hypotheses	26
Chapter 2: Neuroimaging findings in Borderline Personality Disorder	29
2.1. Introduction	32
2.2. Structural neuroimaging studies	32
2.3. Functional neuroimaging studies	35
2.3.1. Resting state functional connectivity	36
2.3.2. Emotion processing and emotion regulation	36
2.3.3. Self-Injury and altered pain processing.....	39
2.3.4. Cognitive disturbances and dissociation	40
2.3.5. Behavioral dysregulation and impulsivity.....	42
2.3.6. Interpersonal disturbances.....	44
2.4. Conclusion	46

Chapter 3: Dissociation and Alterations in Brain Function and Structure.....	49
3.1. Introduction	52
3.1.1. Etiological models: Trauma and dissociation.....	53
3.1.2. Neurobiological models	55
3.1.2.1. Cortico-limbic-disconnection model.....	55
3.1.2.3. Research in dissociative identity disorder (DID)	59
3.1.2.4. Research on structural alterations	59
3.1.3. Interim summary	61
3.2. Dissociation in Borderline Personality Disorder	61
3.2.1. Clinical expressions of dissociation in BPD	61
3.2.2. Neuroimaging research on dissociation in BPD	62
3.2.2.1. Brain function during rest: PET, SPECT, and RS-fMRI studies	62
3.2.2.2. Neurochemical alterations: MRS studies	67
3.2.2.3. Task-related fMRI studies.....	67
3.2.2.5. Structural neuroimaging studies in BPD	70
3.2.3. Interim summary	70
3.3. Overall discussion.....	71
Chapter 4: Amygdala and Anterior Cingulate Resting-state Functional Connectivity	
in Borderline Personality Disorder – Associations with Trait Dissociation..	75
4.1. Introduction	78
4.2. Methods and Materials	80
4.2.1. Participants	80
4.2.2. Procedure.....	80
4.2.3. FMRI data acquisition and analysis	82
4.3. Results	84
4.3.1. Amygdala connectivity (medial temporal lobe network).....	84
4.3.2. Dorsal ACC connectivity (salience network).....	85
4.3.3. Ventral ACC connectivity (default mode network)	86
4.3.4. Exploratory analysis: Trait dissociation and functional connectivity ...	86
4.3.5. Effects of global signal regression	87
4.3.6. Subgroup analysis	87
4.4. Discussion.....	87

Chapter 5: Amygdala and dorsal Anterior Cingulate Connectivity during an Emotional Working Memory Task in BPD – The Role of Dissociation	96
5.1. Introduction	98
5.2. Methods	101
5.2.1. Sample	101
5.2.2. Emotional Working Memory Task (EWMT).....	104
5.2.3. Procedure.....	105
5.2.4. Scanning protocol.....	105
5.2.5. Data analysis	106
5.2.5.1. Psychophysiological interaction (PPI) analysis	106
5.2.5.2. Regression analyses.....	108
5.3. Results	108
5.3.1. Amygdala connectivity:	109
5.3.2. Dorsal anterior cingulate (dACC) connectivity:	111
5.3.2.5. Regression analyses.....	112
5.4. Discussion.....	113
Chapter 6: Dissociation in Borderline Personality Disorder: Disturbed cognitive and emotional inhibition and its neural correlates.....	128
6.1. Introduction	130
6.2. Methods	132
6.2.1. Sample	132
6.2.2. Dissociation induction	134
6.2.3. Emotional Stroop Task.....	135
6.2.4. fMRI scan protocol and data analysis	135
6.2.4.1. Statistical analysis of behavioral data.	136
6.2.4.2. Statistical analysis of fMRI data.	136
6.2.4.3. Post-hoc analysis for trauma, depression, anxiety, tension	137
6.3. Results	138
6.3.1. Manipulation check: dissociation induction.....	138
6.3.2. Emotional Stroop Task.....	139
6.3.4. fMRI during the Emotional Stroop Task.....	141
6.3.5. Post-hoc analysis for trauma, depression, anxiety, and tension	145
6.4. Discussion.....	145

Chapter 7: Reduced Amygdala activity and Emotional Distractibility during Dissociative States in Borderline Personality Disorder	154
7.1. Introduction	156
7.2. Methods	158
7.2.1. Sample	158
7.2.2. Emotional Working Memory Task (EWMT).....	161
7.2.3. Procedure.....	161
7.2.4. FMRI scan protocol.....	162
7.2.5. Statistical analysis	162
7.2.5.1. Manipulation check	162
7.2.5.2. Behavioral (WM) data.....	162
7.2.5.3. Fmri data	163
7.2.5.4. Region of interest (ROI) and whole-brain (WB) analysis:	163
7.2.5.5. Psychophysiological interaction analysis (PPI) analysis.....	164
7.3. Results	164
7.3.1. Dissociation induction:	164
7.3.2. Behavioral data.....	164
7.3.2.1. Errors during the EWMT	165
7.3.2.2. Reaction times during the EWMT	165
7.3.3. FMRI data	165
7.3.3.1. ROI analysis	165
7.3.3.2. Whole-Brain analysis	166
7.3.3.3. PPI analysis	168
7.4. Discussion:.....	169
Chapter 8: General discussion.....	175
8.1. Summary.....	177
8.1.1. Previous neuroimaging research in BPD (Chapter 2).....	177
8.1.2. Neurobiological models on dissociation (Chapter 3).....	178
8.1.3. Present neuroimaging studies (Chapters 4 - 7).....	178
8.2. Integration and discussion of present findings	180
8.2.1. Behavioral findings in BPD	180
8.2.2. Neuroimaging findings in BPD.....	182
8.2.3. The role of dissociation in altered brain function in BPD.....	182

8.3. Strength and limitations	186
8.3.1. Sample characteristics	187
8.3.2. Task-characteristics	188
8.3.3. Neuroimaging data analysis techniques	189
8.4. Implications for future research.....	190
8.5. Clinical implications.....	191
8.6. Conclusion	192
References.....	194
Nederlandse samenvatting.....	232
Acknowledgements	237
Curriculum vitae.....	239
List of publications	240

List of Tables

Table 3.1. Studies on links between brain function/structure and dissociation in BPD .	63
Table 4.1. Demographic and clinical variables in healthy controls and BPD patients ..	81
Table 4.2. Resting-state functional connectivity results: Between groups effects	84
Table 4.3. DES associations with amygdala RSFC in BPD	87
Table S4.1. RSFC results without global signal regression	92
Table S4.2. DES associations with amygdala RSFC without global signal regression.....	92
Table 5.1. Demographic and clinical variables in healthy controls and BPD patients ..	103
Table S5.1. Main effects and interaction effect for bilateral amygdala connectivity	119
Table S5.2. 2x2 Full Factorial Model of task-related amygdala connectivity	120
Table S5.3. Group differences in amygdala connectivity during emotional distraction .	121
Table S5.4. Main effects and interaction effects for bilateral dACC connectivity	122
Table S5.5. 2x2 Full Factorial Model of task-related bilateral dACC connectivity	123
Table S5.6. Group differences in dACC connectivity during emotional distraction	124
Table S5.7. Regression: Reaction times as predictor of amygdala connectivity	125
Table S5.8. Regression: State dissociation as predictor of amygdala connectivity	125
Table 6.1. Demographic and clinical variables	133
Table 6.2. Behavioral data in the emotional Stroop task and related memory tasks	139
Table 6.3. Group differences in neural activation in the Emotional Stroop Task	141
Table 6.4. Neural activation in response to emotional vs. neutral words per group	143
Table 6.5. Group differences in neural activation to emotional versus neutral words ..	144
Table S6.1. Length, valence and frequency of the word stimuli	151
Table S6.2. Behavioral data of negative compared to positive stimuli	152
Table S6.3. Neural activation in response to negative vs.. positive words in the EST ...	153
Table 7.1. Demographic variables, dissociation, arousal, clinical characteristics	160
Table 7.2. Brain activity during the EWMT	167
Table S7.1. PPI Analysis: Amygdala connectivity during negative distractors	173
Table 8.1. Methodological characteristics and results of studies in this thesis	179

List of Figures

Figure 4.1. Group differences in resting-state functional connectivity (RSFC).	85
Figure 4.2. Correlations between dissociation scores and amygdala RSFC in BPD	86
Figure S4.1. Group main effects for RSFC with the three seeds	93
Figure S4.2. Group differences in RSFC of the seeds without global signal regression..	94
Figure S4.3. Correlations without global signal regression.	94
Figure S4.4. Subgroup analysis for BPD patients with and without comorbid PTSD	95
Figure 5.1. Design of the Emotional Working Memory Task (EWMT)	105
Figure 5.2. Results for the main effect of valence on amygdala connectivity.	109
Figure 5.3. Group differences in amygdala connectivity during negative distractors ...	110
Figure 5.4. Group differences in dACC connectivity during negative distractors	112
Figure 5.5. Regression analysis: Dissociation as regressor for bilateral amygdala FC.	113
Figure S5.1. Results for the main effect of group on amygdala connectivity	126
Figure S5.2. Results for the main effect of group on dACC connectivity.	126
Figure S5.3. Results for the interaction effect on dACC connectivity	127
Figure S5.4. Regression analysis: Reaction times as regressor for amygdala FC	127
Figure 6.1. Study design–overview	134
Figure 6.2. Group differences in mean reaction times and accuracy in the EST.	140
Figure 6.3. Neural activation in response to the Emotional Stroop Task.	142
Figure 7.1. Working memory performance during the EWMT	165
Figure 7.2. Percent signal change in the bilateral amygdala during the EWMT	166
Figure 7.3. Results of the Psychophysiological Interaction analysis	166
Figure S7.1. Specific types of errors during the Emotional Working Memory Task	174
Figure 8.1. Brain regions that may be implicated in dissociation in BPD	186

List of most commonly used Abbreviations

ACC	Anterior Cingulate Cortex
BOLD	Blood Oxygen Level-Dependent
BPD	Borderline Personality Disorder
DACC	Dorsal Anterior Cingulate Cortex
D(D)D	Depersonalization(/Derealization) Disorder
DID	Dissociative Identity Disorder
DLPFC	Dorsolateral Prefrontal Cortex
DMPFC	Dorsomedial Prefrontal Cortex
e.g.	for example
EST	Emotional Stroop Task
EWMT	Emotional Working Memory Task
FMRI	Functional Magnetic Resonance Imaging
HC	Healthy controls
IAPS	International Affective Picture System
i.e.	that is
MDD	Major Depressive Disorder
ms.	Milliseconds
OFC	Orbitofrontal Cortex
PD	Personality Disorder
PPI	Psychophysiological Interaction Analysis
PTSD	Posttraumatic Stress Disorder
RT	Reaction times
SSRI	Selective serotonin reuptake inhibitors.
VLPFC	Ventrolateral Prefrontal Cortex
VMPFC	Ventromedial Prefrontal Cortex
WM	Working Memory