



Universiteit
Leiden
The Netherlands

Exploring the world of non-coding genes in stem cells and autoimmunity.

Messemaker, T.C.

Citation

Messemaker, T. C. (2018, April 3). *Exploring the world of non-coding genes in stem cells and autoimmunity*. Retrieved from <https://hdl.handle.net/1887/61075>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/61075>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/61075> holds various files of this Leiden University dissertation.

Author: Messemaker, T.C.

Title: Exploring the world of non-coding genes in stem cells and autoimmunity

Issue Date: 2018-04-03

Chapter 4

Comment on "Functional Analysis of a Complement Polymorphism (rs17611) Associated with Rheumatoid Arthritis"

J Immunol. 2015 Jul 1;195(1):3-4. doi: 10.4049/jimmunol.1500822.

Messemaeker T, Toes RE, Mikkers HM, Kurreeman F

Comment on “Functional Analysis of a Complement Polymorphism (rs17611) Associated with Rheumatoid Arthritis”

In their published article, Giles *et al.* (1) explored functional consequences of rs17611 in the TRAF1-C5 region and propose a mechanism for its contribution to rheumatoid arthritis (RA) pathology, as this SNP would associate with RA. To substantiate the claim that the SNP associates with RA, the authors referred to the following studies: Refs. 2 and 3.

However, the data presented by Kurreeman *et al.* do not indicate a significant association ($p = 0.19$) in 544 subjects (Ref. 2, see Table I). In addition, haplotype block analyses of this risk locus shows that block 2 is significantly associated with RA, while block 3 (which includes rs17611) is not (Ref. 2, see Fig. 1). Similar data by Plenge *et al.* (3) across 3372 subjects support this lack of association. Recent genome-wide association studies data across 55,000 European subjects (4) provide accurate estimations of risk across hundreds of SNPs in this locus.




Figure 1. Regional association plot for the TRAF1-C5 region. The x-axis shows the chromosomal position of the all queried SNPs located on chromosome 9p33p34 over a region of 500 kb. The y-axis displays the $-\log_{10}(p\text{-value})$ of associations with RA. The p values were a result of analysis on the European population as part of the Okada *et al.* paper (4): 14,361 RA cases and 43,923 controls from 18 studies of Europeans descent. Pairwise LD values (r^2) were calculated from individuals of the 1000 Genomes Project (CEU) relative to the highest associating SNP rs10985070 using SNAP (6). N.D., LD for SNP is not determined.

Again, no signal of association for rs17611 has been demonstrated in the available published datasets originating from this study (Fig. 1), which also shows that rs17611 is in relatively low linkage disequilibrium (LD) with the lead SNP rs10985070 ($r^2 = 0.4$). Moreover, independent secondary association signals were not observed in the TRAF1-C5 locus in a study by Eyre *et al.* (5), excluding the possibility that rs17611 could represent a second hit at this locus. On the basis of the published genetic data, we believe caution should be taken in implicating the effects of rs17611 in relation to the immunological mechanism underlying the genetic risk of TRAF1-C5 to RA.

References

1. Giles J. L., E. Choy, C. van den Berg, B. P. Morgan, C. L. Harris. 2015. Functional analysis of a complement polymorphism (rs17611) associated with rheumatoid arthritis. *J. Immunol.* 194: 3029–3034.
2. Kurreeman F. A. S., L. Padyukov, R. B. Marques, S. J. Schrodi, M. Seddighzadeh, G. Stoeken-Rijsbergen, A. H. M. van der Helm-van Mil, C. F. Allaart, W. Verduyn, J. Houwing-Duistermaat, *et al.* 2007. A candidate gene approach identifies the TRAF1/C5 region as a risk factor for rheumatoid arthritis. *PLoS Med.* 4: e278.
3. Plenge R. M., M. Seielstad, L. Padyukov, A. T. Lee, E. F. Remmers, B. Ding, A. Liew, H. Khalili, A. Chandrasekaran, L. R. Davies, *et al.* 2007. TRAF1-C5 as a risk locus for rheumatoid arthritis—a genomewide study. *N. Engl. J. Med.* 357: 1199–1209.
4. Okada, Y., Wu, D., Trynka, G., Raj, T., Terao, C., Ikari, K., Kochi, Y., Ohmura, K., Suzuki, A., Yoshida, S., *et al.* 2013. Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* 506: 376–381.
5. Eyre S., J. Bowes, D. Diogo, A. Lee, A. Barton, P. Martin, A. Zernakova, E. Stahl, S. Viatte, K. McAllister, *et al.* Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate; Wellcome Trust Case Control Consortium. 2012. High-density genetic mapping identifies new susceptibility loci for rheumatoid arthritis. *Nat. Genet.* 44: 1336–1340.
6. Johnson A. D., R. E. Handsaker, S. L. Pulit, M. M. Nizzari, C. J. O'Donnell, P. I. W. de Bakker. 2008. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 24: 2938–2939.

Response to Comment on “Functional Analysis of a Complement Polymorphism (rs17611) Associated with Rheumatoid Arthritis”

Joanna L. Giles, Ernest Choy, Carmen van den Berg, B. Paul Morgan and Claire L. Harris

In their *Comment*, Messemaker *et al.* discuss genetic data that show that block 2 within the TRAF1-C5 locus is a risk for rheumatoid arthritis (RA), while block 3 is

not associated. We acknowledge these data, which illustrate that, when taken in isolation, block 3 does not impact disease risk. It is clear that the genetic data are robust (1); however, there is no question that the rs17611 single nucleotide polymorphism (SNP) in C5 results in an amino acid change in C5 that alters its rate of cleavage by elastase: an enzyme present at high levels in neutrophil-driven or neutrophil-associated diseases such as RA. We believe that it would be important and interesting to further analyze whether this functional polymorphism in block 3, which affects the proinflammatory capacity of C5, has any impact on the risk haplotype in block 2. While genome-wide association studies have identified a number of independent secondary association signals, these are not exclusive (2); the strong functional data that we demonstrate with the C5-V802I variant suggest that potential interactions should be specifically tested in future studies to confirm or exclude them.

In our article (3), we reference other papers where the specific rs17611 SNP has been linked to different diseases. Chai *et al.* (4) show an association ($p < 0.007$) of rs17611 (and its linked set) with periodontal disease; Woehrl *et al.* (5) show an association ($p < 0.002$) of rs17611 with outcome in pneumococcal meningitis; Hoke *et al.* (6) show an association ($p < 0.01$) of rs17611 with adverse cardiovascular outcome; and Greisenegger *et al.* (7) show an association ($p < 0.005$) of rs17611 with risk for ischemic stroke. Together, these linkages provided a strong rationale to explore the functional consequences of the rs17611 SNP on C5 activities, the major focus of our paper.

First, we show that the rs17611 SNP is associated with clear differences in C5 turnover and elevated levels of the proinflammatory product C5a in healthy individuals and RA patients; our finding warrants further investigation of the impact of this polymorphism on risk associated with block 2. We go on to define the mechanism by which the single amino acid change provokes these differences. The genetic data do not detract from the importance of our demonstration that the polymorphism impacts C5 turnover and increases plasma levels of the proinflammatory molecule C5a, and of our description of the atypical route by which this is achieved (3).

References

1. Kurreeman F. A., L. Padyukov, R. B. Marques, S. J. Schrodj, M. Seddighzadeh, G. Stoeken-Rijsbergen, A. H. van der Helm-van Mil, C. F. Allaart, W. Verduyn, J. Houwing-Duistermaat, *et al.* 2007. A

- candidate gene approach identifies the TRAF1/C5 region as a risk factor for rheumatoid arthritis. *PLoS Med.* 4: e278.
2. Eyre S., J. Bowes, D. Diogo, A. Lee, A. Barton, P. Martin, A. Zernakova, E. Stahl, S. Viatte, K. McAllister, *et al*, Biologics in Rheumatoid Arthritis Genetics and Genomics Study Syndicate; Wellcome Trust Case Control Consortium. 2012. High-density genetic mapping identifies new susceptibility loci for rheumatoid arthritis. *Nat. Genet.* 44: 1336–1340.
 3. Giles J. L., E. Choy, C. van den Berg, B. P. Morgan, C. L. Harris. 2015. Functional analysis of a complement polymorphism (rs17611) associated with rheumatoid arthritis. *J. Immunol.* 194: 3029–3034.
 4. Chai L., Y. Q. Song, K. Y. Zee, W. K. Leung. 2010. Single nucleotide polymorphisms of complement component 5 and periodontitis. *J. Periodontal Res.* 45: 301–308.
 5. Woehrl B., M. C. Brouwer, C. Murr, S. G. Heckenberg, F. Baas, H. W. Pfister, A. H. Zwinderman, B. P. Morgan, S. R. Barnum, A. van der Ende, *et al*. 2011. Complement component 5 contributes to poor disease outcome in humans and mice with pneumococcal meningitis. *J. Clin. Invest.* 121: 3943–3953.
 6. Hoke M., W. Speidl, M. Schillinger, E. Minar, S. Zehetmayer, M. Schönherr, O. Wagner, C. Mannhalter. 2012. Polymorphism of the complement 5 gene and cardiovascular outcome in patients with atherosclerosis. *Eur. J. Clin. Invest.* 42: 921–926.
 7. Greisenegger S., S. Zehetmayer, P. Bauer, G. Endler, J. Ferrari, W. Lang, M. Janisiw, L. Steiner, S. Cheng, W. Lalouschek, C. Mannhalter. 2009. Polymorphisms in inflammatory genes and the risk of ischemic stroke and transient ischemic attack: results of a multilocus genotyping assay. *Clin. Chem.* 55: 134–138.

