



Universiteit
Leiden
The Netherlands

The diagnostic and therapeutic management of pulmonary embolism

Hulle, T. van der

Citation

Hulle, T. van der. (2018, January 10). *The diagnostic and therapeutic management of pulmonary embolism*. Retrieved from <https://hdl.handle.net/1887/61127>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/61127>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The following handle holds various files of this Leiden University dissertation:

<http://hdl.handle.net/1887/61127>

Author: Hulle, T. van der

Title: The diagnostic and therapeutic management of pulmonary embolism

Issue Date: 2018-01-10



CHAPTER 2

Recent developments in the
diagnosis and treatment of
pulmonary embolism

T. van der Hulle, C.E.A. Dronkers, F.A. Klok, M.V. Huisman

Journal of Internal Medicine 2016;279:16-29

ABSTRACT

Due to the nonspecific symptoms of the condition, a diagnosis of acute pulmonary embolism (PE) is frequently considered. However, PE will only be confirmed in 10–20% of patients. Because the imaging test of choice, computed tomography pulmonary angiography (CTPA), is costly and associated with radiation exposure and other complications, a validated diagnostic algorithm consisting of a clinical decision rule and D-dimer test should be used to safely exclude PE in 20–30% of patients without the need for CTPA. Recently, the age-adjusted D-dimer threshold has been validated, and this has increased the proportion of patients at older age in which PE can be excluded without CTPA. Initial therapeutic management of PE depends on the risk of short-term PE-related mortality. Haemodynamically unstable patients should be closely monitored and receive thrombolytic therapy unless contraindicated because of an unacceptably high bleeding risk, whereas patients with low-risk PE may be safely discharged early from hospital or receive only outpatient treatment. The PESI score and Hestia decision rule are available to select patients in whom early discharge or outpatient treatment will be safe, although the safety of these strategies should be confirmed in additional studies. Standard PE therapy consists of low molecular weight heparin (LMWH) followed by vitamin K antagonists (VKAs). Recently, several nonvitamin K-dependent oral anticoagulants have been shown to be as effective as LMWH/VKAs, and maybe safer. Determining the optimal duration of treatment for a first unprovoked PE remains a challenge, although clinical prediction rules for estimating the risk of recurrence of venous thromboembolism and anticoagulation-associated haemorrhage are under investigation. Using these prediction rules may lead to both more standardized and more individualized long-term treatment of PE.

INTRODUCTION

Acute pulmonary embolism (PE) is a relatively common disease with an annual rate of 1–2 per 1000 patients [1, 2]. The clinical presentation of acute PE is nonspecific and highly variable, ranging from incidentally diagnosed asymptomatic thrombi to sudden death [3]. As a result, a clinical suspicion of PE is frequently raised, whilst the diagnosis is only confirmed in 10–20% of patients [4, 5]. Diagnostic algorithms have been developed to ensure reliable and efficient management of patients with clinically suspected PE.

Once acute PE is diagnosed, prompt initiation of anticoagulant therapy is indicated to prevent thrombus extension and recurrent (fatal) PE. However, the risk of such an adverse outcome is highly variable, ranging from <1% to >15% [6]. Management decisions including early discharge or prescription of thrombolytic therapy should preferably be based on reproducible risk stratification of the individual patient [6, 7]. For the long-term treatment of acute PE, the decision to continue or stop anticoagulant therapy, after an initial period of 3 months, depends on the balance between the risk of recurrent venous thromboembolism (VTE) and of anticoagulant-associated haemorrhage.

Here, we will present an up-to-date overview of the therapeutic management of established PE, discussing conventional medical therapy, the recently introduced non-vitamin K-dependent oral anticoagulants (NOACs), outpatient treatment, thrombolytic therapy and the duration of treatment. In particular, we will focus on recent advances as a follow-up to a review of a similar topic published in this journal in 2010 [8].

DIAGNOSTIC MANAGEMENT OF CLINICALLY SUSPECTED ACUTE PE

Symptoms that may suggest the presence of acute PE are the sudden onset of dyspnoea, pleuritic chest pain, haemoptysis, extremity swelling suggestive of deep vein thrombosis (DVT) and syncope [3]. Of note, these symptoms are nonspecific for PE and may also be present in many other acute cardiopulmonary conditions. In large diagnostic management studies, the PE prevalence amongst patients with a clinical suspicion of PE ranged from 10% to 30% [4, 5, 9]. PE can only be diagnosed using an imaging test, at present usually computed tomography pulmonary angiography (CTPA), which is associated with high healthcare costs, radiation exposure and a risk of complications such as contrast-induced nephropathy and allergic reactions [10–12]. Therefore, diagnostic management algorithms have been developed to exclude PE without the need for imaging tests in a proportion of patients. These algorithms start with a clinical decision rule (CDR), which predicts the clinical pretest probability of PE, followed by a D-dimer blood test and/or a CTPA.

CDRs and D-dimer testing

Various CDRs have been standardized and validated for use in clinical practice, of these, the Wells rule and revised Geneva score are the most widely validated for PE (**Table 1**) [13, 14]. Originally, different items were awarded different scores, but for practical reasons, both rules have been simplified by assigning only one point to each item, without lowering their diagnostic accuracy [15, 16]. The CDRs are used to categorize patients into either 'PE unlikely' or 'PE likely' groups, with PE prevalences of 12–17% and 37–43%, respectively, in European cohorts [4, 5]. The accuracy of the original and simplified versions of the Wells rule and revised Geneva score were shown to be similar, and the choice of a specific rule may therefore depend on local preference [4].

Table 1. Clinical decision rules.

Item	Original version	Simplified version
Wells rule		
Previous PE or DVT	1.5	1
Heart rate >100 beats/min	1.5	1
Surgery or immobilization within 4 weeks	1.5	1
Haemoptysis	1	1
Active malignancy	1	1
Clinical signs of DVT	3	1
Alternative diagnosis less likely than PE	3	1
Clinical probability categories		
PE unlikely	≤4	≤1
PE likely	>4	>1
Revised Geneva score		
Previous DVT or PE	3	1
Heart rate 75–94 beats/min	3	1
Heart rate ≥95 beats/min	5	2
Surgery or fracture within 1 month	2	1
Haemoptysis	2	1
Active malignancy	2	1
Unilateral lower limb pain	3	1
Pain on lower limb deep vein palpation and unilateral oedema	4	1
Age >65 years	1	1
Clinical probability categories		
PE unlikely	≤ 5	≤2
PE likely	> 5	>2

Note: DVT, deep vein thrombosis; PE, pulmonary embolism.

The D-dimer level in plasma is a marker of fibrinolysis and is elevated in the presence of acute VTE as well as in many other clinical conditions. The sensitivity of D-dimer testing for the latest generation of assays is very high (95–100%), whereas the specificity is moderate (43–93%) [17]. Accordingly, using the standard threshold of $<500 \mu\text{g/L}$, the negative predictive value of D-dimer testing allows for the exclusion of PE in patients with a PE unlikely clinical probability [4, 5, 9]. By applying this strategy, PE can be ruled out in 25–46% of patients without performing an imaging test, with a 3-month VTE incidence of 0.04–0.96% after anticoagulant therapy is withheld [18].

Recently, attempts have been made to increase the number of patients in whom PE can be ruled out without imaging tests. As D-dimer levels increase with age, an age-dependent D-dimer cut-off level was proposed: $\text{age} \times 10 \mu\text{g/L}$ in patients over 50 years of age [19]. The safety of applying this age-dependent D-dimer cut-off was recently confirmed in a large prospective outcome trial [9]. The 3-month VTE incidence in patients in the PE unlikely category and with a D-dimer level $>500 \mu\text{g/L}$ but below the age-adjusted threshold was only 0.3% (95% confidence interval (CI) 0.1–1.7). The absolute increase in the proportion of patients above 50 years old that could be managed without CTPA was 11.6%. In particular, patients aged 75 years and older could be more frequently managed safely without CTPA.

Imaging tests

The historical gold standard imaging test for diagnosing PE is pulmonary angiography, which is associated with a 3-month VTE incidence after a negative result of 1.7% (95% CI 1.0–2.7) [20]. Because pulmonary angiography is an invasive procedure, this test was first replaced by ventilation-perfusion lung scanning (V/Q scanning) and later by CTPA. Although it has been demonstrated that V/Q scanning is safe for excluding PE, with a 3-month VTE incidence of only 0.9% (upper limit of 95% CI 2.3) after a normal result, its major disadvantage is an inconclusive result in 28–46% of patients, in whom the PE prevalence is still 10–30% and who thus require an additional imaging test [21, 22]. CTPA results in an inconclusive test result in only 0.9–4.6% of patients [23]. Furthermore, CTPA is more widely available with faster acquisition and the possibility of establishing an alternative diagnosis, although the value of the latter has recently been questioned [24]. A possible advantage of V/Q scanning is the lower radiation exposure compared to CTPA (1.1 vs. 2–6 mSv), which may be relevant for specific patient groups, particularly young (particularly female) patients and pregnant women. V/Q scanning may also be considered for patients with a known allergy to contrast medium [11, 12].

The sensitivity of multidetector row high-resolution CTPA is excellent and several outcome studies have consistently demonstrated the safety of withholding anticoagulant therapy in patients with a negative CTPA alone [23]. In a meta-analysis, the pooled 3-month VTE incidence after a negative CTPA alone in patients in whom CTPA was


indicated based on a validated CDR and/or D-dimer test was 1.2% (95% CI 0.8–1.8), compared to 1.1% (95% CI 0.6–2.0) after negative CTPA and subsequently negative lower limb compression ultrasonography, indicating that the safety could not be further improved by excluding asymptomatic DVT after a negative CTPA [23]. In addition to ionizing radiation, other potential concerns related to the easily available CTPA as the first-line imaging test are an increased incidence of contrast-induced nephropathy, a risk of unsuspected findings requiring additional medical attention and an increased incidence of isolated subsegmental PE, of which the clinical relevance has become increasingly under debate [10, 25–27].

Integrated approach

The first step in the diagnostic management of patients with clinically suspected acute PE is to determine whether the patient has signs and symptoms of haemodynamic shock. If so, patients should be immediately referred for additional imaging tests and thrombolytic therapy should be started without waiting for a CDR result or D-dimer level.

In patients who are haemodynamically stable, diagnostic management starts with the assessment of the clinical probability of PE using one of the validated CDRs (**Table 1**). In patients with a PE unlikely clinical probability, a D-dimer test is indicated and PE can be ruled out without further imaging tests if the level is $<500 \mu\text{g/L}$ or below the age-adjusted threshold for patients over the age of 50 years. In the remaining patients with either a PE unlikely CDR score in combination with an elevated D-dimer level or a PE likely CDR score, CTPA is indicated to determine the presence of PE (**Figure 1**).

Figure 1. Integrated approach for clinically suspected acute pulmonary embolism in haemodynamically stable patients.



Note: CDR, clinical decision rule; CTPA, computed tomography pulmonary angiography; PE, pulmonary embolism.

Correct use of a validated diagnostic algorithm is required to ensure its safety; this is highlighted by several recent studies showing that a validated diagnostic algorithm is frequently not or incorrectly applied. This leads to the unnecessary use of CTPA and, even more relevant, to a higher 3-month VTE incidence in patients in whom anticoagulant therapy is withheld [28]. A promising intervention to improve adherence may be the implementation of computerized clinical decision support for clinicians [29]. Future studies should focus on further simplification of the diagnostic algorithms to improve

the compliance and consequently the safety and efficiency of diagnostic management in daily clinical practice. Future studies should also investigate whether increasing the D-dimer threshold, for instance with variable D-dimer thresholds depending on the pretest probability, may further improve the efficiency without reducing the safety of the diagnostic algorithm [30].

TREATMENT OF ACUTE PE

Risk stratification of acute PE

Patients with acute PE should be stratified according to the short-term PE-related mortality risk. Risk stratification starts with identifying patients in haemodynamic shock, who are classified as having high-risk or massive PE with an estimated 30-day PE-related mortality risk of >15% [7].

For the remaining haemodynamically stable patients, the ability of clinical prediction rules and several tests to distinguish patients with a low risk (<1%) from those with an intermediate risk (3–15%) of an adverse outcome has been investigated [7].

The Hestia decision rule consists of a set of criteria that can be used to select patients with low-risk PE who are candidates for early discharge or outpatient treatment (**Table 2**) [31]. The major strength of the Hestia decision rule is its clinician friendliness, as it consists of only 11 easy-to-use bedside criteria that should all be negative before early discharge or outpatient treatment can be considered. The Pulmonary Embolism Severity Index (PESI), which combines several clinical signs, symptoms and comorbidities (**Table 3**), is also available and is able to identify patients at low risk of short-term PE-related mortality [6]. To overcome its complexity, a simplified version (sPESI) has been proposed [32].

In addition to the Hestia decision rule and the PESI, several laboratory biomarkers have been shown to predict adverse outcome in haemodynamically stable patients. High concentrations of brain-type natriuretic peptides (BNPs) or the N-terminal of the prohormone of BNP (NT-proBNP) that are elevated during PE-associated right ventricular (RV) overload are strongly associated with mortality in acute PE with an OR of 7.6 (95% CI 3.4–17), whereas normal levels identify patients with a low risk of short-term PE-related mortality (17% vs. 1.7%) [33]. In addition, elevated plasma troponin I or (high-sensitivity) troponin T levels are associated with a high risk of short-term mortality (18% vs. 2.3%) in haemodynamically stable patients with an OR of 5.90 (95% CI 2.7–13) [34].

Both echocardiography and CTPA images can be used to detect RV dysfunction, which is also associated with an elevated risk of short-term PE-related mortality in patients who are haemodynamically stable. At least 25% of patients with PE have signs of RV dysfunction on echocardiography, such as RV dilatation, an increased RV/left ventricular (LV) diameter ratio, hypokinesia of the free RV wall, increased tricuspid regurgitation

Table 2. Hestia decision rule.

Haemodynamically unstable? ^a
Thrombolysis or embolectomy necessary?
High risk of bleeding? ^b
Oxygen supply to maintain oxygen saturation >90% for more than 24 h?
Pulmonary embolism diagnosed during anticoagulant treatment?
Intravenous pain medication for more 24 h?
Medical or social reason for treatment in hospital for more than 24 h?
Creatinine clearance <30 mL/min? ^c
Severe liver impairment? ^d
Pregnant?
Documented history of heparin-induced thrombocytopenia?
Interpretation
If the answer to at least one of the above questions is 'YES', the patient cannot be treated as an outpatient

Note: ^aIncludes, at the discretion of the physician, systolic blood pressure <100 mmHg with heart rate >100 beats/min; condition requiring admission to an intensive care unit; ^bgastrointestinal bleeding in the preceding 14 days, recent stroke (within 4 weeks), recent operation (within 2 weeks), bleeding disorder or thrombocytopenia (platelet count <75 × 10⁹/L) and/or uncontrolled hypertension (systolic blood pressure >180 mmHg or diastolic blood pressure > 110 mmHg); ^ccreatinine clearance calculated according to the Cockcroft–Gault formula; ^dat the discretion of the treating physician.

jet velocity or decreased tricuspid annulus plane systolic excursion. Echocardiographic signs of RV dysfunction in haemodynamically stable patients with PE are associated with an OR of 2.4 (95% CI 1.3–4.3) for short-term mortality [35]. The RV/LV dimensional ratio can also be assessed on axial or reconstructed four-chamber CTPA views, for which thresholds of ≥0.9 or ≥1.0 are generally used. CTPA-assessed RV dysfunction in haemodynamically stable patients is associated with an increased risk of short-term mortality (7.8% vs. 5.1%) as well as with short-term PE-related mortality (2.2% vs. 0.2%) with ORs of 1.8 (95% CI 1.3–2.6) and 7.4 (95% CI 1.4–40), respectively [36]. Interestingly, several studies have shown that models combining prediction rules, biomarkers and imaging tests improve the predictive accuracy compared with the individual tests, although at present, none of these combinations can be recommended to guide management in daily clinical practice [37, 38].

Management of high-risk patients

Thrombolytic therapy is able to rapidly resolve thromboembolic obstruction enabling prompt reduction in pulmonary artery pressure and improvement in RV function. The results of one study suggested that a clinical and echocardiographic improvement can be observed in more than 90% of patients who receive thrombolysis [39]. Thrombolytic therapy is generally recommended for high-risk patients with PE with overt haemodynamic instability without a high risk of bleeding complications, although there is a lack

Table 3. PESI score.

Parameter	Original version	Simplified version
Age	Age in years	1 point (if age >80 years)
Male sex	+10 points	
Cancer	+30 points	1 point
Chronic heart failure	+10 points	1 point
Chronic pulmonary disease	+10 points	
Heart rate ≥ 110 beats/min	+20 points	1 point
Systolic blood pressure <100 mmHg	+30 points	1 point
Respiratory rate >30 breaths/min	+20 points	
Temperature <36 °C	+20 points	
Altered mental status ^a	+60 points	
Arterial oxygen saturation <90% ^b	+20 points	1 point
Risk strata		
	Class I: ≤ 65 points very low 30-day mortality risk 0–1.6%	0 points = 30-day mortality risk 1.0% (95% CI 0–2.1)
	Class II: 66–85 points low mortality risk 1.7–3.5%	≥ 1 point(s) = 30-day mortality risk 10.9% (95% CI 8.5–13.2)
	Class III: 86–105 points moderate mortality risk 3.2–7.1%	
	Class IV: 106–125 points high mortality risk 4.0–11.4%	
	Class V: >125 points very high mortality risk 10.0–24.5%	

Note: PESI, Pulmonary Embolism Severity Index. ^aDefined as disorientation, lethargy, stupor or coma. ^bWith and without the administration of supplemental oxygen.

of convincing supporting evidence from large randomized controlled trials. A meta-analysis of randomized trials including haemodynamically unstable patients with PE demonstrated an OR of 0.53 (2.2% vs. 3.9%; 95% CI 0.32–0.88) for recurrent PE and death in patients who received thrombolytic therapy [40]. In addition, in an epidemiological study, haemodynamically unstable patients with PE who received thrombolytic therapy had a relative risk of 0.2 (95% CI 0.19–0.22) for PE-related in-hospital death compared to those who had not received thrombolytic therapy [41]. Of major concern, it was shown in a recent meta-analysis that thrombolytic therapy substantially increased the risk of major bleeding complications with an OR of 2.73 (9.2% vs. 3.4%; 95% CI 1.91–3.91) compared to standard anticoagulant treatment [40]. Percutaneous catheter-directed therapy and surgical embolectomy are alternatives to thrombolytic therapy, for example

in the case of a contraindication to thrombolysis or after thrombolytic therapy has failed or is deemed inadequate as first-line treatment, although these interventions have not been investigated in clinical outcome studies [42].

Management of nonhigh-risk patients

Amongst the nonhigh-risk haemodynamically stable patients, two key questions with regard to therapeutic management remain to be answered: (i) Do patients with intermediate-risk PE benefit from thrombolytic therapy? (ii) Which patients with low risk of adverse outcome are suitable candidates for outpatient treatment or early discharge?

The benefit of administering thrombolytic drugs to haemodynamically stable patients with intermediate-risk PE has been debated for many years. Individual small randomized trials have consistently demonstrated that thrombolytic therapy reduces the risk of haemodynamic deterioration and recurrent PE, at the cost of a higher risk of major haemorrhage without an effect on overall mortality [40, 43]. In 2014, the results of the randomized double-blind PEITHO trial were published. This study investigated the effects of a single dose of tenecteplase (varying from 30 to 50 mg depending on body weight) versus placebo in 1006 patients with an intermediate risk of PE [44]. Intermediate-risk PE was defined as a PESI score of III–V, elevated biomarkers of cardiac injury and signs of RV dysfunction on echocardiography or CTPA. The study demonstrated a significant reduction in the primary efficacy outcome (a composite of all-cause death or haemodynamic deterioration within 7 days of randomization), with an OR of 0.44 (2.6% vs. 5.6%; 95% CI 0.23–0.88). However, 7-day mortality was not different between the two groups: 1.2% of patients who received tenecteplase and 1.8% of patients who received placebo (OR 0.65; 95% CI 0.23–1.85). Of interest, patients who were randomly assigned to receive tenecteplase had a significantly higher risk of major extracranial bleeding and (predominantly haemorrhagic) stroke within 7 days with ORs of 5.6 (6.3% vs. 1.2%; 95% CI 2.3–13.4) and 12 (2.4% vs. 0.2%; 95% CI 1.6–93.4), respectively. It should be noted that only 23 patients (4.6%) from the placebo cohort eventually received rescue thrombolysis during follow-up because of in-hospital haemodynamic deterioration, and two of these patients died within 7 days.

In conclusion, thrombolytic therapy in haemodynamically stable intermediate-risk patients with PE reduces the risk of the composite endpoint consisting of haemodynamic deterioration and death at the cost of an increase in the incidence of major haemorrhage. Therefore, thrombolytic therapy cannot be recommended for haemodynamically stable intermediate-risk patients with PE. These patients should receive standard anticoagulant therapy and close monitoring, whilst thrombolytic therapy should be reserved for patients with haemodynamic deterioration during the first days of treatment.

One randomized trial and two cohort studies evaluating outpatient treatment or early discharge for patients with PE have been completed and the results published. In the

largest study reported to date, there were initially 14 exclusion criteria (**Table 4**) and thereafter patients with a PESI class I or II were selected [45]. Patients were randomly assigned to either conventional in-hospital treatment or outpatient treatment (discharge after a mean of 0.5 days). Of the 1557 patients screened, 470 (30%) were potentially eligible for early discharge (<24 h) of whom 344 were eventually enrolled in the study. Only one of the 171 (0.6%) evaluable patients receiving outpatient treatment developed a nonfatal recurrent VTE during the 3-month follow-up, and two patients (1.2%) had a major haemorrhage within 14 days. Amongst the 168 evaluable patients treated in hospital, neither recurrent VTE nor major haemorrhage occurred within 14 days. Based on these results, the authors concluded that outpatient treatment was noninferior to in-hospital treatment [45]. A major limitation of this strategy is the complexity of scoring the 11 (items) of the PESI score combined with the many other exclusion criteria that were used in the study. Although this limitation may be partly overcome using the sPESI, it should be noted that the simplified score has not been validated in a prospective management study.

Table 4. Exclusion criteria in a randomized trial investigating outpatient treatment amongst PE patients with a PESI class I or II (from Aujesky et al. [45]).

Arterial hypoxaemia (oxygen saturation on room air <90% measured by pulse oximetry or a partial pressure of oxygen <60 mmHg on arterial blood gas analysis)
Systolic blood pressure <100 mmHg
Chest pain necessitating parenteral administration of opioids
Active bleeding or high risk of bleeding (defined as stroke during the preceding 10 days)
Gastrointestinal bleeding during the preceding 14 days or platelet count <75 × 10 ⁹ /L
Severe renal failure (creatinine clearance <30 mL/min based on the Cockcroft–Gault equation)
Extreme obesity (body mass >150 kg), history of heparin-induced thrombocytopenia or allergy to heparins
Therapeutic oral anticoagulation at the time of diagnosis of PE (INR ≥2.0)
Any barriers to treatment adherence or follow-up (e.g. current alcohol abuse, illicit drug use, psychosis, dementia or homelessness)
Pregnancy
Imprisonment
Diagnosis of pulmonary embolism >23 h before the time of screening (to avoid enrolling already stabilized patients)
Previous enrolment in the trial

Note: INR, international normalized ratio; PE, pulmonary embolism; PESI, Pulmonary Embolism Severity Index.

The Hestia decision rule has also been evaluated as a selection tool for outpatient treatment (**Table 2**) [31]. In a prospective cohort study, 297 (51%) of the 581 consecutive patients with acute PE were selected for outpatient treatment of whom six (2.0%) experienced a nonfatal recurrent VTE. Two patients (0.7%) experienced a major haemor-

rhage, one (0.3%) of whom suffered fatal intracranial bleeding. The major advantages of this strategy are the clinician friendliness and the high degree of simplicity.

NT-proBNP has also been evaluated in a cohort study as a tool for selecting patients for outpatient treatment [46]. Haemodynamically stable patients with a low NT-proBNP concentration (threshold <500 pg/mL) were discharged within 24 h of diagnosis. Of the 351 patients in this cohort, 152 (43%) were treated as outpatients and none developed a recurrent VTE or major haemorrhage or died during 3 months of follow-up [46]. In the Vesta study, the results of which will be presented at the congress of the International Society of Thrombosis and Haemostasis in June 2015, a total of 550 patients were randomly assigned to the Hestia decision rule combined with an NT-proBNP test or the Hestia decision rule alone (NTR2603). Of the 275 patients managed according to the Hestia decision rule in combination with an NT-proBNP test, 34 (12%) had a concentration of NT-proBNP >500 ng/L and were managed as inpatients. None of these patients suffered from the primary outcome (30-day adverse outcome defined as PE or bleeding-related mortality, cardiopulmonary resuscitation or admission to the intensive care unit). Of the 275 patients managed according to the Hestia decision rule alone, three (1.1%; 95% CI 0.2–3.2) experienced the primary outcome, all of whom had a normal NT-proBNP level, which were determined in a post hoc analysis. Therefore, the findings of this study confirm the safety of selecting patients who can be treated as outpatients and suggest that NT-proBNP measurements are of limited additional value.

In conclusion, it has been shown that outpatient treatment of patients with PE based on the Hestia decision rule or a combination of several exclusion criteria and the PESI score is safe. Of note, neither the PESI nor the sPESI rule has been evaluated as a sole test to select patients for outpatient treatment. The combination of a clinical prediction rule, laboratory biomarkers and/or findings on imaging tests to further optimize the identification of patients who can be safely managed in the outpatient setting remains to be evaluated.

Initial anticoagulant therapy (first 3 months)

Acute PE requires initial treatment with a direct-onset anticoagulant drug to prevent the extension of thrombosis or fatal recurrent VTE [7, 47]. Weight-adjusted subcutaneous low molecular weight heparin (LMWH) is the treatment of choice for the large majority of patients. Intravenous unfractionated heparin is reserved for patients with severe renal impairment (creatinine clearance <20 – 30 mL/min), patients with a high risk of haemorrhage including those receiving thrombolytic therapy, haemodynamically unstable patients and individuals who are extremely overweight or underweight. Fondaparinux is an alternative to LMWH and unfractionated heparin in patients with (a history of) heparin-induced thrombocytopenia [7, 48]. Monitoring of the anticoagulant effect of LMWH by determining the antifactor Xa activity is not generally recommended, although it can

be considered in specific circumstances (i.e. in patients with moderate renal impairment or during pregnancy). The target range measured 4 hours after administration is 1.0–2.0 IU/mL for once-daily administration and 0.6–1.0 IU/mL for twice-daily administration of LMWH [7].

Patients with acute PE should be treated for at least 3 months. For decades, vitamin K antagonists (VKAs) have been the long-term treatment of choice and must be started in parallel with one of the parenterally administered direct-onset anticoagulant drugs for at least 5 days and until the international normalized ratio (INR) has reached a therapeutic range (between 2.0 and 3.0) on 2 consecutive days. VKA therapy is highly effective with a rate of recurrent VTE of 3–4% during the initial 3 months of therapy [49]. It has been estimated that the rate of major haemorrhage during the initial 3 months of VKA therapy is 1–2% [49]. A major disadvantage of VKA therapy is the need for frequent laboratory monitoring of the INR with tailored dosing, as a result of the variable pharmacokinetic and pharmacodynamic parameters, and interactions with food and drugs [50].

In recent years, several NOACs have become available for anticoagulant therapy (**Table 5**). These drugs directly and specifically inhibit either thrombin (factor II) or factor Xa. The major advantage of this new drug class is the simplification of PE treatment, due to the rapid onset of action, the predictable anticoagulant effect and a low potential for drug and food interactions [50]. Rivaroxaban and apixaban were investigated with an initial higher dose for 3 weeks and 7 days, respectively, without the need for preceding LMWH therapy, whereas dabigatran and edoxaban were administered after a mean period of 10 days of treatment with LMWH. These drugs can be prescribed in fixed doses and routine laboratory monitoring is not required. A noninferior efficacy compared to standard treatment with VKA has been demonstrated for all NOACs individually in phase III trials [51–55] and confirmed in a meta-analysis: the risk ratio for recurrent VTE was 0.9 (95% CI 0.7–1.1) [56]. Moreover, this meta-analysis demonstrated a beneficial safety profile of NOACs in terms of bleeding complications with a risk ratio for major haemorrhage of 0.6 (95% CI 0.4–0.9); however, it should be noted that the absolute risk of major haemorrhage in real-world daily care may be higher than reported in the randomized controlled trials which applied strict patient selection criteria [57]. Based on these results and the practical advantages of NOACs, these drugs are likely to replace VKAs as the treatment of choice for the majority of patients with PE in the future.

A potential concern with regard to the NOACs is the current unavailability of specific antidotes, although the relevance of this in clinical practice may be limited due to the lower incidence of major haemorrhage as well as the shorter half-life of these drugs compared to VKAs [50]. Moreover, specific antidotes are currently under clinical investigation and are likely to become available in the coming years. The humanized monoclonal antibody idarucizumab, which is specifically designed as an antidote for dabigatran, directly binds to dabigatran with a higher affinity than thrombin and is currently being

Table 5. Long-term anticoagulant therapy regimens.

Drug	Standard dose	Comments
VKA	<i>Individually based on INR</i>	- Target range INR: 2.0–3.0
<i>Oral administration</i>		
LMWH^a	<i>Enoxaparin</i>	- Treatment of choice for cancer-associated PE for at least the first 6 months, evidence for the period beyond the first 6 months is lacking and a switch to a VKA may be considered - After the first month a dose reduction to 80% of the initial dose can be considered - Contraindicated if creatinine clearance <30 mL/min
	<i>1.0 mg/kg b.i.d.</i>	
<i>Subcutaneous injection</i>	<i>1.5 mg/kg o.d.</i>	
	<i>Tinzaparin</i>	
	<i>175 U/kg o.d.</i>	
	<i>Nadroparin</i>	
	<i>86 IU/kg b.i.d.</i>	
	<i>171/kg o.d.</i>	
Dabigatran^a	150 mg b.i.d.	- At least 5 days combined with a direct-action anticoagulant therapy (i.e. LMWH) - Almost completely renal clearance - Contraindicated if creatinine clearance <30 mL/min
<i>Oral administration</i>		
Rivaroxaban^a	20 mg b.i.d.	- First 3 weeks, a higher dose of 15 mg b.i.d. - Contraindicated if creatinine clearance <30 mL/min
<i>Oral administration</i>		
Apixaban^a	5 mg b.i.d.	- First 7 days, a higher dose of 10 mg b.i.d. - Contraindicated if creatinine clearance <25 mL/min
<i>Oral administration</i>		
Edoxaban^a	60 mg o.d.	- At least 5 days combined with a direct-action anticoagulant therapy (i.e. LMWH) - Contraindicated if creatinine clearance <30 mL/min
<i>Oral administration</i>		

Note: VKA, vitamin K antagonist; LMWH, low molecular weight heparin; b.i.d., twice daily; o.d., once daily; INR, international normalized ratio; PE, pulmonary embolism. ^aApproval may vary between countries.

tested in a phase III trial (NCT02104947). Andexanet alpha is a recombinant modified form of factor Xa that directly binds factor Xa inhibitors without procoagulant activity and is a potential universal antidote for the anti-Xa inhibitors as well as for LMWH. Two phase III trials have recently been initiated to evaluate the effects of andexanet alpha in healthy volunteers (NCT02220725, NCT02207725) and a study in patients with a major haemorrhage whilst treated with a factor Xa inhibitor is planned (NCT02329327).

For the specific category of patients with cancer-associated PE, long-term LMWHs are the treatment of choice for at least the first 6 months of therapy, based on a higher efficacy [hazard ratio (HR) of recurrent VTE 0.48; 95% CI 0.30–0.77] in combination with a comparable risk of bleeding complications compared to VKA treatment [58]. Although a meta-analysis of patients with cancer included in the NOAC trials suggested a favourable efficacy and similar safety profile compared to VKAs, dedicated clinical trials comparing these drugs to LMWHs in patients with cancer are required [59]. Therefore, at present, NOACs cannot be recommended for the treatment of cancer-associated VTE. Finally, an inferior vena cava filter is a temporary alternative for patients with an absolute contraindication to anticoagulant therapy [7].

Long-term anticoagulant therapy (after the first 3 months)

To determine the optimal duration of treatment after the initial 3 months, the perceived risk of anticoagulant therapy-associated haemorrhage should be weighed against the risk of recurrent VTE in every patient individually.

It has been estimated that the risk of major haemorrhage after the initial 3 months of therapy is 2.74 per 100 patient-years, but the risk can vary widely depending on patient characteristics [60]. Known risk factors for bleeding complications are older age, previous gastrointestinal bleeding, previous stroke, chronic renal or liver disease, alcohol abuse, concomitant antiplatelet therapy, presence of serious comorbidities and poor control of anticoagulant therapy [7]. Several clinical prediction rules derived in patients with VTE as well as those derived and validated in patients with atrial fibrillation have been investigated. However, none of these prediction rules is currently validated to use in daily clinical practice as a result of low reproducibility, a low c-statistic indicating a limited predictive value and/or the lack of validation studies [61]. Moreover, it should be noted that the case-fatality rate of anticoagulant therapy-associated major haemorrhage (13.4%; 95% CI 9.4–17.4) has been reported to be higher than that of recurrent VTE (3.6%; 95% CI 1.9–5.7), and therefore, the clinical impact of major haemorrhage may be considered to be higher than that of recurrent VTE [49, 60].

Regarding the risk of recurrent VTE, it should be emphasized that extending the duration of anticoagulant therapy only postpones a potential recurrent VTE without diminishing the VTE recurrence risk after anticoagulant therapy is stopped [62]. In patients with PE related to a transient provoking factor (e.g. recent surgery, immobilization, pregnancy or oral contraceptive use), a recurrence risk of approximately 2.5% per year is usually considered low enough to discontinue anticoagulant therapy after 3 months [7, 47]. On the other hand, the recurrence risk in patients with an active malignancy (20.7% during the first 12 months of therapy) [63], the antiphospholipid syndrome [64] or a previous episode of VTE (20.7% after 4 years of follow-up after initial 6 month therapy) [65] is considered to be high enough for guidelines to suggest that these patients should be treated for an extended or indefinite duration [7].

Because most recurrences occur shortly after cessation of anticoagulant therapy, it has been proposed that patients with a second VTE that occurred after a long interval following the first VTE may have a relatively low recurrence risk and therefore may benefit from a limited duration of treatment. However, in a recent cohort study, the incidence rate of a third VTE after an interval of >1 year following cessation of treatment for the first VTE was 12 per 100 patient-years (95% CI 7.4–19.0) and 5.6 per 100 patient-years (95% CI 2.2–12.0) in patients with an unprovoked and a provoked second VTE, respectively [66]. Therefore, differentiation based on the interval between the first and second VTE cannot be recommended for patients with an unprovoked second VTE, whereas a limited duration of treatment can be considered in patients with a provoked second VTE.

In patients with a first unprovoked PE not associated with cancer or the antiphospholipid syndrome, which is associated with a recurrence risk of at least 4.5% per year, determining the optimal duration of treatment is usually more complicated. Standard screening for inherited and acquired thrombophilia is not recommended as studies have failed to demonstrate a benefit and the predictive value for a recurrent VTE of most thrombophilic factors is relatively low [67]. Measuring the D-dimer level as a general marker of coagulation activity may be valuable for predicting the recurrence risk, although results from prospective studies were inconsistent with incidence rates varying from 3.0% to 6.7% per year after anticoagulant therapy was stopped in patients with repeated negative D-dimer test results [68, 69]. The Vienna prediction model is a recently externally validated nomogram consisting of the variables sex, location of the VTE and the D-dimer concentration and is currently being evaluated in a prospective management study [70, 71]. In this ongoing study, patients with a low-risk score (<180 points; expected 12-month recurrence risk 4.4%; 95% CI 2.7–6.2) receive anticoagulant therapy for 3–7 months, whilst therapy is continued in the remaining patients (NCT01972243). The “men continue and HER DOO2 prediction rule” consists of four variables: post-thrombotic signs, D-dimer concentration of ≥ 250 $\mu\text{g/L}$ (whilst on anticoagulant therapy), body mass index ≥ 30 kg/m^2 and age ≥ 60 years. In an observational study, this score differentiated between a low-risk category with an annual risk of 1.6% (≤ 1 point; 95% CI 0.3–4.6) and a higher-risk group with an annual risk of 14.1% (≥ 2 points; 95% CI 10.9–17.3) only in women and is currently being investigated in a prospective management study (NCT00967304) [72]. Finally, the DASH score has been derived from a patient-level meta-analysis and consists of four variables: an abnormal postcoagulation D-dimer concentration, age <50 years, male sex and hormonal therapy. This score was able to differentiate between groups with an annual VTE incidence of 3.1% (95% CI 2.4–3.9) and 9.3% (95% CI 8.1–10.8) [73]. However, in the absence of results from prospective management studies, none of these scores can currently be recommended for daily clinical practice.

The major concern with the use of anticoagulant therapy for an indefinite period is the persistent risk of major haemorrhage. Therefore, alternative strategies that may be associated with a lower risk of haemorrhage have been investigated, including low-intensity VKA therapy with a target INR of 1.5–1.9 and aspirin. Low-intensity VKA therapy was less effective compared to conventional-intensity VKA therapy, with a similar risk of major haemorrhage [74]. Therefore, low-intensity VKA therapy cannot be recommended for the secondary prevention of VTE. More recently, two randomized trials have been conducted to investigate the effect of aspirin after standard anticoagulant therapy for long-term secondary VTE prevention. In the WARFASA trial, a significant reduction in VTE recurrence risk was observed from 11.2% per year for placebo to 6.6% for aspirin (OR 0.58; 95% CI 0.36–0.93), whilst the risk of major haemorrhage was similar (1/197 for

placebo vs. 1/205 for aspirin) [75]. The ASPIRE trial demonstrated only a nonsignificant decrease in the risk of VTE recurrence from 6.5% per year for placebo to 4.8% per year for aspirin (HR 0.74; 95% CI 0.52–1.05) together with a nonsignificant increase in the risk of major haemorrhage from 0.6% to 1.1% per year (HR 1.73; 95% CI 0.72–4.11) [76]. Although these results suggest that aspirin may reduce the VTE recurrence risk, it is clear that it does not provide reliable protection against recurrent VTE in comparison with long-term oral anticoagulation. Therefore, long-term secondary prevention with aspirin is not recommended by current treatment guidelines, although it may be considered as an alternative for extended treatment in patients who are unable to tolerate any type of oral anticoagulants [77].

The effects of dabigatran, apixaban and rivaroxaban on long-term secondary prevention after initial management for VTE (either DVT or PE) have also been investigated. Only dabigatran has been compared to warfarin and was shown to be noninferior (HR 1.44; 95% CI 0.78–2.64) for recurrent VTE with a nonsignificantly lower risk of major haemorrhage (HR 0.52; 95% CI 0.52–1.02) [78]. As expected, dabigatran was more effective than placebo (HR 0.08; 95% CI 0.02–0.25), whilst major haemorrhage occurred in only two of 684 patients (0.3%) versus none of 659 patients receiving placebo. Two doses of apixaban (2.5 and 5 mg twice daily) were compared to placebo for a period of 12 months after initial treatment [79]. These doses of apixaban were shown to be similarly effective with a recurrence risk of 1.7% in both groups compared to 8.8% in patients who received placebo. Remarkably, both doses were not associated with a higher risk of major haemorrhage compared to placebo. Finally, rivaroxaban for an additional 6–12 months after initial treatment demonstrated superior efficacy compared to placebo (8/602 vs. 42/594; HR 0.18; 95%CI 0.09–0.39) for recurrent VTE, whilst nonfatal major haemorrhage occurred in only four of 598 patients (0.7%) treated with rivaroxaban [80]. Although all studies were performed in highly selected groups of patients who completed initial therapy without complications and only dabigatran has been compared to VKA therapy, these findings together with the results from initial management studies suggest that NOACs are as effective as VKA therapy and have a lower risk of major haemorrhage than long-term VKA therapy. Therefore, the introduction of NOACs is likely to change the risk–benefit ratio of long-term anticoagulant therapy and may influence clinicians to extend the duration of treatment more often in selected patient groups.

CONCLUSIONS

The standardized diagnostic management of clinically suspected acute PE allows the exclusion of PE without imaging tests in approximately 30% of patients; this proportion can be further increased using the recently validated age-adjusted D-dimer threshold.

Future studies should focus on further limiting the number of required imaging tests as well as improving the implementation of standardized management in daily clinical practice, as nonadherence decreases safety and efficiency.

PE management is becoming increasingly more stratified by developing criteria to select patients who can be safely treated in the outpatient setting, and thrombolytic therapy should be reserved for patients with haemodynamic instability or those with haemodynamic deterioration during the first days after the diagnosis of PE. The introduction of NOACs simplifies anticoagulant therapy and the associated risk of major haemorrhage seems to be lower than that of VKA therapy. Nevertheless, clinical prediction rules to estimate the risk of anticoagulant therapy-associated major haemorrhage as well as clinical prediction rules to estimate the risk of recurrent VTE are required to come to a more tailored long-term management of PE.

REFERENCES

1. Naess IA, Christiansen SC, Romundstad P, Cannegieter SC, Rosendaal FR, Hammerstrom J. Incidence and mortality of venous thrombosis: a population-based study. *J Thromb Haemost* 2007; 5: 692–9.
2. Silverstein MD, Heit JA, Mohr DN, Petterson TM, O'Fallon WM, Melton LJ III. Trends in the incidence of deep vein thrombosis and pulmonary embolism: a 25-year population-based study. *Arch Intern Med* 1998; 158: 585–93.
3. Stein PD, Beemath A, Matta F et al. Clinical characteristics of patients with acute pulmonary embolism: data from PIOPED II. *Am J Med* 2007; 120: 871–9.
4. Douma RA, Mos IC, Erkens PM et al. Performance of 4 clinical decision rules in the diagnostic management of acute pulmonary embolism: a prospective cohort study. *Ann Intern Med* 2011; 154: 709–18.
5. van Belle A, Büller HR, Huisman MV et al. Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography. *JAMA* 2006; 295: 172–9.
6. Aujesky D, Obrosky DS, Stone RA et al. Derivation and validation of a prognostic model for pulmonary embolism. *Am J Respir Crit Care Med* 2005; 172: 1041–6.
7. Konstantinides SV, Torbicki A, Agnelli G et al. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J* 2014; 35: 3033–80.
8. Bounameaux H. Contemporary management of pulmonary embolism: the answers to ten questions. *J Intern Med* 2010; 268: 218–31.
9. Righini M, Van Es J, Den Exter PL et al. Age-adjusted D-dimer cutoff levels to rule out pulmonary embolism: the ADJUST-PE study. *JAMA* 2014; 311: 1117–24.
10. Kooiman J, Klok FA, Mos IC et al. Incidence and predictors of contrast-induced nephropathy following CT-angiography for clinically suspected acute pulmonary embolism. *J Thromb Haemost* 2010; 8: 409–11.
11. O'Neill J, Murchison JT, Wright L, Williams J. Effect of the introduction of helical CT on radiation dose in the investigation of pulmonary embolism. *Br J Radiol* 2005; 78: 46–50.
12. Sarma A, Heilbrun ME, Conner KE, Stevens SM, Woller SC, Elliott CG. Radiation and chest CT scan examinations: what do we know? *Chest* 2012; 142: 750–60.
13. Le Gal G, Righini M, Roy PM et al. Prediction of pulmonary embolism in the emergency department: the revised Geneva score. *Ann Intern Med* 2006; 144: 165–71.
14. Wells PS, Anderson DR, Rodger M et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000; 83: 416–20.
15. Gibson NS, Söhne M, Kruip MJ et al. Further validation and simplification of the Wells clinical decision rule in pulmonary embolism. *Thromb Haemost* 2008; 99: 229–34.
16. Klok FA, Mos IC, Nijkeuter M et al. Simplification of the revised Geneva score for assessing clinical probability of pulmonary embolism. *Arch Intern Med* 2008; 168: 2131–6.
17. Stein PD, Hull RD, Patel KC et al. D-dimer for the exclusion of acute venous thrombosis and pulmonary embolism: a systematic review. *Ann Intern Med* 2004; 140: 589–602.
18. Pasha SM, Klok FA, Snoep JD et al. Safety of excluding acute pulmonary embolism based on an unlikely clinical probability by the Wells rule and normal D-dimer concentration: a meta-analysis. *Thromb Res* 2010; 125: e123–7.


19. Douma RA, Le Gal G, Söhne M et al. Potential of an age adjusted D-dimer cut-off value to improve the exclusion of pulmonary embolism in older patients: a retrospective analysis of three large cohorts. *BMJ* 2010; 340: c1475.
20. van Beek EJ, Brouwerst EM, Song B, Stein PD, Oudkerk M. Clinical validity of a normal pulmonary angiogram in patients with suspected pulmonary embolism – a critical review. *Clin Radiol* 2001; 56: 838–42.
21. Anderson DR, Kahn SR, Rodger MA et al. Computed tomographic pulmonary angiography vs ventilation-perfusion lung scanning in patients with suspected pulmonary embolism: a randomized controlled trial. *JAMA* 2007; 298:2743–53.
22. Kruip MJ, Leclercq MG, van der Heul C, Prins MH, Büller HR. Diagnostic strategies for excluding pulmonary embolism in clinical outcome studies. A systematic review. *Ann Intern Med* 2003; 138: 941–51.
23. Mos IC, Klok FA, Kroft LJ, de Roos A, Dekkers OM, Huisman MV. Safety of ruling out acute pulmonary embolism by normal computed tomography pulmonary angiography in patients with an indication for computed tomography: systematic review and meta-analysis. *J Thromb Haemost* 2009; 7: 1491–8.
24. Van Es J, Douma RA, Schreuder SM et al. Clinical impact of findings supporting an alternative diagnosis on CT pulmonary angiography in patients with suspected pulmonary embolism. *Chest* 2013; 144: 1893–9.
25. Carrier M, Righini M, Wells PS et al. Subsegmental pulmonary embolism diagnosed by computed tomography: incidence and clinical implications. A systematic review and meta-analysis of the management outcome studies. *J Thromb Haemost* 2010; 8: 1716–22.
26. Den Exter PL, Van Es J, Klok FA et al. Risk profile and clinical outcome of symptomatic subsegmental acute pulmonary embolism. *Blood* 2013; 122: 1144–9.
27. Donato AA, Khoche S, Santora J, Wagner B. Clinical outcomes in patients with isolated subsegmental pulmonary emboli diagnosed by multidetector CT pulmonary angiography. *Thromb Res* 2010; 126: e266–70.
28. Roy PM, Meyer G, Vielle B et al. Appropriateness of diagnostic management and outcomes of suspected pulmonary embolism. *Ann Intern Med* 2006; 144: 157–64.
29. Jimenez D, Resano S, Otero R et al. Computerised clinical decision support for suspected PE. *Thorax* 2015; 70: 909–11.
30. van der Hulle T, Den Exter PL, Erkens PG et al. Variable D-dimer thresholds for diagnosis of clinically suspected acute pulmonary embolism. *J Thromb Haemost* 2013; 11: 1986–92.
31. Zondag W, Mos IC, Creemers-Schild D et al. Outpatient treatment in patients with acute pulmonary embolism: the Hestia Study. *J Thromb Haemost* 2011; 9: 1500–7.
32. Jimenez D, Aujesky D, Moores L et al. Simplification of the pulmonary embolism severity index for prognostication in patients with acute symptomatic pulmonary embolism. *Arch Intern Med* 2010; 170: 1383–9.
33. Klok FA, Mos IC, Huisman MV. Brain-type natriuretic peptide levels in the prediction of adverse outcome in patients with pulmonary embolism: a systematic review and meta-analysis. *Am J Respir Crit Care Med* 2008; 178: 425–30.
34. Becattini C, Vedovati MC, Agnelli G. Prognostic value of troponins in acute pulmonary embolism: a meta-analysis. *Circulation* 2007; 116: 427–33.
35. Coutance G, Cauderlier E, Ehtisham J, Hamon M, Hamon M. The prognostic value of markers of right ventricular dysfunction in pulmonary embolism: a meta-analysis. *Crit Care* 2011; 15: R103.

36. Trujillo-Santos J, Den Exter PL, Gomez V et al. Computed tomography-assessed right ventricular dysfunction and risk stratification of patients with acute non-massive pulmonary embolism: systematic review and meta-analysis. *J Thromb Haemost* 2013; 11: 1823–32.
37. Becattini C, Casazza F, Forgione C et al. Acute pulmonary embolism: external validation of an integrated risk stratification model. *Chest* 2013; 144: 1539–45.
38. Vanni S, Nazerian P, Pepe G et al. Comparison of two prognostic models for acute pulmonary embolism: clinical vs. right ventricular dysfunction-guided approach. *J Thromb Haemost* 2011; 9: 1916–23.
39. Meneveau N, Seronde MF, Blonde MC et al. Management of unsuccessful thrombolysis in acute massive pulmonary embolism. *Chest* 2006; 129: 1043–50.
40. Chatterjee S, Chakraborty A, Weinberg I et al. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA* 2014; 311: 2414–21.
41. Stein PD, Matta F. Thrombolytic therapy in unstable patients with acute pulmonary embolism: saves lives but underused. *Am J Med* 2012; 125: 465–70.
42. Engelberger RP, Kucher N. Ultrasound-assisted thrombolysis for acute pulmonary embolism: a systematic review. *Eur Heart J* 2014; 35: 758–64.
43. Riera-Mestre A, Becattini C, Giustozzi M, Agnelli G. Thrombolysis in hemodynamically stable patients with acute pulmonary embolism: a meta-analysis. *Thromb Res* 2014; 134: 1265–71.
44. Meyer G, Vicaut E, Danays T et al. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med* 2014; 370: 1402–11.
45. Aujesky D, Roy PM, Verschuren F et al. Outpatient versus inpatient treatment for patients with acute pulmonary embolism: an international, open-label, randomised, non-inferiority trial. *Lancet* 2011; 378: 41–8.
46. Agterof MJ, Schutgens RE, Snijder RJ et al. Out of hospital treatment of acute pulmonary embolism in patients with a low NT-proBNP level. *J Thromb Haemost* 2010; 8: 1235–41.
47. Kearon C, Akl EA, Comerota AJ et al. Antithrombotic therapy for VTE disease: antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012; 141(2 Suppl): e419S–94S.
48. Warkentin TE, Maurer BT, Aster RH. Heparin-induced thrombocytopenia associated with fondaparinux. *N Engl J Med* 2007; 356: 2653–5.
49. Carrier M, Le Gal G, Wells PS, Rodger MA. Systematic review: case-fatality rates of recurrent venous thromboembolism and major bleeding events among patients treated for venous thromboembolism. *Ann Intern Med* 2010; 152: 578–89.
50. Ahrens I, Lip GY, Peter K. New oral anticoagulant drugs in cardiovascular disease. *Thromb Haemost* 2010; 104: 49–60.
51. Agnelli G, Büller HR, Cohen A et al. Oral apixaban for the treatment of acute venous thromboembolism. *N Engl J Med* 2013; 369: 799–808.
52. Büller HR, Prins MH, Lensin AW et al. Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. *N Engl J Med* 2012; 366: 1287–97.
53. Büller HR, Decousus H, Grosso MA et al. Edoxaban versus warfarin for the treatment of symptomatic venous thromboembolism. *N Engl J Med* 2013; 369: 1406–15.
54. Schulman S, Kearon C, Kakkar AK et al. Dabigatran versus warfarin in the treatment of acute venous thromboembolism. *N Engl J Med* 2009; 361: 2342–52.
55. Schulman S, Kakkar AK, Goldhaber SZ et al. Treatment of acute venous thromboembolism with dabigatran or warfarin and pooled analysis. *Circulation* 2014; 129: 764–72.

56. van der Hulle T, Kooiman J, Den Exter PL, Dekkers OM, Klok FA, Huisman MV. Effectiveness and safety of novel oral anticoagulants as compared with vitamin K antagonists in the treatment of acute symptomatic venous thromboembolism: a systematic review and meta-analysis. *J Thromb Haemost* 2014; 12: 320–8.
57. Beyer-Westendorf J, Forster K, Pannach S et al. Rates, management, and outcome of rivaroxaban bleeding in daily care: results from the Dresden NOAC registry. *Blood* 2014; 124: 955–62.
58. Lee AY, Levine MN, Baker RI et al. Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. *N Engl J Med* 2003; 349: 146–53.
59. van der Hulle T, Den Exter PL, Kooiman J, van der Hoeven JJ, Huisman MV, Klok FA. Meta-analysis of the efficacy and safety of new oral anticoagulants in patients with cancer-associated acute venous thromboembolism. *J Thromb Haemost* 2014; 12: 1116–20.
60. Linkins LA, Choi PT, Douketis JD. Clinical impact of bleeding in patients taking oral anticoagulant therapy for venous thromboembolism: a meta-analysis. *Ann Intern Med* 2003; 139: 893–900.
61. Klok FA, Kooiman J, Huisman MV, Konstantinides S, Lankeit M. Predicting anticoagulant-related bleeding in patients with venous thromboembolism: a clinically oriented review. *Eur Respir J* 2015; 45: 201–10.
62. van Dongen CJ, Vink R, Hutten BA, Büller HR, Prins MH. The incidence of recurrent venous thromboembolism after treatment with vitamin K antagonists in relation to time since first event: a meta-analysis. *Arch Intern Med* 2003; 163:1285–93.
63. Prandoni P, Lensing AW, Piccioli A et al. Recurrent venous thromboembolism and bleeding complications during anticoagulant treatment in patients with cancer and venous thrombosis. *Blood* 2002; 100: 3484–8.
64. Khamashta MA, Cuadrado MJ, Mujic F, Taub NA, Hunt BJ, Hughes GR. The management of thrombosis in the antiphospholipid-antibody syndrome. *N Engl J Med* 1995; 332: 993–7.
65. Schulman S, Granqvist S, Holmstrom M et al. The duration of oral anticoagulant therapy after a second episode of venous thromboembolism. The Duration of Anticoagulation Trial Study Group. *N Engl J Med* 1997; 336: 393–8.
66. van der Hulle T, Tan M, Den Exter PL et al. Recurrence risk after anticoagulant treatment of limited duration for late, second venous thromboembolism. *Haematologica* 2015; 100: 188–93.
67. Kyrle PA, Eichinger S. Clinical scores to predict recurrence risk of venous thromboembolism. *Thromb Haemost* 2012; 108: 1061–4.
68. Kearon C, Spencer FA, O’Keeffe D et al. D-dimer testing to select patients with a first unprovoked venous thromboembolism who can stop anticoagulant therapy: a cohort study. *Ann Intern Med* 2015; 162: 27–34.
69. Palareti G, Cosmi B, Legnani C et al. D-dimer to guide the duration of anticoagulation in patients with venous thromboembolism: a management study. *Blood* 2014; 124: 196–203.
70. Eichinger S, Heinze G, Jandeck LM, Kyrle PA. Risk assessment of recurrence in patients with unprovoked deep vein thrombosis or pulmonary embolism: the Vienna prediction model. *Circulation* 2010; 121: 1630–6.
71. Marcucci M, Iorio A, Douketis JD et al. Risk of recurrence after a first unprovoked venous thromboembolism: external validation of the Vienna Prediction Model using pooled individual patient data. *J Thromb Haemost* 2015; 13:775–81.
72. Rodger MA, Kahn SR, Wells PS et al. Identifying unprovoked thromboembolism patients at low risk for recurrence who can discontinue anticoagulant therapy. *CMAJ* 2008; 179: 417–26.

73. Tosetto A, Iorio A, Marcucci M et al. Predicting disease recurrence in patients with previous unprovoked venous thromboembolism: a proposed prediction score (DASH). *J Thromb Haemost* 2012; 10: 1019–25.
74. Kearon C, Ginsberg JS, Kovacs MJ et al. Comparison of low-intensity warfarin therapy with conventional-intensity warfarin therapy for long-term prevention of recurrent venous thromboembolism. *N Engl J Med* 2003; 349: 631–9.
75. Becattini C, Agnelli G, Schenone A et al. Aspirin for preventing the recurrence of venous thromboembolism. *N Engl J Med* 2012; 366: 1959–67.
76. Brighton TA, Eikelboom JW, Mann K et al. Low-dose aspirin for preventing recurrent venous thromboembolism. *N Engl J Med* 2012; 367: 1979–87.
77. Konstantinides S, Torbicki A. Management of venous thrombo-embolism: an update. *Eur Heart J* 2014; 35: 2855–63.
78. Schulman S, Kearon C, Kakkar AK et al. Extended use of dabigatran, warfarin, or placebo in venous thromboembolism. *N Engl J Med* 2013; 368: 709–18.
79. Agnelli G, Büller HR, Cohen A et al. Apixaban for extended treatment of venous thromboembolism. *N Engl J Med* 2013; 368: 699–708.
80. Bauersachs R, Berkowitz SD, Brenner B et al. Oral rivaroxaban for symptomatic venous thromboembolism. *N Engl J Med* 2010; 363: 2499–510.





PART 1

DIAGNOSTIC MANAGEMENT OF ACUTE PULMONARY EMBOLISM

