



## Impact of allosteric modulation: Exploring the binding kinetics of glutamate and other orthosteric ligands of the metabotropic glutamate receptor 2

Maarten L.J. Doornbos<sup>a</sup>, Sophie C. Vermond<sup>a</sup>, Hilde Lavreysen<sup>b</sup>, Gary Tresadern<sup>b</sup>,  
Adriaan P. IJzerman<sup>a</sup>, Laura H. Heitman<sup>a,\*</sup>

<sup>a</sup> Division of Drug Discovery and Safety, Leiden Academic Centre for Drug Research (LACDR), Leiden University, P.O. Box 9502, 2300RA Leiden, The Netherlands

<sup>b</sup> Janssen Research and Development, Turnhoutseweg 30, 2340 Beerse, Belgium

### ARTICLE INFO

#### Keywords:

mGlu<sub>2</sub> receptor  
Binding kinetics  
Glutamate  
Orthosteric  
Allosteric modulation

### ABSTRACT

While many orthosteric ligands have been developed for the mGlu<sub>2</sub> receptor, little is known about their target binding kinetics and how these relate to those of the endogenous agonist glutamate. Here, the kinetic rate constants, i.e.  $k_{on}$  and  $k_{off}$ , of glutamate were determined for the first time followed by those of the synthetic agonist LY354740 and antagonist LY341495. To increase the understanding of the binding mechanism and impact of allosteric modulation thereon, kinetic experiments were repeated in the presence of allosteric modulators. Functional assays were performed to further study the interplay between the orthosteric and allosteric binding sites, including an impedance-based morphology assay.

We found that dissociation rate constants of orthosteric mGlu<sub>2</sub> ligands were all within a small 6-fold range, whereas association rate constants were ranging over more than three orders of magnitude and correlated to both affinity and potency. The latter showed that target engagement of orthosteric mGlu<sub>2</sub> ligands is  $k_{on}$ -driven *in vitro*. Moreover, only the off-rates of the two agonists were decreased by a positive allosteric modulator (PAM), thereby increasing their affinity. Interestingly, a PAM increased the duration of a glutamate-induced cellular response. A negative allosteric modulator (NAM) increased both on- and off-rate of glutamate without changing its affinity, while it did not affect these parameters for LY354740, indicating probe-dependency.

In conclusion, we found that affinity- or potency-based orthosteric ligand optimization primarily results in ligands with high  $k_{on}$  values. Moreover, positive allosteric modulators alter the binding kinetics of orthosteric agonists mainly by decreasing  $k_{off}$ , which we were able to correlate to a lengthened cellular response. Together, this study shows the importance of studying binding kinetics in early drug discovery, as this may provide important insights towards improved efficacy *in vivo*.

### 1. Introduction

Glutamate is the most important excitatory neurotransmitter in the central nervous system where it modulates synaptic responses via activation of ionotropic glutamate receptors and metabotropic glutamate (mGlu) receptors [1]. mGlu receptors are class C G protein-coupled receptors (GPCRs) that structurally consist of a large extracellular glutamate binding domain, the so-called Venus Flytrap (VFT) domain, which is connected via a cysteine-rich domain to the typical seven-transmembrane (7TM) domain [2]. The mGlu<sub>2</sub> receptor, which is expressed presynaptically in the periphery of the synapse, is of interest in drug discovery as it negatively modulates the release of glutamate into the synapse [3]. Hence, mGlu<sub>2</sub> receptor activation can reduce glutamate hyperfunction in diseases like schizophrenia and anxiety [4,5],

whereas mGlu<sub>2</sub> receptor blockade can be beneficial for glutamate hypofunction in depression and impaired cognition [6,7].

A variety of glutamate-like ligands targeting the orthosteric binding site in the VFT domain was developed including the reference agonist LY354740 and antagonist LY341495 [8,9]. Until the recent disclosure of the mGlu<sub>2</sub> selective agonist LY2812223 [10], development of orthosteric ligands presented challenges for receptor subtype selectivity and therefore discovery efforts shifted to allosteric modulators that bind in a less conserved pocket in the 7TM domain [11]. Allosteric modulators enhance or inhibit the potency and/or efficacy of the endogenous/orthosteric agonist with little or no intrinsic activity [12]. Two positive allosteric modulators (PAMs) have advanced into clinical trials: AZD8529 [13,14] and JNJ-40411813/ADX71149 [15,16]. Reference PAMs in the field include BINA [17], JNJ-40068782 [18] and

\* Corresponding author.

E-mail address: [l.h.heitman@lacdr.leidenuniv.nl](mailto:l.h.heitman@lacdr.leidenuniv.nl) (L.H. Heitman).

<https://doi.org/10.1016/j.bcp.2018.07.014>

Received 9 May 2018; Accepted 14 July 2018

Available online 17 July 2018

0006-2952/ © 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

JNJ-46281222 [19]. A number of negative allosteric modulators (NAMs) have been characterized *in vivo*, including a recent series of Janssen, RO4491533 and decolurant of which the latter has advanced into clinical trials [7,20–22].

Over the last decade it has become increasingly clear that *in vivo* efficacy is not only depending on optimized *in vitro* affinity and efficacy parameters, but also on optimized kinetics of both receptor binding and activation [23]. The dissociation rate constant  $k_{off}$  and its derived parameter residence time ( $RT = 1/k_{off}$ ) have received increasing attention since the *in vivo* efficacy of multiple marketed GPCR drugs was shown to be related to long RT at the target, such as the long-acting M<sub>3</sub> receptor antagonist tiotropium [24,25]. Although most kinetic studies have emphasized dissociation rate constants, association rate constants have also been described to be important for fast onset of drug action, a high receptor occupancy and even a longer duration of action [26,27]. The importance of  $k_{on}$  was further underscored for its potential in drug safety by Sykes *et al.* (2017) who showed that extrapyramidal side effects induced by dopamine D<sub>2</sub> receptor antagonists are linked to  $k_{on}$  rather than  $k_{off}$  as had been the general hypothesis so far [28].

We have recently shown the importance of  $k_{on}$  and  $k_{off}$  for mGlu<sub>2</sub> PAMs, where their affinity was  $k_{on}$ -driven and their  $k_{off}$  was linked to *in vivo* efficacy [29]. Except for this study, no previous studies have extensively focused on binding kinetics of mGlu<sub>2</sub> ligands. Understanding of binding kinetics may also be helpful in drug discovery of novel orthosteric mGlu<sub>2</sub> ligands. Moreover, appreciating the kinetic binding parameters of the endogenous agonist is important when designing orthosteric ligands as they have to compete for the same binding site [30]. Furthermore, determination of alterations of the kinetic parameters of the endogenous ligand induced by an allosteric modulator will provide valuable mechanistic insights. Since kinetic binding parameters of the endogenous mGlu<sub>2</sub> agonist glutamate have not been quantified before we set up a kinetic radioligand binding assay to enable quantification of kinetic parameters for orthosteric ligands. The major mechanism of action of many allosteric modulators is to modulate endogenous agonist affinity by affecting its kinetic binding parameters, as affinity is determined by  $K_D = k_{off}/k_{on}$ . Therefore, kinetic binding experiments with orthosteric ligands were also performed in the presence of a PAM or NAM. Additionally, we performed functional assays, i.e. [<sup>35</sup>S]GTPγS binding assays measuring G protein activation and a label-free biosensor assay that measures changes in cell morphology representing a more integral cellular response. These functional assays were used to further study the interplay between the orthosteric and allosteric binding sites, and its effect on the level of functional efficacy. Importantly, the cell morphology assay enabled recording in real-time, thereby providing the opportunity to evaluate functional receptor-induced responses over time in addition to the time-dependent binding kinetics assays.

This work provides insights on the binding kinetics of orthosteric ligands at the mGlu<sub>2</sub> receptor and modulation thereof by PAMs or NAMs. As such, it contributes to increased molecular understanding which may strengthen future drug discovery projects focusing on the development of both orthosteric ligands or allosteric modulators for the mGlu<sub>2</sub> receptor as well as for other GPCRs.

## 2. Materials and methods

### 2.1. Chemicals and reagents

LY354740, JNJ-46281222, JNJ-40068782, BINA, RO4491533, and [<sup>3</sup>H]JNJ-46281222 were synthesized at Janssen Research and Development. LY341495 was from Tocris BioScience (Bristol, UK). [<sup>3</sup>H]LY341495 was obtained from American Radiolabeled Companies (St. Louis, MO, USA) and [<sup>35</sup>S]GTPγS from PerkinElmer (Groningen, The Netherlands). Dulbecco's modified Eagle's medium (DMEM), glutamate, GDP and Glutamate-pyruvate transaminase (GPT) were from Sigma Aldrich (St. Louis, MO, USA). Penicillin, streptomycin, L-Proline

and G418 were obtained from Duchefa Biochemie (Haarlem, The Netherlands). Fetal calf serum (FCS) was from Biowest (Nuaille, France). CHO-K1 cells stably expressing the wildtype (WT) hmGlu<sub>2</sub> receptor (CHO-K1\_hmGlu<sub>2</sub>) were from Janssen Research and Development. Other chemicals were from standard commercial sources.

### 2.2. Cell culture

CHO-K1\_hmGlu<sub>2</sub> cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) fetal calf serum, 200 IU·mL<sup>-1</sup> penicillin, 200 μg·mL<sup>-1</sup> streptomycin, 30.5 μg·mL<sup>-1</sup> L-proline and 400 μg·mL<sup>-1</sup> G418 at 37 °C and 5% CO<sub>2</sub>. Cells were subcultured twice every week at a ratio of 1:10.

### 2.3. Membrane preparation

Membrane preparation was performed as previously described [19].

### 2.4. Radioligand binding assays

#### 2.4.1. [<sup>3</sup>H]LY341495 binding assays

Membranes were thawed and subsequently homogenized using an Ultra Turrax homogenizer at 24,000 rpm (IKA-Werke GmbH & Co.KG, Staufen, Germany). Assay buffer (50 mM Tris-HCl pH 7.4, 2 mM CaCl<sub>2</sub>, 10 mM MgCl<sub>2</sub>) was used to dilute the samples to a total reaction volume of 100 μl containing 5 μg membrane protein and 4 nM [<sup>3</sup>H]LY341495. Incubations were performed at 0 °C. Nonspecific binding was determined using 1 mM glutamate and DMSO concentrations were ≤ 0.25%.

Displacement experiments were carried out using radioligand and a competing ligand at multiple concentrations. Samples were incubated for 60 min, after which receptor-bound radioactivity was determined.

Association experiments were performed by incubation of radioligand in the absence or presence of allosteric modulator at 1 μM with membrane aliquots. The amount of receptor-bound radioligand was determined at different time points up to 60 min.


Dissociation experiments were carried out by a 60 min pre-incubation of radioligand and membrane aliquots in the absence or presence of allosteric modulator at 1 μM. The amount of receptor-bound radioligand was determined after dissociation at different time points up to 60 min which was initiated by addition of 5 μl assay buffer containing LY341495 (final concentration 10 μM).

Competition association experiments were performed by incubation of radioligand, competing ligand at its IC<sub>50</sub> concentration in the absence or presence of allosteric modulator at 1 μM with membrane aliquots. The amount of receptor-bound radioligand was determined at different time points up to 60 min.

For all assays, incubations were terminated by rapid filtration over GF/C filter plates (PerkinElmer) using a PerkinElmer 96w Filtermate harvester. Subsequently filters were washed five times using ice-cold wash buffer (50 mM Tris-HCl pH 7.4). Filter-bound reactivity was determined using liquid scintillation spectrometry on a Microbeta 2450<sup>2</sup> microplate counter (PerkinElmer).

#### 2.4.2. [<sup>35</sup>S]GTPγS binding assays

Membrane homogenates (10 μg) were diluted in assay buffer (50 mM Tris-HCl pH 7.4, 100 mM NaCl, 3 mM MgCl<sub>2</sub>) supplemented with 5 μg saponin and 10 μM GDP to a total volume of 80 μl containing increasing concentrations of ligand of interest in the absence or presence of a glutamate concentration equivalent to its EC<sub>80</sub> value (60 μM), if indicated. Basal and maximal receptor activity was determined in the presence of only assay buffer or 1 mM glutamate, respectively. After a 30 min pre-incubation at 25 °C, 20 μl [<sup>35</sup>S]GTPγS (final concentration 0.3 nM) was added. Reactions were stopped after 90 min incubation at 25 °C. Filtration was performed and filter-bound radioactivity determined as described under '<sup>3</sup>H]LY341495 binding assays' except that



**Fig. 1.** Affinity of orthosteric ligands and the effect of allosteric modulators thereon. [<sup>3</sup>H]LY341495 displacement by orthosteric agonists A) glutamate and B) LY354740 and antagonist C) LY341495 in the absence or presence of PAMs or NAM. D) [<sup>3</sup>H]LY341495 binding in the presence of increasing concentrations of the PAM JNJ-46281222 or NAM RO4491533.  $K_i$  values obtained from these graphs are described in Table 1. Data represent the mean  $\pm$  SEM of three individual experiments performed in duplicate. If not shown, error bars are within the symbol.

GF/B filter plates were used and the wash-buffer consisted of 50 mM TRIS-HCl and 5 mM MgCl<sub>2</sub>.

## 2.5. Impedance-based morphology assays

Impedance-based morphology assays were performed using the real-time cell analyser (RTCA) xCELLigence SP system (ACEA Biosciences, San Diego, CA, USA) [31,32], as previously described [33]. The system measures electrical impedance generated by adherence of cells to gold-coated electrodes at the bottom of 96 wells PET E-plates (obtained from Bioké, Leiden, the Netherlands). Changes in impedance (Z) are measured continuously and are displayed as Cell Index (CI), which is defined as  $(Z_i - Z_0) \Omega/15 \Omega$ .  $Z_i$  is the impedance at a given time and  $Z_0$  is the baseline impedance measured at the start of the experiment in the absence of cells. Baseline impedance was determined using 45  $\mu$ l culture medium (as described under ‘cell culture’) per well in a 96 well E-plate. 40,000 CHO-K1\_hmGlu<sub>2</sub> cells were added in a volume of 50  $\mu$ l per well. After resting at room temperature for 30 min, the plate was mounted in the recording station within a humidified 37 °C, 5% CO<sub>2</sub> incubator. Impedance was measured every 15 min overnight. 18 h after cell seeding, wells were stimulated with increasing concentrations of glutamate in the absence or presence of 1  $\mu$ M PAM JNJ-46281222, resulting in final well volumes of 100  $\mu$ l. DMSO concentrations were 0.025% and constant between wells.

## 2.6. Data analysis

Data analyses were performed using Prism 7.00 (GraphPad software, San Diego, CA, USA).  $pIC_{50}$  values in radioligand displacement assays were obtained by non-linear regression curve fitting into a sigmoidal concentration-response curve using the equation:  $Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{-(X - \text{LogIC}_{50})})$ .  $pK_i$  values were obtained from  $pIC_{50}$  values using the Cheng-Prusoff equation [34]. Dissociation rate constant  $k_{off}$  was obtained using an exponential decay analysis of radioligand binding. Association rate constant  $k_{on}$  was determined using the equation  $k_{on} = (k_{obs} - k_{off}) / [L]$ , in which L is the concentration of radioligand and  $k_{obs}$  was determined using an exponential association analysis of radioligand binding.

Association and dissociation rate constants for unlabelled mGlu<sub>2</sub>

PAMs were determined by nonlinear regression analysis of competition association data as described by Motulsky and Mahan [35].

$$\begin{aligned}
 K_A &= k_1 [L] \cdot 10^{-9} + k_2 \\
 K_B &= k_3 [I] \cdot 10^{-9} + k_4 \\
 S &= \sqrt{(K_A - K_B)^2 + 4 \cdot k_1 \cdot k_3 \cdot L \cdot I \cdot 10^{-18}} \\
 K_F &= 0.5(K_A + K_B + S) \\
 K_S &= 0.5(K_A + K_B - S) \\
 Q &= \frac{B_{\max} \cdot k_1 \cdot L \cdot 10^{-9}}{K_F - K_S} \\
 Y &= Q \cdot \left( \frac{k_4 \cdot (K_F - K_S)}{K_F \cdot K_S} + \frac{k_4 - K_F}{K_F} e^{(-K_F \cdot X)} - \frac{k_4 - K_S}{K_S} e^{(-K_S \cdot X)} \right)
 \end{aligned}$$

$pEC_{50}$  and  $pIC_{50}$  values in the [<sup>35</sup>S]GTP $\gamma$ S binding assays were determined using non-linear regression curve fitting into a sigmoidal concentration-response curve using the equation:  $Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{-(\text{LogEC}_{50} - X) \cdot \text{Hill Slope}})$ . The same equation was used to determine  $pEC_{50}$  values from the impedance-based morphology assay. Baseline-corrected  $\Delta$ CI levels at indicated time points were used to obtain these concentration-response curves. Data are shown as mean  $\pm$  SEM of at least three individual experiments performed in duplicate. Statistical analyses were performed as indicated. If p-values were below 0.05, observed differences were considered statistically significant.

## 3. Results

### 3.1. Affinity of orthosteric ligands in the absence or presence of allosteric modulators

The affinities of orthosteric agonists glutamate and LY354740 and antagonist LY341495 were determined by [<sup>3</sup>H]LY341495 displacement assays in the absence and presence of PAMs JNJ-46281222 (1  $\mu$ M), BINA (10  $\mu$ M) and JNJ-40068782 (1  $\mu$ M) or NAM RO4491533 (1  $\mu$ M) (Fig. 1A–C, Table 1). The affinity of both the endogenous agonist glutamate and the synthetic agonist LY354740 was significantly increased in the presence of all PAMs. Specifically, in presence of JNJ-46281222, glutamate affinity was increased by 7-fold ( $pK_i$   $4.52 \pm 0.04$  to  $5.40 \pm 0.08$  in the absence and presence of JNJ-46281222, respectively), while the affinity of LY354740 was increased by 5-fold ( $pK_i$

**Table 1**

Affinity ( $pK_i$ ) of orthosteric ligands in the absence and presence of allosteric modulators determined in [ $^3$ H]LY341495 displacement assays.

	Glutamate	LY354740	LY341495
Vehicle	4.52 ± 0.04	6.79 ± 0.01	8.39 ± 0.09
+ 1 μM JNJ-46281222	5.40 ± 0.08***	7.51 ± 0.15*	8.54 ± 0.02
+ 10 μM BINA	5.30 ± 0.09***	7.38 ± 0.03*	8.57 ± 0.08
+ 1 μM JNJ-40068782	5.23 ± 0.04***	7.43 ± 0.19*	8.59 ± 0.05
+ 1 μM RO4491533	4.56 ± 0.03	6.93 ± 0.15	8.65 ± 0.10

Values represent the mean ± SEM of three individual experiments performed in duplicate. Statistical analyses were performed using a one-way ANOVA with Dunnett's post-test. \* < 0.05, \*\*\* < 0.001.

6.79 ± 0.01 to 7.51 ± 0.15 in the absence and presence of JNJ-46281222, respectively). The affinity of antagonist LY341495 ( $pK_i$  8.39 ± 0.09) was not affected by any of the PAMs. The presence of 1 μM NAM RO4491533 did not affect the affinity of glutamate, LY354740 or LY341495, which shows that the presence of the NAM does not inhibit the orthosteric agonists and antagonist from binding to the receptor. Since JNJ-46281222 induced the highest shift in agonist affinity it was used as representative PAM for further experimentation, along with RO4491533 as a representative NAM. Of note, both JNJ-46281222 and RO4491533 were not able to displace the radiolabelled orthosteric antagonist [ $^3$ H]LY341495 by themselves (Fig. 1D), which indicates that they bind to an allosteric site at the mGlu<sub>2</sub> receptor.

### 3.2. Binding kinetics of orthosteric ligands

We set-up radioligand binding assays allowing the determination of binding kinetics of orthosteric ligands in the absence or presence of allosteric modulators. The kinetic binding parameters  $k_{on}$  and  $k_{off}$  of [ $^3$ H]LY341495 were determined using classical (direct) association and dissociation assays (Fig. 2A, Table 2). [ $^3$ H]LY341495 associated rapidly to the mGlu<sub>2</sub> receptor, where complete association was reached within 5 min, resulting in a  $k_{obs}$  value of  $0.015 ± 0.0007 s^{-1}$ . Dissociation induced by 10 μM unlabelled LY341495 yielded a  $k_{off}$  value of  $0.0066 ± 0.0001 s^{-1}$ . Based on these  $k_{obs}$  and  $k_{off}$  values and the concentration of the radioligand used,  $k_{on}$  was calculated as

**Table 2**

Kinetic binding parameters ( $k_{on}$ ,  $k_{off}$ ) of antagonist LY341495 in the absence or presence of JNJ-46281222 (PAM) or RO4491533 (NAM), determined by direct [ $^3$ H]LY341495 association and dissociation assays.


Assay	$K_D$ (nM) <sup>a</sup>	$k_{on}$ ( $M^{-1} s^{-1}$ )	$k_{off}$ ( $s^{-1}$ )
LY341495	2.9 ± 0.23	$(2.2 ± 0.17) × 10^6$	0.0066 ± 0.0001
+ 1 μM PAM	2.9 ± 0.24	$(2.2 ± 0.67) × 10^6$	0.0063 ± 0.0003
+ 1 μM NAM	3.2 ± 0.17	$(2.1 ± 0.65) × 10^6$	0.0068 ± 0.0002

<sup>a</sup> Kinetic  $K_D$  values, defined by  $K_D = k_{off}/k_{on}$ . Values represent the mean ± SEM of three individual experiments performed in duplicate.

$2.2 ± 0.17 × 10^6 M^{-1} s^{-1}$ . To obtain kinetic binding parameters for unlabelled orthosteric ligands, a competition association assay was performed (Fig. 2B–D). The assay was first validated using unlabelled LY341495, for which  $k_{on}$  and  $k_{off}$  values were comparable to those obtained in the classical association and dissociation assays (Table 3). Subsequently, the kinetic binding parameters of glutamate and LY354740 were assessed (Fig. 2C,D; Table 3). Compared to LY341495, glutamate had a significantly lower  $k_{on}$  value of  $1.6 ± 0.3 × 10^3 M^{-1} s^{-1}$  and a faster dissociation rate ( $k_{off}$   $0.036 ± 0.008 s^{-1}$ ). Interestingly, LY354740 had a  $k_{off}$  value similar to that of glutamate, but a 40-fold higher  $k_{on}$  value ( $k_{off}$   $0.045 ± 0.010 min^{-1}$  and  $k_{on}$   $7.1 ± 2.9 × 10^4 M^{-1} s^{-1}$ , respectively). Of note, the  $K_D$  values, i.e. calculated based on their  $k_{on}$  and  $k_{off}$  values, for glutamate, LY354740 and LY341495, were in good agreement with the  $K_i$  values obtained from equilibrium displacement assays (compare Tables 1 and 3).

### 3.3. The effect of allosteric modulators on the binding kinetics of orthosteric ligands

The kinetic binding parameters  $k_{on}$  and  $k_{off}$  of glutamate, LY354740 and LY341495 were determined in the presence of PAM JNJ-46281222 or NAM RO4491533 at 1 μM (Fig. 2B–D; Tables 2 and 3). The  $k_{on}$  value for glutamate in the presence of 1 μM JNJ-46281222 was slightly but not significantly increased to  $2.4 ± 0.18 × 10^3 M^{-1} s^{-1}$  compared to its value obtained in the absence of PAM. The  $k_{off}$  value on the other hand was significantly decreased by 3-fold to  $0.012 ± 0.001 min^{-1}$ .



**Fig. 2.** Binding kinetics of orthosteric ligands and the effects of allosteric modulators thereon. A) Association and dissociation kinetics of 4 nM [ $^3$ H]LY341495 at the mGlu<sub>2</sub> receptor at 0 °C. Competition association of the agonists B) glutamate and C) LY354740 and D) antagonist LY341495 in the absence or presence of PAM JNJ-46281222 or NAM RO4491533 at 1 μM. Parameters obtained from these graphs are described in Tables 2 and 3. Data represent the mean ± SEM of at least three individual experiments performed in duplicate.

**Table 3**

Kinetic binding parameters ( $k_{on}$ ,  $k_{off}$ ) for agonists glutamate and LY354740 and antagonist LY341495 in the absence or presence of JNJ-46281222 (PAM) or RO4491533 (NAM) obtained from competition association assays using [ $^3$ H]LY341495.

	$K_D$ (nM) <sup>a</sup> (pK <sub>D</sub> )	$k_{on}$ (M <sup>-1</sup> s <sup>-1</sup> )	$k_{off}$ (s <sup>-1</sup> )
Glutamate	23000 ± 6900 (4.65)	(1.6 ± 0.3) × 10 <sup>3</sup>	0.036 ± 0.008
+ 1 μM PAM	4700 ± 600 <sup>*</sup> (5.33)	(2.4 ± 0.18) × 10 <sup>3</sup>	0.012 ± 0.001 <sup>*</sup>
+ 1 μM NAM	23500 ± 2400 (4.63)	(5.2 ± 0.37) × 10 <sup>3</sup> ***	0.12 ± 0.009***
LY354740	630 ± 290 (6.20)	(7.1 ± 2.9) × 10 <sup>4</sup>	0.045 ± 0.010
+ 1 μM PAM	180 ± 20 (6.74)	(7.3 ± 0.56) × 10 <sup>4</sup>	0.013 ± 0.001 <sup>*</sup>
+ 1 μM NAM	550 ± 200 (6.26)	(8.5 ± 2.1) × 10 <sup>4</sup>	0.047 ± 0.013
LY341495	3.5 ± 0.46 (8.45)	(2.4 ± 0.18) × 10 <sup>6</sup>	0.0087 ± 0.0009
+ 1 μM PAM	2.8 ± 0.40 (8.56)	(3.5 ± 0.46) × 10 <sup>6</sup>	0.0096 ± 0.0006
+ 1 μM NAM	3.4 ± 1.2 (8.46)	(3.9 ± 0.93) × 10 <sup>6</sup>	0.013 ± 0.003

<sup>a</sup> Kinetic  $K_D$  values, defined by  $K_D = k_{off}/k_{on}$ . Values represent the mean ± SEM of three individual experiments performed in duplicate. Statistical analyses were performed using a one-way ANOVA with Dunnett's post-test. \* < 0.05, \*\* < 0.01, \*\*\* < 0.001.

Together, this resulted in an approximately 5-fold increased 'kinetic' affinity ( $K_D$ ) for glutamate from 23 ± 6.9 to 4.7 ± 0.6 μM, which was also observed in the equilibrium displacement assays ( $K_i$  values, compare Tables 1 and 3). In the presence of 1 μM RO4491533, the  $k_{on}$  value of glutamate was significantly increased by 3-fold to 5.2 ± 0.37 × 10<sup>3</sup> M<sup>-1</sup> s<sup>-1</sup>, while the  $k_{off}$  value was also significantly increased by 3-fold to 0.12 ± 0.009 s<sup>-1</sup>. As a result, the kinetic  $K_D$  value did not change compared to glutamate in the absence of NAM (24 ± 2.4 μM and 23 ± 6.9 μM in absence or presence of NAM, respectively).

The  $k_{on}$  value of LY354740 was left unchanged in the presence of 1 μM JNJ-46281222 (7.3 ± 0.56 × 10<sup>4</sup> M<sup>-1</sup> s<sup>-1</sup>), whereas the  $k_{off}$  value was decreased by 3-fold to 0.013 ± 0.001 s<sup>-1</sup>. Together this led to an increased kinetic  $K_D$  value of 180 ± 20 nM of the agonist LY354740 in the presence of PAM JNJ-46281222. Interestingly, in contrast to glutamate, no significant shifts in  $k_{on}$  or  $k_{off}$  values were seen for the agonist LY354740 in the presence of 1 μM RO4491533, and therefore the kinetic  $K_D$  was also unaffected (630 ± 290 and 550 ± 200 nM in absence or presence of NAM, respectively).

Lastly, both  $k_{on}$  and  $k_{off}$  values of the antagonist LY341495 were not significantly altered by the presence of PAM or NAM. This was the case for both the direct [ $^3$ H]LY341495 association and dissociation assays (Fig. 2A, Table 2), as well as for the competition association assays (Fig. 2D, Table 3). Hence,  $K_D$  values were similar to those obtained in the absence of allosteric modulator.

### 3.4. Potency and efficacy of orthosteric ligands, effects of allosteric modulators on LY354740 potency and efficacy

To evaluate the effects of JNJ-46281222 and RO4491533 on the functional responses induced by orthosteric ligands, a functional [ $^{35}$ S]GTPγS binding assay was used that measures compound-induced G protein activation. Firstly, concentration-response curves of the agonists glutamate and LY354740 were made (Fig. 3A), which led to pEC<sub>50</sub> values of 4.95 ± 0.01 and 6.94 ± 0.04, respectively. The maximum


**Table 4**

Effect of allosteric modulators JNJ-46281222 (PAM) and RO4491533 (NAM) on LY354740-induced [ $^{35}$ S]GTPγS binding.

	LY354740	
	pEC <sub>50</sub>	$E_{max}$ (%) <sup>a</sup>
LY354740	6.94 ± 0.04	83.5 ± 1.2
+ 1 nM JNJ-46281222	7.04 ± 0.10	117 ± 4.2
+ 3 nM JNJ-46281222	7.14 ± 0.08	154 ± 11**
+ 10 nM JNJ-46281222	7.35 ± 0.10**	178 ± 16***
+ 30 nM JNJ-46281222	7.50 ± 0.07****	215 ± 22****
+ 100 nM JNJ-46281222	7.69 ± 0.07****	222 ± 22****
+ 1 nM RO4491533	6.91 ± 0.07	74.5 ± 1.3**
+ 3 nM RO4491533	6.87 ± 0.11	55.5 ± 2.5****
+ 10 nM RO4491533	6.69 ± 0.05	30.0 ± 1.6****

<sup>a</sup> Expressed as percentage of [ $^{35}$ S]GTPγS binding induced by 1 mM glutamate (set at 100%). Values represent the mean ± SEM of three individual experiments performed in duplicate. Statistical analyses were performed using a one-way ANOVA with Dunnett's post-test. \*\* < 0.01, \*\*\* < 0.001, \*\*\*\* < 0.0001.

level of [ $^{35}$ S]GTPγS binding induced by the synthetic agonist LY354740 was 83.5 ± 1.2% compared to the maximum response induced by the endogenous agonist glutamate (100% at 1 mM) (Table 4). The potency of the antagonist LY341495 was determined in the presence of an EC<sub>80</sub> concentration of glutamate (60 μM). LY341495 inhibited glutamate-induced [ $^{35}$ S]GTPγS binding with a pIC<sub>50</sub> value of 7.40 ± 0.04 (Fig. 3A). Secondly, the effects of increasing concentrations of JNJ-46281222 or RO4491533 on the concentration-response curves of LY354740 were assessed (Fig. 3B, Table 4). Increasing concentrations of the PAM JNJ-46281222 induced a concentration-dependent increase in  $E_{max}$  up to approximately 220% at a concentration of 30 nM or higher. Moreover, the potency of LY354740 was increased significantly from 6.94 ± 0.04 in the absence of PAM to 7.69 ± 0.07 in the presence of 100 nM JNJ-46281222. In contrast, when LY354740 was treated with




**Fig. 3.** Intrinsic agonist potency and efficacy and the effect of allosteric modulators thereon. A) Concentration-response curves of glutamate- and LY354740-induced [ $^{35}$ S]GTPγS binding and concentration-response curve of inhibition of glutamate induced (EC<sub>80</sub>: 60 μM) [ $^{35}$ S]GTPγS binding by LY341495. Effects of increasing concentrations of B) JNJ-46281222 and C) RO4491533 on concentration-response curves of LY354740 in the [ $^{35}$ S]GTPγS binding assay. Parameters obtained from these graphs are described in Table 4. Data are expressed as the percentage of maximal response induced by 1 mM glutamate (100%) and represent the mean ± SEM of three individual experiments performed in duplicate. If not shown, error bars are within the symbol.

increasing concentrations of the NAM RO4491533, the  $E_{max}$  values of the concentration-response curves were reduced concentration-dependently. At a concentration of 30 nM RO4491533 or higher, LY354740-induced [ $^{35}$ S]GTP $\gamma$ S binding was completely abolished (Fig. 3C, Table 4). Interestingly, the potency of LY354740 did not change significantly in the presence of increasing NAM concentrations.

### 3.5. Effect of PAM JNJ-46281222 on duration of glutamate-induced cellular response using an impedance-based morphology assay

To gain further insights in the functional impact of allosteric modulation, receptor activation by glutamate was evaluated using an impedance-based morphology assay (i.e. xCELLigence). This method can record receptor-specific cellular responses in real-time, and thus compound-induced changes in cellular dynamics can be measured over time. Glutamate-induced responses were recorded at increasing concentrations in the absence and presence of 1  $\mu$ M PAM JNJ-46281222 resulting in a concentration-dependent increase in impedance, depicted as Cell Index (Fig. 4A,B). Similar to the [ $^{35}$ S]GTP $\gamma$ S assay, the glutamate potency was significantly ( $p < 0.05$ , student's  $t$ -test) increased from  $pEC_{50}$  5.27  $\pm$  0.19 to 5.99  $\pm$  0.12 in the absence and presence of PAM, respectively. Furthermore, the PAM-induced shift in potency for glutamate in the morphology assay was comparable to the shift for LY354740 in the [ $^{35}$ S]GTP $\gamma$ S assay (Table 4). Interestingly, the duration of the glutamate-induced response in the presence of PAM was also increased from approximately 45 to over 60 min. To further evaluate this effect, concentration-response curves of glutamate in the absence and presence of the PAM were obtained at two different time points after stimulation, i.e. 15 and 45 min (Fig. 4C,D). Comparison of these concentration-response curves yielded a 2-fold decrease in efficacy for the curve from the later time point (Fig. 4C), congruent with the almost loss of glutamate signal in the absence of PAM at 45 min (Fig. 4A). In contrast, the glutamate efficacy in the presence of PAM did not change significantly, when comparing the concentration-response curves of 15 and 45 min after stimulation. This is in line with the observation that the duration of the glutamate-induced response is prolonged by the PAM to approximately 60 min, and thus a decrease in cellular impedance was not yet observed at 45 min (Fig. 4B). Of note, when comparing the potencies for each condition, i.e. glutamate in the absence or presence of PAM, these were not significantly different at the two time points selected (Fig. 4C,D).



**Fig. 4.** Effect of PAM JNJ-46281222 on the duration of glutamate-induced signalling, as determined by an impedance-based morphology assay. A,B) 18 h after seeding, CHO-K1<sub>hmGlu<sub>2</sub></sub> cells (40,000 cells/well) were stimulated by increasing concentrations of the agonist glutamate in absence (A) or presence of 1  $\mu$ M of the PAM JNJ-46281222 (B). Medium with 0.025% DMSO was used as vehicle control. DMSO concentrations were 0.025% in all cases. A representative example is shown of a baseline-corrected response, the so-called  $\Delta$  cell index ( $\Delta$ CI), which was repeated at least three times in duplicate. C,D) Concentration-response curves were obtained from the  $\Delta$ CI values at 15 or 45 min after stimulation.  $pEC_{50}$  values are mentioned in the results section. xCELLigence traces (A, B) are from a representative experiment. Curves (C, D) represent mean  $\pm$  SEM of at least three individual experiments performed in duplicate.


### 3.6. Correlations and kinetic map

To compare the parameters obtained from the different radioligand binding assays, correlation plots were made (Fig. 5A–C). As mentioned above, the affinity values obtained from [ $^3$ H]LY341495 equilibrium displacement assays ( $K_i$ ) and [ $^3$ H]LY341495 competition association assays ( $K_D$ ) were in very good agreement, as exemplified by a high linear correlation (Fig. 5A,  $R^2 = 0.95$ ), further corroborating the robustness of the latter assay. A significant correlation was also found between affinity ( $K_i$ ) and association rate constants ( $k_{on}$ ) (Fig. 5B,  $R^2 = 0.99$ ). Such a correlation, however, was not found between affinity ( $K_i$ ) and dissociation rate constants ( $k_{off}$ ) (Fig. 5C,  $R^2 = 0.29$ ), indicating that affinities of these orthosteric ligands are predominantly  $k_{on}$ -driven.

A kinetic map was made, to further compare the kinetic and affinity parameters (Fig. 5D). In this map  $k_{on}$  (x-axis),  $k_{off}$  (y-axis) and  $K_D$  (diagonal lines) values were plotted.  $k_{on}$  and  $K_D$  values ranged over more than three orders of magnitude, whereas  $k_{off}$  values were only spread within a single order of magnitude and therefore appeared at a similar horizontal level in the kinetic map (Fig. 5D).

As shown in Table 3,  $k_{on}$  and  $k_{off}$  values of glutamate were affected by both PAM and NAM, which was exemplified in the kinetic map by a spread in symbols. Specifically, in the presence of PAM, the  $k_{off}$  of glutamate was decreased by 3-fold. In the presence of NAM both  $k_{on}$  and  $k_{off}$  were increased 3-fold resulting in the same affinity as illustrated by a diagonal shift of the NAM symbol, i.e. a shift along the line of similar  $K_D$ . In the presence of PAM, the  $k_{off}$  of LY354740 was similarly affected compared to glutamate, i.e. in both cases this resulted in a downward shift on the kinetic map, whereas in the presence of NAM no significant shifts were observed, resulting in nearly overlapping symbols. Since the  $k_{on}$  and  $k_{off}$  values of LY34195 were not affected by PAM or NAM all three symbols are overlapping in the kinetic map.

To compare the functional potency of agonists in the absence or presence of PAM to the kinetic parameters  $k_{on}$  and  $k_{off}$  further correlation plots were made (Fig. 5E–G). Firstly, the potencies ( $EC_{50}$ ) obtained in [ $^{35}$ S]GTP $\gamma$ S assays of agonists glutamate and LY354740 were shown to be strongly correlated (Fig. 5E,  $R^2 = 0.99$ ) to the affinity obtained from competition association assays ( $K_D$ ), showing that potencies are driven by the agonist binding affinities although the absolute values differed by approximately one log unit. As the affinities were shown to be correlated to on-rates, a correlation was also found



**Fig. 5.** Correlations between affinity, potency and kinetic parameters. (A,B,C) Correlation between affinity of glutamate (green), LY354740 (blue) and LY341495 (red) determined in [ $^3$ H]LY341495 equilibrium displacement assays ( $K_i$ ) and affinity determined based on kinetic parameters  $k_{on}$  and  $k_{off}$  obtained from [ $^3$ H]LY341495 competition association assays ( $K_D$ ) (A); affinity ( $K_i$ ) and association rate constant ( $k_{on}$ ) (B); affinity ( $K_i$ ) and dissociation rate constant ( $k_{off}$ ) (C). D) Kinetic map with the association rate constant ( $k_{on}$ ) plotted on the x-axis and the dissociation rate constant ( $k_{off}$ ) on the y-axis. Identical affinity ( $K_D$ ) values may result from different combinations of  $k_{on}$  and  $k_{off}$  ( $K_D = k_{off}/k_{on}$ , diagonal dashed lines). E,F,G) Correlation between potency of glutamate and LY354740 determined in [ $^{35}$ S]GPT $\gamma$ S assays ( $EC_{50}$ ) and affinity determined based on kinetic parameters  $k_{on}$  and  $k_{off}$  obtained from [ $^3$ H]LY341495 competition association assays ( $K_D$ ) (E). The  $EC_{50}$  of glutamate in the presence of 1  $\mu$ M PAM is from [19]; potency ( $EC_{50}$ ) and association rate constant ( $k_{on}$ ) (F); potency ( $EC_{50}$ ) and dissociation rate constant ( $k_{off}$ ) (G). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

between agonist potencies and  $k_{on}$  values (Fig. 5F,  $R^2 = 0.84$ ), showing that a high agonist potency is obtained from a high on-rate. No correlation was found between agonist potency and dissociation rate constant ( $k_{off}$ ) (Fig. 5G,  $R^2 = 0.28$ ). However, in the presence of PAM the agonist  $k_{off}$  was decreased for both glutamate and LY354740, which then was the driver for an increased agonist potency, as was also observed for agonist affinity in the presence of PAM.

#### 4. Discussion

Traditionally, affinity and potency are the main parameters studied in *in vitro* drug discovery. In addition, a ligand's target binding kinetic parameters are nowadays commonly appreciated as valuable information for the early phases of drug discovery [25]. For the development of

novel and effective orthosteric mGlu $_2$  ligands it is valuable to know their kinetic binding parameters, but also to understand how these relate to the binding kinetics of the endogenous agonist glutamate [30]. Moreover, a variety of high affinity and selective PAMs and NAMs have been developed that modulate glutamate potency, efficacy and/or affinity. As for orthosteric ligands, it is equally useful to know the allosteric modulator's binding kinetics [29]. Likewise it is also relevant to know how a modulator affects the kinetic binding parameters of the endogenous ligand glutamate. Hence, in the present study we aimed to increase the understanding of binding kinetics of orthosteric mGlu $_2$  ligands both on their own and upon modulation by an allosteric ligand.

The orthosteric ligands used in this study were the endogenous agonist glutamate and reference orthosteric ligands LY354740 (agonist) and LY341495 (antagonist). Radioligand displacement experiments

(Fig. 1, Table 1) showed that these ligands bind the same orthosteric binding site, which is in line with previous observations [36,37]. The kinetic parameters  $k_{on}$  and  $k_{off}$  of glutamate were quantified for the first time, using competition association experiments. To allow the set-up of this assay kinetic [ $^3$ H]LY341495 binding experiments were performed which showed that this ligand associates to its binding site within four minutes and dissociates in approximately ten minutes (Fig. 2A). For the set-up of the competition association assay, the association and dissociation of the radioligand should ideally be slower allowing more data points on the steep part of the curves [38]. This is often achieved by a temperature reduction. However, in these assays the temperature could not be lowered, since it had to be set at 0 °C already to allow quantification of kinetic measurements using this commercially available radioligand. Still we could produce robust data between the different binding assays indicating their validity. To obtain values for  $k_{on}$  and  $k_{off}$  of unlabelled ligands in the absence and presence of allosteric modulators, the Motulsky and Mahan model was used [35,38]. This model requires input for the values of  $k_{on}$  and  $k_{off}$  ( $k_1$  and  $k_2$  in the model) of the labelled radioligand (i.e. [ $^3$ H]LY341495). Importantly, these values should remain constant throughout the experiments, as was the case here for [ $^3$ H]LY341495 which provided similar values for  $k_{on}$  and  $k_{off}$  both in the absence and presence of allosteric modulators (Table 2). Due to the nature of the radioligand (i.e. fast receptor association) there is not much resolution in the association phase of the curves, making it difficult to observe differences in the rates by eye. However, by analysis of the data with the Motulsky and Mahan model we were able to obtain robust binding kinetic parameters between different experiments.

We found that glutamate dissociation ( $0.036\text{ s}^{-1}$ , Table 3) is fast in comparison to the series of mGlu<sub>2</sub> PAMs we studied before, which dissociate on a minute-range at 28 °C ( $k_{off}$  values between  $0.00033$  and  $0.0040\text{ s}^{-1}$ ), which would be even slower at 0 °C [29]. The observation of fast glutamate binding kinetics is in line with studies using FRET sensors that found mGlu<sub>1</sub> receptor conformational changes upon glutamate binding within seconds [39,40]. Fast receptor dissociation of glutamate relates to its physiological role as a neurotransmitter, where short bursts of glutamate at high concentrations are released into the synapse where it should evoke its function during that burst only [3]. Similarly, fast off-rates were obtained for other endogenous neurotransmitters as exemplified by 2-AG and anandamide on the CB<sub>2</sub> receptor ( $k_{off} = 0.053\text{ s}^{-1}$  and  $0.012\text{ s}^{-1}$  at 25 °C, respectively) [41] and acetylcholine on the M<sub>3</sub> receptor ( $k_{off} = 0.093\text{ s}^{-1}$  at 37 °C) [42]. Comparison of these values is troublesome, since for practical reasons experiments were performed at different temperatures. Still, they all share off-rates on the second to minute scale, indicating fast receptor dissociation particularly when comparing to synthetic ligands at the same receptors [41,42].

It is generally acknowledged that allosteric modulators may change the affinity and/or potency of agonists by modulating their  $k_{on}$  and/or  $k_{off}$  values [43]. Generally this has been determined by radioligand dissociation experiments in the presence of allosteric modulators that enable quantification of modulated off-rates [44]. Limitations of such assays are that it is impossible to detect effects on a radioligand's on-rate and that the orthosteric/endogenous ligand of interest needs to be radiolabelled. This is not suitable for most endogenous agonists including glutamate due to a too low target affinity, or too high non-specific binding. Therefore, we used the competition association assay for quantification of both  $k_{on}$  and  $k_{off}$  values of unlabelled orthosteric ligands in the absence or presence of PAM or NAM, as was recently published for an adenosine A<sub>1</sub> receptor PAM [38]. The presence of PAM JNJ-46281222 at 1  $\mu\text{M}$  significantly reduced the  $k_{off}$  of glutamate and LY354740, resulting in an increased affinity for both these agonists. This was in line with the results from the radioligand displacement assay (Table 1) and can be considered a typical PAM effect [43]. A similar effect was also seen in the functional [ $^{35}\text{S}$ ]GTP $\gamma$ S binding assay where the potency of LY354740 was increased by the PAM JNJ-

46281222 by almost 6-fold and its efficacy was more than doubled in line with earlier results using glutamate [19]. Furthermore, a decreased  $k_{off}$ , i.e. increased residence time, for glutamate in the presence of 1  $\mu\text{M}$  PAM JNJ-46281222 results in a prolonged receptor occupancy, which correlated to a prolonged cellular response in the morphology assay (Fig. 4). This functional assay is performed on whole cells under physiologically more relevant conditions (i.e. in culture medium, at 37 °C, in a CO<sub>2</sub> incubator) and is therefore considered to be more translational than classical functional assays [31,32].

Furthermore, the assay can be performed in real-time, which enabled translation from kinetic binding parameters towards functional efficacy over time. Previous studies have also used this assay to study the link between receptor binding kinetics, functional activation kinetics and duration of signalling for agonist-induced responses on the dopamine D<sub>2</sub> and neurokinin 1 receptors [45,46]. Of note, we have previously shown that culture medium contains 100  $\mu\text{M}$  endogenous glutamate [47], and therefore the enzyme glutamate-pyruvate transaminase (GPT) is often used to reduce this level. In the current study we were interested in the effect of a PAM on glutamate-responses, as this is the endogenous ligand and thus most relevant for PAM drug discovery. Therefore, GPT could not be used as this would deplete exogenous glutamate in addition to endogenous glutamate.

The NAM RO4491533 at 1  $\mu\text{M}$  positively modulated both  $k_{on}$  and  $k_{off}$  of glutamate (Table 3), resulting in an unchanged affinity. Of note, these effects on glutamate kinetics cannot be observed in the classical radioligand displacement assays at equilibrium conditions, and would therefore be missed (Table 1). On the other hand, RO4491533 was not able to change  $k_{on}$  and  $k_{off}$  of LY354740, which shows that the effects of this NAM on binding kinetics of agonists were probe-dependent [12]. This finding highlights the importance of using endogenous agonists in studies of allosteric modulation, since results on other (synthetic) agonists may provide different conclusions as a result of probe-dependency. The observation that a high 1  $\mu\text{M}$  concentration of NAM RO4491533 did not modulate the affinity of both agonists indicated that these ligands still bind the orthosteric binding site in the presence of this NAM, which is therefore behaving as a silent or neutral allosteric ligand (NAL) concerning orthosteric ligand binding [11]. On the contrary, RO4491533 displayed a strong negative cooperativity in the functional [ $^{35}\text{S}$ ]GTP $\gamma$ S binding assay as its concentration-dependently decreased the efficacy of the agonist LY354740 without changing its potency, eventually resulting in abolished LY354740 efficacy (Fig. 3; Table 4), similar to its effects on glutamate activity [22]. This mode of negative allosteric modulation, i.e. only affecting efficacy, has been found across the class C GPCR family. NAMs such as CPCCOEt (mGlu<sub>1</sub>) and MPEP (mGlu<sub>5</sub>) also abolished agonist efficacy without modulation of binding affinity [48,49]. The absence of cooperativity between LY341495 and both PAM and NAM was further illustrated by the observation that  $k_{on}$  and  $k_{off}$  of LY341495 were not affected by any allosteric modulator (Table 3).

The kinetic parameters of glutamate, LY354740 and LY341495 were most different in  $k_{on}$ , ranging over more than three orders of magnitude (from  $1.6 \pm 0.3 \times 10^3$  to  $2.4 \pm 0.18 \times 10^6\text{ M}^{-1}\text{ min}^{-1}$ ) as did their affinity values, whereas  $k_{off}$  values were all within a 6-fold range (from  $0.0087 \pm 0.0009$  to  $0.045 \pm 0.010$ ). The correlation plots Fig. 5 show that  $k_{on}$  is strongly correlated to the agonist affinity and potency, which indicates that target engagement of orthosteric mGlu<sub>2</sub> ligands is  $k_{on}$ -driven. Recent simulation studies have emphasized the role of high  $k_{on}$  values for receptor occupancy and drug dosing [26,27]. A mechanism responsible for these effects may be receptor rebinding, which is described as the binding of newly dissociated ligand from the local environment of the receptor [50]. The interstitial localization of the mGlu<sub>2</sub> receptor may result in more rebinding due to less diffusion and therefore higher local concentrations. However, rebinding to the mGlu<sub>2</sub> receptor seems less likely for glutamate due to its low  $k_{on}$  value (both in the absence and presence of a PAM – Table 3) and the active lowering of local glutamate concentrations by glutamate transporters expressed on

the neuronal cell membrane and surrounding glial cells [51]. In contrast, rebinding may play a more prominent role in binding of synthetic ligands which have higher  $k_{on}$  values and are not actively transported away from their site of action, resulting in higher *in vivo* receptor occupancy [50]. The plots in Fig. 5 furthermore showed that whereas between different PAMs a higher affinity is obtained from a higher  $k_{on}$ , the higher agonist affinity and potency obtained in the presence of a PAM results from a lowered  $k_{off}$  value as has been shown to be a common mechanism of action for many PAMs at different GPCRs [43]. As shown in Fig. 5,  $k_{off}$  is not correlated to both affinity and potency. This is similar to our recent work on mGlu<sub>2</sub> PAMs and this may therefore be a receptor-specific property [29]. Furthermore, this earlier study provided a first indication that PAM  $k_{off}$  values may be correlated to *in vivo* efficacy, as measured by effects on sleep-wake architecture in rats, more specifically suppression of Rapid Eye Movement (REM) sleep [52]. In the light of the current observation that the presence of a PAM prolonged glutamate-induced cellular responses by decreasing its  $k_{off}$ , it may be speculated that  $k_{off}$  values of both the PAM and the endogenous agonist are important for the duration of action and thus also for *in vivo* efficacy. As such, this information is valuable for the design of novel orthosteric and allosteric ligands in early drug discovery.

In conclusion, the set-up of a competition association radioligand binding assay enabled quantification of the parameters of binding kinetics for glutamate for the first time.  $k_{off}$  values of the orthosteric ligands were within a single order of magnitude, whereas  $k_{on}$  values were spread over more than three orders of magnitude and were strongly correlated to affinity, indicating that mGlu<sub>2</sub> target engagement is driven by  $k_{on}$  rather than  $k_{off}$ . Binding kinetics of agonists were modulated by the PAM, showing a decrease in  $k_{off}$  of both agonists and a prolonged functional response for glutamate. The NAM altered  $k_{on}$  and  $k_{off}$  of glutamate without changing glutamate's affinity, but did not induce such alterations for agonist LY354740, which indicates probe-dependency. These results show that affinity or potency-only optimization of orthosteric ligands will result in a high  $k_{on}$  value but not necessarily optimized  $k_{off}$  values, which is essential for optimal *in vivo* efficacy, as shown by previous studies. Together, this work contributes to an increased understanding of the molecular processes that underlie the mechanism of GPCR allosteric modulation, specifically how allosteric modulators affect the kinetic parameters of the endogenous agonist. Therefore, this study further emphasizes the need for evaluation of binding kinetics during drug discovery of both orthosteric and allosteric drug candidates for the mGlu<sub>2</sub> receptor as well as for other GPCRs.

## Acknowledgements

This project was financially supported by Vlaams Agentschap Innoveren & Ondernemen project number 120491.

## Conflicts of interests

None.

## List of author contributions

*Participated in research design:* Doornbos, Lavreysen, Tresadern, IJzerman, Heitman.

*Conducted experiments:* Doornbos, Vermond.

*Performed data analysis:* Doornbos, Vermond.

*Wrote or contributed to the writing of the manuscript:* Doornbos, Lavreysen, Tresadern, IJzerman, Heitman.

## References

[1] J.N.C. Kew, J.A. Kemp, Ionotropic and metabotropic glutamate receptor structure and pharmacology, *Psychopharmacology (Berl)* 179 (2005) 4–29, <https://doi.org/10.1007/s00213-005-2200-z>.

- [2] J.-P. Pin, R. Duvoisin, The metabotropic glutamate receptors: structure and functions, *Neuropharmacology* 34 (1995) 1–26 (accessed November 10, 2014), [https://doi.org/10.1016/0028-3908\(94\)00129-G](https://doi.org/10.1016/0028-3908(94)00129-G).
- [3] F. Nicoletti, J. Bockaert, G.L. Collingridge, P.J. Conn, F. Ferraguti, D.D. Schoepp, J.T. Wroblewski, J.P. Pin, Metabotropic glutamate receptors: from the workbench to the bedside, *Neuropharmacology* 60 (2011) 1017–1041, <https://doi.org/10.1016/j.neuropharm.2010.10.022>.
- [4] E. Dunayevich, J. Erickson, L. Levine, R. Landblom, D.D. Schoepp, G.D. Tollefson, Efficacy and tolerability of an mGlu<sub>2/3</sub> agonist in the treatment of generalized anxiety disorder, *Neuropsychopharmacology* 33 (2008) 1603–1610, <https://doi.org/10.1038/sj.npp.1301531>.
- [5] S.T. Patil, L. Zhang, F. Martenyi, S.L. Lowe, K.A. Jackson, B.V. Andreev, A.S. Avedisova, L.M. Bardenstein, I.Y. Gurovich, M.A. Morozova, S.N. Mosolov, N.G. Neznanov, A.M. Reznik, A.B. Smulevich, V.A. Tochilov, B.G. Johnson, J.A. Monn, D.D. Schoepp, Activation of mGlu<sub>2/3</sub> receptors as a new approach to treat schizophrenia: a randomized Phase 2 clinical trial, *Nat. Med.* 13 (2007) 1102–1107, <https://doi.org/10.1038/nm1632>.
- [6] A.M. Feyissa, W.L. Woolverton, J.J. Miguel-Hidalgo, Z. Wang, P.B. Kyle, G. Hasler, C.A. Stockmeier, A.H. Iyo, B. Karolewicz, Elevated level of metabotropic glutamate receptor 2/3 in the prefrontal cortex in major depression, *Prog. Neuropsychopharmacol. Biol. Psychiatry* 34 (2010) 279–283, <https://doi.org/10.1016/j.pnpbp.2009.11.018>.
- [7] C. Goeldner, T.M. Ballard, F. Knoflach, J. Wichmann, S. Gatti, D. Umbricht, Cognitive impairment in major depression and the mGlu<sub>2</sub> receptor as a therapeutic target, *Neuropharmacology* 64 (2013) 337–346, <https://doi.org/10.1016/j.neuropharm.2012.08.001>.
- [8] A. Kingston, P. Ornstein, R. Wright, B. Johnson, N. Mayne, J. Burnett, R. Belagaje, S. Wu, D. Schoepp, LY341495 is a nanomolar potent and selective antagonist of group II metabotropic glutamate receptors, *Neuropharmacology* 37 (1998) 1–12, [https://doi.org/10.1016/S0028-3908\(97\)00191-3](https://doi.org/10.1016/S0028-3908(97)00191-3).
- [9] H. Schaffhauser, J.G. Richards, J. Cartmell, S. Chaboz, J.A. Kemp, A. Klingelschmidt, J. Messer, H. Stadler, T. Woltering, V. Mutel, In vitro binding characteristics of a new selective group II metabotropic glutamate receptor radioligand, [3H]LY354740, in rat brain, *Mol. Pharmacol.* 53 (1998) 228–233, <https://doi.org/10.1124/mol.53.2.228>.
- [10] J.A. Monn, L. Prieto, L. Taboada, J. Hao, M.R. Reinhard, S.S. Henry, C.D. Beadle, L. Walton, T. Man, H. Rudyk, B. Clark, D. Tupper, S.R. Baker, C. Lamas, C. Montero, A. Marcos, J. Blanco, M. Bures, D.K. Clawson, S. Atwell, F. Lu, J. Wang, M. Russell, B.A. Heinz, X. Wang, J.H. Carter, B.G. Getman, J.T. Catlow, S. Swanson, B.G. Johnson, D.B. Shaw, D.L. McKinzie, Synthesis and Pharmacological Characterization of C4-(Thiotriazolyl)-substituted-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylates. Identification of (1R,2S,4R,5R,6R)-2-Amino-4-(1H-1,2,4-triazol-3-ylsulfanyl)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid (LY2812), *J. Med. Chem.* (2015) 7526–7548, <https://doi.org/10.1021/jm501612y>.
- [11] P.J. Conn, A. Christopoulos, C.W. Lindsley, Allosteric modulators of GPCRs: a novel approach for the treatment of CNS disorders, *Nat. Rev. Drug Discov.* 8 (2009) 41–54, <https://doi.org/10.1038/nrd2760>.
- [12] P. Keov, P.M. Sexton, A. Christopoulos, Allosteric modulation of G protein-coupled receptors: a pharmacological perspective, *Neuropharmacology* 60 (2011) 24–35, <https://doi.org/10.1016/j.neuropharm.2010.07.010>.
- [13] Homepage: <https://clinicaltrials.gov/show/NCT00986531>. The Effects AZD8529 on Cognition and Negative Symptoms in Schizophrenics (accessed April 12, 2018).
- [14] Homepage: <https://clinicaltrials.gov/show/NCT02401022>. The Study of AZD8529 for Smoking Cessation in Female Smokers (accessed April 12, 2018).
- [15] H. Salih, I. Anghelescu, I. Kezic, V. Sinha, E. Hoeben, L. Van Nueten, H. De Smedt, P. De Boer, Pharmacokinetic and pharmacodynamic characterisation of JNJ-40411813, a positive allosteric modulator of mGluR2, in two randomised, double-blind phase-I studies, *J. Psychopharmacol.* 29 (2015) 414–425, <https://doi.org/10.1177/0269881115573403>.
- [16] J.M. Kent, E. Daly, I. Kezic, R. Lane, P. Lim, H. De Smedt, P. De Boer, L. Van Nueten, W.C. Drevets, M. Ceusters, Efficacy and safety of an adjunctive mGlu<sub>2</sub> receptor positive allosteric modulator to a SSRI/SNRI in anxious depression, *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 67 (2016) 66–73, <https://doi.org/10.1016/j.pnpbp.2016.01.009>.
- [17] R. Galici, N.G. Echemendia, A.L. Rodriguez, P.J. Conn, A selective allosteric potentiator of metabotropic glutamate (mGlu) 2 receptors has effects similar to an orthosteric mGlu<sub>2/3</sub> receptor agonist in mouse models predictive of antipsychotic activity, *J. Pharmacol. Exp. Ther.* 315 (2005) 1181–1187, <https://doi.org/10.1124/jpet.105.091074>.
- [18] H. Lavreysen, X. Langlois, A. Ahnaou, W. Drinkenburg, P. te Riele, I. Biesmans, I. Van der Linden, L. Peeters, A. Megens, C. Wintmolders, J.M. Cid, A.A. Trabanco, J.I. Andrés, F.M. Dautzenberg, R. Lütjens, G. Macdonald, J.R. Atack, Pharmacological characterization of JNJ-40068782, a new potent, selective, and systemically active positive allosteric modulator of the mGlu<sub>2</sub> receptor and its radioligand [3H]JNJ-40068782, *J. Pharmacol. Exp. Ther.* 346 (2013) 514–527, <https://doi.org/10.1124/jpet.113.204990>.
- [19] M.L.J. Doornbos, L. Pérez-Benito, G. Tresadern, T. Mulder-Krieger, I. Biesmans, A.A. Trabanco, J.M. Cid, H. Lavreysen, A.P. IJzerman, L.H. Heitman, Molecular mechanism of positive allosteric modulation of the metabotropic glutamate receptor 2 by JNJ-46281222, *Br. J. Pharmacol.* 173 (2016) 588–600, <https://doi.org/10.1111/bph.13390>.
- [20] Homepage: <https://clinicaltrials.gov/show/NCT01457677>. ARTDeCo Study: A study of RO4995819 in patients with major depressive disorder and inadequate response to ongoing antidepressant treatment (accessed April 12, 2018).
- [21] M. Van Gool, S.A. Alonso De Diego, O. Delgado, A.A. Trabanco, F. Jourdan, G.J. Macdonald, M. Somers, L. Ver Donck, 1,3,5-Trisubstituted pyrazoles as potent

- negative allosteric modulators of the mGlu2/3 receptors, *ChemMedChem* 12 (2017) 905–912, <https://doi.org/10.1002/cmdc.201700101>.
- [22] B. Campo, M. Kalinichev, N. Lambeng, M. El Yacoubi, I. Royer-Urios, M. Schneider, C. Legrand, D. Parron, F. Girard, A. Bessif, S. Poli, J.-M. Vaugeois, E. Le Poul, S. Celanire, Characterization of an mGluR2/3 negative allosteric modulator in rodent models of depression, *J. Neurogenet.* 25 (2011) 152–166, <https://doi.org/10.3109/01677063.2011.627485>.
- [23] D.C. Swinney, B.A. Haubrich, I. Van Liefde, G. Vauquelin, The role of binding kinetics in GPCR drug discovery, *Curr. Top. Med. Chem.* 15 (2015) 2504–2522 <http://www.ncbi.nlm.nih.gov/pubmed/26126905>.
- [24] M.R. Dowling, S.J. Charlton, Quantifying the association and dissociation rates of unlabelled antagonists at the muscarinic M3 receptor, *Br. J. Pharmacol.* 148 (2006) 927–937, <https://doi.org/10.1038/sj.bjp.0706819>.
- [25] R.A. Copeland, The drug–target residence time model: a 10-year retrospective, *Nat. Rev. Drug Discov.* 15 (2016) 87–95, <https://doi.org/10.1038/nrd.2015.18>.
- [26] G. Vauquelin, Effects of target binding kinetics on in vivo drug efficacy: koff, kon and rebinding, *Br. J. Pharmacol.* 173 (2016) 2319–2334, <https://doi.org/10.1111/bph.13504>.
- [27] W.E.A. de Witte, M. Danhof, P.H. van der Graaf, E.C.M. de Lange, In vivo target residence time and kinetic selectivity: the association rate constant as determinant, *Trends Pharmacol. Sci.* 37 (2016) 831–842, <https://doi.org/10.1016/j.tips.2016.06.008>.
- [28] D.A. Sykes, H. Moore, L. Stott, N. Holliday, J.A. Javitch, J.R. Lane, S.J. Charlton, Extrapyramidal side effects of antipsychotics are linked to their association kinetics at dopamine D2 receptors, *Nat. Commun.* 8 (2017) 763, <https://doi.org/10.1038/s41467-017-00716-z>.
- [29] M.L.J. Doornbos, J.M. Cid, J. Haubrich, A. Nunes, J.W. van de Sande, S.C. Vermond, T. Mulder-Krieger, A.A. Trabanco, A. Ahnaou, W.H. Drinkenburg, H. Lavreysen, L.H. Heitman, A.P. IJzerman, G. Tresadern, Discovery and kinetic profiling of 7-Aryl-1,2,4-triazolo[4,3-a]pyridines: positive allosteric modulators of the metabotropic glutamate receptor 2, *J. Med. Chem.* 60 (2017) 6704–6720, <https://doi.org/10.1021/acs.jmedchem.7b00669>.
- [30] I. Nederpelt, D. Bleeker, B. Tuijt, A.P. IJzerman, L.H. Heitman, Kinetic binding and activation profiles of endogenous tachykinins targeting the NK1 receptor, *Biochem. Pharmacol.* 118 (2016) 88–95, <https://doi.org/10.1016/j.bcp.2016.08.004>.
- [31] B. Xi, N. Yu, X. Wang, X. Xu, Y.A. Abassi, The application of cell-based label-free technology in drug discovery, *Biotechnol. J.* 3 (2008) 484–495, <https://doi.org/10.1002/biot.200800020>.
- [32] N. Yu, J.M. Atienza, J. Bernard, S. Blanc, J. Zhu, X. Wang, X. Xu, Y. Abassi, Real-time monitoring of morphological changes in living cells by electronic cell sensors arrays: an approach to study G protein-coupled receptors, *Anal. Chem.* 78 (2006) 35–43.
- [33] J.M. Hillger, J. Schoop, D.I. Boomsma, P. Eline Slagboom, A.P. IJzerman, L.H. Heitman, Whole-cell biosensor for label-free detection of GPCR-mediated drug responses in personal cell lines, *Biosens. Bioelectron.* 74 (2015) 233–242, <https://doi.org/10.1016/j.bios.2015.06.031>.
- [34] Y. Cheng, W.H. Prusoff, Relationship between the inhibition constant ( $K_i$ ) and the concentration of inhibitor which causes 50 per cent inhibition ( $I_{50}$ ) of an enzymatic reaction, *Biochem. Pharmacol.* 22 (1973) 3099–3108 [https://doi.org/10.1016/0006-2952\(73\)90196-2](https://doi.org/10.1016/0006-2952(73)90196-2).
- [35] H.J. Motulsky, L.C. Mahan, The kinetics of competitive radioligand binding predicted by the law of mass action, *Mol. Pharmacol.* 25 (1984) 1–9 <http://www.ncbi.nlm.nih.gov/pubmed/6708928>.
- [36] J.A. Monn, L. Prieto, L. Taboada, C. Pedregal, J. Hao, M.R. Reinhard, S.S. Henry, P.J. Goldsmith, C.D. Beadle, L. Walton, T. Man, H. Rudyk, B. Clark, D. Tupper, S.R. Baker, C. Lamas, C. Montero, A. Marcos, J. Blanco, M. Bures, D.K. Clawson, S. Atwell, F. Lu, J. Wang, M. Russell, B.A. Heinz, X. Wang, J.H. Carter, C. Xiang, J.T. Catlow, S. Swanson, H. Sanger, L.M. Broad, M.P. Johnson, K.L. Knopp, R.M. Simmons, B.G. Johnson, D.B. Shaw, D.L. McKinzie, Synthesis and pharmacological characterization of C4-disubstituted analogs of 1 S , 2 S , 5 R , 6 S -2-amino-bicyclo[3.1.0]hexane-2,6-dicarboxylate: identification of a potent, selective metabotropic glutamate receptor agonist and determination of agonist-bou, *J. Med. Chem.* 58 (2015) 1776–1794, <https://doi.org/10.1021/jm501612y>.
- [37] R.A. Wright, M.B. Arnold, W.J. Wheeler, P.L. Ornstein, D.D. Schoepp, [3H] LY341495 binding to group II metabotropic glutamate receptors in rat brain, *J. Pharmacol. Exp. Ther.* 298 (2001) 453–460 <http://www.ncbi.nlm.nih.gov/pubmed/11454905>.
- [38] D. Guo, S.N. Venhorst, A. Massink, J.P.D. van Veldhoven, G. Vauquelin, A.P. IJzerman, L.H. Heitman, Molecular mechanism of allosteric modulation at GPCRs: insight from a binding kinetics study at the human A1 adenosine receptor, *Br. J. Pharmacol.* 171 (2014) 5295–5312, <https://doi.org/10.1111/bph.12836>.
- [39] M. Tateyama, H. Abe, H. Nakata, O. Saito, Y. Kubo, Ligand-induced rearrangement of the dimeric metabotropic glutamate receptor 1alpha, *Nat. Struct. Mol. Biol.* 11 (2004) 637–642, <https://doi.org/10.1038/nsmb770>.
- [40] V. Hlavackova, U. Zabel, D. Frankova, J. Bätz, C. Hoffmann, L. Prezeau, J.-P. Pin, J. Blahos, M.J. Lohse, Sequential inter- and intrasubunit rearrangements during activation of dimeric metabotropic glutamate receptor 1, *Sci. Signal.* 5 (2012) ra59, <https://doi.org/10.1126/scisignal.2002720>.
- [41] A. Martella, H. Sijben, A. Rufer, J. Fingerle, U. Grether, C. Ullmer, T. Hartung, A. IJzerman, M. van der Stelt, L. Heitman, A novel selective inverse agonist of the CB2 receptor as a radiolabeled tool compound for kinetic binding studies, *Mol. Pharmacol.* (2017) mol.117.108605. doi:10.1124/mol.117.108605.
- [42] D.A. Sykes, M.R. Dowling, S.J. Charlton, Exploring the mechanism of agonist efficacy: a relationship between efficacy and agonist dissociation rate at the muscarinic M3 receptor, *Mol. Pharmacol.* 76 (2009) 543–551, <https://doi.org/10.1124/mol.108.054452>.
- [43] A. Christopoulos, T. Kenakin, G protein-coupled receptor allosterism and complexing, *Pharmacol. Rev.* 54 (2002) 323–374 <http://www.ncbi.nlm.nih.gov/pubmed/12037145>.
- [44] J.R. Lane, L.T. May, R.G. Parton, P.M. Sexton, A. Christopoulos, A kinetic view of GPCR allostery and biased agonism, *Nat. Chem. Biol.* 13 (2017) 929–937, <https://doi.org/10.1038/nchembio.2431>.
- [45] I. Nederpelt, M. Kuzikov, W.E.A. de Witte, P. Schnider, B. Tuijt, S. Gul, A.P. IJzerman, E.C.M. de Lange, L.H. Heitman, From receptor binding kinetics to signal transduction; a missing link in predicting in vivo drug-action, *Sci. Rep.* 7 (2017) 14169, <https://doi.org/10.1038/s41598-017-14257-4>.
- [46] C. Klein Herenbrink, D.A. Sykes, P. Donthamsetti, M. Canals, T. Coudrat, J. Shonberg, P.J. Scammells, B. Capuano, P.M. Sexton, S.J. Charlton, J.A. Javitch, A. Christopoulos, J.R. Lane, The role of kinetic context in apparent biased agonism at GPCRs, *Nat. Commun.* 7 (2016) 1–14, <https://doi.org/10.1038/ncomms10842>.
- [47] M.L.J. Doornbos, I. Van der Linden, L. Vereyken, G. Tresadern, A.P. IJzerman, H. Lavreysen, L.H. Heitman, Constitutive activity of the metabotropic glutamate receptor 2 explored with a whole-cell label-free biosensor, *Biochem. Pharmacol.* 152 (2018) 201–210, <https://doi.org/10.1016/j.bcp.2018.03.026>.
- [48] S. Litschig, F. Gasparini, D. Rueegg, N. Stoehr, P.J. Flor, I. Vranesic, L. Prézeau, J.P. Pin, C. Thomsen, R. Kuhn, CPCCOEt, a noncompetitive metabotropic glutamate receptor 1 antagonist, inhibits receptor signaling without affecting glutamate binding, *Mol. Pharmacol.* 55 (1999) 453–461 <http://www.ncbi.nlm.nih.gov/pubmed/10051528>.
- [49] S.J. Bradley, C.J. Langmead, J.M. Watson, R.A.J. Challiss, Quantitative analysis reveals multiple mechanisms of allosteric modulation of the mGlu5 receptor in rat astroglia, *Mol. Pharmacol.* 79 (2011) 874–885, <https://doi.org/10.1124/mol.110.068882>.
- [50] G. Vauquelin, S.J. Charlton, Long-lasting target binding and rebinding as mechanisms to prolong in vivo drug action, *Br. J. Pharmacol.* 161 (2010) 488–508, <https://doi.org/10.1111/j.1476-5381.2010.00936.x>.
- [51] P. Marcaggi, D. Attwell, Role of glial amino acid transporters in synaptic transmission and brain energetics, *Glia* 47 (2004) 217–225, <https://doi.org/10.1002/glia.20027>.
- [52] A. Ahnaou, H. Lavreysen, G. Tresadern, J.M. Cid, W.H. Drinkenburg, mGlu2 receptor agonism, but not positive allosteric modulation, elicits rapid tolerance towards their primary efficacy on sleep measures in rats, *PLoS One* 10 (2015) e0144017, <https://doi.org/10.1371/journal.pone.0144017>.