



Universiteit
Leiden
The Netherlands

"Linking lipids to acetylation" novel roles of PI(5)P and PIP4K in SIRT1 regulation and development

Elouarrat, D.

Citation

Elouarrat, D. (2013, September 24). *"Linking lipids to acetylation" novel roles of PI(5)P and PIP4K in SIRT1 regulation and development*. Retrieved from <https://hdl.handle.net/1887/21945>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/21945>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/21945> holds various files of this Leiden University dissertation.

Author: Elouarrat, Dalila

Title: "Linking lipids to acetylation" novel roles of PI(5)P and PIP4K in SIRT1 regulation and development

Issue Date: 2013-09-24

Chapter 1

GENERAL INTRODUCTION

PREFACE

Every cell is able to communicate by sending, receiving or responding to signals in its environment. In order to trigger a physiological response, these signals must be transmitted across the cell membrane. Cells have developed a complex array of mechanisms to sense changes in the extracellular environment and to transduce these changes into intracellular signals (signal transduction) that can generate a cellular response. Signal transduction can be initiated by a ligand binding to a specific receptor at the plasma membrane. These ligands (primary messengers) bind to a diverse family of receptors, such G protein-coupled receptors, tyrosine kinase receptor or serine-threonine kinase receptors. Upon ligand binding, receptors activate diverse signaling pathways giving rise to change in abundance of intracellular signaling molecules, called second messengers. One important group of second messengers is the phosphoinositide family.

PHOSPHOINOSITIDES

Phosphoinositides are a distinct class of phospholipids, consisting of an inositol head group that is linked, via a phosphate diester bond, to the 1, 2-diacylglycerol (DAG) moiety. The hydrophobic DAG moiety remains buried in the membrane (can maintain the phosphoinositide as part of the membrane), whereas the inositol head group is soluble and can interact with components (proteins) in the cytosol. Phosphoinositides can be hydrolysed by phospholipase C family enzymes to generate two important second messengers; DAG, which can stimulate protein kinase C (PKC) activity, and inositol-(1,4,5)trisphosphate (IP3), which triggers calcium release from intracellular stores (Berridge, 1987; Hirasawa, 1985). Phosphatidylinositol (PI) can also be sequentially phosphorylated at the 3', 4' and 5' positions of the inositol ring to generate seven different (biologically active) phosphoinositides, namely PI(3)P, PI(4)P, PI(5)P, PI(3,4)P₂, PI(3,5)P₂, PI(4,5)P₂ and PI(3,4,5)P₃ (Lietha, 2001). The (de)phosphorylation of phosphoinositides and its derivatives is catalyzed by distinct lipid kinases and phosphatases, each specific for a given hydroxyl or phosphate group at the inositol ring (Sasaki *et al.*, 2009). In this way, phosphoinositide-specific kinases and phosphatases can generate various bioactive phosphoinositides in specific subcellular compartments (Figure 1). This subcellular distribution is regulated by specific protein-protein interactions unique to each kinase or phosphatase (Kutateladze, 2010). This allows the generation of a spatially regulated phosphoinositide signaling system.

Phosphoinositides can regulate numerous cellular processes, including ion channel function, protein localization and enzymatic activity, which in turn can affect many downstream processes such as vesicular transport, cell polarity and migration, gene transcription, cytoskeletal dynamics and cell proliferation and survival (Di Paolo *et al.*, 2006; Irvine, 2005; McCrea *et al.*, 2009). Thus, phosphoinositides function (exert their roles) either as precursors of second messengers (DAG and IP3), or by interacting with target proteins that contain specific phosphoinositide-interacting domains (PIDs). Over the past decades, a number of protein domains have been identified that can bind to phosphoinositides with varying degrees of specificity and affinity (Figure 1). These include the pleckstrin homology domain (PH), Phox domain (PX), epsin N-terminal homology domain (ENTH), FYVE domain and lysine/arginine-rich peptide sequences (Kutateladze, 2010; Lemmon, 2008). The interaction of phosphoinositides with these domains is key to the function of many signaling proteins and networks.

NUCLEAR PHOSPHOINOSITIDES

Most of our knowledge of phosphoinositide signaling is derived from receptor-mediated signaling pathways in the cytosol, yet evidence has emerged on the existence of phosphoinositide pathways in the nucleus. Many studies have shown that nuclear phosphoinositide metabolism and its regulation are independent of their cytoplasmic/plasma membrane counterparts (Irvine, 2003). Nuclear phosphoinositides have been shown to play key roles in a wide range of nuclear events, including cell proliferation and differentiation, DNA repair, mRNA splicing but also chromatin structure and transcription as will be described in **chapter 3** of this thesis.




Figure 1. Pathways of phosphoinositide synthesis and degradation. The phosphatidylinositol (PI) molecule includes inositol 1-phosphate bound via its phosphate group to 1,2-diacylglycerol (DAG) moiety. The structure and interconversion are shown for all seven known phosphoinositides namely PI(3)P, PI(4)P, PI(5)P, PI(3,4)P₂, PI(3,5)P₂, PI(4,5)P₂ and PI(3,4,5)P₂ in eukaryotes. Grey arrows represent less well-characterized pathways or pathways that have not been established yet. The phospholipid-binding domains that recognize specific phosphoinositides are indicated in different colours. PI(4,5)P₂ and other phosphoinositides can be hydrolysed by phospholipase C (PLC) enzymes to generate inositol 1,4,5-trisphosphate and diacylglycerol.

PHOSPHATIDYLINOSITOL PHOSPHATE KINASES (PIPKS)

The generation of phosphatidylinositol 4,5-bisphosphate $PI(4,5)P_2$ by phosphatidylinositol phosphate kinases (PIPK) is central to phosphoinositide signaling not only in the cytosol but also within the nucleus (Bunce, 2006). $PI(4,5)P_2$ can be synthesized by two different classes of PIPKs that utilize different substrates (van den Bout *et al.*, 2009). Phosphatidylinositol-4-phosphate 5 kinase (PIP5K) phosphorylates $PI(4)P$ on the 5 position on the inositol ring, while phosphatidylinositol-5-phosphate 4 kinase (PIP4K) phosphorylates $PI(5)P$ on the 4 position (Loijens *et al.*, 1996; Rameh *et al.*, 1997; Roberts *et al.*, 2005). Although $PI(3)P$ can act as a substrate, it is much less efficiently catalyzed than $PI(5)P$ (Morris *et al.*, 2000). Several isoforms of (consisting of α , β , γ isoforms) PIP5Ks and PIP4Ks have been found to localize in the nucleus. Despite synthesizing the same product, PIP5Ks and PIP4Ks appear functionally nonredundant. For instance, genetic deletion of PIP5K can be rescued by expression of PIP5Ks but not PIP4Ks, suggesting that PIP4Ks do not increase the cellular levels of $PI(4,5)P_2$ to the same extent as do the PIP5Ks. Because the cellular levels of $PI(5)P$ is present at much lower levels than $PI(4)P$ in the cell, the amount of $PI(4,5)P_2$ synthesized by PIP4K relative to PIP5K is likely to be much smaller (Clarke *et al.*, 2001; Morris *et al.*, 2000; Rameh *et al.*, 1997; Roberts *et al.*, 2005). Therefore, it is thought that PIP4Ks primarily function to remove the substrate $PI(5)P$ and thereby regulate its potential signaling functions (as will be detailed below).

PHOSPHATIDYLINOSITOL 5-PHOSPHATE 4-KINASE (PIP4K)

The three different PIP4K isoforms, encoded by distinct genes, are found at different cellular locations. The PIP4K α isoform is predominantly localized in the cytosol, although there is also a significant amount in the nucleus, whereas PIP4K γ localizes to the Golgi and intracellular vesicles (Boronenkov *et al.*, 1995; Clarke *et al.*, 2008). PIP4K β is predominantly nuclear; its 16 amino acid α -helix insertion (not present in the other PIP4K isoforms) is essential for nuclear localization (Ciruela *et al.*, 2000). The relative expression levels of PIP4K isoforms can differ between various tissues. The ratio of PIP4K β compared to PIP4K α and PIP4K γ expression is particularly high in skeletal muscle, where there is approximately 10-fold more PIP4K β than PIP4K α and PIP4K γ (Lamia *et al.*, 2004). Furthermore, in situ hybridization studies and RT-PCR in mice revealed similar expression of PIP4K α and β in the brain, whereas PIP4K α is expressed at higher levels than PIP4K β and γ in spleen. PIP4K γ is particularly highly expressed in the kidney compared to PIP4K α and β (Clarke *et al.*, 2008; Volpicelli-Daley *et al.*, 2010). These studies indicate that the differential expression and subcellular localization of each PIP4K isoform may serve specific functions (depending on cell and tissue context) of each organism (Clarke, 2010). The three isoforms of PIP4Ks also differ greatly in catalytic activity: PIP4K α has the highest activity, followed by PIP4K β which has 2000-fold less activity towards $PI(5)P$ compared to PIP4K α , whereas PIP4K γ is generally thought to be inactive (Bultsma *et al.*, 2010; Clarke *et al.*, 2008; Wang *et al.*, 2010). However, recent evidence indicates that distinct isoforms can affect one another's sub-cellular localization (Clarke *et al.*, 2012). For instance, the high activity PIP4K α isoform can interact with PIP4K β possibly by forming a heterodimer. The interaction and targeting of different isoforms suggest a mechanism by which low activity isoforms like

PIP4K β can regulate PI(5)P levels in a given cellular compartment (Bultsma *et al.*, 2010). Finally, it should be mentioned that the physiological roles of PIP4Ks remain poorly understood. PIP4K β -deleted mice are mildly growth retarded and hypersensitive to insulin (Lamia *et al.*, 2004). However, knockout or knockdown phenotypes of PIP4K α and PIP4K γ have not been reported to date. **Chapter 5** examines the role of PIP4Ks in vertebrate embryonic development using zebrafish as a model.

PHOSPHATIDYLINOSITOL 5-PHOSPHATE (PI(5)P) SIGNALING

Function

PI(5)P is the last identified member of the seven known phosphoinositides. Its late discovery is probably due to the fact that PI(5)P is low abundant (1-5%) compared to PI(4)P (Rameh *et al.*, 1997; Sarkes *et al.*, 2010). Therefore, PI(5)P has long been considered to be just an intermediate metabolite in the synthesis of other phosphoinositides. PI(5)P resides in the plasma membrane, intracellular membranes and the nucleus. Interestingly, several physiological and pathological stimuli alter cellular PI(5)P levels. For instance, the levels of PI(5)P are very low in resting cells, but they rise upon thrombin and insulin stimulation in human platelets or by osmotic stress in plant cells (Meijer, 2001; Morris *et al.*, 2000; Ndamukong *et al.*, 2010; Sbrissa *et al.*, 2004). During bacterial infection with *Shigella flexneri* or *Salmonella typhimurium*, the levels of PI(5)P are elevated and were shown to facilitate infection (Niebuhr *et al.*, 2002; Terebiznik, 2002). Moreover, a nuclear pool of PI(5)P has been found to increase when cells progress through the cell cycle and in response to oxidative stress and ultraviolet (UV) irradiation (Clarke *et al.*, 2001; Jones *et al.*, 2006; Jones *et al.*, 2012; Keune *et al.*, 2012).

As described above, phosphoinositides exert their function through interaction with specific phosphoinositide-binding domains. Identification of the binding of PI(5)P to ING2 (Inhibitor of Growth Protein 2) through the plant homeobox domain (PHD) motif defined this domain as the first PI(5)P phosphoinositide-binding domain in the nucleus (Gozani *et al.*, 2003). ING2 is a candidate tumor suppressor that induces growth arrest and apoptosis in a p53-dependent manner (Nagashima, 2001; Shi, 2005). Mutations resulting in loss of interaction with PI(5)P affect localization and the activity of ING2, leading to a decrease in ING2-mediated apoptosis and p53 acetylation. Another example of PI(5)P-PHD domain interaction is the relocalization of the histone H3 lysine 4 methylase ATX1 from the nucleus to the cytoplasm in response to dehydration stress (Alvarez-Venegas *et al.*, 2006; Ndamukong *et al.*, 2010b). Furthermore, the activity of the cullin 3 (CUL3) ubiquitination complex appears to be stimulated by nuclear PI(5)P (Bunce, 2008). Together, these studies indicate that nuclear PI(5)P levels play important roles in regulating nuclear protein function, especially in nuclear events such as chromatin remodeling, as discussed in **Chapter 3**.

PI(5)P signaling functions in the cytosol have also been reported (Grainger, 2012). In the cytosol, overexpression of PIP4K β resulted in a decrease in Akt activity in response to insulin (Carricaburu, 2003). Conversely, cells expressing IpgD, a bacterial PI(4,5)P₂-4-phosphatase that generates PI(5)P, had higher levels of basal and insulin-stimulated Akt phosphorylation (Pendaries, 2006). Interestingly, deletion of PIP4K β in mice also leads to increased insulin-induced Akt activation in muscle (Lamia *et al.*, 2004). Additionally, it has recently been shown

that PI(5)P-dependent Akt activation occurs via translocation of the EGF receptor (Ramel *et al.*, 2011). Next to insulin, H_2O_2 -induced PI(5)P levels also regulate the activity of Akt, which in turn is important for cell growth inhibition by H_2O_2 (Jones *et al.*, 2012). Furthermore, PI(5)P has been shown to play a role in cell morphology by mediating actin rearrangements downstream of the insulin receptor (Sbrissa *et al.*, 2004). More recently, PI(5)P was shown to promote cell migration through the activity of specific kinases and phosphatases involved in metabolizing PI(3,5)P₂ (Oppelt *et al.*, 2012).

Regulation

How is the abundance of PI(5)P regulated in distinct cellular compartments? Several pathways have the potential to generate PI(5)P, but their relative contributions to PI(5)P regulation *in vivo* are unclear (Figure 2). PI(3)P and PI(4)P phosphoinositides are produced by phosphorylation of PI by their specific kinases; however, no such kinase has been identified for PI(5)P so far (Lecompte *et al.*, 2008). Instead, it appears that PI(5)P production is regulated by specific phosphatases that dephosphorylate PI(4,5)P₂. This pathway was first identified by




Figure 2. Regulation of PI(5)P and its distinct roles in nuclear and cytosolic signaling. PI(5)P can be synthesized by MTM/MTMR phosphatases via dephosphorylation of PI(3,5)P₂ and bacterial IpgD and mammalian Type I,II PIP-4-phosphatases that convert PI(4,5)P₂ to PI(4,5)P₂. PIP4K can phosphorylate PI(5)P to PI(4,5)P₂ which is important for the physiological regulation of PI(5)P and downstream signaling. PI(5)P binds to PHD domain present on ING2, ATX1 and other nuclear proteins involved in nuclear signaling. Furthermore PI(5)P levels plays an important role in other cellular functions such as cell migration, insulin responses etc.

the $\text{PI}(4,5)\text{P}_2$ -4-phosphatase *IpgD*, responsible for the profound increase in $\text{PI}(5)\text{P}$ observed in *S.Flexneri*-infected cells (Niebuhr *et al.*, 2002). Subsequently, two human homologous $\text{PI}(4,5)\text{P}_2$ -4-phosphatases have been characterized that can translocate to the nucleus upon stress induction (Zou, 2007). The abundance of $\text{PI}(5)\text{P}$ can also be increased by the stress-activated lipid kinase *PIKfyve*, whose main function is to produce $\text{PI}(3,5)\text{P}_2$ from $\text{PI}(3)\text{P}$ (Tolias *et al.*, 1998). Overexpression of *PIKfyve* increases cellular $\text{PI}(5)\text{P}$ (Sbrissa, 2002). Mice lacking one copy of the gene have reduced $\text{PI}(5)\text{P}$ levels (Ikononov, 2011). However, $\text{PI}(5)\text{P}$ can also be generated by the major product of *PIKfyve*, $\text{PI}(3,5)\text{P}_2$, by myotubularin 3-phosphatases (Schaletzky, 2003). For example, *MTM1* overexpression is required for enhanced $\text{PI}(5)\text{P}$ production in osmotically stressed muscle (Tronchere *et al.*, 2004). However, the ability of myotubularin to dephosphorylate $\text{PI}(5)\text{P}$ and their operation *in vivo* is not yet established.

Unlike the dephosphorylation to $\text{PI}(5)\text{P}$, the $\text{PI}(5)\text{P}$ removal via $\text{PI}(5)\text{P}$ phosphorylation is much better established. *PIP4Ks* plays an important role in the regulation of $\text{PI}(5)\text{P}$ levels in the nucleus, where stressing cells activates p38 MAP kinase, which phosphorylates *PIP4K β* on Ser³²⁶ (Jones *et al.*, 2006). This decreases its *PIP4K* activity leading to an increase in the nuclear abundance of $\text{PI}(5)\text{P}$. A possible explanation of how *PIP4K β* , (which has low catalytic activity) can regulate nuclear $\text{PI}(5)\text{P}$ levels comes from the observation that *PIP4K α* and *PIP4K β* can form heterodimers (Bultsma *et al.*, 2010). Thereby *PIP4K β* can target *PIP4K α* activity to the nucleus. How phosphorylation by the p38 pathway regulates the *PIP4K α* /*PIP4K β* complex and its activity is currently not clear. *PIP4K β* is heavily post-translationally modified: in addition to being phosphorylated, it is also extensively acetylated. Consequently, these modifications might regulate *PIP4K α* /*PIP4K β* complex formation. In **Chapter 4**, we describe acetylated residues on *PIP4K β* and we identify it as a target of the *SIRT1* deacetylase (see below). In conclusion, the metabolic pathways governing the levels and the signaling roles of $\text{PI}(5)\text{P}$ are still not well understood, but recent data reveal its function as lipid mediator.

PROTEIN ACETYLATION

Acetylation refers to the addition of an acetyl group on lysine residues present within both histone and non-histone proteins. Acetylation is mediated by histone acetyl transferases (HATs). Since a large number of nonhistone proteins are targeted by HATs, these enzymes are also called K-acetyltransferases (KATs) (Berndsen, 2008). Acetylation of histones directly influences chromatin remodeling, by reducing the positive charge on lysine residues and by decreasing the affinity of histones for negatively charged DNA. This allows a localized “unraveling” of chromatin, making it more accessible for the binding of coactivators and the basal transcription machinery (Jiang, 2009).

The removal of acetyl groups is catalyzed by histone deacetylases (HDACs) and is associated with chromatin condensation and transcriptional repression. HDACs exert their repressive function on transcription either by condensing the chromatin or as components of large multiprotein complexes, by recruiting inhibitory factors to gene promoter regions (Jiang, 2009). Transcriptional regulation exerted by HDACs determines many cellular processes including cell cycle progression, apoptosis, autophagy, response to diverse types of stress, differentiation, and development (Norris *et al.*, 2009). Mammalian HDACs can be classified

into one of four different classes based on their amino acid sequence and structure. Sirtuins, or class III HDACs, have no homology to class I, II, or IV HDACs (TSA-sensitive enzymes and are localized in both the nucleus and the cytoplasm). Their deacetylase activity depends on the cofactor NAD⁺, rather than zinc, and they are localized in the nucleus, cytoplasm, or mitochondria and can deacetylate histone and nonhistone proteins (Imai *et al.*, 2000).

SIRTUINS

Sirtuins are highly conserved from bacteria to human and SIR2 (silent mating-type information regulation 2) was originally shown to extend lifespan in budding yeast (Kaeberlein, 1999). In mammals, there are seven sirtuins (SIRT1-7) all containing a conserved NAD-binding and catalytic domain, termed the sirtuin core domain, but they differ in their N and C-terminal domains (Frye, 2000). Sirtuins show significant sequence and functional differences from other classes of HDACs in that they carry out deacetylation via a two-step reaction that consumes NAD⁺ and releases, O-acetyl-ADP-ribose (AADPR), the deacetylated substrate and nicotinamide (used as sirtuin inhibitor) (Houtkooper, 2010). Sirtuins, although relatively similar to each other, have divergent biological functions which can be due to distinct cell-type-specific subcellular localisation of each member of the family (Finkel *et al.*, 2009). SIRT1 is located in both the nucleus and the cytoplasm, SIRT2 in the cytoplasm, SIRT3, 4, and 5 are mitochondrial, and SIRT6 and 7 are nuclear (Michan *et al.*, 2007). Apart from intracellular localization, SIRT1, 3, and 5 differ from SIRT2, 4, and 6 in the type of reaction they catalyse. SIRT1, 3, and 5 are NAD⁺-dependent deacetylases catalyzing the deacetylation of histones and nonhistone proteins, whereas SIRT6 is a NAD⁺-dependent ADP ribosyltransferase (ART) mediating mitochondrial protein ribosylation; SIRT2 and 4 exert both NAD⁺-dependent HDAC and ART activities (Westphal, 2007). The enzymatic activity of SIRT7 as well as its specific substrates has not yet been determined.

SIRT1 FUNCTIONS

SIRT1 is the most evolutionarily conserved sirtuin among the seven mammalian homologs, and it has been shown to play crucial roles in complex physiological processes, including metabolism, cancer, and aging (Sebastian *et al.*, 2012). SIRT1 participates in various cellular functions ranging from differentiation and development to metabolism and cell survival by deacetylating diverse substrates, summarized in Figure 3 (Nakagawa *et al.*, 2011). SIRT1 can deacetylate histone H4K16, H3K9 and H1K26 and thereby mediate heterochromatin formation (Liu *et al.*, 2013). Through its enzymatic activity, SIRT1 has also the capacity to regulate the activity of various transcription factors and other regulatory proteins. For example, SIRT1 regulates energy by inducing gluconeogenic while repressing glycolytic gene expressions through deacetylation of PGC-1 α and PPAR γ (Picard, 2004; Rodgers, 2005). Deacetylation of NF κ B, AP1 and Foxp3 by SIRT1 modulates the inflammation and immune pathway (Kong, 2012). And the deacetylation of DNA repair proteins Ku70, NBS1, WRN and XPA by SIRT1 regulates genomic stability (Fan, 2010; Li *et al.*, 2008; Yuan, 2007). SIRT1 regulates cell growth, apoptosis and stress response by deacetylating tumor suppressor protein like forkhead box protein FOXO1,3 and 4, Hif-1 α , HSF1, Rb and survivin (Brunet *et al.*, 2004; Lim, 2010; Luo, 2008; Motta *et al.*, 2004; Wang *et al.*, 2008;




Figure 3. *SIRT1 pathway overview. SIRT1 is an NAD⁺-dependent histone deacetylase that catalyzes the removal of acetyl (Ac) groups from a number of non-histone targets. The downstream effects of target deacetylation include changes in chromatin structure, cellular metabolism as well as cell survival and DNA repair. Several factors are involved in the regulation of SIRT1 function. SIRT1 function can be controlled availability of NAD⁺, posttranslational modifications, changes in expression or protein-protein interactions.*

Wong, 2007). The first and most widely known substrate of SIRT1 is tumor suppressor p53, which has critical roles in the cell-cycle regulation and apoptosis (Luo et al., 2001; McCubrey, 2012).

p53 is a short-lived protein that is sustained at low levels under normal physiological conditions. However, p53 is maintained at a relative high level by posttranslational modifications in response to various stresses (Gu, 2012). The acetylation of p53 in response to DNA damage by HAT CBP/p300 was shown to acetylate p53 at K373/K382 and K320. Lysine acetylation at these sites promotes both its DNA binding ability and its transcriptional activity, and include p21, Bax and puma as p53 targets (Smith, 2002). In mice harboring a mutant p53 allele in which all the acetylation sites have been replaced a deficiency in gene transcription in response to DNA damage was shown, which illustrates that acetylation is indispensable for p53 function (Chao et al., 2006; Tang et al., 2008). Conversely, SIRT1 binds to and deacetylates p53, thereby negatively regulating p53-mediated transcriptional activation. SIRT1 functions by deacetylating of p53 which prevents cellular senescence and apoptosis caused by DNA damage and stress (Luo et al., 2001; Vaziri et al., 2001).

SIRT1 REGULATION

Although much attention has been focused on the identification of the cellular targets and functional networks controlled by SIRT1, the mechanisms that regulate SIRT1 activity by biological stimuli have only recently begun to emerge. As an enzyme, the activity of SIRT1 can

be controlled by the availability of its substrates, post-translational modifications, interactions with other proteins and changes in its expression levels (Figure 3) (Revollo *et al.*, 2013).

The basal intracellular NAD⁺ levels are maintained relatively constant by the NAD⁺ biosynthetic and salvage pathways (Houtkooper, 2010). Proper functioning of these pathways is important for the activation of SIRT1 catalytic activity since the availability of NAD⁺ in cells is a limiting step (Revollo *et al.*, 2004). Glucose deprivation and metabolic changes by foreexample calorie restriction can cause fluctuations in NAD⁺ levels. Since SIRT1 associate with chromatin and its function is NAD⁺-dependent, SIRT1 can couple changes in the cellular metabolic state and NAD⁺ levels to transcription (Chalkiadaki, 2012).

Furthermore, SIRT1 enzymatic activity can be regulated by several posttranslational modifications. First, prosurvival dual specificity tyrosine phosphorylation-regulated kinases (DYRKs), DYRK1A and DYRK3, have been shown to phosphorylate SIRT1 in response to genotoxic stress (Guo, 2010). This PTM substantially enhances SIRT1 deacetylase activity towards acetylated p53 and protects cells from genotoxic stress-induced apoptosis (Guo, 2012). Another serine residue located at the highly conserved core domain of SIRT1, Ser 434, has been shown to be a phosphorylation target of the cyclic AMP/protein kinase A (cAMP/PKA) signaling pathway (Gerhart-Hines *et al.*, 2011). This PTM rapidly enhances the intrinsic deacetylase activity of SIRT1 independently of cellular NAD⁺ levels. Finally, C-Jun N-terminal kinase (JNK)1, phosphorylates SIRT1 on several serines, particularly under stressful cellular conditions (Nasrin, 2009). Intriguingly, these phosphorylations appear to increase the deacetylase activity of SIRT1 towards one of its substrates, histone H3, but have no effect towards another substrate, p53. This suggests that some or all of these PTMs can alter the activity of SIRT1 in a substrate-specific manner. SIRT1 is also sumoylated at K734, which in turn increases its activity and nitrosylation at Cys 387 and 390 in the catalytic core reduces SIRT1's ability to deacetylate PGC-1 α (Kornberg, 2010; Yang, 2007).

In addition to posttranslational modification, alteration in expression levels can also regulate SIRT1 activity. The "tumor suppressor hypermethylated in cancer" protein (HIC1) has been shown to inhibit SIRT1 expression by forming a repressive complex with SIRT1 on its own promoter (Chen, 2005; Zhang *et al.*, 2007). Conversely, through a DNA damage dependent matter transcription factor E2F1 can induce SIRT1 expression (Wang *et al.*, 2006). Members of the FOXO transcription factors also regulated the expression of SIRT1, like FOXO1, which induces SIRT1 expression by binding to FOXO1 response elements in the SIRT1 promoter (Nemoto, 2004; Xiong, 2011). The SIRT1 promoter also contains several cyclic AMP response element-binding protein (CREB) binding sites and through CREB, a transcription factor whose activation is mediated by PKA in response to low nutrient availability, the expression of SIRT1 is induced (Fusco, 2012; Noriega, 2011). The abundance of SIRT1 is also controlled by translational events and RNA stability. For instance, the Hu antigen (HuR) an mRNA binding protein that binds the 3'UTR of SIRT1 mRNA and plays a major role in stabilizing SIRT1 mRNA transcript (Abdelmohsen, 2007; Yamakuchi, 2012). Furthermore, SIRT1 expression and its activity are also under the influence of several miRNA's as reported by several studies.

Finally, protein-protein interactions also play key roles in the regulation of SIRT1. For instance, interaction with active regulator of SIRT1 (AROS) enhances the activity of SIRT1 towards acetylated p53. AROS appears to bind to the N terminus of SIRT1, but little is known

about how this interaction increases the SIRT1 activity (Kim *et al.*, 2007). In contrast to AROS, deleted in breast cancer 1 (DBC1) is a negative regulator of SIRT1 (Kim *et al.*, 2008; Zhao *et al.*, 2008). The ‘essential for SIRT1 activity’ (ESA) domain (a 25-amino-acid region in the C terminus of SIRT1), which intramolecularly interacts with the SIRT1 deacetylase core, was also shown to activate its enzymatic activity (Kang *et al.*, 2011). DBC1 also interacts with the SIRT1 deacetylase core, potentially displacing ESA, decreasing the accessibility of the core domain to protein substrates, and thereby inactivating SIRT1. Moreover, cellular stress results in phosphorylation of DBC1 at Thr 454, which appears to create a second binding site for SIRT1 to augment the DBC1–SIRT1 interaction (Yuan, 2012). However, for most of studies where SIRT1 activity was shown to be modulated by protein-protein interactions, it is not clear yet whether allosteric changes or substrate access to catalytic core are responsible for the decrease in SIRT1 activity.

These studies show that SIRT1 is temporally and spatially regulated in response to various environmental cues. In **Chapter 2**, we describe a novel aspect of the regulation of SIRT1 by its interaction with and activation by phosphoinositides.

THESIS OUTLINE

The studies described in this thesis focus on the novel role of the PI(5)P/PIP4K signaling axis in regulating acetylation and understanding their physiological functions. **Chapter 1** is a general introduction giving an overview of the molecules and pathways involved in phosphoinositide signaling and regulation of SIRT1 activity. **Chapter 2** identifies PI(5)P as novel lipid regulator of SIRT1 activity and that SIRT1 is subject to acute regulation in response to lipid signaling. **Chapter 3** reviews the latest insights of phospholipids in the nucleus and their emerging role as important modulators of chromatin and gene transcription. Next, **chapter 4** explores the posttranslational modifications on PIP4K β and describes its biochemical interaction and deacetylation by SIRT1 deacetylase. **Chapter 5** provides insight on the physiological roles of PIP4Ks and provides the first description of how lack of PIP4K α expression and activity affects zebrafish development. In **chapter 6**, the results presented in this thesis are summarized and discussed

REFERENCE LIST

1. Abdelmohsen K, Pullmann R Jr, Lal A, Kim HH, Galban S, Yang X, *et al.* Phosphorylation of HuR by Chk2 regulates SIRT1 expression. *Molecular Cell* 2007;25:543-57.
2. Alvarez-Venegas R, Sadler M, Hlavacka A, Baluska F, Xia Y, Lu G, *et al.* The Arabidopsis homolog of trithorax, ATX1, binds phosphatidylinositol 5-phosphate, and the two regulate a common set of target genes. *Proc Natl Acad Sci U S A* 2006;103:6049-54.
3. Berndsen C. Catalysis and substrate selection by histone/protein lysine acetyltransferases. *Current opinion in structural biology* 2008;18:682-9.
4. Berridge MJ. Inositol trisphosphate and diacylglycerol: two interacting second messengers. *Annual review of biochemistry* 1987;56:159-93.
5. Boronenkov I, Anderson R. The Sequence of Phosphatidylinositol-4-phosphate 5-Kinase Defines a Novel Family of Lipid Kinases. *Journal of Biological Chemistry* 1995;270:2881-4.
6. Brunet A, Sweeney L, Sturgill JF, Chua K, Greer P, Lin Y, *et al.* Stress-Dependent Regulation of FOXO Transcription Factors by the SIRT1 Deacetylase. *Science* 2004;303:2011-5.
7. Bultsma Y, Keune W, Divecha N. PIP4K α interacts with and modulates nuclear

- localization of the high-activity PI(5)P-4-kinase isoform PIP4K α . *Biochemical Journal* 2010;430:223-35.
8. Bunce. Nuclear PI(4,5)P(2): a new place for an old signal. *Biochimica et biophysica acta* 2006;1761:560-9.
 9. Bunce. Coordinated activation of the nuclear ubiquitin ligase Cul3-SPOP by the generation of phosphatidylinositol 5-phosphate. *Journal of Biological Chemistry* 2008;283:8678-86.
 10. Carricaburu V, Lamia KA, Lo E, Favereaux L, Payraastre B, Cantley LC, et al. The phosphatidylinositol (PI)-5-phosphate 4-kinase type II enzyme controls insulin signaling by regulating PI-3,4,5-trisphosphate degradation. *Proc Natl Acad Sci U S A* 2003;100:9867-72.
 11. Chalkiadaki A. Sirtuins mediate mammalian metabolic responses to nutrient availability. *Nature reviews Endocrinology* 2012;8:287-96.
 12. Chao C, Wu Z, Mazur SJ, Borges H, Rossi M, Lin T, et al. Acetylation of mouse p53 at lysine 317 negatively regulates p53 apoptotic activities after DNA damage. *Mol Cell Biol* 2006;26:6859-69.
 13. Chen. Tumor suppressor HIC1 directly regulates SIRT1 to modulate p53-dependent DNA-damage responses. *Cell* 2005;123:437-48.
 14. Ciruela A, Hinchliffe KA, Divecha N, Irvine RF. Nuclear targeting of the beta isoform of type II phosphatidylinositol phosphate kinase (phosphatidylinositol 5-phosphate 4-kinase) by its alpha-helix 7. *Biochemical Journal* 2000;346:587-91.
 15. Clarke J, Emson P, Irvine R. Localization of phosphatidylinositol phosphate kinase IIgamma in kidney to a membrane trafficking compartment within specialized cells of the nephron. *American Journal of Physiology - Renal Physiology* 2008;295:F1422-F1430.
 16. Clarke J, Irvine R. The activity, evolution and association of phosphatidylinositol 5-phosphate 4-kinases. *Advances in Biological Regulation* 2012;52:40-5.
 17. Clarke JH, Letcher AJ, D'santos CS, Halstead JR, Irvine RF, Divecha N. Inositol lipids are regulated during cell cycle progression in the nuclei of murine erythroleukaemia cells. *Biochemical Journal* 2001b;357:905-10.
 18. Clarke JH, Letcher AJ, D'santos CS, Halstead JR, Irvine RF, Divecha N. Inositol lipids are regulated during cell cycle progression in the nuclei of murine erythroleukaemia cells. *Biochem J* 2001a;357:905-10.
 19. Clarke. Localization, regulation and function of type II phosphatidylinositol 5-phosphate 4-kinases. *Adv Enzyme Regul* 2010;50:12-8.
 20. Di Paolo G, De Camilli P. Phosphoinositides in cell regulation and membrane dynamics. *Nature* 2006;443:651-7.
 21. Fan W. SIRT1 regulates UV-induced DNA repair through deacetylating XPA. *Molecular Cell* 2010;39:247-58.
 22. Finkel T, Deng C, Mostoslavsky R. Recent progress in the biology and physiology of sirtuins. *Nature* 2009;460:587-91.
 23. Frye RA. Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins. *Biochemical and biophysical research communications* 2000;273:793-8.
 24. Fusco S, Ripoli C, Podda MV, Ranieri SC, Leone L, Toietta G, et al.
 25. A role for neuronal cAMP responsive-element binding (CREB)-1 in brain responses to calorie restriction. *Proc Natl Acad Sci U S A* 2012;109:621-6.
 26. Gerhart-Hines Z, Dominy J, Blittler S, Jedrychowski M, Banks A, Lim J, et al. The cAMP/PKA Pathway Rapidly Activates SIRT1 to Promote Fatty Acid Oxidation Independently of Changes in NAD⁺. *Molecular Cell* 2011;44:851-63.
 27. Gozani O, Karuman P, Jones DR, Ivanov D, Cha J, Lugovskoy AA, et al. The PHD finger of the chromatin-associated protein ING2 functions as a nuclear phosphoinositide receptor. *Cell* 2003;114:99-111.
 28. Grainger D, Tavelis C, Ryan AJ, Hinchliffe KA. The emerging role of PI(5)P: another signalling phosphoinositide takes its place. *Biochemical Society transactions* 2012;40:257-61.
 29. Gu B. Surf the post-translational modification network of p53 regulation. *International journal of biological sciences* 2012;8:672-84.
 30. Guo X, Kesimer M, Tolun G, Zheng X, Xu Q, Lu J. The NAD(+)-dependent protein deacetylase activity of SIRT1 is regulated by its oligomeric status. *Scientific Reports* 2012;2:640.
 31. Guo. DYRK1A and DYRK3 promote cell survival through phosphorylation and activation of SIRT1. *Journal of Biological Chemistry* 2010;285:13223-32.

32. Hirasawa K. Phosphatidylinositol turnover in receptor mechanism and signal transduction. *Annual Review of Pharmacology and Toxicology* 1985;25:147-70.
33. Houtkooper R, Cantó C, Wanders RJ, Auwerx J. The secret life of NAD⁺: an old metabolite controlling new metabolic signaling pathways. *Endocrine reviews* 2010;31:194-223.
34. Ikononov O, Sbrissa D, Delvecchio K, Xie Y, Jin JP, Rappolee D, et al. The phosphoinositide kinase PIKfyve is vital in early embryonic development: preimplantation lethality of PIKfyve^{-/-} embryos but normality of PIKfyve^{+/-} mice. *Journal of Biological Chemistry* 2011;286:13404-13.
35. Imai S, Armstrong C, Kaerberlein M, Guarente L. Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* 2000;403:795-800.
36. Irvine R. Inositide evolution- towards turtle domination? *The Journal of Physiology* 2005;566:295-300.
37. Irvine R. Nuclear lipid signalling. *Nat Rev Mol Cell Biol* 2003;4:349-61.
38. Jiang C. Nucleosome positioning and gene regulation: advances through genomics. *Nature reviews Genetics* 2009;10:161-72.
39. Jones D, Bultsma Y, Keune W, Halstead J, Elouarrat D, Mohammed S, et al. Nuclear PI(5)P as a Transducer of Stress Signaling: An In Vivo Role for PIP4Kbeta. *Molecular Cell* 2006;23:685-95.
40. Jones D, Foulger R, Keune W, Bultsma Y, Divecha N. PI(5)P is an oxidative stress-induced second messenger that regulates PKB activation. *The FASEB Journal* 2012.
41. Kaerberlein M. The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes & development* 1999;13:2570-80.
42. Kang H, Suh J, Jung Y, Jung J, Kim M, Chung J. Peptide Switch Is Essential for Sirt1 Deacetylase Activity. *Molecular Cell* 2011;44:203-13.
43. Keune W, Jones D, Bultsma Y, Sommer L, Zhou X, Lu K, et al. Regulation of Phosphatidylinositol-5-Phosphate Signaling by Pin1 Determines Sensitivity to Oxidative Stress. *Sci Signal* 2012;5:ra86.
44. Kim E, Kho J, Kang M, Um S. Active Regulator of SIRT1 Cooperates with SIRT1 and Facilitates Suppression of p53 Activity. *Molecular Cell* 2007;28:277-90.
45. Kim J, Chen J, Lou Z. DBC1 is a negative regulator of SIRT1. *Nature* 2008;451:583-6.
46. Kong. Sirtuin 1 in immune regulation and autoimmunity. *Immunology and cell biology* 2012;90:6-13.
47. Kornberg. GAPDH mediates nitrosylation of nuclear proteins. *Nat Cell Biol* 2010;12:1094-100.
48. Kutateladze T. Translation of the phosphoinositide code by PI effectors. *Nature chemical biology* 2010;6:507-13.
49. Lamia K, Peroni O, Kim Y, Rameh L, Kahn B, Cantley L. Increased Insulin Sensitivity and Reduced Adiposity in Phosphatidylinositol 5-Phosphate 4-Kinase ^{-/-} Mice. *Mol Cell Biol* 2004;24:5080-7.
50. Lecompte O, Poch O, Laporte J. PI(5)Pregulation through evolution: roles in membrane trafficking? *Trends in Biochemical Sciences* 2008;33:453-60.
51. Lemmon MA. Membrane recognition by phospholipid-binding domains. *Nat Rev Mol Cell Biol* 2008;9:99-111.
52. Li K, Casta A, Wang R, Lozada E, Fan W, Kane S, et al. Regulation of WRN Protein Cellular Localization and Enzymatic Activities by SIRT1-mediated Deacetylation. *Journal of Biological Chemistry* 2008;283:7590-8.
53. Lietha D. Phosphoinositides, The Seven Species: Conversion and Cellular Roles. eLS. John Wiley & Sons, Ltd; 2001.
54. Lim J, Lee YM, Chun YS, Chen J, Kim JE, Park JW. Sirtuin 1 modulates cellular responses to hypoxia by deacetylating hypoxia-inducible factor 1alpha. *Molecular Cell* 2010;38:864-78.
55. Liu T, McCall C. Deacetylation by SIRT1 Reprograms Inflammation and Cancer. *Genes Cancer* 2013.
56. Loijens J, Boronenkov I, Parker G, Anderson R. The phosphatidylinositol 4-phosphate 5-kinase family. *Adv Enzyme Regul* 1996;36:115-40.
57. Luo J. SIRTing through breast cancer is just a survivin' game. *Molecular Cell* 2008;32:159-60.
58. Luo J, Nikolaev A, Imai S, Chen D, Su F, Shiloh A, et al. Negative Control of p53 by Sir2alpha Promotes Cell Survival under Stress. *Cell* 2001;107:137-48.
59. McCrea HJ, De Camilli P. Mutations in phosphoinositide metabolizing enzymes and human disease. *Physiology (Bethesda)* 2009;24:8-16.

60. McCubrey J. Recent discoveries in the cycling, growing and aging of the p53 field. *Aging* 2012;4:887-93.
61. Michan S, Sinclair D. Sirtuins in mammals: insights into their biological function. *Biochem J* 2007;404:1-13.
62. Morris J, Hinchliffe K, Ciruela A, Letcher A, Irvine R. Thrombin stimulation of platelets causes an increase in phosphatidylinositol 5-phosphate revealed by mass assay. *FEBS Letters* 2000;475:57-60.
63. Motta M, Divecha N, Lemieux M, Kamel C, Chen D, Gu W, et al. Mammalian SIRT1 Represses Forkhead Transcription Factors. *Cell* 2004;116:551-63.
64. Nakagawa T, Guarente L. Sirtuins at a glance. *Journal of Cell Science* 2011;124:833-8.
65. Nasrin N, Kaushik VK, Fortier E, Wall D, Pearson KJ, de Cabo R, et al. JNK1 phosphorylates SIRT1 and promotes its enzymatic activity. *PLoS One* 2009;4:e8414.
66. Ndamukong I, Jones D, Lapko H, Divecha N, Avramova Z. Phosphatidylinositol 5-Phosphate Links Dehydration Stress to the Activity of ARABIDOPSIS TRITHORAX-LIKE Factor ATX1. *PLoS One* 2010;5:e13396.
67. Nemoto. Nutrient availability regulates SIRT1 through a forkhead-dependent pathway. *Science* 2004;306:2105-8.
68. Niebuhr K, Giuriato S, Pedron T, Philpott D, Gaits F, Sable J, et al. Conversion of PtdIns(4,5) P2 into PtdIns(5)P by the *S. flexneri* effector IpgD reorganizes host cell morphology. *EMBO J* 2002;21:5069-78.
69. Noriega L, Feige JN, Canto C, Yamamoto H, Yu J, Herman MA, et al. CREB and ChREBP oppositely regulate SIRT1 expression in response to energy availability. *EMBO reports* 2011;12:1069-76.
70. Norris KL, Lee JY, Yao TP. Acetylation goes global: the emergence of acetylation biology. *Sci Signal* 2009;2:e76.
71. Oppelt A, Lobert V, Haglund K, Mackey A, Rameh L, Liestol K, et al. Production of phosphatidylinositol 5-phosphate via PIKfyve and MTMR3 regulates cell migration. *EMBO Rep* 2012.
72. Pendaries C, Tronchère H, Arbibe L, Mounier J, Gozani O, Cantley L, et al. PI(5)P activates the host cell PI3-kinase/Akt pathway during *Shigella flexneri* infection. *EMBO journal* 2006;25:1024-34.
73. Picard F, Kurtev M, Chung N, Topark-Ngarm A, Senawong T, Machado De Oliveira R et al. Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR-gamma. *Nature* 2004;429:771-6.
74. Rameh L, Tolias K, Duckworth B, Cantley L. A new pathway for synthesis of phosphatidylinositol-4,5-bisphosphate. *Nature* 1997;390:192-6.
75. Ramel D, Lagarrigue F, Pons V, Mounier J, Dupuis-Coronas S, Chicanne G, et al. *Shigella flexneri* Infection Generates the Lipid PI5P to Alter Endocytosis and Prevent Termination of EGFR Signaling. *Sci Signal* 2011;4:ra61.
76. Revollo J, Li X. The ways and means that fine tune Sirt1 activity. *Trends in Biochemical Sciences* 2013;38:160-7.
77. Revollo JR, Grimm AA, Imai S. The NAD biosynthesis pathway mediated by nicotinamide phosphoribosyltransferase regulates Sir2 activity in mammalian cells. *J Biol Chem* 2004;279:50754-63.
78. Roberts H, Clarke J, Letcher A, Irvine R, Hinchliffe K. Effects of lipid kinase expression and cellular stimuli on phosphatidylinositol 5-phosphate levels in mammalian cell lines. *FEBS Letters* 2005;579:2868-72.
79. Rodgers J, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1alpha and SIRT1. *Nature* 2005;434:113-8.
80. Sarkes D, Rameh L. A novel HPLC-based approach makes possible the spatial characterization of cellular PI(5)P and other phosphoinositides. *Biochemical Journal* 2010;428:375-84.
81. Sasaki T, Takasuga S, Sasaki J, Kofuji S, Eguchi S, Yamazaki M, et al. Mammalian phosphoinositide kinases and phosphatases. *Progress in lipid research* 2009;48:307-43.
82. Sbrissa D, Ikononov O, Strakova J, Shisheva A. Role for a Novel Signaling Intermediate, Phosphatidylinositol 5-Phosphate, in Insulin-Regulated F-Actin Stress Fiber Breakdown and GLUT4 Translocation. *Endocrinology* 2004;145:4853-65.
83. Sbrissa D, Ikononov OC, Deeb R, Shisheva A. Phosphatidylinositol 5-phosphate biosynthesis is linked to PIKfyve and is involved in osmotic

- response pathway in mammalian cells. *Journal of Biological Chemistry* 2002;277:47276-84.
84. Schaletzky J, Dove SK, Short B, Lorenzo O, Clague MJ, Barr FA. Phosphatidylinositol-5-phosphate activation and conserved substrate specificity of the myotubularin phosphatidylinositol 3-phosphatases. *Current Biology* 2003;13:504-9.
 85. Sebastian C, Satterstrom FK, Haigis M, Mostoslavsky R. From Sirtuin Biology to Human Diseases: An Update. *Journal of Biological Chemistry* 2012;287:42444-52.
 86. Shi X. The fellowships of the INGs. *Journal of cellular biochemistry* 2005;96:1127-36.
 87. Smith J. Human Sir2 and the silencing of p53 activity. *Trends in Cell Biology* 2002;12:404-6.
 88. Tang Y, Zhao W, Chen Y, Zhao Y, Gu W. Acetylation Is Indispensable for p53 Activation. *Cell* 2008;133:612-26.
 89. Terebiznik M, Vieira OV, Marcus SL, Slade A, Yip CM, Trimble WS, et al. Elimination of host cell PtdIns(4,5)P(2) by bacterial SigD promotes membrane fission during invasion by Salmonella. *Nat Cell Biol* 2002;4:766-73.
 90. Tolias K, Rameh L, Ishihara H, Shibasaki Y, Chen J, Prestwich G, et al. Type I Phosphatidylinositol-4-phosphate 5-Kinases Synthesize the Novel Lipids Phosphatidylinositol 3,5-Bisphosphate and Phosphatidylinositol 5-Phosphate. *Journal of Biological Chemistry* 1998;273:18040-6.
 91. Tronchere H, Laporte J, Pendariest C, Chaussade C, Liaubet L, Pirola L, et al. Production of phosphatidylinositol 5-phosphate by the phosphoinositide 3-phosphatase myotubularin in mammalian cells. *J Biol Chem* 2004;279:7404-312.
 92. van den Bout I, Divecha N. PIP5K-driven PtdIns(4,5)P₂ synthesis: regulation and cellular functions. *Journal of Cell Science* 2009;122:3837-50.
 93. Vaziri H, Dessain SK, Ng Eaton E, Imai SI, Frye RA, Pandita TK, et al. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* 2001;107:149-59.
 94. Volpicelli-Daley L, Lucast L, Gong L, Liu L, Sasaki J, Sasaki T, et al. Phosphatidylinositol-4-Phosphate 5-Kinases and Phosphatidylinositol 4,5-Bisphosphate Synthesis in the Brain. *Journal of Biological Chemistry* 2010;285:28708-14.
 95. Wang C, Chen L, Hou X, Li Z, Kabra N, Ma Y, et al. Interactions between E2F1 and SirT1 regulate apoptotic response to DNA damage. *Nat Cell Biol* 2006;8:1025-31.
 96. Wang M, Bond N, Letcher A, Richardson J, Lilley K, Irvine R, et al. Genomic tagging reveals a random association of endogenous PI(5)P 4-kinases alpha and beta and a partial nuclear localization of the alpha isoform. *Biochemical Journal* 2010;430:215-21.
 97. Westphal CH. A therapeutic role for sirtuins in diseases of aging? *Trends in Biochemical Sciences* 2007;32:555-60.
 98. Wong S. Deacetylation of the retinoblastoma tumour suppressor protein by SIRT1. *Biochemical Journal* 2007;407:451-60.
 99. Xiong S, Salazar G, Patrushev N, Alexander RW, et al. FoxO1 mediates an autofeedback loop regulating SIRT1 expression. *Journal of Biological Chemistry* 2011;286:5289-99.
 100. Yamakuchi M. MicroRNA Regulation of SIRT1. *Frontiers in physiology* 2012;3:68.
 101. Yang Y, Fu W, Chen J, Olashaw N, Zhang X, Nicosia SV, et al. SIRT1 sumoylation regulates its deacetylase activity and cellular response to genotoxic stress. *Nat Cell Biol* 2007;9:1253-62.
 102. Yuan J, Luo K, Liu T, Lou Z. Regulation of SIRT1 activity by genotoxic stress. *Genes & development* 2012;26:791-6.
 103. Yuan Z, Wang SY, Fleuriet C, Leprince D, Rocheleau JV, Piston DW, et al. SIRT1 regulates the function of the Nijmegen breakage syndrome protein. *Molecular Cell* 2007;27:149-62.
 104. Zhang Q, Wang SY, Fleuriet C, Leprince D, Rocheleau JV, Piston DW, et al. Metabolic regulation of SIRT1 transcription via a HIC1:CtBP corepressor complex. *Proc Natl Acad Sci U S A* 2007;104:829-33.
 105. Zhao W, Kruse JP, Tang Y, Jung SY, Qin J, Gu W. Negative regulation of the deacetylase SIRT1 by DBC1. *Nature* 2008;451:587-90.
 106. Zou J, Marjanovic J, Kisseleva MV, Wilson M, Majerus PW. Type I phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced apoptosis. *Proc Natl Acad Sci U S A* 2007;104:16834-9.

