



Universiteit
Leiden
The Netherlands

Clinical significance of T-cell clonality in mycosis fungoides and other cutaneous T-cell lymphomas

Muche, J.M.

Citation

Muche, J. M. (2010, May 20). *Clinical significance of T-cell clonality in mycosis fungoides and other cutaneous T-cell lymphomas*. Retrieved from <https://hdl.handle.net/1887/15546>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/15546>

Note: To cite this publication please use the final published version (if applicable).

8.2 Bibliography

Articles.

Lukowsky A, Muche JM, Möbs M, Assaf C, Humme D, Hummel M, Sterry W, Steinhoff M. Evaluation of T-cell clonality in archival skin biopsy samples of cutaneous T-cell lymphomas using the Biomed-2 PCR protocol. *Diagn Mol Pathol* 2009 (accepted)

Steinhoff M, Schöpp S, Assaf C, Muche JM, Beyer M, Sterry W, Lukowsky A. Prevalence of genetically defined tumor cells in CD7 as well as CD26 positive and negative circulating T-cell subsets in Sézary syndrome. *Leuk Res.* 2009; 33(1):88-99.

Gellrich S, Muche JM, Wilks A, Jasch KC, Voit C, Fischer T, Audring H, Sterry W. Systemic eight-cycle anti-CD20 monoclonal antibody (rituximab) therapy in primary cutaneous B-cell lymphomas--an applicational observation. *Br J Dermatol.* 2005; 153: 167-73

Muche JM, Toppe E, Sterry W, Haas N. Palpable arciform migratory erythema in an HIV patient, a CD8+ pseudolymphoma. *J Cutan Pathol.* 2004;31: 379-82.

Gellrich S, Wernicke M, Wilks A, Lukowsky A, Muche JM, Jasch KC, Audring H, Mason D, Sterry W. The cell infiltrate in lymphomatoid papulosis comprises a mixture of polyclonal large atypical cells (CD30-positive) and smaller monoclonal T-cells (CD30-negative). *J Invest Dermatol.* 2004; 122: 859-61

Muche JM, Karenko L, Gellrich S, Karhu R, Kytola S, Kahkonen M, Lukowsky A, Sterry W, Ranki A. Cellular coincidence of clonal T cell receptor rearrangements and complex clonal chromosomal aberrations-a hallmark of malignancy in cutaneous T cell lymphoma. *J Invest Dermatol.* 2004; 122: 574-8.

Fischer TC, Gellrich S, Muche JM, Sherev T, Audring H, Neitzel H, Walden P, Sterry W, Tonnies H. Genomic aberrations and survival in cutaneous T cell lymphomas. *J Invest Dermatol.* 2004; 122:579-86

Bohmeyer J, Stadler R, Kremer A, Nashan D, Muche JM, Gellrich S, Luger T, Sterry W. Bexarotene--an alternative therapy for progressive cutaneous T-cell lymphoma? First experiences. *J Dtsch Dermatol Ges.* 2003; 10: 785-9

Kallinich T, Muche JM, Qin S, Sterry W, Audring H, Kroczeck RA. Chemokine receptor expression on neoplastic and reactive T cells in the skin at different stages of mycosis fungoides. *J Invest Dermatol.* 2003; 121: 1045-52

Tumenjargal S, Gellrich S, Linnemann T, Muche JM, Lukowsky A, Audring H, Wiesmuller KH, Sterry W, Walden P. Anti-tumor immune responses and tumor regression induced with mimotopes of a tumor-associated T cell epitope. *Eur J Immunol.* 2003; 33: 3175-85

Muche JM, Sterry W, Gellrich S, Rzany B, Audring H, Lukowsky A. Peripheral blood T-cell clonality in mycosis fungoides and nonlymphoma controls. *Diagn Mol Pathol.* 2003; 12: 142-50

Rosler MJ, Rappl G, Muche JM, Hasselmann DO, Sterry W, Tilgen W, Reinhold U. No evidence of

skin infection with *Chlamydia pneumoniae* in patients with cutaneous T cell lymphoma. *Clin Microbiol Infect.* 2003; 9: 721-3

Kolde G, Muche JM, Schulze P, Fischer P, Lichey J. Infliximab: a promising new treatment option for ulcerated necrobiosis lipoidica. *Dermatology.* 2003; 206: 180-1

Haas N, Hauptmann S, Paralikoudi D, Muche JM, Kolde G. Interferon-gamma treatment induces granulomatous tissue reaction in a case of localized cutaneous leishmaniasis. *Am J Dermatopathol.* 2002; 24: 319-23

Lukowsky A, Richter S, Dijkstal K, Sterry W, Muche JM. A T-cell receptor gamma polymerase chain reaction assay using capillary electrophoresis for the diagnosis of cutaneous T-cell lymphomas. *Diagn Mol Pathol.* 2002; 11: 59-66

Rappl G, Abken H, Muche JM, Sterry W, Tilgen W, Andre S, Kaltner H, Ugurel S, Gabius HJ, Reinhold U. CD4+CD7- leukemic T cells from patients with Sezary syndrome are protected from galectin-1-triggered T cell death. *Leukemia.* 2002; 16: 840-5

Gellrich S, Rutz S, Golembowski S, Jacobs C, von Zimmermann M, Lorenz P, Audring H, Muche JM, Sterry W, Jahn S. Primary cutaneous follicle center cell lymphomas and large B cell lymphomas of the leg descend from germinal center cells. A single cell polymerase chain reaction analysis. *J Invest Dermatol.* 2001; 117: 1512-20

Hombach A, Muche JM, Gerken M, Gellrich S, Heuser C, Pohl C, Sterry W, Abken H. T cells engrafted with a recombinant anti-CD30 receptor target autologous CD30(+) cutaneous lymphoma cells. *Gene Ther.* 2001 8: 891-5

Gellrich S, Muche JM, Pelzer K, Audring H, Sterry W. [Anti-CD20 antibodies in primary cutaneous B-cell lymphoma. Initial results in dermatologic patients] *Hautarzt.* 2001; 52:205-10. German.

Ahnhudt C, Muche JM, Dijkstal K, Sterry W, Lukowsky A. An approach to the sensitivity of temperature-gradient gel electrophoresis in the detection of clonally expanded T-cells in cutaneous T-cell lymphoma. *Electrophoresis.* 2001; 22: 33-8

Rappl G, Muche JM, Abken H, Sterry W, Tilgen W, Ugurel S, Reinhold U. CD4(+)CD7(-) T cells compose the dominant T-cell clone in the peripheral blood of patients with Sezary syndrome. *J Am Acad Dermatol.* 2001; 44: 456-61

Jung M, Muche JM, Lukowsky A, Jung K, Loening SA. Dimethyl sulfoxide as additive in ready-to-use reaction mixtures for real-time polymerase chain reaction analysis with SYBR Green I dye. *Anal Biochem.* 2001; 289: 292-5

Gellrich S, Lukowsky A, Schilling T, Rutz S, Muche JM, Jahn S, Audring H, Sterry W. Microanatomical compartments of clonal and reactive T cells in mycosis fungoides: molecular demonstration by single cell polymerase chain reaction of T cell receptor gene rearrangements. *J Invest Dermatol.* 2000; 115: 620-4

Muche JM, Lukowsky A, Ahnhudt C, Gellrich S, Sterry W. Peripheral blood T cell clonality in mycosis fungoides -an independent prognostic marker? *J Invest Dermatol.* 2000; 115: 504-5

Lukowsky A, Muche JM, Sterry W, Audring H. Detection of expanded T cell clones in skin biopsy samples of patients with lichen sclerosus et atrophicus by T cell receptor-gamma polymerase chain reaction assays. *J Invest Dermatol.* 2000; 115: 254-9

Jung M, Muche JM, Jung K, Loening SA. Methods in molecular biology: minor errors in primer citations with major consequences: how can we minimize these mistakes? *Br J Cancer.* 2000; 82: 1611-3

Muche JM, Lukowsky A, Heim J, Friedrich M, Audring H, Sterry W. Demonstration of frequent occurrence of clonal T cells in the peripheral blood but not in the skin of patients with small plaque parapsoriasis. *Blood.* 1999; 94:1409-17

Linnemann T, Brock C, Sparbier K, Muche JM, Mielke A, Lukowsky A, Sterry W, Kaltoft K, Wiesmuller KH, Walden P. Identification of epitopes for CTCL-specific cytotoxic T lymphocytes. *Adv Exp Med Biol.* 1998; 451: 231-5

Worm M, Muche JM, Schulze P, Sterry W, Kolde G. Hypocomplementaemic urticarial vasculitis: successful treatment with cyclophosphamide-dexamethasone pulse therapy. *Br J Dermatol.* 1998; 139: 704-7

Lukowsky A, Audring H, Heiduk U, Muche JM, Richter S, Sterry W. [Detection of monoclonal T-cells with TCR gamma PCR in mycosis fungoides] *Hautarzt.* 1998 ; 49: 641-5. German.

Scheller U, Muche JM, Sterry W, Lukowsky A. Detection of clonal T cells in cutaneous T cell lymphoma by polymerase chain reaction: comparison of mutation detection enhancement-polyacrylamide gel electrophoresis, temperature gradient gel electrophoresis and fragment analysis of sequencing gels. *Electrophoresis.* 1998; 19: 653-8

Rutz S, Siegel P, Muche JM, Sterry W, Jahn S. Nucleotide sequence comparison of the IgE constant region in patients with atopic dermatitis and non-atopic individuals. *Exp Clin Immunogenet.* 1997; 14:166-72

Muche JM, Lukowsky A, Asadullah K, Gellrich S, Sterry W. Demonstration of frequent occurrence of clonal T cells in the peripheral blood of patients with primary cutaneous T-cell lymphoma. *Blood.* 1997; 90:1636-42

Chapters in books and reviews.

Muche JM. Cutaneous Malignant Lymphoma. Value and Limits of the Detection of Clonal T-Cells. *Akt Dermatol.* 2004; 30: 559-565. German.

Gellrich S, Muche JM, Jahn S, Sterry W. Studying primary cutaneous lymphomas by micromanipulation and single cell PCR – A review. *Giornale Italiano di Dermatologia e venerologia* 2002; 137: 231-238

Muche JM, Sterry W. Vaccination therapy for cutaneous T-cell lymphoma. *Clin Exp Dermatol.* 2002; 27: 602-7

- Muche JM, Sterry W: Erythroderma. In: Bologna, Jorizzo, Rapini. Dermatology. 2003; Mosby
- Sterry W, Muche JM: Other systemic lymphomas with skin infiltrations. In: Freedberg et al. Fitzpatrick's Dermatology in General Medicine. 6th Edition. 2003; McGraw-Hill
- Muche JM, Gellrich S, Sterry W: Therapie kutaner Lymphome. In: Bruhn, Kneba, Löffler: Antineoplastische Therapie. Schattauer Verlag Stuttgart, 3., überarb. u. erw. Aufl 2002
- Muche JM, Sterry W: Tumor biology in cutaneous malignant lymphomas. J Dermatol. 2001; 28: 606-9
- Muche JM, Gellrich S, Sterry W: Treatment of cutaneous T cell lymphomas. Sem Cut Med Surg. 2000; 19: 142-148
- Gellrich S, Muche JM, Audring H, Sterry W: Kutane B-Zell-Lymphome. Hautarzt. 2000; 51: 363-375
- Muche JM, Sterry W: Lymphome, primär kutane. In: Pschyrembel Therapeutisches Wörterbuch, 2. Auflage, de Gruyter Berlin, 2000, pp537
- Asadullah K, Muche JM, Friedrich M, Sterry W: Zirkulierende maligne und benigne Lymphozyten in Frühstadien kutaner Lymphome. In: Plettenberg et al.: Dermatologie an der Schwelle zum neuen Jahrtausend. 463-466, 1999
- Jahn S, Muche JM, Lukowsky A, Sterry W: Malignant cutaneous lymphomas - Pathogenesis, clinical course, and molecularbiological diagnosis. Zeitschrift für Hautkrankheiten. 1997; 72: 15-22

8.3 Acknowledgement

This thesis is the result of collaboration of and support from many persons. Some of them deserve special acknowledgement:

Ansgar Lukowsky. Nicht der Doktorvater meiner Deutschen Promotion und nicht der Promotor meiner Niederländischen "Prüfschrift". Und doch der Vater all meiner Kenntnisse und Fertigkeiten im molekularbiologischen Labor.

Rein Willemze. Voor het begeleiden van het hele traject van dit proefschrift. Maar vooral voor zijn manier om ambitie te wekken. De ambitie om de resultaten uit de onderliggende studies te integreren tot een sluitende visie op het geheel.

Wolfram Sterry. Für meine Kenntnisse über kutane Lymphome und für das Ermöglichen der Untersuchungen, die dieser Arbeit zugrunde liegen.

Annamari Ranki and Leena Karenko. For the most determined and at the same time most pleasant collaboration I've ever had.

Sigbert Jahn, Sylke Gellrich, Cordula Ahnhudt, Uta Scheller und Tanja Fischer. Das verschworene Team von den Labors in der 5. Etage.

Ulrike Heiduk, Sarina Richter, Katja Schreiber und Katharina Dijkstal. Die guten Feen im Labor.

Peter Steijlen. Voor het idee aan dit werk te beginnen.

Claudia, Felix und Amelie. Für die leibliche Versorgung, das Entbehren unseres Computers und das Ertragen meiner unausgeschlafenen Morgenstunden.

Mein Vater. Für die regelmäßige Erinnerung an die Fertigstellung dieser Arbeit.