



Universiteit
Leiden
The Netherlands

Multimodality imaging of coronary artery bypass grafts

Salm, L.P.

Citation

Salm, L. P. (2006, November 7). *Multimodality imaging of coronary artery bypass grafts*. Retrieved from <https://hdl.handle.net/1887/4978>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4978>

Note: To cite this publication please use the final published version (if applicable).



PART I

PREFACE

CHAPTER 1

Introduction

CORONARY ARTERY BYPASS GRAFT SURGERY

Coronary heart disease (CHD) is the most common cause of death worldwide, accounting for nearly 2 million deaths each year in Europe alone (1). CHD is most commonly due to obstruction of the coronary arteries by an atheromatous plaque (2). Manifestations of CHD include unstable and chronic angina pectoris, and myocardial infarction. Therapeutic options for CHD are directed towards prevention of further progression of disease or symptom-relief. This is achieved by obtaining revascularization either by percutaneous coronary intervention or coronary artery bypass graft surgery (CABG), if pharmacological treatment alone is insufficient.

In CABG, revascularization of the myocardium is obtained by rerouting or bypassing the bloodstream around the coronary obstruction. Garrett et al. performed the first human CABG in 1967 as a "bailout" procedure (3). Since then, the surgical technique is further developed and refined. At first, graft material was harvested from the saphenous veins in the lower limbs. Subsequently, arteries were also used as bypass grafts, such as left and right internal mammary arteries (IMA), radial artery and gastro-epiploic artery.

CABG was demonstrated to be an effective treatment for CHD in short- and intermediate-term (4;5). However, graft disease occurs frequently, beginning with accelerated atherosclerosis early after surgery in vein grafts (6;7). Arterial grafts have a superior outcome compared with vein grafts, still obstruction and occlusion do occur (8;9). In a recent study, comparing the outcome of vein and IMA grafts, patency at 10 years was 61% for vein grafts compared with 85% for IMA grafts ($p < 0.001$) (10).

The gold standard for evaluating bypass grafts is coronary angiography. Although coronary angiography offers a rapid and complete exploration of the native coronary arteries and grafts, it is an invasive procedure, requiring arterial puncture, x-ray exposition, and hospitalization. Complications include ventricular arrhythmias, myocardial infarction, cardiac perforation, necessary emergency CABG, and death, even though the risk is small (11;12). Furthermore, coronary angiography does not provide a functional assessment of the myocardial region, perfused by the respective coronary artery or graft. A noninvasive anatomical and functional examination of the coronary arteries and bypass grafts would be preferable to detect insufficiency and identify a target for re-intervention.

NONINVASIVE IMAGING OF CORONARY ARTERY BYPASS GRAFTS

Several noninvasive imaging modalities are available in order to detect bypass graft insufficiency. Using cardiovascular magnetic resonance (CMR), anatomy and function of bypass grafts may be assessed. MR angiography allows an accurate evaluation of patency in both arterial and vein grafts (13-15). The degree of graft stenosis can, thus far, only be evaluated in vein graft bodies, excluding the recipient coronary arteries (16).

A functional parameter that may be evaluated by CMR is the blood flow velocity. Blood flow velocity of native coronary arteries and bypass grafts can be derived invasively at rest and during pharmacological stress using the Doppler flow wire (17). Subsequent calculation of the coronary flow velocity reserve (CFVR) by dividing the stress by the rest parameter provides insight in the physiological condition of the vessel (18;19). Previous studies have shown that flow velocity measurements may be used to evaluate the

hemodynamic consequences of stenoses in native coronary arteries (20;21), and bypass grafts (22), although studies focusing on bypass grafts are few. Using the phase-contrast MR technique, it is feasible to perform flow velocity measurements noninvasively by CMR in both vein (15;23) and arterial grafts (24). The value of CMR flow velocity measurements in predicting the presence of a stenosis in vein grafts was successfully researched (25;26). CMR flow velocity measurements of arterial grafts remain challenging because of the small luminal diameter of the internal mammary arteries, and metal clip artefacts. In earlier studies demonstrating the feasibility to measure flow in left IMA grafts, CMR sequences were limited by low spatial and temporal resolution (24;27;28).

With computed tomography (CT) angiography, assessment of bypass graft patency was extensively investigated (29-31). When it became possible to use multiple detectors in a single spiral CT acquisition, graft patency could be assessed with very high accuracy (32;33). The evaluation of stenoses in grafts by multi-detector row CT is still a challenge because of motion artefacts, metal clip artefacts or beam hardening from calcium deposits (32-34). Since a cardiac CT examination to evaluate native coronary arteries and bypass grafts requires scanning of the entire heart, reconstruction of the left ventricular (LV) function is also achievable. Several studies showed that 4-detector row CT allowed accurate assessment of global and regional LV function (35;36). Validation of the next generation CT equipment in assessing LV function has yet to be conducted.

Single-photon emission computed tomography (SPECT) perfusion imaging is a well-established technique by which to detect CHD by evaluating regional myocardial perfusion at rest and during stress. Excellent sensitivities and specificities of SPECT to detect stenoses in native coronary arteries (37;38) and bypass grafts (39) were demonstrated. More recently, for SPECT imaging the emphasis has shifted from detection of CHD to hemodynamic evaluation of stenoses. Comparative studies between SPECT imaging and invasive assessment using the Doppler flow wire have been performed, demonstrating a good agreement (ranging from 72% to 96%) between these two techniques for the evaluation of the hemodynamic consequence of an intermediate native coronary artery stenosis (21;40;41). Studies evaluating the hemodynamic consequences of bypass graft lesions are lacking.

OUTLINE OF THE THESIS

The aim of the thesis was to describe multiple modalities to examine coronary artery bypass grafts, and to further develop noninvasive imaging techniques to detect or exclude stenoses in native coronary arteries and bypass grafts in patients who experienced recurrent chest pain after CABG.

Part I of the thesis provides an introduction to noninvasive imaging. In chapter 2, noninvasive imaging of coronary artery bypass grafts by CMR and CT is reviewed in detail.

Part II of the thesis focuses on CMR flow velocity imaging in vein and arterial grafts. In chapter 3, a model to detect or exclude stenoses in vein grafts by CMR volumetric flow measurements is described. In chapter 4, two analysis methods for the CMR flow velocity maps, i.e. velocity and volumetric flow, are compared, using flow velocity

maps of vein grafts. In chapter 5, the functional significance of bypass graft stenoses is investigated by SPECT perfusion imaging and CMR, and compared with findings at coronary angiography. In chapter 6, a novel CMR phase-contrast sequence to measure flow velocity in arterial and vein grafts is introduced and validated.

Part III of the thesis concentrates on multi-detector row CT imaging of coronary artery bypass grafts. In chapter 7, a comprehensive assessment by 16-detector row CT of patients after CABG is investigated. In chapter 8, global and regional LV function evaluated by 16-detector row CT is compared with evaluation by echocardiography and CMR.

Part IV focuses on the hemodynamic consequences of vein graft lesions. In chapter 9, the relative merits of invasive Doppler flow velocity measurements and SPECT perfusion imaging are studied in assessing the hemodynamic significance of stenoses in vein grafts, and the results are related to coronary angiography.

REFERENCES

1. British Heart Foundation. European Cardiovascular Disease Statistics. 2000.
2. Morrow DA, Gersh BJ, Braunwald E. Chronic coronary artery disease. In: Braunwald's heart disease. A textbook of cardiovascular medicine. 2005.
3. Garrett HE, Dennis EW, DeBakey ME. Aortocoronary bypass with saphenous vein graft. Seven-year follow-up. *Jama* 1973;223:792-4.
4. Yusuf S, Zucker D, Peduzzi P et al. Effect of coronary artery bypass graft surgery on survival: overview of 10-year results from randomised trials by the Coronary Artery Bypass Graft Surgery Trialists Collaboration. *Lancet* 1994;344:563-70.
5. Davis KB, Chaitman B, Ryan T, Bittner V, Kennedy JW. Comparison of 15-year survival for men and women after initial medical or surgical treatment for coronary artery disease: a CASS registry study. *Coronary Artery Surgery Study*. *J Am Coll Cardiol* 1995;25:1000-9.
6. Motwani JG, Topol EJ. Aortocoronary saphenous vein graft disease: pathogenesis, predisposition, and prevention. *Circulation* 1998;97:916-31.
7. Fitzgibbon GM, Kafka HP, Leach AJ, Keon WJ, Hooper GD, Burton JR. Coronary bypass graft fate and patient outcome: angiographic follow-up of 5,065 grafts related to survival and reoperation in 1,388 patients during 25 years. *J Am Coll Cardiol* 1996;28:616-26.
8. Loop FD, Lytle BW, Cosgrove DM et al. Influence of the internal-mammary-artery graft on 10-year survival and other cardiac events. *N Engl J Med* 1986;314:1-6.
9. Cameron A, Kemp HG, Jr., Green GE. Bypass surgery with the internal mammary artery graft: 15 year follow-up. *Circulation* 1986;74:III30-III36.
10. Goldman S, Zadina K, Moritz T et al. Long-term patency of saphenous vein and left internal mammary artery grafts after coronary artery bypass surgery. Results from a department of Veteran Affairs Cooperative study. *J Am Coll Cardiol* 2004;44:2149-56.
11. Krone RJ, Johnson L, Noto T. Five year trends in cardiac catheterization: a report from the Registry of the Society for Cardiac Angiography and Interventions. *Cathet Cardiovasc Diagn* 1996;39:31-5.
12. Wyman RM, Safian RD, Portway V, Skillman JJ, McKay RG, Baim DS. Current complications of diagnostic and therapeutic cardiac catheterization. *J Am Coll Cardiol* 1988;12:1400-6.
13. Bunce NH, Lorenz CH, John AS, Lesser JR, Mohiaddin RH, Pennell DJ. Coronary artery bypass graft patency: assessment with true fast imaging with steady-state precession versus gadolinium-enhanced MR angiography. *Radiology* 2003;227:440-6.
14. Wintersperger BJ, Engelmann MG, von Smekal A et al. Patency of coronary bypass grafts: assessment with breath-hold contrast-enhanced MR angiography--value of a non-electrocardiographically triggered technique. *Radiology* 1998;208:345-51.
15. Galjee MA, van Rossum AC, Doesburg T, van Eenige MJ, Visser CA. Value of magnetic resonance imaging in assessing patency and function of coronary artery bypass grafts. An angiographically controlled study. *Circulation* 1996;93:660-6.
16. Langerak SE, Vliegen HW, de Roos A et al. Detection of Vein Graft Disease using High-resolution Magnetic Resonance Angiography. *Circulation* 2002;105:328-33.
17. Doucette JW, Corl PD, Payne HM et al. Validation of a Doppler guide wire for intravascular measurement of coronary artery flow velocity. *Circulation* 1992;85:1899-911.
18. Gould KL, Kirkeeide RL, Buchi M. Coronary flow reserve as a physiologic measure of stenosis severity. *J Am Coll Cardiol* 1990;15:459-74.
19. White CW, Wright CB, Doty DB et al. Does visual interpretation of the coronary arteriogram predict the physiologic importance of a coronary stenosis? *N Engl J Med* 1984;310:819-24.
20. Ofili EO, Labovitz AJ, Kern MJ. Coronary flow velocity dynamics in normal and diseased arteries. *Am J Cardiol* 1993;71:3d-9d.
21. Chamuleau SA, Tio RA, de Cock CC et al. Prognostic value of coronary blood flow velocity and myocardial perfusion in intermediate coronary narrowings and multivessel disease. *J Am Coll Cardiol* 2002;39:852-8.
22. White CW. Clinical applications of Doppler coronary flow reserve measurements. *Am J Cardiol* 1993;71:10d-6d.
23. Langerak SE, Kunz P, Vliegen HW et al. Improved MR flow mapping in coronary artery bypass grafts during adenosine-induced stress. *Radiology* 2001;218:540-7.
24. Ishida N, Sakuma H, Cruz BP et al. Mr flow measurement in the internal mammary artery-to-coronary artery bypass graft: comparison with graft stenosis at radiographic angiography. *Radiology* 2001;220:441-7.
25. Langerak SE, Vliegen HW, Jukema JW et al. Value of magnetic resonance imaging for the noninvasive detection of stenosis in coronary artery bypass grafts and recipient coronary arteries. *Circulation* 2003;107:1502-8.

26. Bedaux WL, Hofman MB, Vyt SL, Bronzwaer JG, Visser CA, van Rossum AC. Assessment of coronary artery bypass graft disease using cardiovascular magnetic resonance determination of flow reserve. *J Am Coll Cardiol* 2002;40:1848-55.
27. Debatin JF, Strong JA, Sostman HD et al. MR characterization of blood flow in native and grafted internal mammary arteries. *J Magn Reson Imaging* 1993;3:443-50.
28. Sakuma H, Globits S, O'Sullivan M et al. Breath-hold MR measurements of blood flow velocity in internal mammary arteries and coronary artery bypass grafts. *J Magn Reson Imaging* 1996;6:219-22.
29. Daniel WG, Dohring W, Stender HS, Lichtlen PR. Value and limitations of computed tomography in assessing aortocoronary bypass graft patency. *Circulation* 1983;67:983-7.
30. Bateman TM, Gray RJ, Whiting JS et al. Prospective evaluation of ultrafast cardiac computed tomography for determination of coronary bypass graft patency. *Circulation* 1987;75:1018-24.
31. Stanford W, Brundage BH, MacMillan R et al. Sensitivity and specificity of assessing coronary bypass graft patency with ultrafast computed tomography: results of a multicenter study. *J Am Coll Cardiol* 1988;12:1-7.
32. Ropers D, Ulzheimer S, Wenkel E et al. Investigation of aortocoronary artery bypass grafts by multislice spiral computed tomography with electrocardiographic-gated image reconstruction. *Am J Cardiol* 2001;88:792-5.
33. Nieman K, Pattynama PM, Rensing BJ, van Geuns RJ, de Feyter PJ. Evaluation of patients after coronary artery bypass surgery: CT angiographic assessment of grafts and coronary arteries. *Radiology* 2003;229:749-56.
34. Marano R, Storto ML, Maddestra N, Bonomo L. Non-invasive assessment of coronary artery bypass graft with retrospectively ECG-gated four-row multi-detector spiral computed tomography. *Eur Radiol* 2004;14:1353-62.
35. Dirksen MS, Bax JJ, de Roos A et al. Usefulness of dynamic multislice computed tomography of left ventricular function in unstable angina pectoris and comparison with echocardiography. *Am J Cardiol* 2002;90:1157-60.
36. Juergens KU, Grude M, Maintz D et al. Multi-detector row CT of left ventricular function with dedicated analysis software versus MR imaging: initial experience. *Radiology* 2004;230:403-10.
37. Mahmarijan JJ, Boyce TM, Goldberg RK, Cocanougher MK, Roberts R, Verani MS. Quantitative exercise thallium-201 single photon emission computed tomography for the enhanced diagnosis of ischemic heart disease. *J Am Coll Cardiol* 1990;15:318-29.
38. Maddahi J, Kiat H, Van Train KF et al. Myocardial perfusion imaging with technetium-99m sestamibi SPECT in the evaluation of coronary artery disease. *Am J Cardiol* 1990;66:55E-62E.
39. Lakkis NM, Mahmarijan JJ, Verani MS. Exercise thallium-201 single photon emission computed tomography for evaluation of coronary artery bypass graft patency. *Am J Cardiol* 1995;76:107-11.
40. Deychak YA, Segal J, Reiner JS et al. Doppler guide wire flow-velocity indexes measured distal to coronary stenoses associated with reversible thallium perfusion defects. *Am Heart J* 1995;129:219-27.
41. Verberne HJ, Piek JJ, van Liebergen RA, Koch KT, Schroeder-Tanka JM, van Royen EA. Functional assessment of coronary artery stenosis by doppler derived absolute and relative coronary blood flow velocity reserve in comparison with (99m)Tc MIBI SPECT. *Heart* 1999;82:509-14.