



Universiteit
Leiden
The Netherlands

Primary cilia on endothelial cells : component of the shear stress sensor localized to athero-prone flow areas

Heiden, K. van der

Citation

Heiden, K. van der. (2008, September 11). *Primary cilia on endothelial cells : component of the shear stress sensor localized to athero-prone flow areas*. Retrieved from <https://hdl.handle.net/1887/13093>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13093>

Note: To cite this publication please use the final published version (if applicable).

Chapter 7

Deciphering the Endothelial Shear Stress Sensor

Robert E. Poelmann; Kim Van der Heiden; Adriana C. Gittenberger-de Groot; Beerend P. Hierck.

Department of Anatomy and Embryology, Leiden University Medical Center, Leiden, The Netherlands.

Circulation. 2008;117:1124-1126. Adapted for this thesis.

The single non-motile primary cilium, protruding several microns from the apical surface, contains cytoskeletal elements in a specific fashion. It consists of 9 circularly arranged microtubule doublets, as revealed by transmission electron microscopy¹, but lacks the central ones characteristic for motile cilia and flagellae. The primary cilium is anchored to the basal body and is thereby connected to the cytoskeletal apparatus. Sorokin¹ concluded that primary cilia were probably vestigial remnants of motile cilia. We now know that ciliary functions are abundant. Examples include processes involving Hedgehog and Wnt signaling² and determining left-right asymmetry³. Cilia functioning as sensory antennas in insect ears and the human retina are well established⁴. Because of the widespread presence of primary cilia, it does not come as a surprise that many diseases, often syndromic, are caused by ciliary dysfunction. Several diseases are related to disruption of intraflagellar transport in the case of mutation in, for example, Polaris, *Kif3a* or various Bbs proteins that can lead to such conditions as obesity and Bardet Biedl Syndrome⁵. Other cilium-related proteins involve cell membrane-bound Ca^{2+} channels such as the complex formed by polycystin-1 and -2, encoded from *Pkd1* and *Pkd2*, respectively. Mutations cause polycystic kidney diseases^{6,7}.

Primary cilia were first described in the cardiovascular system in human embryos and adults more than 20 years ago⁸. The geometry of the heart and vascular tree strongly influences hemodynamics with repercussions on the pattern of ciliation. Endothelial ciliation is restricted to areas of low and oscillatory blood flow and is absent in areas of high flow and ensuing wall shear stress and shear-responsive gene expression⁹. Endothelial cells in the adult mouse are ciliated in areas of low and disturbed blood flow¹⁰ in the inner curve of the aortic arch and the branching points of the major arteries. These are areas that are prone to developing atherosclerosis. The ciliated phenotype is typically induced by disturbed or oscillating shear stress. It is noteworthy that patients with polycystic kidney disease present with an increased risk for atherosclerosis and hypertension¹¹.

Nauli *et al.*¹² in the current issue of *Circulation*, proceeding from their thorough and extensive program related to polycystic kidney disease^{6,13}, describe the ability of immortalized and cultured murine endothelial cells to sense low shear ranges using primary cilia. *Polycystin-1*- and *polaris*-deficient cells were used that are unable to transmit flow-related shear stress into Ca^{2+} signaling and nitric oxide (NO) synthesis. They mention that polycystin-1 in the basal body of the cilium is not in itself sufficient for a fluid shear stress response, whereas optimal shear stress does not change cilium structure but rather modifies the responsiveness to high stress by proteolytic modification of polycystin-1. In their article, Nauli *et al.*¹² demonstrate that nearly all immortalized endothelial cells carry a single primary cilium and that mechanical probing of a single cell in a monolayer causes a Ca^{2+} flux throughout the epithelium in a time frame of many seconds. These experiments carried out *in vitro* demonstrate the response potentials of endothelial cells.

If one examines the response time of the endothelial cells to fluid shear stress in terms of a Ca^{2+} flux and NO release, which is effectively 10 to 15 seconds from the trigger, one finds that the activation mechanism *in vivo* needs further exploration. Because the heart rate is 1 to 2 Hz in humans and even 7 to 8 Hz in mice, the pulse frequency is much faster than the endothelial response time. This difference would imply a refractory period for endothelial activation. Furthermore, we have demonstrated that *in vivo*, only a minority of endothelial cells (< 25%) are ciliated, which leaves large areas non-ciliated, nevertheless leading to relatively homogeneous expression patterns of shear responsive genes, such as *Krüppel-like factor-2 (KLF2)*, *Endothelin-1 (ET1)*, or *endothelial NO synthase*^{14,15}. Figure 1 shows the presence of ciliated ventricular and atrial endocardial cells in wild-type and

Pkd1^{-/-} mouse embryos (kindly provided by Dr. D.J.M. Peters, LUMC, The Netherlands), demonstrating the presence of ciliated endothelial cells but also the lack of cilia on most neighboring cells. The intercellular Ca²⁺ exchange, nicely demonstrated by Nauli and colleagues¹², may serve in cilia-poor areas to synchronize the reaction of endothelial cells that are non-ciliated. This effect is most probably facilitated through intercellular ion channels but does not result in activation of NO synthesis and release. The latter aspect appears to be restricted to cells with functional cilia.

Iomini and colleagues¹⁶ demonstrated that endothelial cells lose their primary cilia after 1 to 2 hrs exposure to steady fluid flow. It is obvious that both prolonged steady and pulsed flows, which cause de-ciliation, affect the expression profiles and phenotypes of endothelial cells. These flow profiles typically result in the alignment of cells in the direction of the shear forces and in induction of, for example, *Klf2* and repression of NF-κB-mediated endothelial activation¹⁷. In contrast, disturbed or oscillatory flow induces a ciliated phenotype, induction of *Et1* expression, and endothelial activation. We must realize that sensitivity in ciliary mechanosensation is an important factor, as well, and is a matter of ongoing debate¹⁸.

Changing the shape of the cell and consequently the three-dimensional relations within the cytoskeleton have been described as a basis for mechanosensing and transduction of mechanical signals¹⁹ even in cells without cilia. Cytoskeletal deformation is instrumental in the response to prolonged shear forces by activating membrane proteins through conformational changes²⁰ or through activation of small G-proteins²¹. We have preliminary data that the cilium assists cytoskeletal deformation and acts as a signal amplifier in endothelial cells²².

Combining the now available data from *in vivo* and *in vitro* studies we are still left with some conflicting features: (1) The relation between pulse frequency and endothelial response time, (2) the endothelial heterogeneity in ciliated phenotype *in vivo* and the fact that non-ciliated cells are responsive to shear stress, and (3) the role of the cytoskeleton in mechanosensation. It is tempting to solve the dilemma by considering the shear stress sensing mechanism as a 2-step process: (1) an immediate response that involves ciliary bending, activation of polycystin-1 and -2, a rise in intracellular Ca²⁺, and a concomitant synthesis and release of stored (vasoactive) substances like NO and endothelin, and (2) a prolonged response that is coordinated through cytoskeletal conformational changes and that involves transcriptional activation of shear responsive genes, such as *KLF2*, which drive phenotypic adaptation of endothelial cells to this environmental epigenetic cue. Combining our efforts, we will be able in the near future to arrive at a more detailed plan for mechanosensation in the hemodynamic complexity of the cardiovascular system, both in development and in disease.




Figure 1. Confocal images of the endocardial cell layer of the ventricle (A and C) and the atrium (B and D) of a wild-type (*Wt*) C57BL/6 (A and B) and a *Pkd1*^{-/-} (C and D) mouse embryo of embryonic day 13.5. Only a subset of endocardial cells is ciliated (arrowheads), leaving many neighboring cells unciliated. *Pkd1* deficiency does not affect ciliation. In both *Wt* and *Pkd1*^{-/-} mouse embryos primary cilia are approximately 2 μ m in length and protrude from the endocardial cell into the lumen. Ciliated endocardial cells are only present in areas of disturbed or oscillatory blood flow such as the atrial and ventricular chambers, and are absent from high-shear areas such as the atrioventricular canal and outflow tract. Acetylated α -tubulin (green), and nuclei (red; propidium iodide). Scale bar = 10 μ m.

References

1. Sorokin S. Centrioles and the formation of rudimentary cilia by fibroblasts and smooth muscle cells. *J Cell Biol.* 1962;15:363-377.
2. Eggenschwiler JT, Anderson KV. Cilia and developmental signaling. *Annu Rev Cell Dev Biol.* 2007;23:345-373.
3. Levin M, Palmer AR. Left-right patterning from the inside out: widespread evidence for intracellular control. *BioEssays.* 2007;29:271-287.
4. Singla V, Reiter JF. The primary cilium as the cell's antenna: signaling at a sensory organelle. *Science.* 2006;313:629-633.
5. Satir P, Christensen ST. Overview of structure and function of mammalian cilia. *Annu Rev Physiol.* 2007;69:377-400.
6. Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Lil XG, Elia AEH, Lu WN, Brown EM, Quinn SJ, Ingber DE, Zhou J. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet.* 2003;33:129-137.
7. Weimbs T. Polycystic kidney disease and renal injury repair: common pathways, fluid flow, and the function of polycystin-1. *Am J Physiol Renal Physiol.* 2007;293:F1423-F1432.
8. Bystrevskaya VB, Lichkun VV, Antonov AS, Perov NA. An ultrastructural study of centriolar complexes in adult and embryonic human aortic endothelial cells. *Tissue Cell.* 1988;20:493-503.
9. Van der Heiden K, Groenendijk BCW, Hierck BP, Hogers B, Koerten HK, Mommaas AM, Gittenberger-de Groot AC, Poelmann RE. Monocilia on chicken embryonic endocardium in low shear stress areas. *Dev Dyn.* 2006;235:19-28.
10. Van der Heiden K, Hierck BP, Krams R, de Crom R, Cheng C, Baiker M, Pourquie MJB, Alkemade FE, DeRuiter MC, Gittenberger-de Groot AC, Poelmann RE. Endothelial primary cilia in areas of disturbed flow are at the base of atherosclerosis. *Atherosclerosis.* 2008;196:542-550.
11. Kocaman O, Oflaz H, Yekeler E, Dursun M, Erdogan D, Demirel S, Alisir S, Turgut F, Mercanoglu F, Ecker T. Endothelial dysfunction and increased carotid intima-media thickness in patients with autosomal dominant polycystic kidney disease. *Am J Kidney Dis.* 2004;43:854-860.
12. Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. Endothelial cilia are fluid-shear sensors that regulate calcium signaling and nitric oxide production through polycystin-1. *Circulation.* 2008;117:1161-1171.
13. Nauli SM, Rossetti S, Kolb RJ, Alenghat FJ, Consugar MB, Harris PC, Ingber DE, Loghman-Adham M, Zhou J. Loss of polycystin-1 in human cyst-lining epithelia leads to ciliary dysfunction. *J Am Soc Nephrol.* 2006;17:1015-1025.
14. Groenendijk BCW, Hierck BP, Vrolijk J, Baiker M, Pourquie MJB, Gittenberger-de Groot AC, Poelmann RE. Changes in shear stress-related gene expression after experimentally altered venous return in the chicken embryo. *Circ Res.* 2005;96:1291-1298.
15. Groenendijk BCW, Van der Heiden K, Hierck BP, Poelmann RE. The role of shear stress on *ET-1*, *KLF2*, and *NOS-3* expression in the developing cardiovascular system of chicken embryos in a venous ligation model. *Physiology (Bethesda).* 2007;22:380-389.
16. Iomini C, Tejada K, Mo W, Vaananen H, Piperno G. Primary cilia of human endothelial cells disassemble under laminar shear stress. *J Cell Biol.* 2004;164:811-817.
17. Helderma F, Segers D, de Crom R, Hierck BP, Poelmann RE, Evans PC, Krams R. Effect of shear stress on vascular inflammation and plaque development. *Curr Opin Lipidol.* 2007;18:527-533.
18. Resnick A, Hopfer U. Force-response considerations in ciliary mechanosensation. *Biophys J.* 2007;93:1380-1390.
19. Helmke BP, Davies PF. The cytoskeleton under external fluid mechanical forces: Hemodynamic forces acting on the endothelium. *Ann Biomed Eng.* 2002;30:284-296.
20. Tzima E, Irani-Tehrani M, Kiosses WB, Dejana E, Schultz DA, Engelhardt B, Cao G, DeLisser H, Schwartz MA. A mechanosensory complex that mediates the endothelial cell response to fluid shear stress. *Nature.* 2005;437:426-431.
21. Tzima E. Role of small GTPases in endothelial cytoskeletal dynamics and the shear stress response. *Circ Res.* 2006;98:176-185.
22. Hierck BP, Van der Heiden K, Alkemade FE, van de Pas S, van Thienen JV, Groenendijk BCW, Bax WH, Van der Laarse A, DeRuiter MC, Horrevoets AJG, Poelmann RE. Primary cilia sensitize endothelial cells for fluid shear stress. *Dev Dyn.* 2008;237:725-735.

