



Universiteit
Leiden
The Netherlands

Androgenic switch in barley microspores

De Faria Maraschin, Simone

Citation

De Faria Maraschin, S. (2005, February 9). *Androgenic switch in barley microspores*. Retrieved from <https://hdl.handle.net/1887/606>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/606>

Note: To cite this publication please use the final published version (if applicable).

II. Tissue-specific expression of 14-3-3 isoforms during barley microspore and zygotic embryogenesis

Acta Biol Cracov Bot (2003) 45: 103-106

Simone de Faria Maraschin, Jeanine D. Louwerse, Gerda E.M. Lamers, Herman P. Spaink,

Mei Wang

Abstract

The conserved 14-3-3 proteins have been shown to play regulatory roles in eukaryotic cells, including cell cycle control and differentiation. We were interested in the possible involvement of 14-3-3 proteins in the embryogenic process of barley (*Hordeum vulgare* L.). Barley microspore-derived embryo development was used as a model system. Immunolocalization of three barley 14-3-3 isoforms, 14-3-3A, 14-3-3B and 14-3-3C were carried out using isoform-specific antibodies. In immature microspore-derived embryos, 14-3-3C was specifically expressed underneath the L₁ layer of the shoot apical meristem and in the scutellum. Comparative studies showed that 14-3-3C was also expressed underneath the L₁ layer of the shoot apical meristem and in the scutellum of immature zygotic embryos. We further demonstrated that 14-3-3C expression is restricted to L₂ layer-derived cells of *in vitro* shoot meristematic cultures. Our results reveal that 14-3-3C isoform tissue-specific expression is closely related to defined events during differentiation processes in embryogenesis and *in vitro* meristematic cultures.

Introduction

The 14-3-3 proteins constitute a conserved family of 30 kD proteins present in all eukaryotic organisms studied so far. 14-3-3 dimers bind to phosphorylated motifs in a wide number of target proteins. Several biological functions have been attributed to members of the 14-3-3 protein family, as they are directly involved in the transduction of signals related to stress, cell division and differentiation, activation of transcription and apoptosis, amongst others (Sehnke et al., 2002).

14-3-3 functions in eukaryotic cells are exerted by several isoforms, ranging from 2 in *Saccharomyces cerevisiae* to 15 in *Arabidopsis thaliana* (Rosenquist et al., 2001). Though 14-3-3 activity is thought to be conserved among different 14-3-3 isoforms (van Heusden et al., 1996), they appear to be spatially and temporally regulated during development in both plants and animals, suggesting isoform-specific developmental roles. Tissue-specific expression has been described for the 14-3-3 χ isoform during *Arabidopsis thaliana* development (Daugherty et al., 1996), while the 14-3-3C isoform from barley (*Hordeum vulgare* L.) is known to be specifically expressed in the scutellum and in the L₂ layer of the shoot apical meristem (SAM) of germinating zygotic barley embryos (Testerink et al., 1999). In barley, 3 isoforms have been cloned so far: 14-3-3A, 14-3-3B and 14-3-3C (GenBank X62388, X93170 and Y14200). The developmentally regulated expression pattern of the 14-

3-3C isoform in the SAM described by Testerink et al. (1999) prompted us to further investigate 14-3-3 isoform-specific patterns prior to embryo maturation, in both *in vivo* and *in vitro* embryogenesis. With a set of isoform-specific antibodies, we studied 14-3-3 immunolocalization in immature zygotic embryos and embryo-like structures (ELSs) derived from androgenic microspores. In addition, we have investigated 14-3-3 immunolocalization following cell division of the shoot meristem using *in vitro* shoot meristematic cultures (SMCs).

Materials and Methods

Androgenesis induction and microspore culture

Barley (*Hordeum vulgare* L. cv Igri) androgenesis induction and microspore culture were performed according to Hoekstra et al. (1992). Embryo-like structures (ELSs) ranging 0.5-1.0 mm diameter were harvested from media by forceps after 21 days of culture for immunolocalization studies.

Immature zygotic embryos

In vivo developed immature zygotic embryos were dissected under binocular microscope from 21 DAP (days after pollination) seeds for immunolocalization studies.

Shoot meristematic cultures

Shoot meristematic cultures (SMCs) were obtained from mature grains of barley (*Hordeum vulgare* L. cv. Himalaya) germinated according to Louwerson (2002). The material for immunolocalization was obtained from a 16 month-old culture with high regeneration capacity.

Immunolocalization

Fixation and embedding of 21 days-old ELSs, 21 DAP immature zygotic embryos and 16 month-old SMCs was performed as described by Louwerson (2002). Isoform-specific antibodies raised against the C-terminal part of the 14-3-3A, 14-3-3B and 14-3-3C isoforms were used for *in situ* immunolocalization (Testerink et al., 1999).

Results

14-3-3 immunolocalization was first studied in 21 DAP immature zygotic embryos and in 21 days-old ELSs. Although 21 DAP immature zygotic embryos were at a further developmental stage compared to *in vitro*-developed 21 days-old ELSs, both showed similar 14-3-3 isoform-specific expression patterns (Fig. 1). 14-3-3A was immunolocalized in the scutellum, root and shoot meristems, root cap, leaf primordia, mesocotyl and epidermis, while 14-3-3B was found in all embryogenic tissues (data not shown). 14-3-3C expression was markedly different than that of 14-3-3A and 14-3-3B, being restricted to the scutellum, mesocotyl, coleoptile and to one region of the SAM (Fig. 1). Figure 2 shows a magnified view of the 14-3-3 immunolocalization in the SAM. 14-3-3A signal was detected in the leaf primordia and in some of the cells within the SAM, however no defined pattern could be observed (Fig. 2a,e). While 14-3-3B was ubiquitously expressed (Fig. 2b,f), 14-3-3C could only be detected in a group of cells underneath the L₁ layer of the SAM. No signal was detected in control sections that were incubated only with the secondary antibody (Fig. 2d,h).

L₂-layer-specific 14-3-3C expression in the SAM of mature barley embryos has been previously reported (Testerink et al., 1999). Our results suggest that high levels of 14-3-3C are already present underneath the L₁ layer of the SAM before L₂ layer is morphologically differentiated.




Figure 1. 14-3-3C immunolocalization in 21 DAP zygotic embryos (a) and 21 days-old ELSs (b). Proteins were detected using a secondary alkaline-phosphatase-conjugated antibody followed by incubation in nitroblue tetrazolium, 5-bromo-4-chloro-3-indolyl phosphate (NBT/BCIP) in 5µm sections. Arrowheads indicate 14-3-3C expression underneath L₁ layer of the shoot apical meristem and in scutellum cells. a sam, ct coleoptile, lp leaf primordia, m mesocotyl, r root meristem, rc root cap, sc scutellum.

The question raised is whether 14-3-3C specific expression underneath the L₁ layer of developing SAMs is restricted to L₂ layer formation, or whether it is also a feature of further cell division of the L₂ layer during early organogenesis. To answer this question, we studied 14-3-3C immunolocalization in proliferating barley SMCs.

Figure 3a illustrates shoot formation and an adventitious shoot meristem (boxed) in a cross section of a 16 month-old SMC of barley. We observed the presence of L_2 -derived cells as a thick differentiated layer underneath L_1 and above the corpus.




Figure 2. Magnified view of developing SAM of 21 DAP zygotic embryos (a-d) and 21 days-old ELSs (e-h) showing immunolocalization of 14-3-3 proteins. 14-3-3A (a,e), 14-3-3B (b, f), 14-3-3C (c, g) and control incubated only with secondary antibody (d, h). a SAM, e epidermis, lp leaf primordia.

While 14-3-3A (Fig. 3b) and 14-3-3B (data not shown) were expressed in all layers within the shoot apical meristem, 14-3-3C expression could mainly be observed in L_2 -layer derived cells (Fig. 3c). No 14-3-3 signal was observed when sections were incubated only with the secondary antibody (Fig. 3d).




Figure 3. Structure and 14-3-3 immunolocalization in SMCs. A 5 μ m section of a 16 month-old SMC observed by darkfield (a). The equivalent area in the white box is presented in a higher magnification in (b-d). Proteins were detected using a secondary alkaline-phosphatase-conjugated antibody followed by incubation in NBT/BCIP. 14-3-3A (b), 14-3-3C (c) and control incubated only with secondary antibody (d). e epidermis, L_2 L_2 layer-derived cells, v vascular bundles.

Discussion

The developmental programs that lead to embryo formation are thought to be common to both androgenesis and zygotic embryogenesis (Mordhorst et al., 1997). We describe here that 14-3-3 isoform-specific expression patterns are similar during barley zygotic and microspore-derived embryogenesis. Our results are clear evidence for biochemical similarities among these two processes, indicating that the same spatial expression of members of the 14-3-3 family of regulatory proteins might be needed during the formation of barley embryos.

We demonstrate that L₂-layer specification and cell division from the L₂-layer are both accompanied by specific L₂-layer 14-3-3C expression in immature barley embryos and during early organogenesis of SMCs. The L₂ layer of the shoot apical meristem contributes probably to mesophyll formation in adult plants (Jenik and Irish, 2000). This expression pattern of the 14-3-3 C isoform in the SMC points out to specific functions for this isoform associated with the differentiation and function of the L₂ layer.

In other multicellular organisms, tissue-specific expression of 14-3-3 isoforms during embryogenesis is a known phenomenon. For example, in *Drosophila* 14-3-3ε tissue-specific expression was prior to mitogen-activated protein kinase (MAPK) activation in the same tissue during early embryo development (Tien et al., 1999). 14-3-3ε and 14-3-3ξ have been reported to positively regulate Ras-Raf signaling pathway, leading to MAPK cascade activation in *Drosophila* (Chang and Rubin, 1997; Kockel et al., 1997). Recently, Pnueli et al. (2001) have reported the specific expression of at least one 14-3-3 isoform in the SAM of tomato seedlings (*Lycopersicon esculentum* L.). The authors have shown that the same isoform takes part of a signaling system involved in determining SAM vegetative growth potential, a signaling system that is analogous to Raf-1 in animal cells. Further work is needed to address the possible 14-3-3C targets in the L₂-layer. In this regard, SMCs seem to be a suitable system to study 14-3-3C interacting proteins in the shoot meristem of barley.

Acknowledgments

We thank Sandra van Bergen and Arnoud van Marion for technical assistance.

References

- Chang HC, Rubin GM (1997) 14-3-3 ϵ positively regulates Ras-mediated signalling in *Drosophila*. *Genes Dev* 11: 1132-1139
- Daugherty CJ, Rooney MF, Miller PW, Ferl RJ (1996) Molecular organization and tissue-specific expression of an *Arabidopsis* 14-3-3 gene. *Plant Cell* 8: 1239-1248
- Hoekstra S, van Zijderveld MH, Louwerse JD, Heidekamp F, van der Mark F (1992) Anther and Microspore culture of *Hordeum vulgare* L. cv. Igri. *Plant Sci* 86: 89-96
- Jenik PD, Irish VF (2000) Regulation of cell proliferation patterns by homeotic genes during *Arabidopsis* floral development. *Development* 127: 1267-1276
- Kockel L, Vorbrüggen G, Jäckle H, Mlodzik M, Bohmann D (1997) Requirement for *Drosophila* 14-3-3 ζ in Raf-dependent photoreceptor development. *Genes Dev* 11: 1140-1147
- Louwerse JD (2002) Transformation of barley with the maize transposable element *En/Spm*. Ph.D. thesis, Heriot-Watt University, Edinburgh, Scotland
- Mordhorst AP, Toonen MAJ, de Vries SC (1997) Plant Embryogenesis. *Crit Rev Plant Sci* 16: 535-576
- Pnueli L, Gutfinger T, Hareven D, Ben-Naim O, Ron N, Adir N, Lifschitz E (2001) Tomato SP-interacting proteins define a conserved signaling system that regulates shoot architecture and flowering. *Plant Cell* 13: 2687-2702
- Rosenquist M, Alsterfjord M, Larsson C, Sommarin M (2001) Data mining the *Arabidopsis* genome reveals fifteen 14-3-3 genes. Expression is demonstrated for two out of five novel genes. *Plant Physiol* 127: 142-149
- Sehnke PC, DeLille J, Ferl RJ (2002) Consummating signal transduction: The role of 14-3-3 proteins in the completion of signal-induced transitions in protein activity. *Plant Cell* S339-S354
- Testerink C, van der Meulen RM, Oppedijk BJ, de Boer AH, Heimovaara-Dijkstra S, Kijne JW, Wang M (1999) Differences in spatial expression between 14-3-3 isoforms in germinating barley embryos. *Plant Physiol* 121: 81-87
- Tien AC, Hsei HY, Chien CT (1999) Dynamic expression and cellular localization of the *Drosophila* 14-3-3 ϵ during embryonic development. *Mech Dev* 81: 209-212
- van Heusden GPH, van der Zanden AL, Ferl RJ, Steensma HY (1996) Four *Arabidopsis thaliana* 14-3-3 protein isoforms can complement the lethal yeast *bmh1 bmh2* double disruption. *FEBS Lett* 391: 252-256