



Universiteit
Leiden
The Netherlands

Phenotypic responses to lifelong hypoxia in cichlids

Rutjes, Hendrikus Antonius

Citation

Rutjes, H. A. (2006, October 24). *Phenotypic responses to lifelong hypoxia in cichlids*. Retrieved from <https://hdl.handle.net/1887/4925>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4925>

Note: To cite this publication please use the final published version (if applicable).

REFERENCES

- Almeida-Val,V.M.F., Farias,I.P., Silva,M.N.P., Duncan,W.P., and Val,A.L. (1995) Biochemical adjustments to hypoxia by Amazon cichlids. Brazilian Journal of Medical and Biological Research, 28:1257-1263.
- Almeida-Val,V.M.F., Val,A.L., and Hochachka,P.W. (1993) Hypoxia tolerance in Amazon fishes: Status of an under-explored biological “goldmine”. In: Surviving hypoxia: Mechanisms of control and adaptation. P.W.Hochachka, P.L.Lutz, T.Sick, M.Rosental, and G.Van den Thillart, eds. CRC press, pp. 435-445.
- Almeida-Val,V.M.F., Val,A.L., and Walker,I. (1999) Long- and short-term adaptation of Amazon fishes to varying O₂ levels: intra-specific phenotypic plasticity and interspecific variation. In: Biology of tropical fishes. A.L.Val, and V.M.F.Almeida-Val, eds. INPA, Manaus, pp. 185-206.
- Arunga,J.O. (1981) A case study of the Lake Victoria Nile perch *Lates niloticus* (Mbuta) fishery. Nat.Ac.Adv.Arts Sci.,165-184.
- Babiker,M.M. (1985) The role of organophosphates in adaptation of an obligate water-breathing teleost (*Tilapia nilotica* L.) to hypoxia. Hydrobiologia, 121:59-64.
- Ballintijn,C.M. (1969) Movement pattern and efficiency of the respiratory pump of the carp (*Cyprinus carpio* L.). J.Exp.Biol., 50:593-613.
- Barel,C.D.N. (1993) Concepts of an architectonic approach to transformation morphology. Acta Biotheoretica, 41:345-381.
- Barel,C.D.N., Dorit,R., Greenwood,P.H., Fryer,G., Hughes,N., Jackson,P.B.N., Kawanabe,H., Lowe-McConnel,R.H., Nagoshi,M., Ribbink,A.J., Trewavas,E., Witte,F., and Yamaoka,K. (1985) Destruction of fisheries in Africa’s lakes. Nature, 315:19-20.
- Barel,C.D.N., Van Oijen,M.J.P., and Witte,F. (1977) An introduction to the taxonomy and morphology of the haplochromine Cichlidae from Lake Victoria. A manual to Greenwood’s revision papers. Neth.J.Zool., 27:333-389.
- Barel,C.D.N., Witte,F., and Van Oijen,M.J.P. (1976) The shape of the skeletal elements in the head of a generalised haplochromis species: *H. elegans* Trewavas 1933 (Pisces, Cichlidae). Neth.J.Zool., 26:163-265.
- Basu,S.P. (1959) Active respiration of fish in relation to ambient concentrations of oxygen and carbon dioxide. J.Fish.Res.Bd.Can., 16:175-212.

Beamish,F.W.H. (1963a) Respiration of fishes with special emphasis on standard oxygen consumption II. Influence of weight and temperature on respiration of several species. *Can.J.Zool.*, 42:187.

Beamish,F.W.H. (1963b) Respiration of fishes with special emphasis on standard oxygen consumption III. influence of oxygen. *Can.J.Zool.*, 42:355-366.

Beamish,F.W.H. (1964) Respiration of fishes with special emphasis on standard oxygen consumption III. influence of oxygen. *Can.J.Zool.*, 42:355-366.

Beauchamp,R.S.A. (1958) Utilising the natural resources of Lake Victoria for the benefit of fisheries and agriculture. *Nature*, 181:1634-1636.

Bendib,S., Français,O., Tabeling,P., and Willaime,H. (2001) Analytical study and characterisation of micro-channel and passive micro-diode. 12th micromechanics Europe workshop 16-18 September 2001 Corc, Ireland.

Bouton,N., De Visser,J., and Barel,C.D.N. (2002) Correlating head shape with ecological variables in rock-dwelling haplochromines (Teleostei: Cichlidae) from Lake Victoria. *Biological Journal of the Linnean Society*, 76:39-48.

Chabot,D., and Dutil,J.-D. (1999) Reduced growth of Atlantic cod in non-lethal hypoxic conditions. *Journal of Fish Biology*, 55:472-491.

Chapman,L.J., and Chapman,C.A. (2002) Tropical forest degradation and aquatic ecosystems: our current state of knowledge. In: *Conservation of Freshwater Fishes: Options for the Future*. M.J.Collares-Pereira, I.G.Cowx, and M.M.Coelho, eds. Fishing News Books, an imprint of Blackwell Science, pp. 237-249.

Chapman,L.J., Chapman,C.A., Nordlie,F.G., and Rosenberger,A.E. (2002) Physiological refugia: swamps, hypoxia tolerance and maintenance of fish diversity in the Lake Victoria region. *Comp.Biochem.Physiol.*, 133A:421-437.

Chapman,L.J., Galis,F., and Shin,J. (2000) Phenotypic plasticity and the possible role of genetic assimilation: Hypoxia-induced trade-offs in the morphological traits of an African cichlid. *Ecology Letters*, 3:387-393.

Chapman,L.J., Kaufman,L.S., Chapman,C.A., and McKenzie,F.E. (1995) Hypoxia tolerance in twelve species of East African cichlids: Potential for low oxygen refugia in Lake Victoria. *Conservation Biology*, 9:1274-1288.

- Chapman,L.J., and Liem,K.F. (1995) Papyrus swamps and the respiratory ecology of *Barbus neumayri*. Env.Biol.Fish., 44:183-197.
- Chapman,L.J., Nordlie,F.G., and Seifert,A. (2002) Respiratory oxygen consumption among groups of *Pseudocrenilabrus multicolor victoriae* subjected to different oxygen concentrations during development. Journal of Fish Biology, 61:242-251.
- De Jager,S., and Dekkers,W.J. (1975) Relations between gill structure and activity in fish. Neth.J.Zool., 25:276-308.
- De Jonge,V.N., Elliott,M., and Orive,E. (2002) Causes, historical development, effects and future challenge of a common environmental problem: eutrophication. Hydrobiologia, 475/476:1-19.
- De Visser, J. and Barel, C. D. N. Transformation patterns and functional significance of headshape in cichlid fish (Cichlidae, Teleostei). 55-76. 2000. PhD-Dissertation University of Leiden, Netherlands
- Diaz,R.J., and Rosenberg.R. (1995) Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna. Oceanography and Marine Biology: An Annual Review, 33:245-303.
- Duthie,G.G. (1982) The respiratory metabolism of temperature-adapted flatfish at rest and during swimming activity and the use of anaerobic metabolism at moderate swimming speeds. J.Exp.Biol., 97:359-373.
- Fernandez,M.N., and Rantin,F.T. (1989) Respiratory responses of *Oreochromis niloticus* (Pisces, Cichlididae) to environmental hypoxia under different thermal conditions. Journal of Fish Biology, 35:509-519.
- Fernandez,M.N., and Rantin,F.T. (1994) Relationships between oxygen availability and metabolic cost of breathing in Nile tilapia (*O. niloticus*). Aquaculture consequenses. Aquaculture, 127:346.
- Frey,B.F., Weber,R.E., Van Aardt,W.J., and Fago,A. (1998) The haemoglobin system of the mudfish, *Labeo capensis*; adaptations to temperature and hypoxia. Comp.Biochem.Physiol., 120B:735-742.
- Fryer,G. (1997) Biological implications of a suggested Late Pleistocene desiccation of Lake Victoria. Hydrobiologia, 354:177-182.

Fryer,G. (2001) On the age and origin of the species flock of haplochromine cichlid fishes of Lake Victoria. Proc.R.Soc.Lond.B., 268:1147-1152.

Fryer,G., and Iles,T.D. (1972) The cichlid fishes of the great lakes of Africa: Their Biology and evolution, Oliver and Boyd, Edinburgh.

Galis,F., and Barel,C.D.N. (1980) Comparative functional morphology of the gills of African lacustrine cichlidae (Pisces, Teleostei). Neth.J.Zool., 30:392-430.

Goldschmidt,T., Witte,F., and De Visser,J. (1990) Ecological segregation in zooplanktivorous haplochromine species (Pisces: Cichlidae) from Lake Victoria. Oikos, 58:343-355.

Gracey,A.Y., Troll,J.V., and Somero,G.N. (2001) Hypoxia-induced gene expression profiling in the euryoxic fish *Gillichthys mirabilis*. PNAS, 98:1993-1998.

Gray,I.E. (1954) Comparative study of the gill area of marine fishes. Biol.Bull., 107:219-225.

Greaney,G.S., Place,A.R., Cashon,R.E., Smith,G., and Powers,D.A. (1980) Time course of changes in enzyme activities and blood respiratory properties of killifish during long-term acclimation to hypoxia. Physiol.Zool., 53:136-144.

Greenwood,P.H. (1959) The monotypic genera of cichlid fishes in Lake Victoria part II and a revision of the Lake Victoria *Haplochromis* species (Pisces Cichlidae) part III. Bulletin of the British Museum of Natural History (Zoology), 5:164-177.

Greenwood,P.H. (1960) A revision of the Lake Victoria *Haplochromis* species (Pisces, Cichlidae) part IV. Bulletin of the British Museum of Natural History (Zoology), 6:1-281.

Greenwood,P.H. (1965a) Environmental effects on the pharyngeal mill of a cichlid fish, *Astatoreochromis alluaudi*, and their taxonomic implications. Proc.Linn.Soc.Lond., 176:1-10.

Greenwood,P.H. (1965b) The cichlid fishes of Lake Nabugabo, Uganda. Bull.Br.Mus.Nat.Hist.(Zool.), 12:313-357.

Greenwood,P.H. (1973) A revision of the *Haplochromis* and related species (Pisces: Cichlidae) from Lake George, Uganda. Bull.Br.Mus.Nat.Hist.(Zool.), 25:141-242.

Greenwood,P.H. (1974) The cichlid fishes of Lake Victoria, East Africa: the biology and evolution of a species flock. Bull.Br.Mus.Nat.Hist.(Zool.), suppl.6:1-134.

- Greenwood,P.H. (1981) The haplochromine cichlids of the East African Lakes: Collected papers on their taxonomy, biology and evolution, Kraus International Publications, Munich, Germany.
- Harmsen,E., De Tombe,P.Ph., and De Jong,J.W. (1982) Simultaneous determination of myocardial adenine-nucleotides and creatine-phosphate by high-performance liquid-chromatography. *Journal of Chromatography*, 230:131-136.
- Hattingh,J. (1976) Haemoglobins in *Labeo umbratus*: The influence of temperature and oxygen. *South African Journal of Science*, 72:27-28.
- Hecky,R.E. (1993) Peter Kilham Memorial Lecture: The eutrophication of Lake Victoria. *Verh.Internat.Verein.Limnol.*, 25:39-48.
- Hecky,R.E., and Bugeny,F.W.B. (1992) Hydrology and chemistry of the African Great Lakes and water quality issues: problems and solutions. *Mitt.Internat.Verein.Limnol.*, 23:45-54.
- Hecky,R.E., Bugeny,F.W.B., Ochumba,P.B.O., Talling,J.F., Mugidde,R., Gophen,M., and Kaufman,L.S. (1994) Deoxygenation of the deep water of Lake Victoria, East Africa. *Limnol.Oceanogr.*, 39:1476-1481.
- Hoogerhoud,R.J.C. (1984) A taxonomic reconsideration of the haplochromine genera *Gairochromis* Greenwood, 1980 and *Labrochromis* Regan, 1920 (Pisces, Cichlidae). *Neth.J.Zool.*, 34:539-565.
- Hoogerhoud, R. J. C. Ecological morphology of some cichlid fishes. PhD-Dissertation University of Leiden, Netherlands 1-133. 1986.
- Hoogerhoud,R.J.C., Witte,F., and Barel,C.D.N. (1983) The ecological differentiation of two closely resembling Haplochromis species from Lake Victoria (*H. iris* and *H. hiatus*; pisces, Cichlidae). *Neth.J.Zool.*, 33:283-305.
- Hoppeler,H., Vogt,M., Weibel,E.R., and Lück,M. (2003) Response of skeletal muscle mitochondria to hypoxia. *Exp.Physiol.*, 88:109-119.
- Houston,A.H. (1980) Components of the haematological response of fishes to environmental temperature change. In: Environmental physiology of fishes. M.A.Ali, ed. Plenum, New York, pp. 241-298.
- Houston,A.H., and Cyr,D. (1974) Thermoacclimatory variation in the haemoglobin systems of goldfish (*Carassius auratus*) and rainbow trout (*Salmo giardneri*). *J.Exp.Biol.*, 61:455-461.

Houston,A.H., and Tun,N. (1986) Environmentally -related changes in red cell levels of ionic modulators of hemoglobin-O₂ affinity in rainbow trout, *Salmo gairdneri*. Comp.Biochem.Physiol., 85A:779-783.

Hughes,G.M. (1966) The dimensions of fish gills in relation to their function. Journal of Experimental Biology, 45:177-195.

Hughes,G.M. (1972) Morphometrics of fish gills. Respir.Physiol., 14:1-25.

Hughes,G.M., and Morgan,M. (1973) The structure of fish gills in relation to their respiratory function. Biological Reviews, 48:419-475.

Johnson,T.C., Scholz,C.A., Talbot,M.R., Kelts,K., Ricketts,R.D., Ngobi,G., Beuning,K., Ssemmanda,I., and McGill,J.W. (1996) Late Pleistocene desiccation of Lake Victoria and rapid evolution of cichlid fishes. Science 273:1091-1093.

Johnston,I.A., and Bernard,L.M. (1982a) Routine oxygen consumption and characteristics of the myotomal muscle in tench: Effects of long-term acclimation to hypoxia. Cell Tiss.Res., 227:161-177.

Johnston,I.A., and Bernard,L.M. (1982b) Ultrastructure and metabolism of skeletal muscle fibers in the tench: Effects of long-term acclimation to hypoxia. Cell Tiss.Res., 227:179-199.

Jones,D.R., Randall,D.J., and Jarman,G.M. (1970) A graphical analysis of oxygen transfer in fish. Resp.Physiol., 10:285-298.

Joyce,D.A., Lunt,D.H., Bills,R., Turner,G.F., Katongo,C., Duftner,N., Sturmbauer,C., and Seehausen,O. (2005) An extant cichlid fish radiation emerged in an extinct Pleistocene lake. Nature, 435:90-95.

Karlson,K., Rosenberg,R., and Bonsdorff,E. (2002) Temporal and spatial large-scale effects of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic waters - a review. Oceanography and Marine Biology, 40:427-489.

Kaufman,L.S. (1992) Catastrophic change in species-rich freshwater ecosystems. The lessons of Lake Victoria. Bioscience, 42:846-858.

Keppler,D., and Decker,K. (1988) Glycogen. In: Methods of enzymatic analysis. Volume VI, Metabolites 1: Carbohydrates. H.A.Bergmeyer, J.Bergmeyer, and M.Graßl, eds. VHC Verlagsgesellschaft, Weinheim, pp. 11-18.

Lomholt,J.P., and Johansen,K. (1979) Hypoxia acclimation in carp- how it affects O₂ uptake, ventilation, and O₂ extraction from water. Physiol.Zool., 52:38-49.

- Marinsky,C.A., Houston,A.H., and Murad,A. (1989) Effects of hypoxia on hemoglobin isomorph abundances in rainbow trout, *Salmo giardneri*. Can.J.Zool., 68:884-888.
- Matsuda,N., and Sakai,K. (1999) Technical evaluation of oxygen transfer rates of fish gills and artificial gills. ASAIO J., 45:293-298.
- Mcnatt,R.A., and Rice,J.A. (2004) Hypoxia-induced growth rate reduction in two juvenile estuary-dependant fishes. Journal of Experimental Marine Biology and Ecology, 311:147-156.
- Meyer,A. (1990) Morphometric and allometry in the trophically cichlid fish, *Cichlasoma citrinellum*: Alternative adaptations and ontogenetic changes in shape. J.Zool.Lond., 221: 237-260.
- Mortensen,N.A., Okkels,F., and Bruus,H. (2005) reexamination of Hagen-Poiseuille flow: Shape dependence of the hydraulic resistance in microchannels. Physical Reviews E, 71: 057301-1-057301-4.
- Muir,B.S., and Brown,C.E. (1971) Effects of blood pathway on the blood-pressure drop in fish gills, with special references to tunas. J.Fish.Res.Bd.Can., 28:947-955.
- Muir,B.S., and Hughes,G.M. (1969) Gill dimensions for three species of tunny. J.Exp.Biol., 51:271-285.
- Muusze,B., Marcon,J., Van den Thillart,G.J.E.E.M., and Val,A.L. (1998) Hypoxia tolerance of Amazon fish Respirometry and energy metabolism of the cichlid *Astronotus ocellatus*. Comp.Biochem.Physiol., 120:151-156.
- Nagl,S., Tichy,H., Mayer,W.E., Takezaki,N., Takahata,N., and Klein,J. (2000) The origin and age of haplochromine fishes in Lake Victoria, East Africa. Proc.R.Soc.Lond.B., 267:1049-1061.
- Nikinmaa,M., Cech Jr,J.J., Ryhanen,E.L., and Salama,A. (1987) Red cell function of carp (*Cyprinus carpio*) in acute hypoxia. J.Exp.Biol., 47:53-58.
- Njuguna,S.G. (1991) Water hyacinth: The world's worst aquatic weed infests lakes Naivasha and Victoria. Swara, 14:8-10.
- Ochumba,P.B.O. (1990) Massive fish kills within the Nyanza Gulf of Lake Victoria, Kenya. Hydrobiologia, 208:93-99.

Ochumba, P. B. O., Gophen, M., and Kaufman, L. S. Changes in oxygen availability in the Kenyan portion of Lake Victoria: effects on fisheries and biodiversity. (Proceedings of the Lake Victoria Ecosystem Workshop held in Jinja, Uganda, Aug. 17 - 21, 1992), 33. 1993. New England, U.S.A., Edgerton Research Laboratory of the New England Aquarium, report 93-3. Kaufman, L. S., Ochumba, P. B. O., and Ongutu-Ohwayo, R.

Ochumba,P.B.O., and Kibaara,D.I. (1989) Observations on blue-green algal blooms in the open waters of Lake Victoria, Kenya. Afr.J.Ecol., 27:23-34.

Ongutu-Ohwayo,R. (1990) The decline of the native fishes of lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus*, and the Nile tilapia, *Oreochromis niloticus*. Env.Biol.Fish., 27:81-96.

Ott,M.E., Heisler,N., and Ultsch,G.R. (1980) A re-evaluation of the relationship between temperature and the critical oxygen tension in freshwater fishes. Comp.Biochem.Physiol., 67A:337-340.

Palzenberger,M., and Pohla,H. (1992) Gill surface area of water-breathing freshwater fish. Reviews in Fish Biology and Fisheries, 2:187-216.

Pearson,T.H., and Rosenberg,R. (1992) Energy flow through the Kattegat: A comparative examination of the eutrophication of a coastal marine ecosystem. Netherlands Journal of Sea Research, 28:317-334.

Petersen,J.K., and Petersen,G.I. (1990) Tolerance, behaviour consumption in the sand goby, *Pomatoschistus minutus* (Pallas), exposed to hypoxia. Journal of Fish Biology, 37:921-933.

Pihl,L., Baden,S.P., and Diaz,R.J. (1991) Effects of periodic hypoxia on distribution of demersal fish and crustaceans. Marine Biology, 108:349-360.

Pihl,L., Baden,S.P., Diaz,R.J., and Schaffner,L.C. (1992) Hypoxia-induced structural changes in the diet of bottom-feeding fish and Crustacea. Marine Biology, 112:349-361.

Randall,D.J. (1970) Gas exchange in fish. In: Fish physiology, Volume IV, The nervous system, circulation, and respiration. W.S.Hoar, and D.J.Randall, eds. Academic press, London, pp. 253-286.

Rantin,F.T., Kalinin,A.L., Lass,M.L., and Fernandez,M.N. (1992) Respiratory responses to hypoxia in relation to mode of life of two erythrinid species (*Hoplias malabaricus* and *Hoplias lacerdae*). Journal of Fish Biology, 41:805-812.

- Reddy,M. (2005) Restoration and management of tropical eutrophic lakes, Science Publishers Inc. Enfield (NH) USA.
- Roman,M.R., Gauzens,A.L., Rhinehart,W.K., and White,J.R. (1993) Effects of low oxygen waters on Chesapeake Bay zooplankton. Limnol.Oceanogr., 38:1603-1614.
- Ross,L.G. (2000) Environmental physiology and energetics. In: Tilapias: Biology and exploitation. C.M.Malcolm, Beveridge, J.Brendan, and e.McAndrews, eds. Kluwer academic publishers, Dordrecht/Boston/London, pp. 89-128.
- Schlichting,C.D. and Pigliucci,M. (1998) Phenotypic evolution: a reaction norm perspective. Sinauer.
- Schofield,P.J., and Chapman,L.J. (2000) Hypoxia tolerance of introduced Nile perch: implications for survival of indigenous fishes in the Lake Victoria basin. African Zoology, 35: 35-42.
- Schumann,D., and Piiper,J. (1966) Der Sauerstoffbedarf der Atmung bei Fischen nach Messungen an der narkotisierten Schleie (*Tinca tinca*). Pflugers Arch.Ges.Physiol., 288:15-26.
- Schwantes,A.R., Bartlett,G.R., and Schwantes,M.L.B. (1991) The hemoglobin of *Geophagus brasiliensis* (cichlidae). Comp.Biochem.Physiol., 99B:157-160.
- Seehausen,O. (1996) Lake Victoria rock cichlids: Taxonomy, ecology, and distribution, Verduyn Cichlids, Zevenhuizen, Netherlands.
- Seehausen,O. (2002) Patterns in fish radiation are compatible with pleistocene desiccation of Lake Victoria and 14600 year history for its cichlid species flock. Proc.R.Soc.Lond.B., 269: 491-497.
- Seehausen,O., Koertsier,E., Schneiden,M.V., Chapman,L.J., Chapman,C.A., Van Alphen,J.J.M., and Bills,R. (2003) Nuclear markers reveal unexpected genetic variation and a Congolese-Nilotic origin of the Lake Victoria cichlid species flock. Proc.R.Soc.Lond., 270: 129-137.
- Smith,D.G., and Johnson,D.W. (1977) Oxygen exchange in a simulated trout gill secondary lamella. Am.J.Physiol.Reg.Int.Comp.Physiol., 2:R146-R161.
- Smits,J.D., Anker,G.C., Witte,F., and Barel,C.D.N. (1997) Comparative functional anatomy of the pharyngeal jaw apparatus in two morphs of *Astatoreochromis alluaudi* (Pisces, Cichlididae). Neth.J.Zool., 47:313-347.

- Smits,J.D., Witte,F., and Povel,G.D.E. (1996a) Differences between inter- and intraspecific architectonic adaptations to pharyngeal mollusc crushing in cichlid fishes. *Biological Journal of the Linnean Society*, *59*:367-387.
- Smits,J.D., Witte,F., and Van Veen,F.G. (1996b) Functional changes in the anatomy of the pharyngeal jaw apparatus of *Astatoreochromis alluaudi* (Pisces, Cichlididae), and their effects on adjacent structures. *Biological Journal of the Linnean Society*, *59*:389-409.
- Sollid,J., De Angelis,P., Gundseren,K., and Nilsson,G. (2003) Hypoxia induces adaptive and reversible gross morphological changes in crucian carp gills. *The Journal of Experimental Biology*, *3667*-3673.
- Sollid,J., Weber,R.E., and Nilsson,G.E. (2005) Temperature alters the respiratory surface area of crucian carp *Carassius carassius* and goldfish *Carassius auratus*. *Journal of Experimental Biology*, *208*:1109-1116.
- Stitt,M. (1984) Citrate synthase (Condensing enzyme). In: *Methods of enzymatic analysis*. Volume IV: Enzymes 2: Esterases, Glycosidases, Lyases, Ligases. H.A.Bergmeyer, H.A.Bergmeyer, and M.Graßl, eds. VCH Verlagsgesellschaft, Weinheim, pp. 353-358.
- Thetmeyer,H., Waller,U., Black,K.D., Inselmann,S., and Rosenthal,H. (1999) Growth of european sea bass (*Dicentrarchus labrax* L.) under hypoxic conditions and oscillating oxygen conditions. *Aquaculture*, *174*:355-367.
- Trewavas,E. (1983) Tilapiine fishes of the genera *Sarotherodon*, *Oreochromis* and *Danakilia*, British Museum (Natural History).
- Tun,N., and Houston,A.H. (1986) Temperature, oxygen, photoperiod, and the hemoglobin system of the rainbow trout *Salmo giardneri*. *Can.J.Zool.*, *64*:1883-1888.
- Ultsch,G.R., Jackcon,D.C., and Moalli,R. (1981) Metabolic oxygen conformity among lower vertebrates: The toadfish revisited. *J.Comp.Physiol.*, *142*:439-443.
- Val,A.L. (2000) Organic phosphates in the red blood cells of fish. *Comp.Biochem.Physiol.*, *125A*:417-435.
- Val,A.L., Lessard,J., and Randall,D.J. (1995) Effects of hypoxia on rainbow trout (*Oncorhynchus mykiss*): intraerythrocytic phosphates. *J.Exp.Biol.*, *198*:305-310.
- Van Dam,A.A., and Pauly,D. (1995) Simulation of the effects of oxygen on food consumption and growth of Nile tilapia *Oreochromis niloticus* (L.). *Aquaculture Research*, *26*:427-440.

- Van Dam, L. On the utilisation of oxygen and regulation of breathing in some aquatic animals. 1938. PhD-Dissertation University of Groningen.
- Van den Thillart,G., and Kesbeke,F. (1978) Anaerobic production of carbon dioxide and ammonia by goldfish, *Carassius auratus* (L.). Comp.Biochem.Physiol., 59A:393-400.
- Van den Thillart,G.J.E.E.M., Dalla via,J., Vitali,G., and Cortesi,P. (1994) Influence of long term hypoxia exposure on the energy metabolism of *Solea solea*. I. Critical O₂ levels for aerobic and anaerobic metabolism. Mar.Ecol.Prog.Ser., 104:109-117.
- Van den Thillart,G.J.E.E.M., Kesbeke,F., and Van Waarde,A. (1980) Anaerobic Energy Metabolism of Goldfish, *Carassius auratus* (L.). Influence of hypoxia and anoxia on phosphorylated compounds and glycogen. J.Comp.Physiol., 136:45-52.
- Van den Thillart,G.J.E.E.M., and Smit,H. (1984) Carbohydrate metabolism of goldfish (*Carassius auratus* (L.)). Effects of long-term hypoxia acclimation on enzyme patterns of red muscle, white muscle and liver. J.Comp.Physiol.B., 154:477-786.
- Van den Thillart,G.J.E.E.M., and Van Raaij,M.T.M. (1995) Endogenous fuels: Non-invasive versus invasive approaches. In: Biochemistry and molecular biology of fishes, vol. 4. P.W.Hochachka, and T.P.Mommsen, eds. Elsevier Science B.V..
- Van den Thillart,G.J.E.E.M., and Van Waarde,A. (1985) Teleosts in hypoxia: Aspects of anaerobic metabolism. Molecular Physiology, 8:393-409.
- Van den Thillart,G.J.E.E.M., and Verbeek,R. (1991) Anoxia-induced oxygen debt of Goldfish (*Carassius auratus* L.). Physiol.Zool., 64:525-540.
- Van Ginneken,V.J.T., Addink,A., Van den Thillart,G.J.E.E.M., Körner,F., Noldus,L., and Buma,M. (1997) Metabolic rate and level of activity determined in tilapia (*Oreochromis mossambicus*) by direct and indirect calorimetry and videomonitoring. Thermochimica Acta, 291:1-13.
- Van Ginneken,V.J.T., Vanderschoot,J., and Addink,A.D.F. (1995) Direct calorimetry of aquatic animals: Dynamic response of biological processes. Thermochimica Acta, 249:143-159.
- Van Leeuwen,J.L., and Spoor,C.W. (1987) Biting force in cichlids: a discussion of the planar model of Barel (1983). Neth.J.Zool., 37:307-314.
- Van Oijen,M.J.P., Witte,F., and Witte-Maas,E.L.M. (1981) An introduction to ecological and taxonomical investigations on the haplochromine cichlids from the Mwanza Gulf of Lake Victoria. Neth.J.Zool., 31:149-174.

- Van Raaij,M.T.M., Djemila,S.S.P., Balm,H.M., Steffens,A.B., and Van den Thillart,G.J.E.E.M. (1996) Behavioural strategy and the physiological stress response in rainbow trout exposed to severe hypoxia. *Hormones and Behaviour*, *30*:85-92.
- Van Velzen,J., Bouton,N., and Zandee,R. (1998) A procedure to extract phylogenetic information from morphometric data. *Neth.J.Zool.*, *48*:305-322.
- Vassault,A. (1987) Lactate dehydrogenase, Uv-method with pyruvate and NADH. In: Methods of enzymatic analysis. Volume III: Oxireductases,Transferases. H.A.Bergmeyer, J.Bergmeyer, and M.Graßl, eds. VCH Verlagsgesellschaft, Weinheim, pp. 118-133.
- Verheijen,E., Blust,R., and Decleir,W. (1994) Metabolic rate, hypoxia tolerance and aquatic surface respiration of some lacustrine and riverine African cichlid fishes (Pisces: Cichlidae). *107A*, *2*:411.
- Verheyen,E., Blust,R., and Doumen,C. (1985) The oxygen uptake of *Sarotherodon niloticus* L. and the oxygen binding properties of its blood and hemolysate (Pisces: Cichlididae). *Comp.Biochem.Physiol.*, *81A*:423-426.
- Verheyen,E., Salzburger,W., Snoeks,j., and Meyer,A. (2003) Origin of the superflock of cichlid fishes from Lake Victoria, East Africa. *Science*, *300*:325-329.
- Verschuren,D., Edgington,D.N., Kling,H.J., and Johnson,T.C. (1998) Silica depletion in Lake Victoria: Sedimentary signals at offshore stations. *J.Great Lakes Res.*, *24*:118-130.
- Verschuren,D., Johnson,T.C., Kling,H.J., Edgington,D.N., Leavitt,P.R., Brown,E.T., Talbot,M.R., and Hecky,R.E. (2002) History and timing of human impact on Lake Victoria, East Africa. *Proc.R.Soc.Lond.B.*, *269*:289-294.
- Vianen,G.J., Obels,P.P., Van den Thillart,G.J.E.E.M., and Zaagsma,J. (2002) B-Adrenoceptors mediate inhibition of lipolysis in adipocytes of tilapia (*Oreochromis mossambicus*). *Am.J.Physiol.Endocrinol.Metab.*, *282*:E325.
- Vianen,G.J., Van den Thillart,G.J.E.E.M., Van Kampen,M., Van Heel,T.I., and Steffens,A.B. (2001) Plasma lactate and stress hormones in common carp (*Cyprinus carpio*) and rainbow trout (*Oncorhynchus mykiss*) during stepwise decreasing oxygen levels. *Neth.J.Zool.*, *51*:33-50.
- Virani,N.A., and Rees,B. (2000) Oxygen consumption, blood lactate and inter-individual variation in the gulf killifish, *Fundulus grandis*, during hypoxia and recovery. *Comp.Biochem.Physiol.*, *126*:397-405.

- Wahlefeld,A., and Siedel,J. (1985) Creatine and creatinine. In: Methods of enzymatic analysis, Volume VIII, Metabolites 3: Lipids, amino acids and related compounds. H.A.Bergmeyer, J.Bergmeyer, and M.Graßl, eds. VCH Verlagsgesellschaft, Weinheim, pp. 488-507.
- Wanink,J.H. (1991) Survival in a perturbed environment: The effects of Nile perch introduction on the zooplanktivorous fish community of Lake Victoria. In: Terrestrial and Aquatic ecosystems: Perturbation and Recovery. O.Ravera, ed. Ellis Horwood Ltd., Chichester, UK, pp. 269-275.
- Wanink,J.H., Kashindye,J.J., Goudswaard,K.P.C., and Witte,F. (2001) Dwelling at the oxycline: does increased stratification provide a predation refugium for the Lake Victoria sardine *Rastrineobola argentea*? Freshw.Biol., 46:75-85.
- Wanink,J.H., and Witte,F. (2000a) Rapid morphological changes following niche shift in the zooplanktivorous cyprinid *Rastrineobola argentea* from Lake Victoria. Neth.J.Zool., 50:365-372.
- Wanink,J.H., and Witte,F. (2000b) The use of perturbation as a natural experiment: effects of predator introduction on the community structure of zooplanktivorous fish in Lake Victoria. Advances In Ecol.Res., 31:553-570.
- Wannamaker,C.M., and Rice,J.A. (2000) Effects of hypoxia on movements and behavior of selected estuarine organisms from the southeastern United States. J.Exp.M.Biol.Ecol., 249: 145-163.
- Weber,R.E. (1990) Functional significance and structural basis of multiple hemoglobins with special reference to ectothermic vertebrates. In: Comparative physiology. R.K.H.Kinne, and E.Kinne-Saffran, eds. Karger S., Basel, pp. 57-75.
- Weber,R.E. (1994) Hemoglobin-based O₂ transfer in viviparous animals. Israel Journal of Zoology, 40:541-550.
- Weber,R.E. (1996) Hemoglobin adaptations in Amazonian and temperate fish with special references to hypoxia, allosteric effectors and functional heterogeneity. In: Physiology and biochemistry of the fishes of the amazon. A.L.Val, V.M.F.Almeida-Val, and D.J.Randall, eds. INPA Manaus, pp. 75-90.
- Weber,R.E. (2000) Adaptations for oxygen transport: Lessons from fish hemoglobins. In: Hemoglobin function in vertebrates. Molecular adaptation in extreme and temperate environments. G.d.Prisco, B.Giardina, and R.E.Weber, eds. Springer-Verlag Italia.

- Weber,R.E., and Jensen,F.B. (1988) Functional adaptations in hemoglobins from ectothermic vertebrates. *Ann.Rev.Physiol.*, *50*:161-179.
- Weber,R.E., and Wells,R.M.G. (1989) Hemoglobin structure and function. In: Comparative pulmonary physiology. S.C.Wood, ed. Marcel Dekker Inc., pp. 279-309.
- Welcomme,R.L. (1967) Observations on the biology of the introduced species of tilapia in Lake Victoria. *Rev.Zool.Bot.Afr.*, *76*:249-279.
- Welcomme,R.L. (1988) International introductions of inland fish species. *FAO Fish Tech.Pap.*, *294*:1-318.
- Wendelaar Bonga,S.E. (1997) The stress response in fish. *Physiological Reviews*, *77*:591-625.
- Wilhelm,D.F., and Weber,R.E. (1983) Functional characterisation of hemoglobins from South Brazilian fresh water teleosts-II. Three cichlids (*Crenicichla lepidot A*, *Aequidens portalegrensis* and *Geophagus brasiliensis*). *Comp.Biochem.Physiol.*, *75A*:483-489.
- Witte,F. (1981) Initial results of the ecological survey of the haplochromine cichlid fishes from the Mwanza Gulf of Lake Victoria: Breeding patterns, trophic and species distribution. *Neth.J.Zool.*, *31*:175-202.
- Witte,F. (2002) On research on tilapiine and haplochromine cichlids. *Hydrobiologica*, *470*: 1-3.
- Witte,F., Barel,C.D.N., and Hoogerhoud,R.J.C. (1990) Phenotypic plasticity of anatomical structures and its ecomorphological significance. *Neth.J.Zool.*, *40*:278-298.
- Witte,F., Goldschmidt,T., Wanink,J.H., Van Oijen,M.J.P., Goudswaard,K.P.C., Witte-Maas,E.L.M., and Bouton,N. (1992) The destruction of an endemic species flock: quantitative data on the decline of the haplochromine cichlids of Lake Victoria. *Environmental Biology of Fishes*, *34*:1-28.
- Witte,F., Goudswaard,K.P.C., Katunzi,E.F.B., Mkumbo,O.C., Seehausen,O., and Wanink,J.H. (1999) Lake Victoria's ecological changes and their relationships with the riparian societies. In: Ancient Lakes: Their Cultural and Biological diversity. H.Kawanabe, G.W.Coulter and A.C.Roosevelt ed. Kenobi Productions. Printed in Belgium, pp. 189-202.
- Witte,F., Msuku,B.S., Wanink,J.H., Seehausen,O., Katunzi,E.F.B., Goudswaard,K.P.C., and Goldschmidt,T. (2000) Recovery of cichlid species in Lake Victoria: an examination of factors leading to differential extinction. *Reviews in Fish Biology and Fisheries*, *10*:233-241.

- Witte,F., and Van Oijen,M.J.P. (1995) Biology of haplochromine trophic groups. In: Fish stocks and fisheries of Lake Victoria: A handbook for field observations. F.Witte, and W.L.T.van Densen, eds. Samara Publishing, Cardigan, UK, pp. 321-335.
- Witte,F., Wanink,J.H., Rutjes,H.A., Van der Meer,H.J., and Van den Thillart,G.J.E.E.M. (2005) Eutrophication and its influences on the fish fauna of Lake Victoria. In: Restoration and management of tropical eutrophic lakes. M.Reddy, ed. Science Publishers Inc. Enfield (NH) USA.
- Witte,F., and Witte-Maas,E.L.M. (1981) Haplochromine cleaner fishes: a taxonomic and eco-morphological description of two new species. Revision of the haplochromine species (Teleostei, Cichlidae) from Lake Victoria. Part I. Neth.J.Zool., 31:203-231.
- Witte,F., and Witte-Maas,E.L.M. (1987) Implications for taxonomy and functional morphology of intraspecific variation in haplochromine cichlids of Lake Victoria. In: From form to fishery. PhD-Dissertation University of Leiden.
- Wu,R.S. (2002) Hypoxia: from molecular responses to ecosystem responses. Marine Pollution Bulletin, 45:35-45.
- Wu,R.S., Zhou,B.S., Randall,D.J., Woo,N.Y., and Lam,P.K.S. (2003) Aquatic hypoxia is a disruptor and impairs reproduction. Environ.Sci.Technol., 15:1137-1141.
- Yamamoto,K.-I. (1991) Relationship of respiration to body weight in the carp *Cyprinus carpio* under resting and normoxic conditions. Comp.Biochem.Physiol., 100A:113-116.
- Yamamoto,K.-I. (1992) Relationship of respiration to body weight in the tilapia *Oreochromis niloticus* under resting and normoxic conditions. Comp.Biochem.Physiol., 103A:81-83.
- Zhou,B.S., Wu,R.S.S., Randall,D.J., and Lam,P.K.S. (2001) Bioenergetics and RNA/DNA ratios in the common carp (*Cyprinus carpio*) under hypoxia. J.Comp.Physiol.B., 171:49-57.
- Zhou,B.S., Wu,R.S.S., Randall,D.J., Lam,P.K.S., Ip,Y.K., and Chew,S.F. (2000) Metabolic adjustments in the common carp during prolonged hypoxia. Journal of Fish Biology, 57:1160-1171.