



Universiteit  
Leiden  
The Netherlands

## **Nucleotide excision repair : complexes and complexities : a study of global genome repair in human cells**

Volker, Marcel

### **Citation**

Volker, M. (2006, May 15). *Nucleotide excision repair : complexes and complexities : a study of global genome repair in human cells*. Retrieved from <https://hdl.handle.net/1887/4390>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4390>

**Note:** To cite this publication please use the final published version (if applicable).

# **Nucleotide excision repair – complexes and complexities**

**A study of global genome repair in human cells**



**Nucleotide excision repair – complexes and complexities**  
**A study of global genome repair in human cells**

**proefschrift**

ter verkrijging van  
de graad van Doctor aan de Universiteit Leiden,  
op gezag van de Rector Magnificus Dr. D. D. Breimer,  
hoogleraar aan de faculteit der Wiskunde en  
Natuurwetenschappen en die der Geneeskunde,  
volgens besluit van het College voor Promoties  
te verdedigen op maandag 15 mei 2006  
klokke 16.15 uur

door

**Marcel Volker**

Geboren te Leiderdorp,  
in 1976

## **Promotiecommissie**

Promotores	Prof. Dr. L.H.F. Mullenders Prof. Dr. Ir. A.A. van Zeeland
Referent	Dr. W. Vermeulen (Erasmus Universiteit Rotterdam)
Overige leden	Prof. Dr. H. van Steeg Prof. Dr. H. Tanke

*All we have to decide is what to do with the time that is given us*

**J.R.R. Tolkien** - the Lord of the Rings

**aan mijn vader**

**voor mijn moeder**

## Contents

<b>Aim &amp; outline of the thesis</b>	9
<b>1 Cellular responses to DNA damage</b>	15
1.1 Cell cycle arrest	15
1.2 DNA repair mechanisms	15
Direct reversal	15
Base excision repair and single-strand break repair	16
Double-strand break repair	17
Nucleotide excision repair	17
Replication-associated responses	18
Postreplication repair	18
Translesion synthesis	18
Damage avoidance: template switching and daughter-strand gap repair	19
Mismatch repair	19
<b>2 Nucleotide excision repair: background and concepts</b>	23
2.1 Two modes of NER: GGR and TCR	23
Transcription-coupled repair: an outline	23
2.2 Inherited disorders resulting from defective NER	25
Xeroderma pigmentosum (XP)	25
Cockayne syndrome (CS)	25
Trichothiodystrophy (TTD)	26
2.3 Sequential incorporation of factors into the GGR complex	27
2.4 NER substrate specificity: the bipartite recognition model	29
<b>3 Global genome repair proteins and their functions</b>	33
3.1 UV-DDB: a key factor assisting damage recognition	33
The role of CPD repair in cellular survival and mutagenesis	34
3.2 XPC-hHR23B	35
XPC-hHR23B probes for helix destabilisation assisted by UV-DDB	36
3.3 Post-transcriptional regulation of UV-DDB and XPC – regulating the GGR pathway via the damage-recognition step	37
Degradation of DDB2 following UV	37
Increased activity of XPC following UV	38
Other XPC-regulating proteins	39
Regulation at the transcriptional level: a supervising role for p53	39
3.4 TFIIH	40
Possible roles for TFIIH in detecting chemical alterations and in regulating dual incision	41
3.5 XPA	
Damage verification by XPA: regulation of the endonucleases in cooperation with TFIIH?	43
3.6 RPA	44
RPA complements XPA in damage verification	45
3.7 XPG	45
3.8 Order of binding of RPA, XPG and XPA	46
XPG first?	46
RPA first?	47
XPA first?	47

No fixed order of binding of XPG, RPA and XPA?	48
3.9 ERCC1-XPF	48
3.10 DNA resynthesis and ligation	49
<b>4 More interplay between NER and transcription: inhibition of transcription following DNA damage</b>	53
4.1 Transcription inhibition <i>in cis</i> and <i>in trans</i>	53
4.2 A role for TFIIH in transcription inhibition <i>in trans</i> ?	54
4.3 What causes the transcription inhibition <i>in trans</i> ?	55
<b>5 Chromatin, chromatin remodelling and NER</b>	59
5.1 ATP-dependent non-covalent chromatin remodelling	60
SWI/SNF and ISWI	60
CSB	60
5.2 Histone tail modifications	60
Histone acetylation and NER	61
5.3 Non-chromatin remodellers assisting NER <i>in vivo</i>	62
Gadd45	62
HMGN1	62
p53	62
5.4 Post-repair chromatin restoration	63
CAF-1 and Asf1	63
<b>References</b>	65
<b>6 Sequential assembly of the nucleotide excision repair factors <i>in vivo</i></b>	83
<b>7 Local UV-induced DNA damage in cell nuclei results in local transcription inhibition</b>	103
<b>8 Xeroderma pigmentosum group A protein loads as a separate factor onto DNA lesions</b>	113
<b>9 The UV-damaged DNA binding protein mediates efficient targeting of the nucleotide excision repair complex to UV-induced photolesions</b>	135
<b>10 Differential stability of early and late components of the nucleotide excision repair complex <i>in vivo</i></b>	155
<b>Summary</b>	173
<b>Samenvatting voor de leek</b>	179
<b>Abbreviations</b>	184
<b>Curriculum vitae</b>	186
<b>List of publications</b>	187
<b>Appendix: Selected colour figures</b>	





## ***Aim and outline of the thesis***



## Aim and outline of the thesis

Of all exogenous agents that damage genomic DNA and hence present a threat to the integrity of its genetic information, the ultraviolet B (UVB) component of sunlight possesses high clinical relevance because of its abundance. UVB induces predominantly two types of photolesions: cyclobutane pyrimidine dimers (CPD) and pyrimidine 6-4 pyrimidone photoproducts (6-4 photoproducts or 6-4PP). Uniquely, placental mammals rely solely on the nucleotide excision repair (NER) system to repair these photolesions. In addition, NER is capable of removing a wide range of bulky, helix-distorting lesions, including numerous chemical adducts such as those caused by the anticancer drug *cis*-diamminedichloroplatinum(II) (cisplatin), the poison gas nitrogen mustard and polycyclic aromatic hydrocarbons found in burnt food and cigarette smoke. The relevance of the NER pathway, particularly in the removal of UV-induced photolesions, is emphasised by the existence of three inherited disorders caused by mutations in genes encoding NER proteins. Patients suffering from these disorders invariably display hypersensitivity to sunlight, and patients suffering from one disease, xeroderma pigmentosum, combine this with a dramatically increased risk of developing skin cancers.

In the first part of this thesis, the different cellular defence mechanisms against various types of DNA damage are described. Chapter 1 summarises the possible reactions of a cell to DNA damage and briefly sketches the DNA repair pathways. Chapters 2 and 3 concern themselves solely with NER: chapter 2 introduces some background important to understanding NER while chapter 3 provides a detailed step-by-step description of the global genome repair pathway, highlighting the roles of every factor involved. Subsequently chapter 4 elaborates on the inhibition of transcription that is observed after the introduction of NER-types of DNA damage. Chapter 5 puts NER in a nuclear perspective, describing its interplay with chromatin, the natural substrate of any DNA repair pathway.

The research presented in this thesis is focused on various aspects of the NER pathway in live cells. One of the main questions addressed is whether the NER complex assembles on the lesion in a sequential manner or whether it pre-exists as a so-called 'repairosome'. Experimental evidence as described in chapter 6 suggests that a sequential assembly of NER proteins on DNA lesions is most likely. The technique of local UV irradiation used in this study was utilised to address another question: the possible involvement of transcription factor TFIIH in the UV-induced temporary inhibition of transcription. The result of this research, together with a detailed description of the local irradiation method, is presented in chapter 7. The cellular characteristics of XPA, an NER protein involved in damage verification were further investigated in chapter 8. In contrast to what had been reported previously, we found that XPA does not form a cellular complex with the single-strand binding protein RPA. After having investigated the functioning of essential NER proteins, the role of the accessory damage recognition factor UV-DDB in NER was assessed. Using a significantly lower UV dose compared to that used in earlier published research, we discovered a previously ill-recognised role for UV-DDB in 6-4PP repair which is described in chapter 9. Turning the attention to the later stages of the NER reaction, the involvement of DNA polymerases  $\delta$  and  $\epsilon$  and DNA ligase I in the DNA resynthesis step of NER *in vivo* was investigated, while concurrently the stability of the preincision NER complex and the DNA resynthesis complex were assessed. The results of this research are presented in chapter 10.

