



Universiteit
Leiden
The Netherlands

Calcium- and BTB domain protein-modulated PINOID protein kinase directs polar auxin transport

Robert-Boisivon, H.S.

Citation

Robert-Boisivon, H. S. (2008, May 21). *Calcium- and BTB domain protein-modulated PINOID protein kinase directs polar auxin transport*. Retrieved from <https://hdl.handle.net/1887/12863>


Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/12863>


Note: To cite this publication please use the final published version (if applicable).

Color supplements




Chapter 2, Figure 3, page 38.

TOUCH3 is a negative regulator of PINOID *in vivo*.




Chapter 2, Figure 4, page 40.

TOUCH3 and PINOID co-localization is auxin-dependent.



Chapter 2, Figure 6, page 42.

TOUCH3 and auxin cause PINOID to dissociate from the plasma membrane.




Chapter 3, Figure 2, page 60.
pbp1-1 loss-of-function enhances *pid-14* embryo phenotypes.




Chapter 3, Figure 3, page 63.

pbp1-1 loss-of-function partially rescues *pid-14* inflorescences.




Chapter 3, Figure 5, page 65.

PBP1 and PBP1H act redundantly on root growth, embryo patterning and leaf phyllotaxis.




Chapter 4, Figure 2, page 81.

The expression pattern of the BT proteins suggests a genetic redundancy among the family members.




Chapter 4, Figure 4, page 86.
The quintuple *bt* loss-of-function is gametophytic lethal.



Chapter 5, Figure 4, page 107.

Overexpression of *BT1* enhances *pid-14* embryo phenotypes and inhibits *35Spro::PID-21* root meristem collapse.



Chapter 5, Figure 6, page 110.

RPS5Apro>>BTB expression results in axilliary branch and inflorescence meristem defects.

