



Universiteit
Leiden
The Netherlands

Biocontrol of tomato foot and root rot by *Pseudomonas* bacteria in stonewool

Validov, S.

Citation

Validov, S. (2007, December 6). *Biocontrol of tomato foot and root rot by Pseudomonas bacteria in stonewool*. Retrieved from <https://hdl.handle.net/1887/12480>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12480>

Note: To cite this publication please use the final published version (if applicable).

Reference List

- Abbas,H.K., Mirocha,C.J., Vesonder,R.F., and Gunther,R. (1990) Acute toxic effects of an isolate of moniliformin-producing *Fusarium oxysporum* and purified moniliformin on rats. Arch. Environ. Contam Toxicol. **19**: 433-436.
- Abbasi,P.A., Miller,S.A., Meulia,T., Hoitink, H.A. and Kim,J.M. (1999) Precise detection and tracing of *Trichoderma hamatum* 382 in compost-amended potting mixes by using molecular markers. Appl. Environ. Microbiol. **65**: 5421–5426.
- Abd-Elsalam,K.A.,A. Asran-Amal,F. Schnieder,Q. Migheli, and J.A.Vetreet (2006) Molecular detection of *Fusarium oxysporum* f. sp *vasinfectum* in cotton roots by PCR and real-time PCR assay. J. Plant Dis. Prot. **113**: 14-19.
- Alabouvette,C. and Couteaudier,Y. (1992) Biological control of Fusarium wilts with non-pathogenic Fusaria. In: Biological control of plant diseases. Plenum press, New York, pp. 415-426.
- Altschul,S.E., Madden,T.L., Schaffer,A.A., Zhang,J., Zhang,Z., Miller,W., and LipmanD.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. NAR **25**: 3389-3402.
- Altschul,S.F., Gish,W., Miller,W., Meyers,E.W., and Lipman,D.J. (1990) Basic Local Alignment Search Tool. J. Mol. Biol. **215**: 403-410.
- Amiet-Charpentier,C., Gadille,P., and Benoit,J.P. (1999) Rhizobacteria microencapsulation: properties of microparticles obtained by spray-drying. J. Microencapsul. **16**: 215-229.
- Amiet-Charpentier,C., Gadille,P., Digat,B., and Benoit,J.P. (1998) Microencapsulation of rhizobacteria by spray-drying: formulation and survival studies. J. Microencapsul. **15**: 639-659.
- Anonymous. (1998) Sichere Biotechnologie. Eingruppierung biologischer Agenzien: Bakterien, BG Chemie, Merkblatt B 006 8/98 ZH 1/346, Jedermann-Verlag Dr. Otto Pfeffer oHG, Heidelberg, Germany.
- Anzai,Y., Kim,H., Park,J.Y., Wakabayashi,H., and Oyaizu,H. (2000) Phylogenetic affiliation of the pseudomonads based on 16S rRNA sequence. Int. J. Syst. Evol. Microbiol. 50 Pt 4: 1563-1589.
- Appel,D.J., and Gordon,T.R., 1996. Relationships among pathogenic and nonpathogenic isolates of *Fusarium oxysporum* based on the partial sequence of the intergenic spacer region of the ribosomal DNA. Mol. Plant-Microbe Interact. **9**: 125-138

- Arie,T.,I. Kaneko, T.Yoshida, M.Noguchi, Y.Nomura, and I.Yamaguchi. 2000. Mating-type genes from asexual phytopathogenic ascomycetes *Fusarium oxysporum* and *Alternaria alternata*. Mol.Plant Microbe Interact. **13**: 1330-1339.
- Attitalla,I.H., Fatehi, J. Levenfors,J. and Brishammar,S. 2004. A rapid molecular method for differentiating two special forms (*lycopersici* and *radicis-lycopersici*) of *Fusarium oxysporum*. Mycol.Res. **108**: 787-794.
- Audenaert,K., Pattery,T., Cornelis,P., and Hofte,M. (2002) Induction of systemic resistance to *Botrytis cinerea* in tomato by *Pseudomonas aeruginosa* 7NSK2: role of salicylic acid, pyochelin, and pyocyanin. Mol. Plant Microbe Interact. **15**: 1147-1156.
- Baayen,R.P.,. Forch,M.G, C. Waalwijk, Bonants,P.J.M., Loffler,H.J.M. and Roebroeck,E.J.A.. 1998. Pathogenic, genetic and molecular characterization of *Fusarium oxysporum* f.sp. *lilii*. Eur. J. Plant Pathol. **104**: 887-894.
- Baayen,R.P., O'Donnell,K., Bonats,P.J.M., Cigelnik,E., Kroon,L.P.N.M. Roebroeck,E.J.A., and Waalwijk,C., 2000. Gene genealogies and AFLP analyses in the *Fusarium oxysporum* complex identify monophyletic and nonmonophyletic formae speciales causing wilt and rot disease, Ecol. Popul. Biol. **90**: 891-900
- Baayen,R.P., O'Donnell,K., Breeuwsman,S., Gerser,D.M., and Wallwijk,C., 2001. Molecular relationships of Fungi within the *Fusarium redolens*- *F. hostae* clade, Mycology **91**: 1037-1043
- Baker,K.F. and Snyder,W.C. eds. (1965) Ecology of soil-borne plant pathogens: prelude to biological control. Univ. of California Press. Berkeley.
- Bano,N. and Musarrat,J. (2003) Characterization of a new *Pseudomonas aeruginosa* strain NJ-15 as a potential biocontrol agent. Curr. Microbiol. **46**: 324-328.
- Bauer,W. D., and Robinson, J. B. (2002) Disruption of bacterial quorum sensing by other organisms. Curr. Opin. Biotechnol. **13**: 234-237
- Bent,A.F. (2002) Crop diseases and strategies for their control. In Plants, genes and Crop Biotechnology Chrispeels M.J. and Savada D.E. (eds) 2nd ed. Sudbury, Canada, Jones and Bartlett Publishers, pp 390-413.
- Berg,H.C. (2003) The rotary motor of bacterial flagella. Annu. Rev. Biochem. **72**: 19-54.
- Berg,G., Eberl,L., and Hartmann,A. (2005) The rhizosphere as a reservoir for opportunistic human pathogenic bacteria. Environ. Microbiol. **7**: 1673-1685.

- Biffen, R.H. 1905. Mendel's laws of inheritance and wheat breeding. J. Agr. Sci., **1**: 4-48
- Bloemberg,G.V. and Lugtenberg,B.J. (2001) Molecular basis of plant growth promotion and biocontrol by rhizobacteria. Curr. Opin. Plant Biol. **4**: 343-350.
- Bloemberg,G.V., Wijffjes,A.H.M., Lamers,G.E.M., Stuurman,N., and Lugtenberg,B.J.J. (2000) Simultaneous imaging of *Pseudomonas fluorescens* WCS365 populations expressing three different autofluorescent proteins in the rhizosphere: New perspectives for studying microbial communities. Mol. Plant-Microbe Interact. **13**: 1170-1176.
- Bolwerk,A., Lagopodi,A.L., Wijffjes,A.H.M. , Lamers,G.E.M. , Chin-A-Woeng,T.F.C., Lugtenberg,B.J.J. and Bloemberg,G.V. (2003). Interactions in the tomato rhizosphere of two *Pseudomonas* biocontrol strains with the phytopathogenic fungus *Fusarium oxysporum* f. sp. *radicis-lycopersici*. Mol. Plant-Microbe Interact. **16**: 983-993.
- Bolwerk,A., Lagopodi,A.L., Lugtenberg,B.J.J., and Bloemberg,G.V. (2005) Visualization of interactions between a pathogenic and a beneficial *Fusarium* strain during biocontrol of tomato foot and root rot. Mol. Plant-Microbe Interact. **18**: 710-721.
- Booth,C. (1970), The genus *Fusarium*. Commonwealth Mycological Institute, Kew, UK.
- Bourgault,A.M. and Lamothe,F. (1988) Evaluation of the KOH test and the antibiotic disk test in routine clinical anaerobic bacteriology. J. Clin. Microbiol. **26**: 2144-2146.
- Brayford,D. (1996) *Fusarium oxysporum* f. sp. *radicis-lycopersici*. Mycopathol. **133**: 61-63.
- Brouwer,M., Lievens,B., Van Hemelrijck,W., Van den,A.G., Cammue,B.P., and Thomma,B.P. (2003) Quantification of disease progression of several microbial pathogens on *Arabidopsis thaliana* using real-time fluorescence PCR. FEMS Microbiol. Lett. **228**: 241-248.
- Brown,M.R. and Foster,J.H. (1970) A simple diagnostic milk medium for *Pseudomonas aeruginosa*. J. Clin. Pathol. **23**: 172-177.
- Bulat,S.A., LuĖbeck, M., Alekhina, I.A., Jensen, D.F., Knudsen, I.M.B. and LuĖbeck, P.S. (2000) Identification of a universally primed-PCR-derived sequence characterized amplified region marker for an antagonistic strain of *Clonostachys rosea* and development of a strain-specific PCR detection assay. Appl. Environ. Microbiol. **66**, 4758–4763.

- Burgess L.W. (1981) General ecology of the fusaria. In: Nelson P.E., Toussoun T.A., Cook R.J., eds. *Fusarium: diseases, biology and taxonomy*. University Park, PA, USA: The Pennsylvania State University Press, 225-235.
- Buyer, J.S. and Sikora, L.J. (1990) Rhizosphere Interactions and Siderophores. *Plant and Soil* **129**: 101-107.
- Cai, G., Rosewich, G.L., Schneider, R.W., Kistler, H.C., Davis, R.M., Elias, K.S., and Miyao, E.M. (2003) Origin of race 3 of *Fusarium oxysporum* f. sp. *lycopersici* at a single site in California. *Ecol. Popul. Biol.* **93**: 1014-1022
- Camacho, M.M. (2001) Molecular characterization of the role of type 4 pili, NDH-1 and PyrR in rhizosphere colonization of *Pseudomonas fluorescens* WCS365. PhD Thesis, Leiden University, Leiden, the Netherlands.
- Carsolio, C., Gutierrez, A., Jimenez, B., van Montagu, M., and Herrera-Estrella, A. (1994) Characterization of ech-42, a *Trichoderma harzianum* endochitinase gene expressed during mycoparasitism. *Proc. Natl. Acad. Sci. U. S. A* **91**: 10903-10907.
- Castric, P.A. (1975) Hydrogen cyanide, a secondary metabolite of *Pseudomonas aeruginosa*. *Can. J. Microbiol.* **21**: 613-618.
- Cazorla, F.M., Duckett, S.B., Bergstrom, E.T., Noreen, S., Odijk, R., Lugtenberg, B.J.J. et al. (2006) Biocontrol of avocado dematophora root rot by antagonistic *Pseudomonas fluorescens* PCL1606 correlates with the production of 2-hexyl 5-propyl resorcinol. *Mol. Plant-Microbe Interact.* **19**: 418-428.
- Chapon, A., Boutin, M., Rimer, D., Delalande, L., Guillerme, A.Y. and Sarniguet, A. (2003) Direct and specific assessment of colonisation of wheat rhizoplane by *Pseudomonas fluorescens* Pf29A. *Eur. J. Plant Pathol.* **109**, 61-70.
- Charest, P.M., Ouellette, G.B. and Pauze, F.J. (1984) Cytological observations of early infection process by *Fusarium oxysporum* f. sp. *radicis-lycopersici* in tomato plants. *Can. J. Botany*, **62**: 1232-1244.
- Chiarini, L., Bevivino, A., Dalmastri, C., Tabacchioni, S., and Visca, P. (2006) Burkholderia cepacia complex species: health hazards and biotechnological potential. *Trends Microbiol.* **14**: 277-286.
- Chin, A.W., Bloemberg, G. V., van der Bij, A.J., van der Drift, K.M.G.F, Schripsema, J., Kroon, B., Scheffer, R.J., Keel, C., Bakker, P.A.H.M., Tichy, H.V., de Bruijn, F.J., Thomas-Oates, J.E. and Lugtenberg, B.J.J. (1998) Biocontrol by phenazine-1-carboxamide-producing *Pseudomonas chlororaphis* PCL1391 of tomato root rot caused by *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Molecular Plant-Microbe Interactions* **11**: 1069-1077.

- Chin-A-Woeng,T.F.C., BloembergG. V., MuldersI. H., DekkersL.C, and LugtenbergB.J. (2000). Root colonization by phenazine-1-carboxamide-producing bacterium *Pseudomonas chlororaphis* PCL1391 is essential for biocontrol of tomato foot and root rot. *Mol.Plant Microbe Interact.* **13**: 1340-1345.
- Chin-A-Woeng,T.F.C., Bloemberg,G.V. and Lugtenberg,B.J.J. (2003). Mechanisms of biological control of phytopathogenic fungi by *Pseudomonas spp.* In: *Plant-Microbe Interactions*, Vol 6. Stacey, G. & Keen, N.T. (eds). The American Phytopathological Society, St. Paul, MN., pp.173-225.
- Chin-A-Woeng,T.F.C., Bloemberg,G.V., and Lugtenberg,B.J.J. (2003) Phenazines and their role in biocontrol by *Pseudomonas* bacteria. *New. Phytol.* **157**: 503-523.
- Chiocchetti,A., Bernardo,I., Daboussi,M.J., Garibaldi,A., Gullino,M.L., Langin,T., Migheli,Q. (1999) Detection of *Fusarium oxysporum* f. sp. *dianthi* in carnation tissue by PCR amplification of transposon insertions. *Phytopathol.* **89**: 1169-1175.
- Clerc,A., Manceau,C., and Nesme,X. (1998) Comparison of randomly amplified polymorphic DNA with amplified fragment length polymorphism to assess genetic diversity and genetic relatedness within genospecies III of *Pseudomonas syringae*. *Appl. Environ. Microbiol.* **64**: 1180-1187.
- Cook,R.J. and Rovira,A.D. (1976) The role of bacteria in biological control of *Gaeumannomyces graminis* by suppressive soil *Soil Biol. Biochem.* **8**, 269-273
- Coroler,L., Elomari,M., Hoste,B., Gillis,M. Izard,D. and H. Leclerc. (1996) *Pseudomonas rhodesiae* sp. nov., a new species isolated from natural mineral waters. *Syst. Appl. Microbiol.* **19**: 600–607.
- Correll,J.C., Gordon,T.R. and Elliott,V.J. (1987). Host Range, Specificity, and Biometrical Measurements of *Leveillula-Taurica* in California. *Plant Disease* **71**: 248-251.
- Cross,J.V. and Polonenko,D.R. (1996) An industry perspective on registration and commercialization of biocontrol agents in Canada. *Can. J. Plant Pathol.* **18**: 446-454.
- Curwin,B.D., Hein,M.J., Sanderson,W.T., Nishioka,M.G., Reynolds,S.J., Ward,E.M., and Alavanja,M.C. (2005) Pesticide contamination inside farm and nonfarm homes. *J. Occup. Environ. Hyg.* **2**: 357-367.
- de Clercq, D., Cognet, S., Pujol, M., Lepoivre, P. and Jijakli, M.H. (2003) Development of a SCAR marker and a semi-selective medium for specific

- quantification of *Pichia anomala* strain K on apple fruit surfaces. *Postharvest Biol. Technol.* **29**, 237–247.
- de Weert, S., H. Vermeiren, I. H. M. Mulders, I. Kuiper, N. Hendrickx, G. V. Bloemberg, J. Vanderleyden, R. de Mot, and B. J. J. Lugtenberg. (2002) Flagella-driven chemotaxis towards exudate components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. *Mol. Plant-Microbe Interact.* **15**: 1173-1180.
- de Weert, S., Kuiper, I., Lagendijk, E. L., Lamers, G. E. M., and Lugtenberg, B. J. J. (2004) Role of chemotaxis toward fusaric acid in colonization of hyphae of *Fusarium oxysporum* f. sp. *radicis-lycopersici* by *Pseudomonas fluorescens* WCS365. *Mol. Plant-Microbe Interact.* **17**: 1185-1191.
- de Weert, S., Dekkers, L. C., Kuiper, I., Bloemberg, G. V., and Lugtenberg, B. J. (2004) Generation of enhanced competitive root-tip-colonizing *Pseudomonas* bacteria through accelerated evolution. *J. Bacteriol.* **186**: 3153-3159.
- de Weger, L. A., van der Vlugt, C. I. M., Wijffjes, A. H. M., Bakker, P. A. H. M., Schippers, B. and Lugtenberg, B. (1987) Flagella of a plant growth stimulating *Pseudomonas fluorescens* strain are required for colonization of potato roots. *J. Bacteriol.* **169**: 2769-2773.
- de Weger, L. A., Jann, B., Jann, K. and Lugtenberg, B. J. J. (1987a) Lipopolysaccharides of *Pseudomonas* spp. that stimulate plant growth: composition and use for strain identification. *J. Bacteriol.* **169**: 1441-1446.
- Dekkers, L. (1997) Isolation and characterization of novel rhizosphere colonization mutants of *Pseudomonas fluorescens* WCS365. Leiden University. Leiden. The Netherlands.
- Dekkers, L. C., Bloemendaal, C. J., de Weger, L. A., Wijffelman, C. A., Spaink, H. P. and Lugtenberg, B. J. J. (1998) A two-component system plays an important role in the root-colonizing ability of *Pseudomonas fluorescens* strain WCS365. *Mol. Plant-Microbe Interact.* **11**: 45-56.
- Dekkers, L. C., Mulders, I. H., Phoelich, C. C., Chin-A-Woeng, T. F. C., Wijffjes, A. H. M., and Lugtenberg, B. J. J. (2000) The *sss* colonization gene of the tomato-*Fusarium oxysporum* f. sp. *radicis-lycopersici* biocontrol strain *Pseudomonas fluorescens* WCS365 can improve root colonization of other wild-type *Pseudomonas* spp. bacteria. *Mol. Plant-Microbe Interact.* **13**: 1177-1183.
- Desaki, Y., Miya, A., Venkatesh, B., Tsuyumu, S., Yamane, H., Kaku, H. et al., (2006) Bacterial lipopolysaccharides induce defense responses associated with programmed cell death in rice cells. *Plant Cell Physiol.* **47**: 1530-1540.

- Ditta,G., Stanfield,S., Corbin,D., and Helinski,D.R. (1980) Broad host range DNA cloning system for gram-negative bacteria: construction of gene bank of *Rhizobium meliloty*. *Proc. Natl. Acad. Sci. USA* **77**: 7347-7351.
- Donaldson,G.C., Ball,L.A., Axelrood,P.E. and Glass,N.L., 1995. Primer sets developed to amplify conserved genes from filamentous ascomycetes are useful in differentiating *Fusarium* species associated with conifers. *Appl. Environ. Microbiol.* **61**, 1331-1340
- Duffy,B.K. and Défago,G. (1999) Environmental factors modulating antibiotic and siderophore biosynthesis by *Pseudomonas fluorescens* biocontrol strains. *Appl. Environ. Microbiol.* **65**: 2429-2438.
- Duffy,B.K. and Défago,G. (1997) Zinc improves biocontrol of *Fusarium* crown and root rot of tomato by *Pseudomonas fluorescens* and represses the production of pathogen metabolites inhibitory to bacterial antibiotic biosynthesis. *Phytopathol.* **87**: 1250-1257.
- Duijff,B.K., Pouhair,D., Olivain,C., Alabouvette,C., and Lemanceau,P. (1998) Implication of systemic induced resistance in the suppression of *Fusarium* wilt of tomato by *Pseudomonas fluorescens* WCS417r and nonpathogenic *Fusarium oxysporum* Fo47. *Eur. J. Plant Pathol.* **104**: 903-910.
- Edel,V., Steinberg,C., Gautheron,G., Recorbet,G., and Alabouvette,C, (2001) Genetic diversity of *Fusarium oxysporum* populations isolated from different soils in France, *FEMS Microbiol. Ecol.* **36**: 61-71
- Egamberdieva,D., Kamilova,F., Validov,S., Gafurova,L., Kucharova,Z., Lugtenberg,B. High incidence of plant growth-stimulating bacteria associated with the rhizosphere of wheat grown on salinated soil in Uzbekistan *Environ. Microbiol. (OnlineEarly Articles)* doi: 10.1111/j.1462-2920.2007.01424.x
- Elias,K.S., Schneider,R.W., and Lear,M.M. 1992. Genetic diversity within and among races and vegetative compatibility groups of *Fusarium oxysporum* f. sp. *lycopersici* as determined by isozyme analysis. *Phytopathology*, **82**: 1421-1427
- Elias,K.S., Zamir,D., Lichtman-Pleban,T., and Katan,T. 1993. Population structure of *Fusarium oxysporum* f. sp. *lycopersici*: restriction fragment length polymorphisms provide genetic evidence that vegetative compatibility group is an indicator of evolutionary origin. *Mol. Plant Microbe Interact.* **6**: 565-572
- Enright,M.R., McInerney,J.O., and Griffin,C.T. (2003) Characterization of endospore-forming bacteria associated with entomopathogenic nematodes, *Heterorhabditis spp.*, and description of *Paenibacillus nematophilus* sp. nov. *Int. J. Syst. Evol. Microbiol.* **53**: 435-441.

- European Commission. (2001) Directive 2001/18/EC of the European Parliament and of the Council of 12 March 2001 on the deliberate release into the environment of genetically modified organisms and repealing Council Directive 90/220/EEC-Comission Declaration Official Journal of the European Communities L106, 17/04/2001, pp. 0001–0039.
- Folman,L.B., Postma,J., and van Veen,J.A. (2001) Ecophysiological characterization of rhizosphere bacterial communities at different root locations and plant developmental stages of cucumber grown on rockwool. *Microb. Ecol.* **42**: 586-597.
- Fontanille,P. and Larroche,C. (2003) Optimization of isonovalal production from alpha-pinene oxide using permeabilized cells of *Pseudomonas rhodesiae* CIP 107491. *Appl. Microbiol. Biotechnol.* **60**: 534-540.
- Fravel,D., Olivian,C., Alabouvette,C. (2003) *Fusarium oxysporum* and its biocontrol. *New Phytologist.*, **157**: 493-502.
- Fravel,D.R., Rhodes,D.J. and Larkin,R.P. (1999) Production and commercialization of biocontrol products In: Integrated Pest and Disease Management in Greenhouse Crops (Albajes, R., Gullino, L.M., van Lenteren, J.C. and Elad, Y., Eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Fravel,D. (1999) Hurdles and bottlenecks on the road to biocontrol of plant pathogens. *Australasian Plant Pathology* **28**: 53-56.
- Fuchs,J.-G, Moenne-Loccoz,Y., and Defago,G. (1997) Nonpathogenic *Fusarium oxysporum* strain Fo47 induces resistance to *Fusarium* wilt in tomato. *Plant Dis.* **81**: 492-496.
- Garret S.D. (1970) Pathogenic root-infection fungi. London, UK: Cambridge University Press
- Gerrits,J.P.L. and Weisbeek,P.G. (1996) Induction of systemic aquired resistance by saprophytic *Pseudomonas spp.* In the model plant *Arabidopsis thaliana*, p.13-14. In: NWO-LNV Priority Program Crop Protection Progress Report. Lunteren, The Netherlands.
- Gordon,S.A., and Weber,R.P. (1951) Colorimetric estimation of indole acetic acid. *Plant Physiol.* **26**: 192-195.
- Gordon,T.R. and Martyn,R.D. 1997. The evolutionary biology of *Fusarium oxysporum*. *Annu.Rev.Phytopathol.* **35**: 111-128

- Gunther,R., Kishore,P.N., Abbas,H.K., and Mirocha,C.J. (1989) Immunosuppressive effects of dietary wortmannin on rats and mice. *Immunopharmacol. Immunotoxicol.* **11**: 559-570.
- Haas,D. and Defago,G. (2005) Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat. Rev. Microbiol.* **3**: 307-319.
- Hahn,F. (2002) Fungal spore detection on tomatoes using spectral Fourier signatures. *Biosyst. Engineer.* **81** : 249-259.
- Han,J., Sun,L., Dong,X., Cai,Z., Sun,X., Yang,H. Wang,Y. and Song, W. (2005) Characterization of a novel plant growth-promoting bacteria strain *Delftia tsuruhatensis* HR4 both as a diazotroph and a potential biocontrol agent against various plant pathogens. *Syst. Appl. Microbiol.* **28**: 66-76.
- Hanks,G.R. (1996) Control of *Fusarium oxysporum* f.sp.*narcissi*, the cause of narcissus basal rot, with thiabendazole and other fungicides. *Crop Protection*, **15**: 549-558
- Harman,G.E., Howell,C.H., Vitero,A., Chet,I., and Lorito,M. (2004) *Trichoderma* species-opportunistic, avirulent plant symbionts. *Nature Rev. Microbiol.* **2**: 43-56.
- Hartmann,A., Gantner,S., Schuegger,R., Steidle,A., Durr,C., Schmid,M., Dazzo,F., Eberl,L. and Langgebartels,C. (2004) N-acyl homoserin lactones of rhizosphere bacteria trigger systemic resistance in tomato plants. In: *Proceedings of the 11th International Congress on Mol. Plant-Microbe Interact.s.* St.-Petersburg, Russia, July 18-26, 2003. Tikhonovich I., Lugtenberg B. and Provorov N. (eds.). pp. 554-556.
- Heath,D. (1995) The chi-squared test for a difference between two population proportions. In: *An introduction to experimental design and statistics for biology.* University College, London, UK, pp. 216-218.
- Hoffland,E., Findenegg,G.R., and Nelemans,J.A. (1989) Solubilization of rock phosphate by rape. *Plant Soil* **113**: 161-165.
- Hong,S., Park,Y., Jeong,Y., Bae,K.S., (2000) Sequence Comparison of Mitochondrial Small Subunit Ribosomal DNA in *Penicillium*, *J. Microbiol.* **38**: 62-65
- Howe,T.G. and Ward,J.M. (1976) The utilization of Tween 80 as carbon source by *Pseudomonas*. *J. Gen. Microbiol.* **92**: 234-235.

- Howie, W.J., Cook, R.J. and Weller, D.M. (1987). Effects of soil matric potential and cell motility on wheat root colonization by fluorescent pseudomonads suppressive to take-all. *Phytopathol.* **77**: 286-292.
- Hunt, M.D., Neuenschwander, U.H., Delaney, T.P., Weymann, K.B., Friedrich, L.B., Lawton, K.A., Steiner, H.Y., Ryals, J.A. (1996) Recent advance in systemic acquired resistance. *Gene* **7**: 89-95.
- Iavicoli, A., Boutet, E., Buchala, A. and Metraux, J.-P. (2003) Induced systemic resistance in *Arabidopsis thaliana* in response to root inoculation with *Pseudomonas fluorescens* CHA0. *Mol. Plant-Microbe Interact.* **10**: 851-858.
- Ishikawa, R., Shirouzu, K., Nakashita, H., Lee, H.Y., Motoyama, T., Yamaguchi, I. et al. (2005) Foliar spray of validamycin A or validoxylamine A controls tomato Fusarium wilt. *Phytopathol.* **95**: 1209-1216.
- Jacobson, D.J. and Gordon, T.R. (1990) Variability of Mitochondrial-Dna As An Indicator of Relationships Between Populations of *Fusarium oxysporum* f.sp. *melonis*. *Mycological Research.* **94**: 734-744.
- Jain, D.K., Thomson, D.K.C., Lee, H., and Trevors, J.T. (1991) A drop-collapsing test for screening surfactant producing microorganisms. *J. Microbiol. Methods* **13**: 271-279.
- Jones, J., Jones, J.P., Stall, R.E., and Zitter, T.A. (1991) Compendium of tomato diseases. American Phytopathological Society, St. Paul, MN.
- Jones, J., Jones, J.P., Stall, R.E., and Zitter, T.A. 1991. Compendium of tomato diseases. American Phytopathological Society, St. Paul, MN.
- Jones, K.A. and Burges, H.D. (1998) Technology of formulation and application. In *Formulation of microbial biopesticides. Beneficial microorganisms, nematodes and seed treatment*. Burges, H.D. (ed). Dordrecht, The Netherlands, Kluwer Academic Publishers, pp. 7-30.
- Kahng, H.Y., Nam, K., Kukor, J.J., Yoon, B.J., Lee, D.H., Oh, D.C. et al. (2002) PAH utilization by *Pseudomonas rhodesiae* KK1 isolated from a former manufactured-gas plant site. *Appl. Microbiol. Biotechnol.* **60**: 475-480.
- Kamilova, F., Kravchenko, L.V., Shaposhnikov, A.I., Azarova, T., Makarova, N., and Lugtenberg, B. (2006-a) Organic acids, sugars, and L-tryptophane in exudates of vegetables growing on stonewool and their effects on activities of rhizosphere bacteria. *Mol. Plant-Microbe Interact.* **19**: 250-256.
- Kamilova, F., Kravchenko, L.V., Shaposhnikov, A.I., Makarova, N., and Lugtenberg, B. (2006-b) Effects of the tomato pathogen *Fusarium oxysporum* f.

- sp radialis-lycopersici* and of the biocontrol bacterium *Pseudomonas fluorescens* WCS365 on the composition of organic acids and sugars in tomato root exudate. *Mol. Plant-Microbe Interact.* **19**: 1121-1126.
- Kamilova,F., Validov,S., Azarova,T., Mulders,I., and Lugtenberg,B. (2005) Enrichment for enhanced competitive plant root tip colonizers selects for a new class of biocontrol bacteria. *Environ. Microbiol.* **7**: 1809-1817.
- Katan,J., Grinstein,A., Greenberger,A., Yarden,O. and DeVay,J.E. (1987). First decade (1976-1986) of soil solarization (solar heating)-A chronological bibliography. *Phytoparasitica* **15**: 229-255.
- Katan,T. (1999) Current status of vegetative compatibility groups in *Fusarium oxysporum*. *Phytoparasitica*. **27**: 51-64.
- Katan,T., Zamir,D., Sarfatti,M. and Katan,J. (1991). Vegetative Compatibility Groups and Subgroups in *Fusarium oxysporum* f.sp. *radialis-lycopersici*. *Phytopathology*. **81**: 255-262.
- Keel,C., Schnider,U., Maurhofer,M., Voisard,C., Laville,J., Burger,U., Wirthner,P., Haas,D., Defago,G. (1992) Suppression of root diseases by *Pseudomonas fluorescens* CHA0: importance of the bacterial secondary metabolite 2,4-diacetyl phloroglucinol. *Mol. Plan-Microbe Interact.* **5**: 4-13.
- Kerenyi,Z., Moretti,A., Waalwijk,C., Olah,B. and Hornok,L. (2004) Mating type sequences in asexually reproducing *Fusarium* species. *Appl.Environ.Microbiol.* **70**: 4419-4423.
- Kim,Y.T., Yoo,S.J. and Kim,H.G. (1997) [Some genetic characteristics of tomato wilt pathogen, *Fusarium oxysporum* f.sp. *lycopersici* in Korea.] *Korean J. Plant Pathol.* **13**: 299-303 (in Korean).
- King, E.O., Ward, M.K. and Raney, D.E. (1954) Two simple media for the demonstration of pyocyanin and fluorescin. *J. Lab. Clin. Med.* **44**: 301-307.
- Kistler, H. C., C. Alabouvette, R. P. Baayen, S. Bentley, D. Brayford, A. Coddington, J. Correll, M. J. Daboussi, K. Elias, D. Fernandez, T. R. Gordon, T. Katan, H. G. Kim, J. F. Leslie, R. D. Martyn, Q. Migheli, N. Y. Moore, K. O'Donnell, R. C. Ploetz, M. A. Rutherford, B. Summerell, C. Waalwijk, and S. Woo. (1998) Systematic numbering of vegetative compatibility groups in the plant pathogenic fungus *Fusarium oxysporum*. *Phytopathol.* **88**: 30-32.
- Kistler,H.C. (1997) Genetic diversity in the plant-pathogenic fungus *Fusarium oxysporum*. *Phytopathology* **87**: 474-479.

- Kloepper, J.W., Schroth, M.N., and Miller, T.D. (1980) Effect of rhizosphere colonization by plant growth-promoting rhizobacteria on potato plant development and yield. *Phytopathol.* **70**: 1078-1082.
- Koch, E. (2001) Authorisation of microbial preparation as plant strengthening products. In: (Koch, E. and Leinonen, P., eds.) *Formulation of microbial inoculants*. Working Group 2 Proceedings of a meeting held in Braunschweig, Germany, 5 and 6 February 1999, organised by the Institute of Technology, FAL, Germany. COST Action 830 "Microbial inoculants for agriculture and environment".
- Koenig, R.L., Ploetz, R.C. and Kistler, H.C. (1997) *Fusarium oxysporum* f. sp. cubense consists of a small number of divergent and globally distributed clonal lineages. *Phytopathol.* **87**, 915-923.
- Kroon, B.A.M. (1992) *Fusarium* wilt of tomato. Aspects of resistance and pathogenecity. PhD Thesis. Amsterdam University, Amsterdam, the Netherlands.
- Kuiper, I., Bloemberg, G. V., and Lugtenberg, B.J.J. (2001-a) Selection of a plant-bacterium pair as a novel tool for rhizostimulation of polycyclic aromatic hydrocarbon-degrading bacteria. *Mol. Plant-Microbe Interact.* **14**: 1197-1205.
- Kuiper, I., Kravchenko, L., Bloemberg, G.V., and Lugtenberg, B.J.J. (2002) *Pseudomonas putida* strain PCL1444, selected for efficient root colonization and naphthalene degradation, effectively utilizes root exudate components. *Mol. Plant-Microbe Interact.* **15**: 734-741.
- Kuiper, I., Bloemberg, G.V., Noreen, S., Thomas-Oates, J.E., and Lugtenberg, B.J.J. (2001b) Increased uptake of putrescine in the rhizosphere inhibits competitive root colonization by *Pseudomonas fluorescens* strain WCS365. *Mol. Plant-Microbe Interact.* **14**: 1096-1104.
- Kunze, G., Zipfel, C., Robatzek, S., Niehaus, K., Boller, T., and Felix, G. (2004) The N terminus of bacterial elongation factor Tu elicits innate immunity in Arabidopsis plants. *Plant Cell* **16**: 3496-3507.
- Lagopodi, A.L., Ram, A.F.J., Lamers, G.E.M., Punt, P.J., Van den Hondel, C.A.M.J., Lugtenberg, B.J.J., and Bloemberg, G.V. (2002) Novel aspects of tomato root colonization and infection by *Fusarium oxysporum* f. sp. *radicis-lycopersici* revealed by confocal laser scanning microscopic analysis using the green fluorescent protein as a marker. *Mol. Plant-Microbe Interact.* **15**: 172-179.
- Lee, S.B., and Taylor, J.W., (1992) Phylogeny of five fungus-like protocistan Phytophthora species, inferred from the internal transcribed spacers of ribosomal DNA. *Mol. Boil. Evol.* **9**: 636-653

- Lee, S.W. and Cooksey, D.A. (2000) Genes expressed in *Pseudomonas putida* during colonization of a plant-pathogenic fungus. *Appl. Environ. Microbiol.* **66**: 2764-2772.
- Lemanceau, P., and Alabouvette, C. (1990). Biological control of *Fusarium* diseases by the association of fluorescent *Pseudomonas* and non-pathogenic *Fusarium*. In: Plant growth promoting rhizobacteria report. Keel C., Knoller B., and Defago G. (eds), Interlaken, Switzerland, pp. 45-50.
- Lemanceau, P., Bakker, P.A.H.M., Dekogel, W.J., Alabouvette, C., and Schippers, B. (1992) Effect of Pseudobactin-358 Production by *Pseudomonas-Putida* Wcs358 on Suppression of *Fusarium*-Wilt of Carnations by Nonpathogenic *Fusarium oxysporum* Fo47. *Applied and Environ. Microbiol.* **58**: 2978-2982.
- Lievens, B., Brouwer, M., Vanachter, A.C.R.C., Levesque, C.A., Cammue, B.P.A. and Thomma, B.P.H.J. (2005) Quantitative assessment of phytopathogenic fungi in various substrates using a DNA macroarray. *Environ. Microbiol.* **7**: 1698-1710.
- Lievens, B., Brouwer, M., Vanachter, A.C., Levesque, C.A., Cammue, B.P., and Thomma, B.P. (2003) Design and development of a DNA array for rapid detection and identification of multiple tomato vascular wilt pathogens. *FEMS Microbiol. Lett.* **223**: 113-122.
- Ligon, J.M., Hill, D.S., Hammer, P.E., Torkewitz, N.R., Hofmann, D., Kempf, H.J., and van Pee, K.H. (2000) Natural products with antifungal activity from *Pseudomonas* biocontrol bacteria. *Pest Manag. Sci.* **56**: 688-695.
- Lorito, M., Farkas, V., Rebuffat, S., Bodo, B., and Kubicek, C.P. (1996) Cell wall synthesis is a major target of mycoparasitic antagonism by *Trichoderma harzianum*. *J. Bacteriol.* **178**: 6382-6385.
- Louws, F.J., Rademaker, J.L.W., Bruijn, F.J. (1999) The three Ds of PCR-based genomic analysis of phyto-bacteria: Diversity, Detection and Disease Diagnosis. *Annu. Rev. Phytopathol.* **37**: 81-125
- Lugtenberg, B.J.J., Dekkers, L.C. (1999). What makes *Pseudomonas* bacteria rhizosphere competent? *Environ. Microbiol.* **1**: 9-13.
- Lugtenberg, B.J.J. and Bloemberg, G.V. (2004) Life in the rhizosphere. In: *Pseudomonas*. V.1. Ramos, J.L. (ed), Kluwer Academic / Plenum Publishers, New York, pp.403-430.
- Lugtenberg, B.J.J., Dekkers, L.C. and Bloemberg, G.V. (2001) Molecular determinants of rhizosphere colonization by *Pseudomonas*. *Annu. Rev. Phytopathol.* **39**: 461-490.

- Mahaffee,W.F., Bauske,E.M., van Vuurde,J.W.L., van der Wolf,J.M., van den Brink,M. and Kloepper,J.W. (1997) Comparative analysis of antibiotic resistance, immunofluorescent colony staining, and a transgenic marker (Bioluminescence) for monitoring the environmental fate of a rhizobacterium. Appl. Environ. Microbiol. **63**, 1617–1622.
- Mari,M., Bertolini,P., and Pratella,G.C. (2003) Non-conventional methods for the control of post-harvest pear diseases. J. Appl. Microbiol. **94**: 761-766.
- Marlatt,M.I., Correll,J.C., Kaufmann,P., and Cooper,P.E. (1996) Two genetically distinct populations of *Fusarium oxysporum* f.sp. *lycopersici* race 3 in the United States. Plant Dis. 80: 1336-1342
- Marschner,H. (1995) Mineral Nutrition of Higher Plants, Ed 2. Academic Press, London.
- Mavrodi,D.V., Mavrodi,O.V., Spadden-Gardener,B.B., Landa,B.B., Weller,D.M., and Thomashow,L.S. (2002) Identification of differences in genome content among pHID-positive *Pseudomonas fluorescens* strains by using PCR-based subtractive hybridization. Appl. Environ. Microbiol. **68**: 5170-5176.
- Milton,D.L., Hardman,A., Camara,M., Chabra,S.R., Bycroft,B.W., Stewart,G.B., and Williams,P. (1997) Quorum sensing in *Vibrio anguillarum*: characterization of vanI/vanR locus and identification of the autoinducer N-(3-oxodecanoyl)-L-homoserine lactone. J. Bacteriol. **179**: 3004-3012.
- Mirocha,C.J., Abbas,H.K., Kommedahl,T., and Jarvis,B.B. (1989) Mycotoxin production by *Fusarium oxysporum* and *Fusarium sporotrichioides* isolated from *Baccharis spp.* from Brazil. Appl. Environ. Microbiol. **55**: 254-255.
- Montesinos,E. (2003) Development, registration and commercialization of microbial pesticides for plant protection. Int. Microbiol. **6**: 245-252.
- Moore,W.C., Brunt,A.A., Price,D., Rees,A.R. and Dickens,J.S.W. (1979) Diseases of bulbs, Second edn. Reference Book HPD1. Her Majesty's Stationery Office, London, 205 pp
- Morello,J.E., Pierson,E.A. and Pierson III,L.S. (2004) Negative cross-communication among wheat rhizosphere bacteria: effect on antibiotic production by the biological control bacterium *Pseudomonas aureofaciens* 30-84. Appl. Environ. Microbiol. **70**: 3103-3109.
- Nam,H.S., Anderson,A.J., Yang,K.Y., Cho,B.H., and Kim,Y.C. (2006) The *dctA* gene of *Pseudomonas chlororaphis* O6 is under RpoN control and is required for effective root colonization and induction of systemic resistance. FEMS Microbiol. Lett. **256**: 98-104.

- Nam,H.S., Spencer,M., Anderson,A.J., Cho,B.H., and Kim,Y.C. (2003) Transcriptional regulation and mutational analysis of a *dctA* gene encoding an organic acid transporter protein from *Pseudomonas chlororaphis* O6. *Gene* **323**: 125-131.
- Nelson,A.J., Elias,K.S., Arevalo,G.E., Darlington,L.C. and Bailey,B.A. (1997) Genetic characterization by RAPD analysis of isolates of *Fusarium oxysporum* f.sp. *erythroxyli* associated with an emerging epidemic in Peru. *Phytopathol.* **87**: 1220~1225.
- Nelson,P.E., Toussoun,T.A. and Cook,R.J. (1981) *Fusarium: Diseases, Biology and Taxonomy*. Pennsylvania State University Press, University Park, PA.
- Notz,R., Maurhofer,M., Dubach,H., Haas,D., and Defago,G. (2002) Fusaric acid-producing strains of *Fusarium oxysporum* alter 2,4- diacetylphloroglucinol biosynthetic gene expression in *Pseudomonas fluorescens* CHA0 in vitro and in rhizosphere of wheat. *Appl. Environ. Microbiol.* **68**: 2229-2235.
- O'Donnell,K., Cigelnik,E. and Nirenberg,H.I. (1998) Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia.* **90**: 465-493.
- O'Donnell,K., Kistler,H.C., Cigelnik,E., and Ploetz,R.C. (1998) Multiple evolutionary origins of the fungus causing Panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proc. Natl. Acad. Sci. USA* **95**: 2044-2049.
- Olivain,C. and Alabouvette,C. (1999) Process of tomato root colonization by a pathogenic strain of *Fusarium oxysporum* f. sp. *lycopersici* in comparison with a non-pathogenic strain. *New Phytologist* **141**: 497-510.
- Olivain,C., Trouvelot,S., Binet,M.N., Cordier,C., Pugin,A., and Alabouvette,C. (2003) Colonization of flax roots and early physiological responses of flax cells inoculated with pathogenic and nonpathogenic strains of *Fusarium oxysporum*. *Appl. Environ. Microbiol.* **69**: 5453-5462.
- Palmfeldt,J., Radstrom,P., and Hahn-Hagerdal,B. (2003) Optimization of initial cell concentration enhances freeze-drying tolerance of *Pseudomonas chlororaphis*. *Cryobiol.* **47**: 21-29.
- Pasquali,M., Acquadro,A., Balmas,V., Migheli,Q., Gullino,M.L., Garibaldi,A. (2003) RAPD characterization of *Fusarium oxysporum* isolates pathogenic on *Argyranthemum frutescens* L. *J. Phytopathol.* **151**: 30-35.

- Pasquali,M., Acquadro,A., Balmas,V., Migheli,Q., Gullino,M.L. and Garibaldi,A. (2004). Development of PCR primers for a new *Fusarium oxysporum* pathogenic on Paris daisy (*Argyranthemum frutescens* L.). Eur. J. Plant Pathol. **110**: 7-11.
- Pasquali,M., Piatti,M. Gullino,L., and Garibaldi,A. (2006). Development of a real-time polymerase chain reaction for the detection of *Fusarium oxysporum* f. sp basilici from basil seed and roots. Eur. J. Plant Pathol. **154**: 632-636.
- Paulitz,T.C. (2000) Population dynamics of biocontrol agents and pathogens in soils and rhizospheres. Eur. J. Plant Pathol. **106**: 401-413.
- Paulitz,T.C. and Bélanger,R.R. (2001) Biological control in greenhouse systems. Annu. Rev. Phytopathol. **39**: 103-133.
- Paulitz,T.C., Park,C.S. and Baker,R. (1987) Biological control of *Fusarium* wilt of cucumber with nonpathogenic isolates of *Fusarium oxysporum*. Can. J. Microbiol. **33**: 349-353.
- Pier,A.C. (1981) Mycotoxins and animal health. Adv. Vet. Sci. Comp Med. **25**: 185-243.
- Pierson,L.S., III and Thomashow,L.S. (1992) Cloning and heterologous expression of the phenazine biosynthetic locus from *Pseudomonas aureofaciens* 30-84. Mol. Plant-Microbe Interact. **5**: 330-339.
- Pierson,L.S., Wood,D.W., Pierson,E.A., and Chancey,S.T. (1998) N-acyl-homoserine lactone-mediated gene regulation in biological control by fluorescent pseudomonads: Current knowledge and future work. Eur. J. Plant Pathol. **104**: 1-9.
- Pieterse,C.M.J., van Wees,S.C.M., Hoffland,E., van Pelt,J.A., and van Loon,L.C. (1996) Systemic resistance in *Arabidopsis* induced by biocontrol bacteria is independent of salicylic acid accumulation and pathogenesis related gene expression. Plant Cell **8**: 1225-1237.
- Piper,K.R., Beck von Bodman,S., and Farrand,S.K. (1993). Conjugation factor of *Agrobacterium tumefaciens* regulates Ti-plasmid transfer by autoinduction. Nature **362**: 448-450.
- Pitt,J.I. (2000) Toxigenic fungi: which are important? Med. Mycol. 38 Suppl **1**: 17-22.
- Postma,J. and Luttikholt,A.J.G. (1996) Colonization of carnation stems by a nonpathogenic isolate of *Fusarium oxysporum* and its effect on *Fusarium oxysporum* f sp *dianthi*. Can. J. Botany **74**: 1841-1851

- Postma,J., Willemsen-de Klein,M.J.E.I., and van Elsas,J.D. (2000) Effect of the indigenous microflora on the development of root and crown rot caused by *Pythium aphanidermatum* in cucumber grown on rockwool. *Phytopathol.* **90**: 125-133.
- Prithiviraj,B., Vikram,A., Kushalappa,A.C., and Yaylayan,V. (2004) Volatile metabolite profiling for the discrimination of onion bulbs infected by *Erwinia carotovora* ssp *carotovora*, *Fusarium oxysporum* and *Botrytis allii*. *Eur. J. Plant Pathol.* **110**: 371-377.
- Puhalla,J.E. (1985) Classification of strains of *Fusarium oxysporum* on the basis of vegetative compatibility. *Can. J. Bot.* **63**: 179-183
- Pujol,M., Badosa,E., Cabrefiga,J., Montesinos,E. (2005) Development of a strain-specific quantitative method for monitoring *Pseudomonas fluorescens* EPS62e, a novel biocontrol agent of fire blight FEMS *Microbiol. Lett.* **249**: 343-352.
- Raupach,G.S., Liu,L., Murphy,J.F., Tuzun,S., Kloepper,J.W. (1996) Induced systemic resistance in cucumber and tomato against cucumber mosaic cucumovirus using plant growth-promoting rhizobacteria (PGPR). *Plant Dis.* **80**: 891-894.
- Rep,M., van der Does,H.C., Meijer,M., van Wijk,R., Houterman,P.M., Dekker,H.L., de Koster,C.G., and Cornelissen,B.J. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol.Microbiol.* **53**: 1373-1383.
- Robinson,R.A. (1997), "The Acceptance of Horizontal Resistance in Crops." *Biotechnology and Development Monitor*, N.**33**, P. 24
- Romero,D., de Vicente,A., Rakotoaly,R.H., Dufour,S.E., Veening,J.W., Arrebola,E. et al. (2007) The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* toward *Podosphaera fusca*. *Mol. Plant-Microbe Interact.* **20**: 430-440.
- Rowe R.C. (1980) Comparative pathogenicity and host ranges of *Fusarium oxysporum* isolates causing crown and root rot of greenhouse and field-grown tomatoes in North America and Japan. *Phytopathol.* **70**: 1143-1148.
- Sakiyama,C.C., Paula,E.M., Pereira,P.C., Borges,A.C., and Silva,D.O. (2001) Characterization of pectin lyase produced by an endophytic strain isolated from coffee cherries. *Lett. Appl. Microbiol.* **33**: 117-121.
- Sambrook,J. and Russel,D. (2001) *Molecular Cloning: A Laboratory Manual*. Cold Spring Laboratory Press, Cold Spring Harbor NY, USA.

- Schaad,N.M., Frederick,R.D. (2002) Real-time PCR and its application for rapid plant disease diagnostics. *Can. J. Plant Pathol.* **24**: 250-258.
- Schena,L., Finetti-Sialer,M. and Gallitelli,D. (2002) Molecular detection of strain L47 of *Aureobasidium pullulans*, a biocontrol agent of postharvest diseases. *Plant Dis.* **86**, 54–60.
- Scher,F.M. and Baker,R. (1980) Mechanism of biological control in a *Fusarium*-suppressive soil. *Phytopathol.* **70**: 412-417.
- Schippers,B., Bakker,A.W. and Bakker,P.A.H.M. (1987). Interactions of deleterious and beneficial microorganisms and the effect of cropping practices. *Annu. Rev. Phytopathol.* **25**: 339-358.
- Schnider-Keel,U., Seematter,A., Maurhofer,M., Blumer,C., Duffy,B., Gigot-Bonnefoy,C., Reimann,C., Notz,R., Défago,G., Haas,D. and Keel,C. (2000) Autoinduction of 2,4-diacetylphloroglucinol biosynthesis in the biocontrol agent *Pseudomonas fluorescens* CHA0 and repression by the bacterial metabolites salicylate and pyoluteorin. *J. Bacteriol.* **182**: 1215-1225.
- Schollenberger,M., Muller,H.M., Ruffe,M., Suchy,S., Planck,S., and Drochner,W. (2005) Survey of *Fusarium* toxins in foodstuffs of plant origin marketed in Germany. *Int. J. Food Microbiol.* **97**: 317-326.
- Schouten,A., van den Berg,G., Edel-Hermann,V., Steinberg,C., Gautheron,N., Alabouvette,C., de Vos,C.H., Lemanceau,P., Raaijmakers,J.M. (2004) Defense responses of *Fusarium oxysporum* to 2,4-diacetylphloroglucinol, a broad-spectrum antibiotic produced by *Pseudomonas fluorescens*. *Mol. Plant-Microbe Interact.* **17**: 1201-1211.
- Schuhegger,R., Ihring,A., Gantner,S., Bahnweg,G., Knappe,C., Vogg,G. et al., (2006) Induction of systemic resistance in tomato by N-acyl-L-homoserine lactone-producing rhizosphere bacteria. *Plant Cell Environ.* **29**: 909-918.
- Shigematsu,T., Yumihara,K., Ueda,Y., Numaguchi,M., Morimura,S., and Kida,K. (2003) *Delftia tsuruhatensis* sp. nov., a terephthalate-assimilating bacterium isolated from activated sludge. *Int. J. Syst. Evol. Microbiol.* **53**: 1479-1483.
- Shimahara,K., and Takiguchi,Y. (1988). Preparation of crustacean chitin. In: *Methods in Enzymology*, **161**, pp. 417-423. W.A.Wood and S.T.Kellog (eds.), Academic Press, Inc., San Diego.
- Shipton,P.J., Cook,R.J., Sitton,J.W. (1973) Occurrence and transfer of a biological factor in soil that suppress take-all of wheat in eastern Washington. *Phytopathol.* **63**: 511-517.

- Simons,M., van der Bij,A.J., Brand,J., de Weger,L.A., Wijffelman,C.A., and Lugtenberg,B.J.J. (1996) Gnotobiotic system for studying rhizosphere colonization by plant growth-promoting *Pseudomonas* bacteria. *Mol.Plant-Microbe Interact.* **9**: 600-607.
- Simons,M., Permentier,H.P., de Weger,L.A., Wijffelman,C.A. and Lugtenberg,B.J.J. (1997) Amino acid synthesis is necessary for tomato root colonisation by *Pseudomonas fluorescens* strain WCS365. *Mol. Plant-Microbe Interact.* **10**: 102-106.
- Skrobek,A., Boss,D., Defago,G., Butt,T.M., and Maurhofer,M. (2006) Evaluation of different biological test systems to assess the toxicity of metabolites from fungal biocontrol agents. *Toxicol. Lett.* **161**: 43-52.
- Smith,I.M., J. Dunez,D.H. Phillips, Lelliott,R.A. and Archer, S.A. eds. (1988) *European handbook of plant diseases*. Blackwell Scientific Publications: Oxford. 583pp.
- Smith, L.M., Tola, E., de Boer, P. and O’Gara F. (1999) Signalling by the fungus *Pythium ultimum* represses expression of two ribosomal RNA operons with key roles in the rhizosphere ecology of *Pseudomonas fluorescens* F113. *Environ. Microbiol.* **1**: 495-502.
- Sneath,P.H., Snokal,R.R. (1973) *Numerical Taxonomy*. San Francisco, USA: Freeman Publications.
- Sokal,R.R., and Rohlf,F.J. (1981) *Biometry*. San Fransisco, USA: Freeman Publications.
- Summerell,B.A., Salleh,B. Leslie,J.F. (2003) A utilitarian approach to *Fusarium* identification. *Plant Disease*. **87**: 117-128
- Taylor,J.W., Geiser,D.M., Burt,A. and Oufopanou,V. (1999) The evolutionary biology and populations genetics underlying fungal strain typing. *Clin. Microbiol. Rev.* **12**: 126-146
- Teunissen,H.A., Rep,M., Houterman,P.M., Cornelissen,B.J. and Haring,M.A. (2003) Construction of a mitotic linkage map of *Fusarium oxysporum* based on Foxy-AFLPs. *Mol.Genet.Genomics* **269**: 215-226.
- Teunissen,H.A., Rep,M., Houterman,P.M., Cornelissen,B.J., and Haring,M.A. (2003) Construction of a mitotic linkage map of *Fusarium oxysporum* based on Foxy-AFLPs. *Mol. Genet. Genomics* **269**: 215-226.
- Thomashow,L.S. and WellerD.M. (1995) Current concepts in the use of introduced bacteria for biological disease control: mechanisms and antifungal

- metabolites. In: Plant-Microbe Interactions. G.Stacey and N.T.Keen (eds.), Chapman & Hall, New York, pp.187-235.
- Thomashow,L.S. and Weller,D.M. (1990) Role of Antibiotics and Siderophores in Biocontrol of Take-All Disease of Wheat. *Plant and Soil* **129**: 93-99.
- Thomashow,L.S., Weller,D.M., Bonsall,R.F., and Pierson,L.S. (1990) Production of the Antibiotic Phenazine-1-Carboxylic Acid by Fluorescent *Pseudomonas* Species in the Rhizosphere of Wheat. *Appl. Environ. Microbiol.* **56**: 908-912.
- Thompson,J.D., Higgins,D.G., Gibson,T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *NAR* **22**: 4673-4680.
- Tjamos,S.E., Flemetakis,E., Paplomatas,E.J., and Katinakis,P. (2005) Induction of resistance to *Verticillium dahliae* in *Arabidopsis thaliana* by the biocontrol agent K-165 and pathogenesis-related proteins gene expression. *Mol. Plant Microbe Interact.* **18**: 555-561.
- Vakalounakis, D. J. and G. A. Fragkiadakis. (1999) Genetic diversity of *Fusarium oxysporum* isolates from cucumber: Differentiation by pathogenicity, vegetative compatibility, and RAPD fingerprinting. *Phytopathol.* **89**: 161-168.
- Vakalounakis,D.J., Fragkiadakis,G.A. (2003) Plant pathobreeding with emphasis in tomato and cucurbits (in Greek). Vakalounakis, Heraklio, Greece.
- Vakalounakis,D.J., Wang,Z., Fragkiadakis,G.A., Skaracis,G.N., Li,D.-B. (2004) Characterization of *Fusarium oxysporum* isolates obtained from cucumber (*Cucumis sativus*) in China by pathogenicity, VCGs and RAPD. *Plant Dis.* **88**: 645-649.
- Validov,S., Kamilova,F., Qi,S., Stephan,D., Wang,J.J., Makarova,N., and Lugtenberg,B. (2007) Selection of bacteria able to control *Fusarium oxysporum* f. sp. *radicis-lycopersici* in stonewool substrate. *J. Appl. Microbiol* **102**: 461-471.
- van den Broek,D., Chin-A-Woeng,T.F., Eijkemans,K., Mulders,I.H., Bloemberg,G.V. and LugtenbergB.J.J. (2003) Biocontrol traits of *Pseudomonas spp.* are regulated by phase variation. *Mol Plant- Microbe Interact* **16**: 1003-1012
- van Loon L.C, Bakker, P.A, Pieterse C.M. (1998) Systemic resistance induced by rhizosphere bacteria. *Annu. Rev. Phytopathol.* **36**: 453-83.

- van Rij,E.T., Wesselink,M., Chin-A-Woeng,T.F.C., Bloemberg,G.V. and Lugtenberg,B.J.J. (2004) Influence of environmental conditions on the production of phenazine-1-carboxamide by *Pseudomonas chlororaphis* PCL1391. *Mol. Plant-Microbe Interact.* **17**: 557-566.
- van Wees,S.C.M., Pieterse,C.M.J., Trijssenaar,A., van 't Westende,Y., Hartog,F., and van Loon,L.C. (1997) Differential induction of systemic resistance in *Arabidopsis* by biocontrol bacteria. *Mol. Plant-Microbe Interact.* **10**: 716–24
- Vanneechoutte,M., Boerlin,R., Tichy,H.-V., Bannerman,E., Jager,B., and Bille,J. (1998) Comparison of PCR-based DNA fingerprinting techniques for identification of *Listeria* species and their use for atypical *Listeria* isolates. *Int. J. Syst. Bacteriol.* **48**: 127-139.
- Vujanovic,V., Hamel,C., Jabaji-Hare,S. and St Arnaud,M. (2002) Development of a selective myclobutanil agar (MBA) medium for the isolation of *Fusarium* species from asparagus fields. *Can. J. Microbiol.* **48**: 841-847
- Walker,T.S., Bais,H.P., Deziel,E., Schweizer,H.P., Rahme,L.G., Fall,R., and Vivanco,J.M. (2004) *Pseudomonas aeruginosa*-plant root interactions. Pathogenicity, biofilm formation, and root exudation. *Plant Physiol.* **134**: 320-331.
- Walsh,G.A., Murphy,R.A., Killeen,G.F., Headon,D.R., and Power,R.F. (1995) Technical note: detection and quantification of supplemental fungal β -glucanase activity in animal feed. *J. Anim. Sci.* **73**: 1074-1076.
- Weisburg,W.G., Barns,S.M., Pelletier,D.A., and Lane,D.J. (1991) 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* **173**: 697-703.
- Weller,D. M. and Thomashow,L.S. (1994) Current challenges in introducing beneficial microorganisms into the rhizosphere., P. 1-17. In: F. O' Gara, D. N. Dowling, and B. Boesten (eds.), *Molecular ecology of rhizosphere organisms*. VCH verlagsgesellschaft mbH., Weinheim, Germany.
- Weller,D.M., Landa,B.B., Mavrodi,O.V., Schroeder,K.L., De La Fuente,L., Bankhead,S.B. et al. (2007) Role of 2,4-diacetylphloroglucinol-producing fluorescent *Pseudomonas spp.* in the defense of plant roots. *Plant Biol.* **9**: 4-20.
- Weller,D.M., Raaijmakers,J.M., Gardener,B.B., and Thomashow,L.S. (2002) Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annu. Rev. Phytopathol.* **40**: 309-348.
- Williams,J.G., Kubelik,A.R., Livak,K.J., Rafalski,J.A., and Tingey,S.V. (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *NAR* **18**: 6531-6535.

- Wilson,M. and Lindow,S.E. (1993) Interactions Between the Biological-Control Agent *Pseudomonas fluorescens* A506 and *Erwinia-Amylovora* in Pear Blossoms. *Phytopathol.* **83**: 117-123.
- Woo,S.L., Zoina,A., Del Sorbo,G., Lorito,M., Nanni,B., Scala,F. and Noviello,C. (1996) Characterization of *Fusarium oxysporum* f. sp. *phaseoli* by pathogenic races, VCGs, RFLPs, and RAPD. *Phytopathol.* **86**: 966-973.
- Working Group "Effect of pesticides from market gardening on water quality", (1988) Invloed van tuinbouwactiviteiten op de waterkwaliteit in de polder Nieuwland en Noordland (Westland). Defland Water Board, Delft.
- Yoo,S.J., Kim,H.G. and Yu,S.H. (1995) [Vegetative compatibility grouping of *Fusarium oxysporum* f.sp. *lycopersici* isolated from Korea.] *Korean J.Plant Pathol.* **11**: 330-337 (in Korean).
- Yoon,B.J., Lee,D.H., Kang,Y.S., Oh,D.C., Kim,S.I., Oh,K.H., and Kahng,H.Y. (2002) Evaluation of carbazole degradation by *Pseudomonas rhodesiae* strain KK1 isolated from soil contaminated with coal tar. *J. Basic Microbiol.* **42**: 434-443.
- Zeng,Q.Y., Wang,X.R. and Blomquist,G. (2003) Development of mitochondrial ssu rDNA-based oligonucleotide probes for specific detection of common airborne fungi. *Molecular and Cellular Probes* **17**: 281-288.
- Zhang,Z.G., Zhang,J.Y., Wang,Y.C. and Zheng,X. B. (2005) Molecular detection of *Fusarium oxysporum* f. sp. *niveum* and *Mycosphaerella melonis* in infected plant tissues and soil *FEMS Microbiol. Lett.* **251**: 357.