



Universiteit
Leiden
The Netherlands

Late-type Giants in the Inner Galaxy

Messineo, M.

Citation

Messineo, M. (2004, June 30). *Late-type Giants in the Inner Galaxy*. Retrieved from <https://hdl.handle.net/1887/512>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/512>

Note: To cite this publication please use the final published version (if applicable).

Late-type Giants in the Inner Galaxy

PROEFSCHRIFT

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van de Rector Magnificus Dr. D.D. Breimer,
hoogleraar in de faculteit der Wiskunde en
Natuurwetenschappen en die der Geneeskunde,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 30 juni 2004
klokke 16.15 uur

door

Maria Messineo
geboren te Petralia Soprana (Italië)
in 1970

Promotiecomissie

Promotor : Prof. dr. H. J. Habing

Referent : Dr. J. Lub

Overige leden : Prof. dr. W. B. Burton

Dr. M.R. Cioni (European Southern Observatory, Garching bei München)

Prof. dr. K. Kuijken

Prof. dr. K. M. Menten (Max-Planck-Institut fuer Radioastronomie, Bonn)

Prof. dr. A. Omont (Institut d'Astrophysique de Paris)

Prof. dr. P. T. de Zeeuw

A Lucia e Leonardo

Front cover: “I Girasoli” by Renato Guttuso. Permission to reproduce this painting was kindly granted by Dr. Fabio Carapezza Guttuso and authorised by SIAE 2004 (Italian Society of Authors and Publishers).

Back cover: ISOGAL image taken at $7\ \mu\text{m}$ (LW5 filter) of a field of $20' \times 20'$ centred on $(l, b) = -0.27^\circ, -0.6^\circ$, field FC-00027-00006.

Contents

1	Introduction	1
1.1	Late-type giants	2
1.1.1	AGB star and variability	4
1.1.2	Circumstellar maser emission	5
1.2	The Milky Way galaxy	6
1.2.1	Stellar line of sight velocity surveys and the importance of maser surveys	9
1.3	Outline of this thesis	10
	Bibliography	12
2	86 GHz SiO maser survey of late-type stars in the Inner Galaxy I.	
	Observational data	13
2.1	Introduction	13
2.2	Source selection	15
2.2.1	ISOGAL selection criteria	17
2.2.2	MSX selection criteria	18
2.3	Observations and data reduction	19
2.3.1	Flux stability	20
2.3.2	Detection criteria	21
2.3.3	Confusion with interstellar H ¹³ CN emission	22
2.3.4	Confusion with interstellar SiO emission	23
2.4	Results	25
2.4.1	Detection rate	25
2.4.2	Variability	29
2.4.3	Line intensity	31
2.4.4	Longitude-velocity diagram	31
2.4.5	Comparison with previous detections	34
2.4.6	Comments on individual objects	36
2.5	Conclusions	38
	Bibliography	41
3	86 GHz SiO maser survey of late-type stars in the Inner Galaxy II.	
	Infrared photometry	43

Contents

3.1	Introduction	43
3.2	Identification of the SiO targets in various infrared catalogues . . .	45
3.2.1	ISO GAL data	46
3.2.2	MSX data	46
3.2.3	ISO GAL-MSX cross-identifications	47
3.2.4	DENIS data	48
3.2.5	2MASS data	51
3.2.6	SiO targets and IRAS identifications	53
3.2.7	SiO targets and visual identifications	53
3.2.8	The table	55
3.3	A comparison between the ISO GAL and MSX samples	55
3.4	Some other remarks on the SiO targets	59
3.4.1	SIMBAD search	59
3.4.2	Is the targeted star the actual SiO emitter?	60
3.5	Variability	60
3.5.1	Variability information from DENIS and 2MASS data	62
3.5.2	Variability flag	63
3.6	The distribution of the SiO targets in the IRAS two-colour diagram	63
3.7	MSX colour-colour diagrams	66
3.8	Conclusion	68
	Bibliography	71
4	86 GHz SiO maser survey of late-type stars in the Inner Galaxy III.	
	Interstellar extinction and colours	73
4.1	Interstellar extinction law	74
4.1.1	Near-infrared interstellar extinction	75
4.1.2	Mid-infrared interstellar extinction	76
4.2	Interstellar extinction of field stars from near-infrared colour-magnitude diagrams	80
4.2.1	Reference red giant branch	85
4.2.2	Determination of extinction value and extinction law in the J, H, K_s CMD	86
4.2.3	Outside the Bulge	89
4.2.4	Dispersion of the extinctions along a line of sight	90
4.3	Near-infrared properties of known Mira stars	90
4.3.1	Colour-magnitude diagram of outer Bulge Mira stars and surrounding field stars	91
4.3.2	Colour-colour diagram of Mira stars	92
4.4	Interstellar extinction and intrinsic colours of the SiO targets	94
4.4.1	“Foreground objects”	97
4.5	Intrinsic colours and mass-loss rates	97
4.6	Conclusion	99
A	SAAO and 2MASS colours and magnitudes	100
A	IRAS and MSX filters	101
	Bibliography	102

5	86 GHz SiO maser survey of late-type stars in the Inner Galaxy IV.	
	Bolometric magnitudes	105
5.1	Introduction	105
5.2	Apparent bolometric magnitudes	107
5.2.1	Variability	109
5.3	SiO targets with $ l < 5^\circ$	110
5.3.1	Luminosities	110
5.3.2	Initial masses and ages	111
5.3.3	Red Supergiant stars ?	112
5.4	Comparison with OH/IR stars	113
5.5	Period–Luminosity relation	115
5.6	Stars in the the Nuclear Disk and a fourth dimension: extinction	117
5.7	Bolometric magnitudes and the $l - v$ diagram	119
5.8	Conclusion	121
A	Bolometric corrections	122
A.1	Bolometric corrections of monitored LPV stars	123
A.2	Bolometric corrections for stars in the inner Galaxy	126
	Bibliography	126
6	Considerations on the dynamics of maser stars in our Galaxy.	129
6.1	Introduction	129
6.2	Available evidence for a Galactic bar	130
6.2.1	Asymmetry in the longitude distribution of maser stars	130
6.2.2	Longitude–velocity diagram	131
6.2.3	Nuclear Disk	133
6.3	A simple dynamical model	134
6.3.1	Geometry	135
6.3.2	Equations of motion	135
6.3.3	Gravitational potential	136
6.3.4	First results	136
6.4	Summary and future plans	136
	Bibliography	137
7	The ISOGAL survey and the completeness analysis	139
7.1	Introduction	139
7.2	Observations	139
7.3	Data processing and analysis	141
7.4	Artificial sources	141
7.5	Concluding remarks	145
	Bibliography	146
	Nederlandse samenvatting	147
	Curriculum Vitae	153
	Nawoord	155

Contents