



Universiteit
Leiden
The Netherlands

Respiratory tract infection: prevention, early detection and attenuation of immune response

Groeneveld, G.H.

Citation

Groeneveld, G. H. (2020, March 11). *Respiratory tract infection: prevention, early detection and attenuation of immune response*. Retrieved from <https://hdl.handle.net/1887/86287>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/86287>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/86287> holds various files of this Leiden University dissertation.

Author: Groeneveld, G.H.

Title: Respiratory tract infection: prevention, early detection and attenuation of immune response

Issue Date: 2020-03-11



4

Clinical factors, C-reactive protein point of care test and chest X-ray in patients with pneumonia: a survey in primary care.

Geert H. Groeneveld,
Robert J. van de Peppel,
Margot W.M. de Waal,
Theo J.M. Verheij,
Jaap T. van Dissel.

Eur J Gen Pract. 2019 Oct;25(4):229-235

ABSTRACT

Background

In patients with an acute lower respiratory tract infection, the decision to prescribe antibiotics is sometimes difficult. C-reactive protein point of care test and chest X-ray are available as additional diagnostic tests, but the usefulness in clinical practice is unknown. To assess the proportion of Dutch general practitioners that use additional diagnostics in patients with an acute lower respiratory tract infection and whether clinical factors and C-reactive protein point of care test affect the behaviour in requesting chest X-rays.

Methods

In 2014, a questionnaire was sent to a random sample of 900 Dutch general practitioners. Outcome parameters are the use of C-reactive protein and chest X-ray, the percentage of GPs who guide their decision in requesting chest X-rays by CRP testing and the expectation regarding presence or absence of pneumonia. In addition, distribution of considerations for requesting chest X-rays were assessed.

Results

Two hundred fifty-five completed questionnaires (29%) were returned. More than half (54%) use the C-reactive protein test, these GPs tend to use less chest X-rays ($p=0.07$). GPs overestimate the chance that pneumonia would be present on the radiograph and 70% consider the detection or exclusion of abnormalities other than pneumonia as the main reasons for requesting a chest X-ray.

Conclusions

GPs report that CRP results affect their behaviour regarding the request of a chest x ray in patients with lower respiratory tract infection and therefore research is needed to substantiate the use of these diagnostic tools for this purpose.

INTRODUCTION

In patients that present with an acute lower respiratory tract infection, the decision whether or not to prescribe antibiotics is sometimes difficult, especially in moderately ill patients [1, 2]. Antibiotics are used more restrictively by Dutch general practitioners (GP) than by their colleagues in other European countries [3]. Nevertheless, there are also large regional differences within the Netherlands [4]. These differences are an expression of the complexity of the consideration of whether or not to prescribe an antibiotic. In general, one can state that patients with acute bronchitis do not need antimicrobial treatment while patients with pneumonia do [5,6]. Unfortunately, for the diagnosis of pneumonia, the use of anamnesis and physical examination alone provide insufficient support [7-9].

Two types of additional (diagnostic) tests for acute lower respiratory tract infection can be used in general practice: the C-reactive protein point of care test (CRP POCT) and the chest X-ray. A low CRP (< 20 mg/l) can exclude pneumonia with reasonable certainty, irrespective of clinical findings, while an elevated CRP (> 100 mg/l) greatly increases the chance of pneumonia warranting antibiotic treatment [8,10]. A recent meta-analysis ascertained that even when clinical variables are taken into account, the CRP test can help to confirm or exclude pneumonia [11]. Different guidelines (e.g. the British and the Dutch guideline) therefore, indicated the use of the CRP test in moderately ill patients [1,12]. Studies that evaluated whether the CRP POCT reduced the number of antibiotic prescriptions showed variable results [13,14].

A chest X-ray can be used to detect pneumonia, but the use of this examination in all individuals in whom a pneumonia is suspected, is not recommended. The chest X-ray is currently only recommended in the Dutch guideline to investigate the cause of lack of recovery, uncertainty about the diagnosis or treatment, or when a condition other than pneumonia is suspected as an explanation for the symptoms [1]. The British guideline does not mention chest X-ray as a diagnostic tool in patients with suspected pneumonia or exacerbations of asthma and COPD. Every year GPs request about 31 chest radiographs per 1000 person-years [15]. Research into the effectiveness of requesting chest X-rays by the GP for certain subgroups of patients with an acute lower respiratory tract infection is lacking. The objective of this study was to assess the use of chest X-ray and the CRP POCT in patients with an acute respiratory tract infection in Dutch primary care. We asked the GPs about their estimates and experiences with this complex situation where evidence for a specific strategy is lacking.

METHODS

Study design and setting

Between May and September 2014 a questionnaire-based cross-sectional study was performed in the Netherlands. The registry from the *Netherlands Institute for Healthcare Research* (NIVEL) contains address information of all GPs in the Netherlands. A random sample of 900 addresses was drawn. The questionnaire (see below) was sent in May 2014 by mail to these family practice addresses.

Construction of the questionnaire

The two main investigators (GHG and RJP) held an exploratory focus group discussion with various GPs in the Leiden region, the Netherlands. In this discussion, open questions were asked about the way in which the GPs use additional diagnostic tests in patients with acute lower respiratory tract infection and in what way the results of the tests affect their treatment policy [16]. An acute lower respiratory tract infection was defined as complaints for less than three weeks.

With the results, a list with open and closed questions was generated and distributed among 15 GPs in the Leiden region via the newsletter of the Leiden Primary Care Research Network. The answers and feedback received via this route contributed to the final quantitative questionnaire.

Quantitative questionnaire

The questionnaire first asks about the number of years of work experience, the number of hours a week that the GP works at the general practice, and an estimate of the number of chest X-ray request in a year for patients with acute lower respiratory tract infection.

Main outcomes are the use of CRP POCT, the percentage of GPs who guide their decision in requesting chest X-rays by CRP testing and the expectation regarding presence of pneumonia on chest X-ray. In addition, indications for use of CRP POCT, clinical parameters and distribution of reasons for requesting chest X-rays (in GPs with and without CRP test available), which other pathology the GP wants to exclude and diagnostic and therapeutic consequences when pneumonia is present or absent were assessed.

The various characteristics and consequences could be scored on five-point Likert scales, with answers varying from “(almost) never”/“Very unimportant” to “(almost) always”/“Very important”. The complete questionnaire is available in the Supplementary Material.

Analysis

The returned questionnaires were anonymized. Descriptive analyses and comparison of proportion with Chi Square test were performed with SPSS (IBM, version 23).

RESULTS

Study population

Twenty-three questionnaires were returned due to outdated address details. In total, after one reminder letter, 255 of the 877 (29%) questionnaires were returned completed in September 2014. The respondents reported a median work experience of 14 years, (interquartile range, IQR, 9 - 22 years) and a median work week of 36 hours (IQR 30 - 41.5 hours) at the general practice.

Chest X-ray

Median reported number of chest X-rays per year for patients with an acute lower respiratory tract infection was 10 (IQR 4-12). The 24 respondents (9%) that never requested a chest X-ray for this indication, could not answer the remaining questions. Median work experience and work week in the respondents who never request a chest X-ray did not differ from respondents who did request chest X-rays.

Table 1 and 2 provide an overview of the reports of GPs regarding considerations and objectives to request a chest X-ray. The majority (70% of all GPs) consider the detection or exclusion of abnormalities other than pneumonia as one of the main reasons for requesting a chest X-ray. The exclusion of malignancy, heart failure, sarcoidosis, and tuberculosis are mentioned repeatedly. If the chest X-ray has been requested to exclude other pathology, the GP will state this in 90% of the cases on the X-ray application form. Factors that play an important role in the decision to request a chest X-ray are mainly age, smoking, and the duration of the complaints.

The expectation of 217 GPs (14 GPs did not answer this question and 24 never request a chest X-ray) to detect a lung infiltrate on the chest X-ray was less than 10% in 13% of GPs, between 10 and 20% in 19% of GPs, and more than 50% in 68% of GPs. If an infiltrate suspect for pneumonia is present, 227 of the 230 GPs (99%; 1 GP did not answer this question and 24 GPs never request a chest X-ray) often, to almost always, prescribe an antibiotic. In the absence of a pneumonia, 4% of GPs often to almost always, prescribe an antibiotic (Figure 1).

Table 1. Questionnaire response from general practitioners: Clinical factors in the consideration to request a chest X-ray in patients with an acute lower respiratory tract infection (n=226*).

| Clinical factors in the consideration to request a chest X-ray | Rating | |
|--|---------------|----------------------------|
| | Important (%) | Neutral or unimportant (%) |
| Smoking | 191 (85) | 35 (15) |
| Duration of the complaints | 186 (82) | 40 (18) |
| Age | 179 (79) | 47 (21) |
| Presence of fever | 98 (43) | 128 (57) |
| Duration of fever | 95 (42) | 131 (58) |
| Response to previous antibiotics | 92 (41) | 134 (59) |
| Producing sputum, and sputum color | 28 (12) | 198 (88) |

*29 respondents never requested chest X-rays and/or did not give an answer to this question.

Table 2. Questionnaire response from general practitioners: Reasons to request a chest X-ray in patients with an acute lower respiratory tract infection (n=228[†]).

| Reasons to request a chest X-ray | Number of times indicated to be the most important (% ^{**}) |
|---|---|
| Detection or exclusion of other lung abnormalities, such as a lung tumor | 159 (69.7) |
| Confirm the diagnosis of pneumonia | 87 (38.2) |
| Exclude the diagnosis of pneumonia | 76 (33.3) |
| Reassuring the patient | 22 (9.6) |
| Uncertainty about further policy | 21 (9.2) |
| As a guide to decide on antibiotic prescription | 18 (7.9) |
| Conditions that GPs want to exclude | Number of times indicated (%), N=190^{***} |
| Lung cancer | 160 (84.2) |
| Heart failure | 46 (24.2) |
| Sarcoidosis | 36 (18.9) |
| Tuberculosis | 24 (12.6) |
| Pneumothorax | 15 (6.9) |
| Other^{****} | 48 (25.2) |

[†]27 respondents never requested chest X-rays and/or did not give an answer to this question.

^{**}Percentages add up to >100% because some respondents gave more than one reason the same score.

^{***} some GPs who did not state the exclusion of other lung abnormalities as the most important reason also answered this question; in addition, several answers could be filled in.

^{****} other disorders included foreign body, pulmonary embolism, and systemic lupus erythematosus and were each mentioned by <5% of all respondents.




Figure 1. Questionnaire response from general practitioners: Policy following the chest X-ray in patients with an acute lower respiratory tract infection (n=230*).

Bi-directional bar chart. On the left the policy followed in case a pneumonia was detected on the chest X-ray, on the right the policy followed in case no pneumonia was detected on the chest X-ray. In the middle, description of the policy.

* 24 respondents never request a chest X-ray and one did not answer this question.

CRP point of care test

The CRP POCT is used by more than half of GPs (54%). A large proportion of them, also use the test to evaluate suspected infections other than pneumonia (Table 3), e.g. diverticulitis, urinary tract infection, or an unknown “other” infection. Eighty percent of all GPs reported that they foresee that CRP POCT can replace chest X-ray as a diagnostic test partially or completely. GPs with CRP test available are more confident than those that do not have this test available (86% versus 71%, p<0.01).

Table 3. Questionnaire response from general practitioners: Use and indications for use of the CRP point of care test (n=246*).

| | Number (%) |
|--|-----------------|
| Respondents that use the CRP point of care test in the general practice | 134 (54) |
| Use only if there is a suspicion of respiratory tract infection | 35 (26) |
| Use in case of suspected respiratory tract or other infection | 83 (62) |
| Hardly ever use the test | 16 (12) |
| In many cases, the CRP point of care test plays a role in the consideration to request a chest X-ray** | 75 (56) |
| Respondents that do not use the CRP point of care test in the general practice | 112 (46) |
| Would like to purchase the test in the future | 85 (76) |
| Would not like to purchase the test in the future | 27 (24) |

* Nine GPs did not answer this question

** Respondents that indicated that this “often” or “(almost) always” plays a role.

Difference between GPs with and without CRP test

GPs with CRP POCT available reported to request less chest X-rays than their colleagues without CRP POCT available (median 6, IQR 3-10 versus median 10, IQR 5-14 respectively; $p=0.07$).

Expectation regarding presence of pneumonia did not differ between GPs with or without CRP POCT available ($p=0.67$).

Presence and colour of sputum was reported to be more important when considering chest X-ray by GPs without than those with CRP POCT available (Figure S1 in the supplementary material).

Guidance whether or not to prescribe antibiotics is reported as reason for requesting chest X-ray less frequently in GPs with CRP than in GPs without CRP. Other reasons were not different (see Figure S2 in the supplementary material).

GPs who do not use CRP POCT reported more frequently than those who do use CRP POCT to start symptom management in case pneumonia is confirmed (neutral to almost always 15% versus 9%; $p=0.05$) or ruled out with chest X-ray (neutral to almost always 57% versus 41%; $p<0.01$). All other policy items did not differ significantly between GP groups.

DISCUSSION

Main findings

This study shows that in 255 Dutch GPs the use of additional diagnostic tools for the suspicion of acute lower respiratory tract infection was diverse. GPs reported to estimate the probability of having a pneumonia as high among the patients for whom they request a chest X-ray. Nearly 70% of GPs request the photo mainly to exclude other pathology. More than half of the GPs had the CRP POCT available in 2014 and the majority used this test to determine whether or not to request a chest X-ray. GPs using CRP POCTs reported to request less chest X-rays than GPs who did not use this test. These latter GPs reported to use chest X-ray more often to guide the decision to prescribe antibiotics. Many GPs also used the CRP POCT for other purposes.

Strengths and limitations

The strengths of this study are the random sample of GPs in the Netherlands and the considerable number of 255 completed surveys that were available for analysis. The inventory based on focus group interview and pilot questionnaires during the pilot study means that

the diversity of ideas, experiences, and behaviors in the target group were well explored. The fact that both GPs with and without a CRP POCT, as well as GPs that vary from never to frequently requesting chest X-rays have responded, makes that the sample has, in any case, included all extremes of diagnostic policy.

A limitation of the study is the potential occurrence of sampling bias. The 'selection' of respondents could be different from that of the GPs who did not respond. Although the absolute number of questionnaires analyzed is considerable, the response rate of 29% is not high. A review by Creavin *et al.* showed a mean response rate of 61% [17]. However, response rate in recent surveys among Dutch GPs is substantially lower [17-19]. Respondents could be more interested in this topic than non-responders and thereby more aware of guidelines and evidence, resulting in more prudent use of diagnostic tools. The years of work experience and the number of working hours of the respondents correspond to the national average, 14.9 years and 31.2 hours per week respectively [20]. Moreover, McFarlane *et al.* demonstrated that higher response rates in a survey of physicians are not associated with lower selection bias [21].

Nonetheless, potential difference in characteristics between GPs who filled in the questionnaire and the ones that did not respond, might still be present. However, the study provides a useful insight into the considerations of the Dutch GP about additional diagnostic tools for acute lower respiratory tract infections.

The short questionnaire brings about that not every possible consideration has been asked. For example, it is not clear in what type of patient the CRP POCT is actually used, if CRP kinetics are taken into consideration and how GPs interpret the results. A previous study showed that most GPs use the CRP POCT in patients who are moderately ill when it is not immediately obvious whether or not the patient needs an antibiotic. In the same study, it was found that the CRP POCT is sometimes used too frequently, even in situations where this test should have no consequences for the policy [22].

This is a survey-based study about opinions and perceptions, which do not necessarily reflect the real management and prescription habits. The survey was completed in 2014. It is possible that with an increase in use, the interpretation of the results will also change slightly.

Interpretation

The expectation about the likelihood to detect a lung infiltrate on the X-ray is high. Two-thirds expect an infiltrate in more than 20% of patients. This estimate does not match

the findings in several primary care studies, where a pneumonia on the chest X-ray was detected in only 5 to 13% [8,11,23].

The chest X-ray is the gold standard for the detection or exclusion of pneumonia, while clinical features, including a low CRP value, can safely exclude pneumonia [11,12]. The added value of the chest X-ray in the detection or exclusion of pneumonia is therefore mainly present in the group of patients with a high probability of the presence of an infiltrate. This mainly concerns patients with clinical characteristics fitting with pneumonia that have a high CRP value. We hypothesize that GPs may request too much chest X-rays because they overestimate the likelihood of pneumonia. With better pre-test (pre-chest X-ray) assessment, for example by using CRP, they could rule out pneumonia more often without chest X-ray. On the other hand, GPs incorrectly withhold some patients from a chest X-ray because they do not adequately determine the group of patients with a high pre-test (pre-chest X-ray) probability, partially because only 54% in our study used CRP test. In addition, given the discrepancy between the pre-test assessment and the actual percentage of pneumonia present on lung images, pneumonia can often be excluded with a chest X-ray. In the latter case, antibiotics are prescribed less frequently.

The lack of evidence is the reason that the chest X-ray is currently not clearly defined in the standard of the Dutch Society of GPs or the British guidelines for the detection or exclusion of pneumonia [1,12]. However, this study shows that GPs already use the results of the CRP test in their decision to request a chest X-ray and/or that they foresee that the CRP test can replace the chest X-ray as a diagnostic tool.

Often the detection or exclusion of a condition other than pneumonia is indicated as the main reason to request a chest X-ray. In a European cohort of nearly 3,000 patients with acute cough who underwent a chest X-ray, a clinically relevant abnormality -other than pneumonia- was found in 3% [24]. Therefore, the chance that a GP will find such aberrations is small. A malignancy can be missed on the chest X-ray, especially if at that time an infiltrate is present in the same area. It is then preferable to repeat the chest X-ray after the pneumonia has been treated [25].

Exact information about availability and use of CRP POCT in European countries is not known. Opong et al. reported that CRP POCT was available in 12 of 14 primary care networks in 13 European countries [26]. There were marked differences in the availability of CRP test between Spain and Denmark [27] and between CRP use in Belgium (3%), the UK (15%) and the Netherlands (48%) in 2012-2013 [28]. The use of CRP has increased in Scandinavian countries between 2004 and 2013 [29].

When comparing Danish primary care versus Spanish primary care, chest X-rays are used more frequently to confirm pneumonia in Spain [27].

Implications

With the frequent use of the CRP POCT to aid in the decision to request a chest radiograph, there appears to be a need for research into a diagnostic algorithm, that would incorporate clinical characteristics and a CRP result, to determine in which patient a chest X-ray has added value.

This study also shows that GPs using the CRP POCT often use this test for other infections than pneumonia. The use of the CRP test is only recommended for patients with acute lower respiratory tract infections or diverticulitis. For both disorders, the use of the CRP test has many limitations [1,30]. Restraint in the use of this test is therefore required until new research proves that either the CRP POCT has added value for other indications, or that the CRP test can replace a chest radiograph.

CONCLUSION

GPs widely use the CRP POCT and often base their decision to request a chest X-ray on the outcome. They overestimate the chance of finding a pneumonia in these patients. Clinical variables in combination with the CRP POCT, could help the GP to request chest radiographs more selectively for patients with acute lower respiratory tract infection. Research is however first needed to substantiate the use of these diagnostic tools for this purpose.

ACKNOWLEDGEMENTS

We thank the GPs from the Leiden region who contributed to the realization of the survey, the GPs who gave their feedback in the pilot study, and all GPs who completed and returned the final questionnaire.

REFERENCES

1. Verheij ThJM, Hopstaken RM, Prins JM, et al. NHG Standard Acute Cough (First review). *Huisarts Wet.* 2011;54:68-92.
2. Rosh AJ, Newman DH. Evidence-based emergency medicine/rational clinical examination abstract. Diagnosing pneumonia by medical history and physical examination. *Ann Emerg Med.* 2005;46:465-7.
3. Adriaenssens N, Coenen S, Versporten A, et al. European Surveillance of Antimicrobial Consumption (ESAC): outpatient antibiotic use in Europe (1997-2009). *J Antimicrob Chemother.* 2011;66 Suppl 6:vi3-12.
4. Volksgezondheidszorg.info. Users antibiotics per municipality [Internet]. Bilthoven: RIVM 2013 [updated 25 maart 2016 ; cited 2019 Mar 21] Available from: <https://www.volksgezondheidszorg.info/onderwerp/genees-en-hulpmiddelen-en-lichaamsmaterialen/regionaal-internationaal/geneesmiddelen#node-gebruikers-antibiotica-gemeente>.
5. Little P, Stuart B, Moore M, et al. Amoxicillin for acute lower-respiratory-tract infection in primary care when pneumonia is not suspected: a 12-country, randomised, placebo-controlled trial. *Lancet Infect Dis.* 2013;13:123-9.
6. Teepe J, Little P, Elshof N, et al. Amoxicillin for clinically unsuspected pneumonia in primary care: subgroup analysis. *Eur Respir J.* 2016;47:327-30.
7. Zaat JOM, Stalman WAB, Assendelft WJJ. Knock, knock, who is there? A systematic literature review of the value of anamnesis and physical examination for suspected pneumonia. *Huisarts Wet.* 1998;41:461-9.
8. Hopstaken RM, Muris JW, Knottnerus JA, et al. Contributions of symptoms, signs, erythrocyte sedimentation rate, and C-reactive protein to a diagnosis of pneumonia in acute lower respiratory tract infection. *Br J Gen Pract.* 2003;53:358-64.
9. Graffelman AW, le Cessie S, Knuistingh Neven A, et al. Can history and exam alone reliably predict pneumonia? *J Fam Pract.* 2007;56:465-70.
10. van Vugt SF, Broekhuizen BD, Lammens C, et al. Use of serum C reactive protein and procalcitonin concentrations in addition to symptoms and signs to predict pneumonia in patients presenting to primary care with acute cough: diagnostic study. *BMJ.* 2013;346:f2450.
11. Minnaard MC, de Groot JA, Hopstaken RM, et al. The added value of C-reactive protein measurement in diagnosing pneumonia in primary care: a meta-analysis of individual patient data. *CMAJ.* 2017;189:E56-E63
12. National Institute for Health and Excellence (NICE). Pneumonia in adults: diagnosis and management 2014 [updated December 2014; cited 2019 May 22]. Available from: <https://www.nice.org.uk/guidance/cg191/chapter/1-Recommendations>
13. Minnaard MC, van de Pol AC, Hopstaken RM, et al. C-reactive protein point-of-care testing and associated antibiotic prescribing. *Fam Pract.* 2016;33(4):408-13.
14. Huang Y, Chen R, Wu T, et al. Association between point-of-care CRP testing and antibiotic prescribing in respiratory tract infections: a systematic review and meta-analysis of primary care studies. *Br J Gen Pract.* 2013;63:e787-94.
15. Speets AM, Kalmijn S, Hoes AW, et al. Frequency of chest radiography and abdominal ultrasound in the Netherlands: 1999-2003. *Eur J Epidemiol.* 2005;20:1031-6.
16. Moser A, Korstjens I. Series: Practical guidance to qualitative research. Part 3: Sampling, data collection and analysis. *Eur J Gen Pract.* 2018;24:9-18.

17. Creavin ST, Creavin AL, Mallen CD. Do GPs respond to postal questionnaire surveys? A comprehensive review of primary care literature. *Fam Pract.* 2011;28:461-7.
18. Wammes JJ, Jeurissen PP, Verhoef LM, et al. Is the role as gatekeeper still feasible? A survey among Dutch general practitioners. *Fam Pract.* 2014;31:538-44.
19. Sinnema H, Franx G, Spijker J, et al. Delivering stepped care for depression in general practice: results of a survey amongst general practitioners in the Netherlands. *Eur J Gen Pract.* 2013;19:221-9.
20. Cijfers uit de registratie van huisartsen [Internet] Utrecht: Stichting NIVEL; 2015 [updated Februari 2015; cited 2019 Mar 21] Available from: <https://nvl004.nivel.nl/nivel-2015/sites/default/files/bestanden/Cijfers-uit-de-registratie-van-huisartsen-peiling-jan-2014.pdf>.
21. McFarlane E, Olmsted MG, Murphy J, et al. Nonresponse bias in a mail survey of physicians. *Eval Health Prof.* 2007;30:170-85.
22. Cals JW, Chappin FH, Hopstaken RM, et al. C-reactive protein point-of-care testing for lower respiratory tract infections: a qualitative evaluation of experiences by GPs. *Fam Pract.* 2010;27:212-8.
23. Speets AM, Hoes AW, van der Graaf Y, et al. Chest radiography and pneumonia in primary care: diagnostic yield and consequences for patient management. *Eur Respir J.* 2006;28:933-8.
24. van Vugt S, Broekhuizen L, Zuithoff N, et al. Incidental chest radiographic findings in adult patients with acute cough. *Ann Fam Med.* 2012;10:510-5.
25. Tang KL, Eurich DT, Minhas-Sandhu JK, et al. Incidence, correlates, and chest radiographic yield of new lung cancer diagnosis in 3398 patients with pneumonia. *Arch Intern Med.* 2011;171:1193-8.
26. Oppong R, Coast J, Hood K, et al. Resource use and costs of treating acute cough/lower respiratory tract infections in 13 European countries: results and challenges. *Eur J Health Econ.* 2011;12:319-29.
27. Christensen SF, Jorgensen LC, Cordoba G, et al. Marked differences in GPs' diagnosis of pneumonia between Denmark and Spain: a cross-sectional study. *Prim Care Respir J.* 2013;22:454-8.
28. Howick J, Cals JW, Jones C, et al. Current and future use of point-of-care tests in primary care: an international survey in Australia, Belgium, The Netherlands, the UK and the USA. *BMJ open.* 2014;4:e005611.
29. Haldrup S, Thomsen RW, Bro F, et al. Microbiological point of care testing before antibiotic prescribing in primary care: considerable variations between practices. *BMC Fam Pract.* 2017;18:9.
30. Berger MY, De Wit NJ, Vogelenzang R, et al. NHG-Standaard Diverticulitis [NHG standard diverticulitis] *Huisarts Wet.* 2011;54:492-9

SUPPLEMENTARY MATERIAL




Figure S1. Bidirectional bar chart: questionnaire response of general practitioners with and without CRP test available, regarding the considerations to request a chest X-ray in patients with an acute lower respiratory tract infection (n=226).

* 29 respondents never requested a chest X ray and/or did not give an answer to this question.




Figure S2. Bidirectional bar chart: questionnaire response of general practitioners with and without CRP test available, about reasons to request a chest X-ray in patients with an acute respiratory tract infection (n=228).

* 27 respondents never requested a chest X ray and/or did not give an answer to this question.

