



Universiteit
Leiden
The Netherlands

Immune parameters affecting maternal tolerance towards the fetus in normal and aberrant pregnancies

Craenmehr, M.H.C.

Citation

Craenmehr, M. H. C. (2020, June 16). *Immune parameters affecting maternal tolerance towards the fetus in normal and aberrant pregnancies*. Retrieved from <https://hdl.handle.net/1887/116771>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/116771>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/116771> holds various files of this Leiden University dissertation.

Author: Craenmehr, M.H.C.

Title: Immune parameters affecting maternal tolerance towards the fetus in normal and aberrant pregnancies

Issue Date: 2020-06-16



Craenmehr, M. H. C.; Nederlof, I.; Cao, M.; Drabbels, J. J. M.; Spruyt-Gerritse, M. J.; Anholts, J. D. H.; Kapsenberg, H. M.; Stegehuis, J. A.; van der Keur, C.; Fasse, E.; Haasnoot, G. W.; van der Hoorn, M. P.; Claas, F. H. J.; Heidt, S.; Eikmans, M., *Int J Mol Sci* 2019, 20, (3).

4 increased HLA-G expression in term placenta of women with a history of recurrent miscarriage despite their genetic predisposition to decreased HLA-G levels

Abstract

HLA-G is an immune modulating molecule present on fetal extravillous trophoblasts at the fetal-maternal interface. Single nucleotide polymorphisms (SNPs) in the 3 prime untranslated region (3'UTR) of the HLA-G gene can affect the level of HLA-G expression, which may be altered in women with recurrent miscarriages (RM). This case-control study included 23 women with a medical history of three or more consecutive miscarriages who delivered a child after uncomplicated pregnancy, and 46 controls with uncomplicated pregnancy. Genomic DNA was isolated to sequence the 3'UTR of HLA-G. Tissue from term placentas was processed to quantify HLA-G protein and mRNA levels. The women with a history of RM had a lower frequency of the HLA-G 3'UTR 14-bp del/del genotype compared to controls (OR 0.28; P = 0.039), which has previously been related to higher soluble HLA-G levels. Yet, HLA-G protein (OR 6.67; P = 0.006) and mRNA (OR 6.33; P = 0.010) expression was increased in term placentas of women with a history of RM compared to controls. In conclusion, during a successful pregnancy HLA-G expression is elevated in term placentas from women with a history of RM compared to controls, despite a genetic predisposition associated with decreased HLA-G levels. These findings suggest that HLA-G upregulation could be a compensatory mechanism in the occurrence of RM to achieve an ongoing pregnancy.

Introduction

About 1 to 2% of couples trying to conceive experience recurrent miscarriages (RM) [1]. Accepted etiological categories for RM include chromosomal abnormalities, uterine anatomic abnormalities and antiphospholipid antibody syndrome. However, a significant proportion of the couples trying to conceive do not know the underlying cause for this recurring problem [2], leaving them with a burden of uncertainty.

During pregnancy, the maternal immune system needs to accept the semi-allogeneic fetal tissue. For this reason, several mechanisms are at play at the fetal-maternal interface. The absence of the human leukocyte antigen (HLA) class I antigens A and B and HLA class II on fetal trophoblast cells helps to prevent allorecognition by T and B cells, whereas the presence of HLA-C, HLA-E, HLA-F and HLA-G provide self-signals to control (natural killer) NK responses [3, 4]. Low levels of HLA-G have been associated with RM [5, 6]. By alternative splicing, the HLA-G pre-mRNA can give rise to seven different isoforms, of which four are membrane-bound (HLA-G1, -G2, -G3 and -G4) and three are soluble (HLA-G5, -G6 and -G7) [7]. Whereas in healthy tissue membrane-bound HLA-G is only expressed on trophoblasts, the soluble form of HLA-G can be detected in various body fluids, such as amniotic fluid, blood and seminal plasma [8-10]. One mechanism leading to the generation of soluble (s)HLA-G1 is the cleavage of membrane-bound HLA-G from the cell surface by the activity of metalloproteinases [11].

Several polymorphisms are present in the 3 prime untranslated region (3'UTR) of the HLA-G gene. Since the 3'UTR is targeted by microRNAs (miRNA) that can negatively influence expression, polymorphisms in this region may have an influence on the efficiency of miRNA binding, and consequently on the level of HLA-G expression and on pregnancy outcome. The 14bp insertion/deletion polymorphism affects the stability of HLA-G mRNA and thereby the expression

of HLA-G [12]: the insertion is associated with low levels of sHLA-G [13]. Presence of the +3187A allele is associated with decreased mRNA stability and decreased HLA-G expression [14]. The presence of a guanine at position +3142 increases the affinity of miR-148a, miR-148b and miR-152, which leads to downregulation of HLA-G expression [15].

We analyzed the HLA-G 3'UTR genotype of women with a history of RM and of a control group of women with no history of RM. We also analyzed the HLA-G 3'UTR genotype of their offspring. The combination of multiple polymorphic sites was used to generate 3-UTR haplotypes. In addition, we studied HLA-G mRNA and protein expression levels in term placentas of women with successful pregnancies in both study groups.

Materials and Methods

Subjects and materials

This case control study included women with a medical history of RM who delivered a child after uncomplicated pregnancy. These women visited the Department of Obstetrics and Gynecology, Leiden University Medical Center (LUMC) between 2012 and 2015, and no underlying cause for RM was found after a full clinical workup according to the local guidelines, which are in line with the international ESHRE guideline. Twenty-three women with a history of at least three miscarriages and an uncomplicated singleton pregnancy were included in this study, of whom placental tissue was stored for research purposes. For the control group, 46 women were included with a history of ≤ 1 miscarriage, of whom placental tissue of a healthy singleton pregnancy was stored for research purposes after delivery at the Department of Obstetrics and Gynecology, LUMC.

For additional experiments we collected products of conception from eight first

trimester miscarriages (GA: 6-10 weeks) and four first trimester elective abortions (GA: 5-10 weeks). The miscarriage material was obtained from women with a history of RM from the Department of Obstetrics and Gynecology in the LUMC. Elective abortion material was received anonymously from an abortion clinic [16].

The protocol was approved by the Ethical committee of the LUMC (P11.196), and all participants gave informed consent for inclusion in the study.

HLA-G polymorphisms and haplotypes

Peripheral blood and umbilical cord blood for both groups was processed to genotype HLA-G in the mothers and children, respectively. Genomic DNA was isolated to sequence a 699/713-bp fragment covering the 3'UTR of exon 8, starting just before the 14-bp insertion/deletion and ending 591-bp downstream of the insertion/deletion. To sequence the haplotype on each of the two alleles, amplification reactions were performed using the generic 3'-primer that was tailed with a M13 sequence to cover the 3'UTR region of HLA-G. The following polymorphisms were identified: the 14-bp insertion/deletion (rs371194629), +3003C/T (rs1707), +3010C/G (rs1710), +3027A/C (rs17179101), +3035C/T (rs17179108), +3142C/G (rs1063320), +3187A/G (rs9380142), +3196C/G (rs1610696), +3422C/T (rs17875408), +3496A/G (rs1233330), and +3509G/T (rs1611139).

UTR haplotypes were composed based on the combination of SNPs. Conversion of sequencing data to UTR haplotypes was carried out by using specialized HLA interpretation software (SBT Engine, GenDX, Utrecht, the Netherlands). The forward primer (GTGATGGGCTGTTAAAGTGTCACC), the reverse primer (GACGTTGTAAAACGACGGCCAGTAGGGGAAGAGGTGTAGGGGTCTG) and an M13 universal primer (GACGTTGTAAAACGACGGCCAGT) were ordered from

Sigma (St. Louis, Missouri USA). The underlining represents the M13 sequence.

Immunohistochemistry

HLA-G and trophoblasts were detected by standard immunohistochemical procedures. After delivery, placental tissues were dissected and fixed in 4% neutral buffered formaldehyde and embedded in paraffin. Section slides of 6 µm were cut, mounted on Superfrost/Plus glass slides (Thermo Scientific, Waltham, Massachusetts, USA), and dried overnight at 37°C. Sections were deparaffinized in xylene and ethanol. Depending on the primary antibody used, unmasking of the antigens was achieved by enzyme digestion with trypsin or incubation with citrate buffer in a microwave. This was followed by endogenous peroxidase blocking in 3% H₂O₂ in methanol for enzymatic staining. All incubations were at room temperature and wash steps in between the incubations were performed in PBS. Slides were pre-incubated with PBS/1% BSA to reduce background staining. Excess buffer was removed and slides were incubated with mouse monoclonal primary antibodies overnight at room temperature. Antibodies were diluted in PBS containing 1% BSA.

For enzymatic and immunofluorescence staining, primary antibodies against the free heavy chain of all HLA-G isoforms (MEM-G2; EXBIO Praha, Czech Republic) and against cytokeratin 8 (CAM5.2; Becton Dickinson, Franklin Lakes, New Jersey, USA) were used. The next day, incubation with secondary antibody (EnVision solution, goat anti-mouse HRP, undiluted; DAKO, Agilent, Santa Clara, California, USA) for enzymatic staining; Goat-anti-mouse IgG1-AF488 A21121 and Goat-anti-mouse IgG2a-AF546 A21133 for immunofluorescence staining; Thermo Scientific) was performed, and substrate was visualized with diaminobenzidine (DAB metal Enhanced substrate kit; 34065; Thermo Scientific) for enzymatic staining. Specimens were counterstained with hematoxylin and mounted in

Micromount Mounting Medium (Leica, Nussloch, Germany) for enzymatic staining and ProLong Gold Antifade Mountant with DAPI (P36931; Thermo Scientific) for immunofluorescence staining.

Quantification of immunohistochemical stainings

We set out to compare the extent of MEMG2 and CAM5.2 staining in term placentas between the study groups. All slides were scanned by a Panoramic Midi scanner (3DHISTECH, Budapest, Hungary). The entire decidua basalis was quantitatively analyzed using the HistoQuant modus in Quant Center software (3DHISTECH). This was done by two investigators (JS and HK) independently for 10 placentas to analyze interobserver variability. For each staining, the same thresholds and training scenarios were used for patient and control slides. We corrected for the selected surface area when calculating the percentage positivity of a staining.

For first trimester material, we could not define the decidua. Therefore, we analyzed only the HLA-G positive parts of the slides. Scoring of the slides was performed by two investigators (JS and ME) independently, blinded for the cause of the abortion. Based on the extent of staining, cases were classified according to a semi-quantitative scoring system, i.e., (1) minimal, (2) moderate, or (3) intense staining. Examples of stainings are shown in Supplementary Fig. S1.

RNA isolation and qPCR

Tissue homogenates from term placentas were processed for mRNA quantification of HLA-G by real-time qPCR. Tissue sections were immersed in ML lysis buffer (Nucleospin miRNA isolation kit from Macherey-Nagel, Düren, Germany) and stored at -20°C until isolation. RNA was extracted using NucleoSpin columns (Macherey-Nagel) and tested for integrity by gel electrophoresis (Experion, Bio-

Rad). RNA quantity was determined on a NanoDrop 2000 Spectrophotometer (Thermo Scientific). RNA was combined with oligo DT (Promega; 0.5 µg), dNTP (Promega; 10 mM), and random nucleotide hexamers (0.5 µg; Promega, Fitchburg, Wisconsin, USA). This mixture was incubated at 65°C for 5 minutes and then put on ice. Complementary DNA synthesis from mRNA was carried out using Superscript III (40 µg/µL RNaseOUT, SuperScriptIII 200 µg/µL, 0.1M DTT; Promega). The reactions were proceeded at 25°C for 5 minutes and 50°C for 60 minutes. Reactions were terminated by increasing the temperature to 70°C for 5 minutes.

PCR assays were carried out using iQ™ SYBR® Green Supermix (Bio-Rad) on a Viia7 Real-time PCR system (Applied Biosystems, Foster City, California, USA). The PCR program consisted of 10 minutes at 95 °C, followed by 40 cycles of 15 seconds at 95 °C and 1 minute at 60 °C. Levels of mRNA transcripts for HLA-G were normalized to the geometric mean signal of reference genes GAPDH and β-actin. The forward (ACCCACTCCTCCACCTTTGAC) and reverse (TCCACCACCCTGTTGCTGTAG) primer for GAPDH; the forward (ACCACACCTTCTACAATGAG) and reverse (TAGCACAGCCTGGATAGC) primer for beta-actin; the forward (GACAGCGACTCGGCGT) and reverse (GTGTTCCGTGTCTCCTCT) primer for HLA-G were ordered from Sigma.

We also studied miRNA levels in the tissue homogenates. For this, RNA template was reverse transcribed into cDNA using the miRCURY LNA™ Universal RT miR PCR kit (Exiqon, Vedbaek, Denmark). LNA™ enhanced primer sets were used targeting the following miRNAs of interest: hsa-miR-148a (MIMAT0000243), hsa-miR-148b (MIMAT0000759), hsa-miR-152 (MIMAT0000438), and hsa-miR-365 (MIMAT0000710). Levels of these miRNAs were normalized to the geometric mean signal of previously described reference genes hsa-miR-16 (MIMAT0000069) and hsa-miR-103 (MIMAT0000101) [17, 18].

All PCR reactions were performed in duplicate. Signals were normalized using the $\Delta\Delta Cq$ method. Quantitative PCR measurements were analyzed using QuantStudio Real-Time PCR System Software (Applied Biosystems). To verify the accuracy of amplification, melting curve analyses were performed at the end of each PCR run.

Statistical analysis

Spearman's correlation analysis and Bland-Altman plotting were performed for the assessments of validity and reproducibility [19]. Differences between groups were tested by Mann-Whitney U tests, chi-square tests or logistic regression analysis. Values of $P < 0.05$ were considered to indicate statistical significance. Association between HLA-G SNPs and RM was studied with binary logistic regression. Per HLA-G genotype the highest prevalence was defined as the reference group. Alleles with a frequency of $<5\%$ were excluded from analysis. For the calculations on the HLA-G genotypes Bonferroni adjustment was used to correct for multiple comparisons. Observed heterozygosity in both groups was computed by the direct counting method. Adherences of genotypic proportions to expectations under Hardy-Weinberg equilibrium were tested separately for each SNP using the PyPop 0.7.0 software (California, USA) [20]. Statistical analyses were performed using GraphPad Prism version 7.02 for Windows (GraphPad Software, California, USA) and SPSS Statistics 23 (IBM SPSS Software, New York, USA).

Results

Patient characteristics

Characteristics of the RM group and control group are listed in Table 1. Groups did not differ in maternal age and gestational age (GA) at delivery. As expected, the women in the RM group had fewer previous live born children compared to the

control women ($P < 0.001$). Of the RM group, 65.2% had no children, compared to 19.6% in the control group.

Table 1. Subject characteristics.

	Pregnancy after RM (n=23)	Uneventful pregnancy (n=46)	P-value*
Maternal age at time of index pregnancy in years	34 (22-39) [#]	33 (20-41) [#]	0.548
Gestational age at time of birth in weeks	39 (37-41) [#]	39 (37-42) [#]	0.109
Gravidity at time of index pregnancy	5 (4-9) [#]	3 (1-7) [#]	<0.001
Parity at time of index pregnancy	0 (0-2) [#]	1 (0-5) [#]	<0.001
Number of previous miscarriages	4 (3-7) [#]	0 (0-1) [#]	<0.001

*Mann-Whitney U Test; [#] median, min-max

HLA-G polymorphisms and haplotypes

We analyzed multiple SNPs to distinguish eight haplotypes of the 3'-UTR in exon 8 of HLA-G. All genotyped SNPs fit the Hardy-Weinberg expected proportions in both groups of women and in their offspring (Supplementary Tables S1 and S2). No differences in frequency for individual SNPs or in haplotype distribution was found between groups (Supplementary Tables S3 and S4), except for the 14-bp indel polymorphism. We found a higher frequency of HLA-G 14-bp ins/del heterozygotes in RM women (65.2%) as compared to control women (36.4%) (OR 3.28; $P = 0.026$) and a lower del/del genotype (17.4% compared to 43.2%) (OR 0.28; $P = 0.039$), whereas the frequencies of ins/ins genotype are very similar in both groups (17.4% vs. 20.5%) (Table 2). Nevertheless the allelic frequencies of deletion and insertion do not differ significantly between RM and controls (Table 2). The 14-bp insertion is known to influence mRNA stability [21], resulting in lower HLA-G expression [13]. The children in both groups did not differ in frequency of individual SNPs (Supplementary Table S6), haplotypes (Supplementary Table S5) and 14-bp indel (Table 3).

Table 2. The 14-bp insertion/deletion in the 3'-UTR region of HLA-G in the women with a history of RM and the control groups.

	RM women (n=23)		Control women (n=44)*		OR	95% CI	P-value[§]
Genotype frequency							
Del/Del	4	17.4%	19	43.2%	0.28	0.08-0.95	0.039
Ins/Del	15	65.2%	16	36.4%	3.28	1.14-9.43	0.026
Ins/Ins	4	17.4%	9	20.5%	0.82	0.22-3.01	0.810
Phenotype frequency							
Ins phenotype	19	82.6%	25	56.8%	3.61	1.05-12.38	0.039
Del phenotype	19	82.6%	35	79.6%	1.22	0.33-4.50	0.810
Allele frequency							
Insertion	23	50.0%	54	38.6%	1.59	0.77-3.26	0.205
Deletion	23	50.0%	54	61.4%	0.63	0.31-1.29	0.205

*In 2 control subjects the 14bp ins/del could not be defined (4%).

[§]Chi-square.

OR, odds ratio; 95% CI, 95% confidence interval; del, deletion; ins, insertion.

Table 3. The 14-bp insertion/deletion in the 3'-UTR region of HLA-G in the offspring of the group with a history of RM and the control group.

	RM offspring (n=23)		Control offspring (n=45)*		OR	95% CI	P-value[§]
Genotype frequency							
Del/Del	8	34.8%	16	34.0%	0.97	1.33-2.77	0.969
Ins/Del	11	47.8%	22	46.8%	0.96	0.35-2.62	0.936
Ins/Ins	4	17.4%	7	14.9%	1.14	0.30-4.39	0.789
Phenotype frequency							
Ins phenotype	15	65.2%	29	61.7%	1.03	0.36-2.97	0.969
Del phenotype	19	82.6%	38	84.4%	0.88	0.23-3.36	0.789
Allele frequency							
Insertion	19	43.3%	36	40.0%	1.06	0.51-2.17	0.875
Deletion	27	58.7%	54	60.0%	0.95	0.46-1.95	0.875

*In 1 control subject the 14bp ins/del could not be defined (2%).

[§]Chi-square.

OR, odds ratio; 95% CI, 95% confidence interval; del, deletion; ins, insertion.

Placental HLA-G expression is elevated in women with history of RM

Trophoblasts were stained in the term placentas by means of immunohistochemistry with an anti-cytokeratin antibody (CAM5.2) (Fig. 1a). On the sequential slides an antibody recognizing the free heavy chain of all HLA-G isoforms (MEM-G2) was applied (Fig. 1b). Expression of HLA-G was confined to the cytokeratin positive cells in the decidua basalis, as determined by double staining experiments by immunofluorescence (Supplementary Fig. S2). We annotated the decidual part of the placental tissues and quantified the extent of staining within these annotations. Spearman’s correlation coefficient of the inter-observer reproducibility for our approach of quantitation was $r=0.79$. In the Bland-Altman plot of inter-observer measurements (Supplementary Fig. S3), most of the values ranged within a mean \pm two SD, meaning that the reproducibility of the measurement is acceptable [19].




Figure 1. Expression of trophoblast cell marker and HLA-G in term placenta. Representative examples of staining for (a) trophoblasts with cytokeratin marker CAM5.2 and (b) all HLA-G iso forms with marker MEM-G2. Decidual parts of the placenta were annotated to specify the area for analysis.

No significant difference was observed in the extent of trophoblast staining between groups (Fig. 2a). However, the extent of decidual HLA-G protein expression was elevated in the placentas of women with a history of RM (median 32.6%) compared to the control group (median 21.9%, $P < 0.0001$) (Fig. 2b).

HLA-G expression was similar in placentas of women who gave birth to their firstborn compared to women who already had a successful previous pregnancy (Supplementary Fig. S4). Using the median expression in the controls, the RM subjects were divided into either low or high HLA-G protein expression groups (Table 4). From RM cases, 87.0% belonged to the high HLA-G protein expression group (OR 6.67, 95% CI: 1.74-25.57; $P = 0.006$).




Figure 2. (a) Percentage positivity for trophoblast staining. No difference was observed in trophoblast staining between women with a history of RM and controls. (b) Percentage positivity for HLA-G staining. A higher HLA-G protein expression was observed in the decidual part of the placenta of women with a history of RM compared to controls. (c) HLA-G mRNA expression was measured in the placentas of women with a history of RM and controls. HLA-G mRNA expression was increased in term placenta of women with a history of RM compared to controls.

Table 4. HLA-G protein expression in the placentas of women with a history of RM and controls.

	RM women (n=23)		Control women (n=44)*		OR	95% CI	P-value
Low HLA-G protein expression	3	13.0%	23	50.0%	6.67	1.74-25.57	0.006[§]
High HLA-G protein expression	20	87.0%	23	50.0%			

[§]logistic regression. OR, odds ratio; 95% CI, 95% confidence interval.

To verify the differences observed at the protein level, we analyzed the mRNA expression of HLA-G in homogenates of term placentas from both groups. For this, we developed primers targeting exon 2 and 3 of the HLA-G gene, so all HLA-G isoforms were recognized. To verify that the primers only recognize HLA-G, and not HLA-C, their specificity was checked by sequencing of the amplicons. The mean Cq value for all placentas was 25.2 ± 2.0 (range 21-33), indicating expression well above background. Similar to what was observed for HLA-G protein expression,

the placentas of women with a history of RM had a 2.3-fold higher HLA-G mRNA expression than women without a history of RM (median relative level 8.2 versus 3.6, $P < 0.005$) (Fig. 2c). RM subjects were divided into either low or high HLA-G mRNA expression groups (Table 5). From RM cases, 86.4% belonged to the high HLA-G mRNA expression group (OR 6.33, 95% CI: 1.56-25.71; $P = 0.010$). No correlation was found between maternal and fetal HLA-G genotype with HLA-G expression.

Table 5. HLA-G mRNA expression in the placentas of women with a history of RM and controls.

	RM women (n=22)*		Control women (n=32)*		OR	95% CI	P-value
Low HLA-G mRNA expression	3	13.6%	16	50.0%	6.33	1.56-25.71	0.010[§]
High HLA-G mRNA expression	19	86.4%	16	50.0%			

*In 1 RM case (4%) and 12 control subjects (26%) mRNA expression could not be defined.

[§]logistic regression. OR, odds ratio; 95% CI, 95% confidence interval.

We wondered whether the higher placental HLA-G expression in the RM group was accompanied by a lower level of miRNAs. Members of the miR-148 family, and miR152 and miR-365 have been identified to target the 3'UTR of HLA-G [22, 23]. Cq values for miR-148a, miR148b and miR-152 ranged between 16 and 29. Cq values for miR-365 ranged between 25 and 39. After normalization for two reference miRNAs, no difference was observed between groups in the levels of miR-152 and miR-365 (Supplementary Fig. S5). Placental miR-148a ($P = 0.0009$) and miR-148b levels ($P = 0.0154$) were higher in the RM group compared to controls. Thus, increased HLA-G expression in the RM group was not accompanied by decreased miRNA levels.

HLA-G expression in first trimester miscarriage material and elective abortions is similar

Since HLA-G protein expression was elevated in term placentas after successful pregnancies, we additionally analyzed HLA-G expression in first trimester placentas, using the same antibody for immunohistochemistry. To this aim, we collected first trimester miscarriage of patients with a history of RM and elective abortion material and stained slides for HLA-G. Since the decidua could not be clearly defined in this early material, HLA-G positive region were selected to semi-quantitatively scored the extent of protein staining. The average score for each group is shown in Fig. 3. No difference in HLA-G protein expression was found between early miscarriages and elective abortions.




Figure 3. Amount of HLA-G staining in first trimester miscarriage and elective abortion material. HLA-G positive parts in the EVT regions of the placental slides were scored to be (1) minimally, (2) moderately or (3) intensely stained.

Discussion

In this study we investigated the HLA-G genotype and HLA-G mRNA and protein expression in term placentas of women with a history of RM and of women with no such history. A homogenous well-defined case group of women with at least three consecutive unexplained RM within 20 weeks of gestation was included. A

lower frequency of the HLA-G 3'UTR 14-bp deletion genotype was observed in the case group, suggesting that genetic predisposition to a low level of HLA-G played a role in the etiology of previous RM. In the current successful pregnancies, a significantly higher HLA-G protein and mRNA expression was found in the placenta of the RM group compared to the control group.

The most studied polymorphism in exon 8 of 3-UTR of the HLA-G gene is the 14 bp indel polymorphism, which has been associated with altered HLA-G expression. The insertion genotype is associated with low levels of sHLA-G [13]. In addition, the fetal 14-bp ins/ins genotype has been associated with lower surface expression of HLA-G on first trimester trophoblast cells than the 14-bp del/del genotype [24]. We did not find any differences in fetal HLA-G 3'UTR haplotypes or individual SNPs between both groups. However, we found a higher frequency of HLA-G 14-bp ins/del heterozygotes in RM women (65.2%) as compared with control women (39.1%), and a lower frequency of HLA-G 14 bp del/del homozygotes (17.4% and 43.2%, respectively). This is consistent with some studies [25-27], but not others [28, 29]. Since several studies have focused on the HLA-G 14-bp polymorphism in RM with controversial or inconclusive results, Wang *et al* performed a meta-analysis [30], which suggested that the HLA-G 14-bp insertion allele was associated with increased risk of RM. In 2014, yet another meta-analysis indicated that there was only an association between the HLA-G 14-bp indel polymorphism and RM in patients with three or more miscarriages [31]. In the present study, we have not addressed polymorphisms in the HLA-G promotor region, but they may be associated to RM, as recently shown [32].

Both the individual SNPs and the most common extended 3-UTR haplotypes of HLA-G were studied in the group of women with a history of RM and controls. HLA-G haplotype distribution and frequencies of individual SNPs in the 3-UTR region of HLA-G were neither significantly different between the groups of women, nor in their offspring (Supplementary Table S3-6). Studies have shown

that individual SNPs in the 3-UTR region of HLA-G are not significantly associated with RM, but that the UTR-4 haplotype seemed to be protective against RM [33, 34]. Similarly we observed a lower incidence of the HLA-G UTR-4 haplotype in women with RM (10.9% in RM women vs 15.9% in control women) (Supplementary Table S3). Remarkably, the HLA-G UTR-4 haplotype was more frequently present in the offspring of women with RM than in the offspring of controls (21.7% vs 13.3% respectively) (Supplementary Table S5) and less frequently in miscarriage material from women with RM (10%, data not shown). Even though these results were not statistically significant, possibly due to limited sample size, collectively they support the idea that this haplotype might have a protective effect in uncomplicated pregnancy.

HLA-G in the placenta is suggested to play a role in the induction of immunological tolerance at the fetal-maternal interface, by functioning as a trophoblast-restricted inhibitory ligand of maternal immune cells. Only a few studies have focused on HLA-G protein expression in the placentas of women with a history of RM, with contradicting results [35, 36]. Remarkably, the present immune-histochemical analysis of term placentas of successful pregnancies showed a significantly higher HLA-G protein expression in women with a history of RM compared to controls, although this RM group had a lower frequency of the 14 bp del/del genotype. This is not in line with results previously found in peripheral blood [25, 37] and suggests that local regulation is involved. HLA-G was mostly confined to the trophoblast areas at the fetal-maternal interface (decidua basalis), as determined by double label immunofluorescence experiments (Supplementary Fig. S2), and the level of HLA-G expression was independent of previous pregnancies.

Since the level of HLA-G expression can depend on the differentiation status of EVT, as determined by *in vitro* studies using isolated primary trophoblasts [38], it is unclear whether the observed differences in HLA-G expression are a direct consequence of transcriptional regulation or a secondary of an altered

differentiation status of the EVT. Possibly, for a successful pregnancy to occur after previous RM, a compensatory mechanism resulting in high HLA-G protein expression is in place. When comparing first trimester miscarriage material of women with a history of RM and material of elective abortions, we did not observe a difference in HLA-G expression between both groups, suggesting that successful pregnancy in women with a history of RM is due to high fetal HLA-G expression in the current pregnancy. Besides HLA-G, other molecules and immune interactions may be involved in the immune-regulation leading to successful pregnancy.


We found a higher miR-148a and miR-148b expression in the term placentas of the RM group despite the elevated HLA-G expression. Apparently, either these miRNAs do not bind or binding does not result in post-transcriptional repression of HLA-G. This leads us to hypothesize that the higher HLA-G protein expression in the RM group may be the result of an epigenetically-regulated compensatory mechanism to achieve an ongoing pregnancy in patients with a history of RM. Alternatively, the higher HLA-G protein expression in the case group may be an epiphenomenon resulting from the previous miscarriages. It is possible that the elevated HLA-G in the term placentas of women with RM is the result of proteolytic cleavage of the membrane bound HLA-G1 isoform resulting from activity of metalloproteases, leading to elevated sHLA-G levels. The antibody recognizing MEMG2 in our immune-histochemical assays does not distinguish membrane bound HLA-G from soluble HLA-G. Previous miscarriages could lead to increased metalloprotease (MMP) levels [39], which in turn lead to increased proteolytic shedding of HLA-G1 [11]. MMP2 and MMP9 mRNA expression was not elevated in term placentas of women with a history of RM compared to controls (Supplementary Fig. S6), but this does not fully exclude the involvement of MMPs since their activity was not tested in the current setting.

In conclusion, whereas women with RM have a genetic predisposition to lower HLA-G levels, HLA-G expression is increased in the placenta of ongoing pregnancies after


RM. This implies that HLA-G upregulation could be a compensatory mechanism in the occurrence of RM to achieve an ongoing pregnancy. Whether the higher HLA-G expression in the ongoing pregnancy after RM is a cause or a consequence of the successful pregnancy remains to be established. Future studies should be concentrated on further establishing the role of HLA-G in complicated pregnancies. Measurement of maternal sHLA-G may provide further insight on the prognosis of the outcome of pregnancies in women with a history of RM.

References


1. Regan, L., M. Backos, and R. Rai, *Recurrent Miscarriage, investigation and treatment of couples*. Royal College of Obstetricians and Gynaecologists - Greentop Guideline 17, 2011.
2. Practice Committee of the American Society for Reproductive, M., *Evaluation and treatment of recurrent pregnancy loss: a committee opinion*. *Fertil Steril*, 2012. **98**(5): p. 1103-11.
3. Kovats, S., et al., *A class I antigen, HLA-G, expressed in human trophoblasts*. *Science*, 1990. **248**(4952): p. 220-3.
4. Ishitani, A., et al., *Protein expression and peptide binding suggest unique and interacting functional roles for HLA-E, F, and G in maternal-placental immune recognition*. *J Immunol*, 2003. **171**(3): p. 1376-84.
5. Kofod, L., et al., *Endometrial immune markers are potential predictors of normal fertility and pregnancy after in vitro fertilization*. *Am J Reprod Immunol*, 2017. **78**(3).
6. Pfeiffer, K.A., et al., *Soluble HLA levels in early pregnancy after in vitro fertilization*. *Hum Immunol*, 2000. **61**(6): p. 559-64.
7. Lee, N., et al., *The membrane-bound and soluble forms of HLA-G bind identical sets of endogenous peptides but differ with respect to TAP association*. *Immunity*, 1995. **3**(5): p. 591-600.
8. Rebmann, V., et al., *Detection of soluble HLA-G molecules in plasma and amniotic fluid*. *Tissue Antigens*, 1999. **53**(1): p. 14-22.
9. Shaikly, V.R., et al., *Analysis of HLA-G in maternal plasma, follicular fluid, and preimplantation embryos reveal an asymmetric pattern of expression*. *J Immunol*, 2008. **180**(6): p. 4330-7.
10. Larsen, M.H., et al., *Human leukocyte antigen-G in the male reproductive system and in seminal plasma*. *Mol Hum Reprod*, 2011. **17**(12): p. 727-38.
11. Rizzo, R., et al., *Matrix metalloproteinase-2 (MMP-2) generates soluble HLA-G1 by cell surface proteolytic shedding*. *Mol Cell Biochem*, 2013. **381**(1-2): p. 243-55.
12. Rousseau, P., et al., *The 14 bp deletion-insertion polymorphism in the 3' UT region of the HLA-G gene influences HLA-G mRNA stability*. *Hum Immunol*, 2003. **64**(11): p. 1005-10.
13. Chen, X.Y., et al., *The 14 bp deletion polymorphisms in HLA-G gene play an important role in the expression of soluble HLA-G in plasma*. *Tissue Antigens*, 2008. **72**(4): p. 335-41.
14. Yie, S.M., et al., *A single base-pair mutation in the 3'-untranslated region of HLA-G mRNA is associated with pre-eclampsia*. *Mol Hum Reprod*, 2008. **14**(11): p. 649-53.
15. Tan, Z., et al., *Allele-specific targeting of microRNAs to HLA-G and risk of asthma*. *Am J Hum Genet*, 2007. **81**(4): p. 829-34.
16. Meuleman, T., et al., *Increased complement C4d deposition at the maternal-fetal interface in unexplained recurrent miscarriage*. *J Reprod Immunol*, 2016. **113**: p. 54-60.
17. Hua, Y.J., et al., *Comparison of normalization methods with microRNA microarray*. *Genomics*, 2008. **92**(2): p. 122-8.
18. Wotschovsky, Z., et al., *Reference genes for the relative quantification of microRNAs in renal cell carcinomas and their metastases*. *Anal Biochem*, 2011. **417**(2): p. 233-41.
19. Bland, J.M. and D.G. Altman, *Statistical methods for assessing agreement between two methods of clinical measurement*. *Lancet*, 1986. **1**(8476): p. 307-10.
20. Lancaster, A.K., et al., *PyPop update--a software pipeline for large-scale multilocus population genomics*. *Tissue Antigens*, 2007. **69 Suppl 1**: p. 192-7.
21. Hviid, T.V., et al., *HLA-G allelic variants are associated with differences in the HLA-G mRNA isoform profile and HLA-G mRNA levels*. *Immunogenetics*, 2003. **55**(2): p. 63-79.
22. Mori, A., et al., *HLA-G expression is regulated by miR-365 in trophoblasts under hypoxic conditions*. *Placenta*, 2016. **45**: p. 37-41.
23. Manaster, I., et al., *MIRNA-mediated control of HLA-G expression and function*. *PLoS One*, 2012. **7**(3): p. e33395.
24. Djuricic, S., et al., *Allelic imbalance modulates surface expression of the tolerance-inducing HLA-G molecule on primary trophoblast cells*. *Mol Hum Reprod*, 2015. **21**(3): p. 281-95.
25. Kalotra, V., et al., *The HLA-G 14 bp insertion/deletion polymorphism and its association with soluble HLA-G levels in women with recurrent miscarriages*. *HLA*, 2018. **91**(3): p. 167-174.
26. Xue, S., et al., *Recurrent spontaneous abortions patients have more -14 bp/+14 bp heterozygotes in the 3'UT region of the HLA-G gene in a Chinese Han population*. *Tissue Antigens*, 2007. **69 Suppl 1**: p. 153-5.
27. Tripathi, P., et al., *Role of 14-bp deletion in the HLA-G gene in the maintenance of pregnancy*. *Tissue Antigens*, 2004. **64**(6): p. 706-10.
28. Vargas, R.G., et al., *Association of HLA-G alleles and 3' UTR 14 bp haplotypes with recurrent miscarriage in Brazilian couples*. *Hum Immunol*, 2011. **72**(6): p. 479-85.
29. Hviid, T.V., et al., *HLA-G polymorphisms in couples with recurrent spontaneous abortions*. *Tissue Antigens*, 2002. **60**(2): p. 122-32.
30. Wang, X., W. Jiang, and D. Zhang, *Association of 14-bp insertion/deletion polymorphism of HLA-G gene with unexplained recurrent spontaneous abortion: a meta-analysis*. *Tissue Antigens*, 2013. **81**(2): p. 108-15.
31. Fan, W., et al., *Relationship between HLA-G polymorphism and susceptibility to recurrent miscarriage: a meta-analysis of non-family-based studies*. *J Assist Reprod Genet*, 2014. **31**(2): p. 173-84.
32. Yazdani, N., et al., *HLA-G regulatory variants and haplotypes with susceptibility to recurrent pregnancy loss*. *Int J Immunogenet*, 2018. **45**(4): p. 181-189.
33. Amodio, G., et al., *Association of genetic variants in the 3'UTR of HLA-G with Recurrent Pregnancy Loss*. *Hum Immunol*, 2016. **77**(10): p. 886-891.
34. Meuleman, T., et al., *Lower frequency of the HLA-G UTR-4 haplotype in women with unexplained recurrent miscarriage*. *J Reprod Immunol*, 2018. **126**: p. 46-52.
35. Emmer, P.M., et al., *Altered phenotype of HLA-G expressing trophoblast and decidual natural killer cells in pathological pregnancies*. *Hum Reprod*, 2002. **17**(4): p. 1072-80.
36. Bhalla, A., et al., *Comparison of the expression of human leukocyte antigen (HLA)-G and HLA-E in women with normal pregnancy and those with recurrent miscarriage*. *Reproduction*, 2006. **131**(3): p. 583-9.
37. Dahl, M., et al., *Human leukocyte antigen (HLA)-G during pregnancy part II: associations between maternal and fetal HLA-G genotypes and soluble HLA-G*. *Hum Immunol*, 2015. **76**(4): p. 260-71.
38. Goldman-Wohl, D.S., et al., *HLA-G expression in extravillous trophoblasts is an intrinsic property of cell differentiation: a lesson learned from ectopic pregnancies*. *Mol Hum Reprod*, 2000. **6**(6): p. 535-40.
39. Jokimaa, V., et al., *Altered expression of genes involved in the production and degradation of endometrial extracellular matrix in patients with unexplained infertility and recurrent miscarriages*. *Mol Hum Reprod*, 2002. **8**(12): p. 1111-6.




Supplementary Figure S1. Examples of minimal (a), moderate (b), and intense (c) HLA-G staining in first trimester placenta.




Supplementary Figure S2. (a) Cytokeratin 8 (CAM5.2, red) and HLA-G (MEM-G2, green) colocalize in the decidua; yellow in merged image indicates overlap of red and green labels. (b) CAM5.2 stains all trophoblasts in the placenta. (c) MEM-G2 staining is limited to the extravillous trophoblasts in the decidual part of the placenta.




Supplementary Figure S3. Bland-Altman plot of interobserver measurements of HLA-G staining. Ninety percent of the values ranged within a mean ± 2 SD deviations, indicating acceptable reproducibility.



Supplementary Figure S4. HLA-G expression in placentas of healthy first pregnancies compared to subsequent pregnancies. Previous pregnancies did not influence placental HLA-G expression in the control group with uncomplicated pregnancies.



Supplementary Figure S5. miRNA expression in term placentas of women with a history of RM and controls. (a-b) miR-148a and miR-148b expression was elevated in placenta of women with a history of RM compared to controls. No difference in miRNA expression was seen for (c) miR-152 and (d) miR-365 (statistics: Mann-Whitney test).



Supplementary Figure S6. MMP2 and MMP9 mRNA expression in term placentas of women with a history of RM and controls. (a) MMP2 mRNA expression was similar between women with a history of RM and controls. (b) MMP9 mRNA expression was similar between women with a history of RM and controls (statistics: Mann-Whitney test).

Supplementary Table S1. Hardy-Weinberg analyses for HLA-G 3'UTR genotypes in women with RM and controls.

SNP	Recurrent miscarriage (n=23)			Controls (n=46)		
	common <i>P</i>	homozygote <i>P</i>	heterozygot <i>P</i>	common <i>P</i>	homozygote <i>P</i>	heterozygot <i>P</i>
14-bp	0.1444	0.3020	0.3020	0.1220	0.3119	0.2870
+3003	#	0.9370	&	0.5888	0.7548	0.6054
+3010	0.7171	0.7637	0.7526	0.1379	0.2975	0.2953
+3027	#	0.9653	&	0.8602	0.9474	0.8626
+3035	0.6481	0.7965	0.6619	0.7823	0.9045	0.7874
+3142	0.7171	0.7637	0.7526	0.2322	0.3997	0.3987
+3187	0.5487	0.6508	0.5836	0.3077	0.3077	0.2440
+3196	0.4238	0.5347	0.4688	0.2315	0.3908	0.2679
+3422	0.9250	0.9520	0.9292	0.5149	0.7170	0.5321
+3496	#	0.8996	&	0.5888	0.7548	0.6054
+3509	0.5855	0.6588	0.6280	0.2938	0.4428	0.3332

All Hardy-Weinberg analyses. *P*, p value. # Too many parameters for chi-square test. & Too few cases to calculate p-value.

Supplementary Table S2. Hardy-Weinberg analyses for HLA-G 3'UTR genotypes in the RM and control offspring.

SNP	Recurrent miscarriage (n=23)			Controls (n=46)		
	common <i>P</i>	homozygote <i>P</i>	heterozygot <i>P</i>	common <i>P</i>	homozygote <i>P</i>	heterozygot <i>P</i>
14-bp	0.9577	0.9647	0.9637	0.9011	0.9341	0.9314
+3003	0.9471	0.9644	0.9504	0.8976	0.9458	0.9013
+3010	0.2058	0.2920	0.2883	0.3181	0.4854	0.4820
+3027	#	0.9653	&	#	0.9877	&
+3035	0.7224	0.8528	0.7319	0.7009	0.8533	0.7094
+3142	0.2921	0.4575	0.4567	0.8841	0.9179	0.9179
+3187	0.8147	0.8691	0.8275	0.7358	0.7894	0.7617
+3196	0.5487	0.6508	0.5836	0.7862	0.8389	0.8038
+3422	0.4882	0.6574	0.5128	0.9110	0.9424	0.9161
+3496	0.9471	0.9644	0.9504	0.8976	0.9458	0.9013
+3509	0.5487	0.6508	0.5836	#	0.8150	0.7861

All Hardy-Weinberg analyses. *P*, p value. # Too many parameters for chi-square test. & Too few cases to calculate p-value.

Supplementary Table S3. Haplotypes of women in the RM group and the control group.

	RM women 2n=46		Controls 2n=88*		OR	95% C.I.		P
						Lower	Upper	
UTR-1	13	28.3%	28	31.8%	0.90	0.413	1.965	0.792
UTR-2	16	34.8%	23	26.1%	1.60	0.742	3.451	0.231
UTR-3	5	10.9%	8	9.1%	1.28	0.394	4.159	0.681
UTR-4	5	10.9%	14	15.9%	0.68	0.229	2.019	0.487
UTR-5	4	8.7%	4	4.5%	2.10	0.499	8.789	0.312
UTR-7	3	6.5%	7	8.0%	0.85	0.209	3.440	0.817
UTR-8	0	0.0%	0	0.0%	x	x	x	x
UTR-18	0	0.0%	3	3.4%	0.00	0.000		0.999
UTR-N	0	0.0%	1	1.1%	0.00	0.000		1.000
	46	100%	88	100%				

All univariate logistic regression analysis. *P*, p value; OR, odds ratio; 95% CI, 95% confidence interval; n.a, not applicable. *In 2 control subjects the UTR haplotype could not be defined (4%). The 3'UTR haplotype nomenclature is consistent with publication by Castelli *et al.*

Supplementary Table S4. HLA-G 3'UTR genotypic polymorphisms in women with recurrent miscarriage and uneventful pregnancy.

	RM women (n=23)		Controls (n=44)		OR	95% CI	P	Pc
	n	%	n	%				
3003	0	0.0%	2	4.5%	n.c.			
CC	4	17.4%	10	22.7%	0.67	0.19-2.45	0.549	1.000
CT	19	82.6%	32	72.7%	ref.			
3010	8	34.8%	12	27.3%	0.94	0.30-3.01	0.923	1.000
CG	12	52.2%	17	38.6%	ref.			
GG	3	13.0%	15	34.1%	0.28	0.07-1.20	0.087	1.000
3027	0	0.0%	0	0.0%	n.c.			
AA	3	13.0%	6	13.6%	0.95	0.22-4.21	0.946	1.000
AC	20	87.0%	38	86.4%	ref.			
3035	16	69.6%	36	81.8%	ref.			
CC	7	30.4%	8	18.2%	1.97	0.61-6.36	0.258	1.000
CT	0	0.0%	0	0.0%	n.c.			
3142	3	13.0%	14	31.8%	0.32	0.08-1.36	0.124	1.000
CC	12	52.2%	18	40.9%	ref.			
CG	8	34.8%	12	27.3%	1.00	0.32-3.17	1.000	1.000
3187	11	47.8%	23	52.3%	ref.			
AA	11	47.8%	14	31.8%	1.64	0.57-4.48	0.362	1.000
AG	1	4.3%	7	15.9%	0.30	0.03-2.74	0.285	1.000
3196	10	43.5%	27	61.4%	ref.			
CC	12	52.2%	12	27.3%	2.70	0.92-7.95	0.071	1.000
CG	1	4.3%	5	11.4%	n.c.			
3422	15	65.2%	33	75.0%	ref.			
CC	7	30.4%	9	20.5%	1.71	0.54-5.46	0.365	1.000
CT	1	4.3%	2	4.5%	n.c.			
3496	0	0.0%	2	4.5%	n.c.			
AA	5	21.7%	10	22.7%	0.89	0.26-3.01	0.850	1.000
AG	18	78.3%	32	72.7%	ref.			
3509	9	39.1%	26	59.1%	ref.			
GG	12	52.2%	13	29.5%	2.67	0.90-7.94	0.078	1.000
GT	2	8.7%	5	11.4%	1.16	0.19-7.04	0.875	1.000
TT								

Data are all n (%). All univariate logistic regression analysis. Per HLA-G genotype the highest prevalence was defined as the reference group. If percentages in a group were below 5%, no calculations were performed. *P*, *p* value; *Pc*, *p* value corrected for multiple comparisons; *OR*, odds ratio; *95% CI*, 95% confidence interval; n.c., not calculated; ref, reference group.

Supplementary Table S5. Haplotypes of the offspring in the RM group and the control group.

	RM offspring 2n=46		Controls 2n=90*		OR	95% C.I.		P
	n	%	n	%		Lower	Upper	
UTR-1	11	23.9%	28	31.1%	0.70	0.309	1.566	0.381
UTR-2	13	28.3%	26	28.9%	0.97	0.441	2.131	0.939
UTR-3	6	13.0%	11	12.2%	1.08	0.371	3.125	0.891
UTR-4	10	21.7%	12	13.3%	1.81	0.714	4.565	0.212
UTR-5	3	6.5%	7	7.8%	0.83	0.204	3.360	0.791
UTR-7	3	6.5%	3	3.3%	2.02	0.392	10.446	0.400
UTR-8	0	0.0%	1	1.1%	0.00	0.000		1.000
UTR-18	0	0.0%	2	2.2%	0.00	0.000		0.999
UTR-N	0	0.0%	0	0.0%	x	x	x	x
	46	100%	90	100%				

All univariate logistic regression analysis. *P*, *p* value; *OR*, odds ratio; *95% CI*, 95% confidence interval; n.a., not applicable. *In 1 control subject the UTR haplotype could not be defined (2%). The 3'UTR haplotype nomenclature is consistent with publication by Castelli *et al.*

Supplementary Table S6. HLA-G 3'UTR genotypic polymorphisms in the offspring of women with recurrent miscarriage and uneventful pregnancy.

	RM offspring (n=23)		Controls (n=45)		OR	95% CI	P	Pc
3003 CC	1	4.3%	1	2.2%	n.c.			
CT	8	34.8%	10	22.2%	1.94	0.63-5.95	0.245	1.000
TT	14	60.9%	34	75.6%	ref.			
3010 CC	5	21.7%	11	24.4%	0.58	0.16-2.02	0.389	1.000
CG	15	65.2%	19	42.2%	ref.			
GG	3	13.0%	15	33.3%	0.25	0.06-1.04	0.057	0.969
3027 AA	0	0.0%	0	0.0%	n.c.			
AC	3	13.0%	3	6.7%	2.10	0.39-11.34	0.389	1.000
CC	20	87.0%	42	93.3%	ref.			
3035 CC	17	73.9%	35	77.8%	ref.			
CT	6	26.1%	10	22.2%	1.24	0.38-3.97	0.722	1.000
TT	0	0.0%	0	0.0%	n.c.			
3142 CC	5	21.7%	11	24.4%	0.71	0.20-2.50	0.598	1.000
CG	14	60.9%	22	48.9%	ref.			
GG	4	17.4%	12	26.7%	0.52	0.14-1.95	0.335	1.000
3187 AA	13	56.5%	20	44.4%	ref.			
AG	9	39.1%	21	46.7%	0.66	0.23-1.88	0.436	1.000
GG	1	4.3%	4	8.9%	n.c.			
3196 CC	11	47.8%	24	53.3%	ref.			
CG	11	47.8%	17	37.8%	1.41	0.50-4.00	0.516	1.000
GG	1	4.3%	4	8.9%	n.c.			
3422 CC	14	60.9%	29	64.4%	ref.			
CT	9	39.1%	14	31.1%	1.33	0.47-3.82	0.594	1.000
TT	0	0.0%	2	4.4%	n.c.			
3496 AA	1	4.3%	1	2.2%	n.c.			
AG	8	34.8%	10	22.2%	1.94	0.63-5.95	0.245	1.000
GG	14	60.9%	34	75.6%	ref.			
3509 GG	11	47.8%	23	51.1%	ref.			
GT	11	47.8%	17	37.8%	1.35	0.48-3.85	0.571	1.000
TT	1	4.3%	4	8.9%	0.52	0.05-5.25	0.581	1.000
CT	0	0.0%	1	2.2%	n.c.			

Data are all n (%). All univariate logistic regression analysis. Per HLA-G genotype the highest prevalence was defined as the reference group. If percentages in a group were below 5%, no calculations were performed. *P*, *p* value; *Pc*, *p* value corrected for multiple comparisons; *OR*, odds ratio; *95% CI*, 95% confidence interval; n.c., not calculated; ref, reference group.