



Universiteit
Leiden
The Netherlands

Hunting for new physics in the primordial Universe

Wang, D.-G.

Citation

Wang, D. -G. (2020, August 27). *Hunting for new physics in the primordial Universe. Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/135951>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/135951>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/135951> holds various files of this Leiden University dissertation.

Author: Wang, D.-G.

Title: Hunting for new physics in the primordial Universe

Issue date: 2020-08-27

3

Shift-symmetric orbital inflation

Abstract: We present a new class of two-field inflationary attractor models, known as ‘*shift-symmetric orbital inflation*’, whose behaviour is strongly multi-field but whose predictions are remarkably close to those of single-field inflation. In these models, the field space metric and potential are such that the inflaton trajectory is along an ‘angular’ isometry direction whose ‘radius’ is constant but *arbitrary*. As a result, the radial (isocurvature) perturbations away from the trajectory are exactly *massless* and they freeze on superhorizon scales. These models are the first exact realization of the ‘ultra-light isocurvature’ scenario, previously described in the literature, where a combined shift symmetry emerges between the curvature and isocurvature perturbations and results in primordial perturbation spectra that are entirely consistent with current observations. Due to the turning trajectory, the radial perturbation sources the tangential (curvature) perturbation and makes it grow linearly in time. As a result, only one degree of freedom (*i.e.* the one from isocurvature modes) is responsible for the primordial observables at the end of inflation, which yields the same phenomenology as in single-field inflation. In particular, isocurvature perturbations and local non-Gaussianity are highly suppressed here, even if the inflationary dynamics is truly multi-field. We comment on the generalization to models with more than two fields.

Keywords: multi-field inflation, cosmological perturbation theory

Based on:

A. Achúcarro, E. Copeland, O. Iarygina, G. Palma, D.-G. Wang, Y. Welling
Shift-symmetric orbital inflation: single field or multi-field?
Phys. Rev. D Rapid Communications, **102**, 021302(R) (2020), [arXiv:1901.03657].

3.1 Introduction

Single field slow roll inflation is the leading explanation for the observations through the CMB [7] that primordial perturbations are very close to Gaussian and adiabatic, yet embedding it in an ultraviolet complete theory such as string theory is notoriously difficult. Moduli fields arising from string compactifications require stabilizing to realize single field inflation [19], and large field excursions test the validity of using four dimensional effective theories¹.

One may therefore wonder whether there are multi-field inflationary scenarios with a phenomenology similar to that of single field inflation. Most of the previous studies on multi-field inflation focus on the slow-roll slow-turn regime, where the isocurvature and curvature perturbations are weakly coupled, therefore one may get single-field like phenomenology unsurprisingly. Meanwhile, models with significant multi-field turning behaviour are commonly considered to be problematic. In the usual understanding of this regime, light fields during inflation may lead to isocurvature perturbations and local non-Gaussianity tightly constrained by current observations. However, it has been suggested recently that inflation with non-stabilized light fields on an axion-dilaton system can be compatible with the latest CMB data [75, 78, 120–124]. In particular, it was pointed out in [78] that, when the perturbations orthogonal to the trajectory are *massless* but efficiently *coupled* to the inflaton, the isocurvature modes are dynamically suppressed. This is the “ultra-light isocurvature” scenario.

In this paper we provide for the first time a family of exact models of inflation in which the multi-field effects are significant, but the phenomenology remains similar to single field inflation. The models combine two ingredients: First, the inflaton trajectory proceeds along an isometry direction of the field space, so it is Orbital Inflation in the sense of [81, 82]. This ensures time independence of the coupling between the radial and tangential inflationary perturbations. Second, the trajectory can have an *arbitrary* radius (within some range described below), and a constant radius is proven to be a neutrally stable attractor. Hence, isocurvature perturbations become exactly massless. The two ingredients, combined, guarantee that the sourcing of the curvature perturbation is sustained over many e-folds of inflationary expansion. The action for the perturbations inherits a symmetry

¹The recent swampland debate highlights the importance of finding viable scenarios for inflation that are not strictly single-field. See, for instance, the discussion in [47] as compared to [24, 137]

between background solutions that is not manifest in the potential or in the Lagrangian. We show that, at the end of inflation, only the isocurvature degree of freedom is responsible for the generation of primordial observables, but perturbations still remain adiabatic and Gaussian. We call this scenario *shift-symmetric orbital inflation*.

Crucially this scenario provides a new direction to explore inflation and a potential resolution to some of the problems faced by the embedding of inflation in string theory. That is, in the construction of inflationary models wherein every modulus is stabilized except for the inflaton, one could be missing less restrictive realizations of inflation compatible with current observational constraints. We set $\hbar = c = 1$ and the reduced Planck mass $M_p \equiv (8\pi G)^{-1/2} = 1$, where G is Newton's constant.

3.2 A toy model

To illustrate the idea, we first consider the following Lagrangian in flat field space with polar coordinates (illustrated in Fig. 3.1)

$$\mathcal{L} = \frac{1}{2} [\rho^2 (\partial\theta)^2 + (\partial\rho)^2] - \frac{1}{2} m^2 \left(\theta^2 - \frac{2}{3\rho^2} \right). \quad (3.1)$$

The potential has a monodromy in the angular coordinate, and although it is unbounded at $\rho \rightarrow 0$, inflation only takes place in the physically consistent regime where $V(\rho, \theta) > 0$. Moreover, as shown in the perturbation analysis below, our study is restricted to radii that cannot be too small. Therefore, we only care about the local form of the potential close to the inflationary trajectory, which we assume is captured well by (3.1). In general, it is difficult to solve the background equations analytically in such a system. However, this model has the following exact neutrally stable solutions at any radius (see Fig. 3.1)

$$\rho = \rho_0, \quad \dot{\theta} = \pm \sqrt{\frac{2}{3}} \frac{m}{\rho_0^2}. \quad (3.2)$$

The Friedmann equation becomes $H^2 = m^2 \theta^2 / 6$ on the attractor, where H is the Hubble parameter, and the first slow-roll parameter is $\epsilon \equiv -\dot{H}/H^2 = \frac{2}{\rho_0^2 \theta^2}$. This trajectory is nongeodesic in field space, with turning effects that depend on the radius κ of the trajectory. Note that here $\kappa = \rho_0$ but, if the field space geometry is curved, κ will be a more general function of ρ_0 .

The situation is reminiscent of circular orbits in a spherically symmetric gravitational field, where the centripetal force stabilizes the radial direction,




Figure 3.1: The toy model potential $V(\rho, \theta)$ given in (3.1) together with a typical inflationary trajectory indicated with the solid black line.

and the inflaton can circle at any radius with the corresponding angular velocity. For the field system on the cosmological background, only the isometric circular orbits appear, and we need to break the shift symmetry of θ in the potential to overcome the Hubble friction. We can label each solution by a continuous parameter c with the corresponding map

$$\rho_c = \rho_0 + c, \quad (\theta_c^2)' = \frac{(\theta_0^2)'}{(1 + c/\kappa)^2}, \quad (3.3)$$

where the prime $'$ denotes a derivative with respect to e-folds $d/dN = d/(Hdt)$. This transformation identifies all the trajectories in (3.2) and hints at the existence of a shift symmetry for the perturbations. In flat gauge, the isocurvature perturbation σ is associated with $\delta\rho$ and the curvature perturbation \mathcal{R} with $\frac{\rho}{\sqrt{2\epsilon}}\delta\theta$, which equals $\frac{1}{4}\rho^2\delta(\theta^2)$ in this toy model. To find the effect of the transformation on the perturbations, we split $\rho = \rho_0 + \sigma$ and $(\theta^2)' = (\theta_0^2)'(1 - \mathcal{R}')$. This allows us to determine how a small c changes σ and \mathcal{R}' . In the long wavelength limit every transformed set of perturbations $(\sigma_c, \mathcal{R}'_c)$ provide a new solution to the equations of motion. This is because homogeneous perturbations map background solutions onto each other. Therefore, we expect the following *symmetry* for linearized pertur-

bations

$$\sigma \rightarrow \sigma + c, \quad \mathcal{R}' \rightarrow \mathcal{R}' + \frac{2}{\kappa}c. \quad (3.4)$$

Given the shift symmetry of σ , the isocurvature perturbation is expected to be massless and freeze after horizon-exit. Meanwhile, the symmetry also indicates that \mathcal{R} has a growing solution that is dictated by the constant σ on superhorizon scales.

To get an intuitive notion of the perturbations behavior, we employ the δN formalism [131–135]. From the Friedmann equation and the exact solution (3.2), the number of e-folds until the end of inflation is $N = \rho^2\theta^2/4 - 1/2$. The curvature perturbation at the end of inflation is

$$\mathcal{R}(k_*) = \delta N \simeq \frac{1}{\sqrt{2\epsilon_*}}(\rho\delta\theta)_* + \frac{2N_*}{\kappa}\delta\rho_*, \quad (3.5)$$

where $(\rho\delta\theta)_*$ and $\delta\rho_*$ are field fluctuations with typical amplitude $\frac{H_*}{2\pi}$ at horizon-exit of the k_* mode. This yields the following power spectrum of curvature perturbations

$$P_{\mathcal{R}}(k_*) \simeq \frac{H_*^2}{4\pi^2} \left(\frac{1}{2\epsilon_*} + \frac{4N_*^2}{\kappa^2} \right). \quad (3.6)$$

Here the first contribution has an adiabatic origin, just like in the single-field models, and the second term corresponds to the conversion from isocurvature to curvature modes on superhorizon scales. When the radius of the trajectory is small enough, namely $8\epsilon_* \ll \kappa^2 \ll 8\epsilon_*N_*^2 \approx 4N_*$, the second term in (3.6) dominates. Then the final power spectrum becomes $P_{\mathcal{R}}(k_*) \simeq H_*^2N_*^2/(\pi^2\kappa^2)$, which is generated by one single degree of freedom – the isocurvature mode.

3.3 Shift-symmetric orbital inflation

To construct generic models with the above properties, we begin with an axion-dilaton system in a non-trivial field manifold (θ, ρ) with kinetic term $K = -\frac{1}{2}(f(\rho)\partial_\mu\theta\partial^\mu\theta + \partial_\mu\rho\partial^\mu\rho)$. This field space, of curvature $\mathbb{R} = f_\rho^2/2f^2 - f_{\rho\rho}/f$, arises generically from UV completions of inflation in quantum gravity or from an effective field theory (EFT) viewpoint. To realize shift-symmetric orbital inflation, we assume the inflationary trajectory to be isometric, *i.e.* along the θ direction at *any* (constant) radius in field space. The potential can be derived by generalizing the Hamilton-Jacobi

formalism [131, 138–140] to a two-field system. It has the general form

$$V = 3H^2 - 2\frac{H_\theta^2}{f(\rho)}, \quad (3.7)$$

where H is a function of θ only, $H_\theta \equiv dH/d\theta$ and $f(\rho) > 0$. The model (3.1) is recovered for $H \propto \theta$ and $f(\rho) = \rho^2$, corresponding to a flat field space parametrized by polar coordinates. This non-linear system admits exact solutions

$$\dot{\theta} = -2\frac{H_\theta}{f}, \quad \rho = \rho_0. \quad (3.8)$$

Thus the inflaton moves in an orbit of constant radius, as ensured by the Hamilton-Jacobi formalism. As in the toy model, this trajectory is not along a geodesic. Here the tangent and normal vectors to the trajectory are $\mathcal{T}^a = 1/\sqrt{f}(1, 0)$ and $\mathcal{N}^a = (0, 1)$, and the radius of the turning trajectory is a constant given by $\kappa = 2f/f_\rho$. It follows that all these trajectories are *neutrally stable*: a small perturbation orthogonal to a given orbital trajectory will bring us to one of the neighbouring trajectories. The attractor behaviour is explicitly demonstrated in the appendix of the original paper.

3.4 Analysis of perturbations

In flat gauge, the comoving curvature perturbation \mathcal{R} is defined as the projection of the field perturbation along the inflationary trajectory $\mathcal{R} = \frac{1}{\sqrt{2\epsilon}}\mathcal{T}_a\delta\phi^a$, and the isocurvature perturbation σ corresponds to the orthogonal projection $\sigma = \mathcal{N}_a\delta\phi^a$. Then for generic multi-field models, the quadratic action of perturbations takes the following form [78]

$$S^{(2)} = \frac{1}{2} \int d^4x a^3 \left[2\epsilon \left(\dot{\mathcal{R}} - \frac{2H}{\kappa}\sigma \right)^2 + \dot{\sigma}^2 - \mu^2\sigma^2 + \dots \right], \quad (3.9)$$

where ellipses stand for the gradient terms $-(\partial_i\sigma)^2 - 2\epsilon(\partial_i\mathcal{R})^2$. The interaction between curvature and isocurvature modes is given by the term $a^3(8\epsilon H/\kappa)\dot{\mathcal{R}}\sigma$. To guarantee perturbative analysis we require that $\sqrt{8\epsilon}/\kappa \ll 1$ [78, 141]. The mass of entropy perturbations is defined as $\mu^2 \equiv V_{NN} + \epsilon H^2 (\mathbb{R} + 6/\kappa^2)$, where the first term is obtained from the standard Hessian of the potential $V_{NN} \equiv \mathcal{N}^a\mathcal{N}^b(V_{ab} - \Gamma_{ab}^c V_c)$, the second and third terms correspond to the field space curvature and turning contributions respectively.

For shift-symmetric orbital inflation, we expect the isocurvature perturbations to be exactly massless, as in the toy model, and this is confirmed by

using (3.8) to show $\mu^2 = 0$. This implies that the quadratic action (3.9) has the combined shift symmetry (3.4), as in the toy model. The power spectra of perturbations in the massless limit can be directly estimated from the coupled evolution of perturbations [78]. When $\mu = 0$, the linearized system simplifies in the superhorizon limit, yielding

$$\mathcal{R}'_k = \frac{2}{\kappa}\sigma_k, \quad \sigma_k = \frac{H_*}{2\pi}, \quad (3.10)$$

where $*$ denotes evaluation at the time of horizon crossing. That is, on superhorizon scales the isocurvature perturbation quickly becomes a constant, and it sources the growth of \mathcal{R} . At the end of inflation, the primordial curvature perturbation can be expressed as $\mathcal{R}_k = \mathcal{R}_* + 2N_*\sigma_k/\kappa$, where the first term is the curvature perturbation amplitude at horizon-exit, and the second term comes from the isocurvature source. Thus these two contributions are uncorrelated with each other, and the dimensionless power spectrum for \mathcal{R} is given by

$$P_{\mathcal{R}} = \frac{H_*^2}{8\pi^2\epsilon_*} (1 + \mathcal{C}), \quad (3.11)$$

where $\mathcal{C} = 8\epsilon_*N_*^2/\kappa^2$ represents the contribution from isocurvature modes. This result agrees with the δN calculation for the toy model given in (3.6). The full calculation via the in-in formalism gives the same answer up to subleading corrections [78]. Note that the power spectrum is completely determined by the isocurvature perturbations if $\mathcal{C} \gg 1$, which corresponds to trajectories with a small radius κ or, equivalently, significant turning effects with $8\epsilon_* \ll \kappa^2 \ll 8\epsilon_*N_*^2$. Thus at the end of inflation, curvature perturbations are highly enhanced compared to the ones at horizon-exit. Meanwhile, the isocurvature power spectrum for $\mathcal{S} \equiv \sigma/\sqrt{2\epsilon}$ remains unchanged as $P_{\mathcal{S}} = \frac{H_*^2}{8\pi^2\epsilon_*}$. Therefore, the amplitude of the isocurvature perturbation is dynamically suppressed, *i.e.* $P_{\mathcal{S}}/P_{\mathcal{R}} \simeq 1/\mathcal{C} \ll 1$. The details of how $P_{\mathcal{S}} \neq 0$ can generate isocurvature components in the CMB are rather model-dependent, and one cannot automatically claim that a suppressed ratio $P_{\mathcal{S}}/P_{\mathcal{R}}$ is compatible with observations. However, if \mathcal{R} and \mathcal{S} contributed similarly to the curvature and isocurvature components in the CMB, the result is compatible with current constraints.

3.5 Phenomenology

We now turn to the observational predictions of shift-symmetric orbital inflation. For any positive \mathcal{C} , from (3.11), the tensor-to-scalar ratio can be

expressed as $r = 16\epsilon_*/(1 + \mathcal{C})$, and the scalar spectral index is $n_s - 1 \equiv \frac{d \ln P_{\mathcal{R}}}{d \ln k} = -2\epsilon_* - \eta_* + (d\mathcal{C}/dN)/(1 + \mathcal{C})$, where we used $d \ln k = dN$. Note that $\frac{\partial N_*}{\partial N} = -1$, since N_* counts the number of e-folds backwards. These predictions depend on the function $H(\theta)$. As in single field inflation, this function determines how slow-roll parameters ϵ and $\eta \equiv \epsilon'/\epsilon$ scale with N_* .

For concreteness, we consider models with $H \sim \theta^p$. Solving (3.8) for $\theta(N)$ yields² $\epsilon_* \simeq p/(2N_*)$ and $\eta_* \simeq 1/N_*$. The predictions for n_s and r are therefore well approximated by

$$n_s - 1 \simeq -\frac{p+1}{N_*} - \frac{4p}{\kappa^2 + 4pN_*}, \quad r \simeq \frac{8p\kappa^2}{N_*\kappa^2 + 4pN_*^2}. \quad (3.12)$$

We plot these results against the Planck 1σ and 2σ contours [7] in Fig. 3.2. N_* is taken to be between 50 and 60, and the radius κ^2 varies between 1 and 10^5 . The purple region is for $p = 1$, corresponding to the toy model (3.1), and we also show the predictions for $p = 0.5$ (red region), $p = 0.2$ (yellow region) and $p = 0.1$ (green region).

Notice that n_s and r only depend on the value of κ and are therefore insensitive to the details of the field metric. When $\kappa \rightarrow \infty$ one recovers the predictions of chaotic inflation with $V \propto \phi^{2p}$. Meanwhile as κ decreases, predictions are pushed downwards and to the left in this $n_s - r$ diagram. Therefore, in the case of power-law potentials only for small p do the predictions remain within the Planck contours. The interesting regime here is still the case with significant turning (small κ or $\mathcal{C} \gg 1$), where the final power spectrum $P_{\mathcal{R}} \simeq \frac{H_*^2 N_*^2}{\pi^2 \kappa^2}$ mainly has an isocurvature origin. Then the tensor-to-scalar ratio is given by $r = 2\kappa^2/N_*^2 = 16\epsilon_*/\mathcal{C}$, which is suppressed. The spectral index reduces to $n_s - 1 = -(p+2)/N_*$ which, for small p , lies in the sweet spot $n_s = 0.9649 \pm 0.0042$.

Another important observable is primordial non-Gaussianity, which is currently bounded by Planck through $f_{\text{NL}}^{\text{loc}} = 0.8 \pm 5$ [109]. There are examples in the literature of how $\mathcal{O}(1)$ local non-Gaussianity can arise in multi-field models, especially when the coupling between isocurvature and curvature modes is large [57, 142–144] - see [145] for a review. There are also examples of how small levels of non-Gaussianity can arise in multifield models [146–148]. However, in most cases a detailed analytic understanding of the size of the non-Gaussianity is lacking because the associated dynamics is non-linear and complicated. This is not the case in

²We note that for $0 < p < 1$ this toy model is not well defined as $\theta \rightarrow 0$, as can be seen in (3.7). This is not a problem as the inflationary period we are interested in occurs before that point is reached. The true underlying potential would have to be completed in some way. This is similar to case with say axion monodromy.




Figure 3.2: The analytical predictions (3.12) for (n_s, r) compared to the *Planck* 1σ and 2σ contours [7]. We show the predictions for wavenumbers which cross the horizon 50–60 e-folds before the end of inflation. The predictions for $n_s - r$ depend on the value of $\kappa \in [1, 1000]$, where the values (1, 2, 4, 8, 16, 32, 64, 128, 256) are depicted with thick lines (from bottom to top).

shift-symmetric orbital inflation, where we find that we can both easily satisfy the Planck constraint and crucially understand its origin analytically. The amplitude of local non-Gaussianity can be determined using the δN formalism. In a generic multi-field inflation model with curved field manifold, we have $f_{\text{NL}}^{\text{loc}} = \frac{5}{6} G^{ab} G^{cd} N_a N_c N_{bd} / (G^{ab} N_a N_b)^2$ [142, 149], where $G_{ab} = \text{diag}\{f(\rho), 1\}$ is the field space metric, N_a and N_{ab} are derivatives of N with respect to the fields (θ, ρ) . To gain some analytical understanding, here we still focus on models with $H \sim \theta^p$, where N can be expressed as $N = f(\rho)\theta^2/4p - p/2$. The amplitude of local non-Gaussianity then follows

$$f_{\text{NL}}^{\text{loc}} = \frac{5}{12} \eta_* \left[1 - \frac{\mathcal{C}^2}{(1 + \mathcal{C})^2} \frac{\kappa^2 \mathbb{R}}{2} \right], \quad (3.13)$$

where we used the relation $\mathcal{C} = 2p^2/(\epsilon_* \kappa^2)$. When $\kappa \rightarrow \infty$, we have $\mathcal{C} \rightarrow 0$ and $\mathcal{C}^2 \kappa^2 \rightarrow 0$. Thus the second term in (3.13) vanishes, which leads to the single field result $f_{\text{NL}}^{\text{loc}} = 5\eta_*/12$ as expected. The enhancement of non-Gaussianity is possible in the intermediate regime $\mathcal{C} \sim \mathcal{O}(1)$, where the

transfer from isocurvature to adiabatic modes is inefficient. In that case, $f_{\text{NL}}^{\text{loc}} \sim -5p\mathbb{R}/12$ can be large if the field space is highly curved.

For the interesting regime with $\mathcal{C} \gg 1$, the δN expansion is dominated by N_ρ and $N_{\rho\rho}$. This then leads to what, at first sight, appears as the counterintuitive result that $f_{\text{NL}}^{\text{loc}}$ is negligible and slow-roll suppressed

$$f_{\text{NL}}^{\text{loc}} \simeq \frac{5}{6} \frac{N_{\rho\rho}}{N_\rho^2} = \frac{5}{12} \eta_* \left(1 - \frac{\kappa^2 \mathbb{R}}{2} \right). \quad (3.14)$$

This is the same as happened in the calculation of the power spectrum: the contribution to the curvature perturbation sourced by the isocurvature modes dominates the final result. The bispectrum is found to be slow-roll suppressed, just like in single field inflation, but there are small corrections from the field space curvature, which violates Maldacena's consistency relation [39, 40]. We have recently confirmed this result via a scaling symmetry approach in [80].

3.6 Discussions

We have proposed a class of multi-field inflationary models that demonstrate a new type of attractor trajectory along the isometry direction in field space. Here the isocurvature modes become massless and freeze on super-horizon scales. Moreover, when the turning effects become significant, the curvature perturbations keep growing after horizon-exit and thus isocurvature modes are dynamically suppressed. As a consequence, these multi-field models yield the single-field-like phenomenology favored by observations.

Additional isocurvature perturbations will either decay if they are massive or freeze if they are light. Therefore, although our computations were done in a simple two-field setting, we expect the conclusions will continue to hold in multi-field extensions with more than two fields, provided that the number of additional light isocurvature fields is not too large.

We have shown and explained how in shift-symmetric orbital inflation, a negligible amount of local non-Gaussianity is produced. Here the isocurvature degree of freedom can be the dominant contribution to the bispectrum, but in such cases f_{NL} is slow-roll suppressed. This result teaches us a generic lesson: that in multi-field models, even if the isocurvature-to-adiabatic conversion is very efficient, the resulting non-Gaussianity can still be suppressed. A large coupling between curvature and isocurvature modes enhances the transfer of non-Gaussianity, but for this transfer to generate large non-Gaussianity, one needs sizable self-interactions affecting

the isocurvature field during horizon crossing [57, 141]. In this class of scenarios, however, the shift symmetry along the radial direction (3.4) has a role in suppressing the self-interactions of the isocurvature field (see [80]). Therefore, it is perfectly fine to study multi-field models with significant and sustained turning trajectories, without worrying about generating large non-Gaussianity.

Our model has important implications on the realization of inflation in UV-complete theories. Contrary to what is usually assumed, and as emphasized in [78], it is not always necessary to stabilize all compactification moduli, or to have a large mass hierarchy between the inflaton and other fields. The most problematic effects usually associated with multi-field effects – the generation of isocurvature perturbations and non-Gaussianity at unacceptable levels – cancel each other in the shift-symmetric orbital scenario ³. From an EFT point of view this can be traced back to the effect of derivative interactions among the curvature and isocurvature perturbations that are absent in single-field inflation. These are unavoidable on curved trajectories and curved field spaces and, therefore, ubiquitous in string compactifications.

³As already emphasized, large isocurvature perturbations at the end of inflation do not necessarily imply large isocurvature components in the CMB, the details of which are rather model dependent. Nevertheless, in this class of models the potentially significant generation of isocurvature modes in the CMB is automatically alleviated by the mechanism at play.

