



Universiteit  
Leiden  
The Netherlands

## **Growing up safely: Attachment-based interventions in child protection cases**

Asdonk, S. van der

### **Citation**

Asdonk, S. van der. (2020, September 15). *Growing up safely: Attachment-based interventions in child protection cases*. Retrieved from <https://hdl.handle.net/1887/136527>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/136527>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/136527> holds various files of this Leiden University dissertation.

**Author:** Asdonk, S. van der

**Title:** Growing up safely: Attachment-based interventions in child protection cases

**Issue date:** 2020-09-15



## Chapter 4

### Effectiveness of an attachment-based intervention for the assessment of parenting capacities in maltreating families: A randomized controlled trial

Sabine van der Asdonk, Whitney D. de Haan,  
Sheila R. van Berkel, Marinus H. van IJzendoorn,  
Ralph C. A. Rippe, Carlo Schuengel, Chris Kuiper,  
Ramón J. L. Lindauer, Mathilde Overbeek, &  
Lenneke R. A. Alink

Published in *Infant Mental Health Journal*, 1-15  
doi:10.1002/imhj.21874

## Abstract

Even though Parenting Capacity Assessments (PCAs) are essential for child protection services to support placement decisions for maltreating families, presently no evidence-based PCA protocols are available. In this randomized controlled trial, we tested the quality of an attachment-based PCA protocol based on Video-feedback Intervention to promote Positive Parenting and Sensitive Discipline (VIPP-SD). We recruited 56 parent-child dyads ( $M^{\text{age}}$  children = 3.48 years) in family residential clinics that conduct PCAs to support placement decisions. After pre-test, families were randomized to receive the Regular Assessment Procedure (RAP) ( $n = 28$ ), or an additional assessment based on VIPP-SD ( $n = 28$ ). An immediate post-test and a 10-month follow-up were conducted. Multilevel models showed that therapists felt equally confident about their recommendations regarding child placement for both groups. Moreover, children in the VIPP-SD group did not show fewer behavior problems and did not experience recurring child maltreatment less often than children in the RAP group. Finally, parents who received VIPP-SD were generally evaluated as less capable than parents who received RAP. Thus, we found no evidence that PCAs incorporating the VIPP-SD protocol outperformed PCAs as usual. Possible reasons for this lack of effectiveness, such as insufficient power due to small sample size, are discussed.

*Keywords:* child maltreatment, parenting capacity, placement decisions, attachment-based intervention, RCT

## Introduction

Child maltreatment constitutes a major public health concern; it affects millions of children worldwide and is associated with a broad spectrum of negative and long-lasting developmental outcomes (Gilbert et al., 2009). When child maltreatment is suspected or substantiated in a family, child protection services may consider out-of-home placement. Essential for deciding whether or not a child should be placed out of home are assessments of parenting capacities (PCAs). Unfortunately, currently no evidence-based methods for PCAs are available. Considering the complexity of placement decisions and their impact on the lives of children and their parents, valid PCA protocols are needed to effectively support placement decisions. One proposal for improved PCA protocols is that parenting capacities should be evaluated based on parents' response to an evidence-based intervention (Harnett, 2007). Building on this proposal and existing theories regarding child maltreatment and its etiology, several researchers have suggested using an attachment-based intervention for this purpose (Cyr & Alink, 2017; Cyr et al., 2012; Lindauer, Bakermans-Kranenburg, Van IJzendoorn, & Schuengel, 2010). Parallel to a recent Canadian study (Cyr, Paquette, Dubois-Comtois, & Lopez, 2015), the current randomized controlled trial (RCT) is among the first to empirically evaluate whether the quality of placement decisions can be improved by structurally evaluating parents' response to an attachment-based intervention.

### Parenting capacity assessments

Although a number of guidelines have been developed for PCAs (e.g., American Psychological Association, 1998; Budd, 2001), empirical evidence on the effectiveness of these assessments is scarce (Vischer, Grietens, Knorth, & Mulder, 2017). In addition, several limitations of PCAs in practice have been reported: These assessments often concern only one time point, do not include observations of parent-child interactions in the home environment, and emphasize parents' weaknesses more than their strengths (Budd, 2001, 2005). In order to improve the quality of these assessments, several researchers have suggested using a more structured and dynamic approach (Cyr et al., 2012; Harnett, 2007; Lindauer et al., 2010). The approach they propose consists of structurally assessing parents' capacity to change relevant parenting behavior by evaluating parents' response to an evidence-based intervention. The intervention should be conducted in a short time period, include systematic observations of the parent-child relationship in the home setting, and focus on the strengths of parents. It is argued that such an approach would be particularly valuable for cases that are equivocal and where an initial (risk) assessment does not demonstrate a clear picture of the child's well-being (Harnett, 2007).

### Focus on attachment-based interventions

Based on the existing knowledge on maltreatment, parents' response to an attachment-based intervention aimed at improving parental sensitivity would provide highly relevant information for a PCA (Cyr & Alink, 2017; Cyr et al., 2012; Lindauer et al., 2010). Parental sensitivity, which refers to parents' ability to notice, interpret, and respond to child signals in an appropriate and prompt manner while adapting to the child's changing developmental needs (Ainsworth, Bell, & Stayton, 1974), is universally considered as an important indicator

of positive child development (Ainsworth et al., 1974; Mesman, Van IJzendoorn, & Bakermans-Kranenburg, 2012) and has often been identified as relevant for PCAs (Cyr & Alink, 2017; Cyr et al., 2012; Lindauer et al., 2010; Teti & Candelaria, 2002; Ward, Brown, & Hyde-Dryden, 2014; White, 2005). Several studies have shown that attachment-based interventions aimed at improving parental sensitivity have positive effects for maltreating parents, or parents at risk for maltreatment, and their children (Bernard et al., 2012; Moss et al., 2011; Negrao, Pereira, Soares, & Mesman, 2014; Steele, Murphy, Bonuck, Meissner, & Steele, 2019). These studies found positive outcomes both at the level of the parent-child relationship (i.e., increased quality of parental sensitivity and the attachment-relationship and less harsh discipline) and at the level of child development (i.e., improved self-regulation skills and fewer behavioral and emotional problems). Besides their focus on improving parental sensitivity, these interventions have in common that they are short-term, include video-feedback, and focus on parents' strengths. The effectiveness of these interventions has been strongly supported by empirical evidence, which increases the informational value of response to intervention or lack thereof (Cyr et al., 2012; Harnett, 2007; Lindauer et al., 2010). A recent Canadian study found that implementing a PCA protocol based on an evidence- and attachment-based video-feedback intervention enabled clinicians to better predict reoccurrences of child maltreatment (Cyr et al., 2015). Although these results are promising, more studies are necessary, (1) to establish these effects more firmly, and (2) to evaluate whether such a protocol could also be effective in other countries with different child protection systems.

### **Evaluating the quality of placement decisions**

The quality of a procedure for PCAs depends on the reliability and validity of subsequent placement decisions. Relating this to the current study, the reliability of the proposed assessment approach has recently been investigated in a vignette study where we demonstrated that providing decision-makers with information about parents' response to an attachment-based intervention can lead to increased agreement on placement decisions (Van der Asdonk et al., 2019). This is an important foundation for the current study, because sufficient reliability is a prerequisite for strong validity. Although validity might be a difficult construct to appropriately evaluate in this context, the validity of placement decisions should at least be reflected in (a) professionals' confidence that their recommendation regarding the child's placement is accurate (face validity) and (b), because the main goal of child protection services is to act in the best interest of children's well-being, an improved quality of life for children (predictive validity). Importantly, several longitudinal studies have shown that reunifications of maltreated children with their parents are often not stable over time and that some parents will abuse or neglect their children again in the future (Biehal, Sinclair, & Wade, 2015; Lutman & Farmer, 2013). This indicates that severe parenting problems may still exist and children's quality of life does not always improve following placement decisions. Moreover, mixed results have been reported regarding children's mental well-being, with some studies showing worse outcomes for children who were reunified with their parents than for children who remained in out-of-home care (Biehal et al., 2015), and other studies finding opposite results (Lloyd & Barth, 2011). These findings do not only emphasize the complexity of placement decisions, but also stress the need for studies that take children's well-being into account when evaluating methods to improve the quality of decisions. Therefore, in the current study we looked at reoccurrences of child maltreatment and children's emotional and behavioral problems as

indicators of their quality of life following placement decisions. In addition, we looked at the severity of parenting problems for birth parents following placement decisions as a proxy of children's well-being.

### **Reasoning biases in decision-making**

One aspect that has been found to compromise the quality of decision-making is related to common reasoning biases in decision-making (Kahneman, Slovic, Slovic, & Tversky, 1982). In a study that investigated professional reasoning in child protection reports, it was shown that professionals can be prone to hold on to their initial judgements about a family, even when they are faced with new and contradictory evidence (Munro, 1999). One way to prevent such intuitive reasoning mistakes might be by providing more concrete, relevant, and objective information for professionals to guide their decision-making. Such concrete information may be produced by a structured, attachment-based assessment protocol (Cyr et al., 2012; Lindauer et al., 2010), because it informs professionals about parents' ability to benefit from an intervention to improve important parenting skills. If this information can indeed reduce reasoning biases in child protection cases, this should be reflected by a higher tendency of professionals to change their initial judgements after receiving additional information provided by the assessment protocol.

### **Current study**

The current RCT tested the effect of evaluating parents' response to an attachment-based intervention on the quality of placement decisions in the Netherlands. For this purpose, we developed a procedure for PCAs based on the Video-feedback Intervention to promote Positive Parenting and Sensitive Discipline (VIPP-SD; Juffer, Bakermans-Kranenburg, & Van IJzendoorn, 2017), an assessment procedure that is similar to the protocol developed by Cyr et al. (2012). VIPP-SD is an evidence-based intervention for improving parental sensitivity, also for families at risk for maltreatment (Juffer et al., 2017). We implemented VIPP-SD in four family residential clinics throughout the Netherlands to which families are referred for a PCA in the context of a decision regarding a possible out-of-home placement or reunification with their child(ren). At the end of the families' assessment period, the therapist provides a recommendation on which the child's subsequent placement is usually based. We hypothesized (1) that recommendations about the necessity of out-of-home placement at the start of families' assessment period were more often modified by therapists after VIPP-SD than after the regular assessment procedure, (2) that therapists felt more confident on their recommendations based on VIPP-SD than on their recommendations based on the regular assessment procedure, (3) that children for whom a recommendation was based on VIPP-SD showed fewer emotional and behavioral problems than children for whom a recommendation was based on the regular assessment procedure, and (4), for the group of children who returned to their parents after the assessment, that there were fewer reoccurrences of child maltreatment in families for whom a recommendation was based on VIPP-SD than in families for whom a recommendation was based on the regular assessment procedure. In addition to these primary research questions, we explored whether the evaluation of parenting capacities differed between families who received VIPP-SD and families in the regular assessment procedure. Finally, for the group of children who returned to their parents after the assessment, we explored whether families for whom a recommendation was made based on VIPP-SD received

less intensive parenting support, indicating less severe parenting problems, after leaving the clinic than families for whom a recommendation was made based on the regular assessment procedure.

## Method

### Participants

Fifty-six families participated in this study. Recruitment took place from May 2015 until December 2017 in four family residential clinics that are located in different regions of the Netherlands. These clinics constitute a unique setting in the Dutch child protection system which enables highly intensive observation and treatment of families for whom a placement decision is being considered (either in the context of an out-of-home placement or a reunification). Families usually reside in these clinics for 24 hours a day on weekdays (and, if necessary, during weekends) for a period of two to three months, during which they are regularly observed by family workers and receive highly intensive support at all levels of the family system. The evaluation of families' trajectory at the clinics is used as a recommendation for the children's court judge or involved family guardian, depending on who referred the family to the clinic. For the current study, families were selected based on the following inclusion criteria: (1) the family was referred to the clinic for an evaluation of their parenting capacities in the context of a decision regarding out-of-home placement or reunification with their child(ren), (2) the child's age was between 6 months and 7 years, (3) the primary caregiver spoke a basic level of Dutch, (4) the primary caregiver did not have a (severe) intellectual disability that affected his or her ability to understand the instructions of the intervention, and (5) the primary caregiver did not have severe mental health problems which required acute intervention. If a family that met our inclusion criteria started their assessment in the clinic, one of the staff members informed the researchers so that they could explain the study to the families. The recruitment goal was set on 71 families. A power analysis in G\*Power 3.0 (Faul, Erdfelder, Lang, & Buchner, 2007) conducted prior to this study indicated that for 60 randomized participants and two-tailed significance tests at  $\alpha = .05$ , power to detect medium effects on primary study outcomes would be .80. The majority of approached families (79%) agreed to participate. We asked the primary caregiver to participate. If there was more than one child in the family, the youngest child between 1 and 7 years was invited to participate. Overall, 41 families (73% of enrolled families) completed the post-test. All families, except for those who indicated they did not want to participate anymore ( $n = 6$ ), were approached again for follow-up. The final follow-up sample consisted of 34 dyads (61% of the original sample). See Figure 4.1 and Appendix 4.A for a more detailed description of the sample flow.

About half of the children (55%) from the original sample ( $N = 56$ ) were boys, and the children were on average 3.48 years old ( $SD = 1.74$ ). Primary caregivers had an average age of 32.32 years ( $SD = 6.43$ ) and were primarily mothers (93%). Most parents were single parents (64%). For 55% of the families, referral to the clinic concerned assessment regarding a possible reunification with the participating child.






Figure 4.1. Flow chart of sample throughout the study.

## **Procedure**

This research was approved by the Dutch Central Committee on Research Involving Human Subjects, the ethical review board of the Institute of Education and Child Studies at Leiden University, and the Ethics Committee for Legal and Criminological Research at Vrije Universiteit Amsterdam. The study is registered in the Netherlands Trial Register (Trial NL7632). The pre-test was conducted as soon as was possible after the parent(s) signed informed consent for the study and consisted of a two-hour appointment in a lab setting at the clinic. In addition, the therapist responsible for the family's recommendation on future placement filled out a short questionnaire about the family. After pre-test, families were randomly assigned to either VIPP-SD ( $n = 28$ ) or the regular assessment procedure ( $n = 28$ ). Randomization was done by one of the researchers with a computer-generated blocked randomization sequence that was stratified for the four clinics, so that for each clinic the families were equally divided over the two conditions. The post-test was similar to the pre-test and took place on average 9.5 weeks after pre-test. Initially, we aimed to conduct two follow-up assessments at 8 and 12 months. However, due to practical issues making it complicated to complete two follow-up assessments with the families (i.e., phone numbers changed, multiple efforts required to reach parents at their homes for one appointment) it was decided to conduct only one follow-up assessment for each family. This assessment took place approximately 10 months after post-test (range: 8 – 16 months) and consisted of a 1-hour home visit during which the primary caregiver filled out questionnaires and a semi-structured interview was conducted by a trained researcher. Four participating children lived in a foster family at the time of the follow-up assessment. To contact the foster family, parental permission was required, which was granted for two of these children. Foster parents were asked to fill out questionnaires through email – preceded by a telephone appointment to provide information on the study and to request informed consent. After pre- and post-test, families received a gift card of 20 euros, and after follow-up they received a small gift.

## **Assessment of parenting capacities**

### **Regular assessment procedure**

The regular assessment procedure (RAP) consisted of care as usual at one of the clinics. Although the four clinics were not completely uniform in their treatment programs, the general structure was similar: all parents received various forms of treatment aimed at improving family dynamics, including observations of parent-child interactions, group sessions with other parents, and individual sessions for the parent(s). Some parents and children additionally received specialized therapy based on their individual needs (e.g., trauma therapy or emotion-regulation training). To limit similarities with VIPP-SD, none of the families in the RAP condition received video feedback. The standard period for family treatments differed between the clinics (range: 8 - 12 weeks). In all clinics, an evaluation was conducted at the end of the assessment period in which the therapist and involved family workers evaluated parents' progress during their treatment in the clinic. This evaluation resulted in a recommendation that was provided to either the involved family guardian, social worker, or children's court judge, depending on who had referred the family to the clinic. To be able to use the content of this evaluation for research purposes, we composed a structured parenting capacity evaluation form and asked the therapists to fill out this form for each family at the end of the

family's assessment period. This form consists of 15 items on a six-point Likert-scale, of which five items concern general aspects of the therapeutic relationship and parents' attitude during the intervention (e.g., "Was the parent open to change his/her behavior?"), and ten items concern changes in parents' behavior following the intervention (e.g., "The parent shows progress in adequately responding to negative child signals, such as crying and resistant or naughty behavior"). The internal consistency of the assessment form was high ( $\alpha$  for all 15 items = .93). In each clinic, there was one therapist (with a Master degree) who was responsible for families' recommendations. The family workers generally had a Bachelor degree and worked directly with the families.

### **VIPP-SD**

We slightly adapted VIPP-SD by adding an explicit evaluation of parenting capacities at the end of the intervention. Thus, VIPP-SD in this study consisted of (1) an intervention and (2) an assessment form. For the intervention part, either VIPP or VIPP-SD (Juffer, Bakermans-Kranenburg, & Van IJzendoorn, 2008) was delivered to the family, depending on the child's age: parents of a 6- to 12-months-old received VIPP ( $n = 3$ ) and parents of a child older than 12 months received VIPP-SD. VIPP focuses on improving parental sensitivity through video feedback and consists of six sessions in which the parent-child dyad is videotaped during common, daily interactions such as playing together or a meal. VIPP-SD additionally focuses on improving sensitive discipline strategies of the parent. For a detailed overview of VIPP(-SD), see Juffer et al. (2008). VIPP-SD was delivered by family workers at the clinics who were trained to be VIPP-interveners for this study. For six parent-child dyads, a trained (assistant) researcher provided VIPP-SD because no trained family worker was available at that time. Each VIPP-SD trajectory was monitored during supervision meetings with one of the trained researchers. After the final session, we asked the intervener to fill out the parenting capacity assessment form to evaluate parents' response to VIPP-SD and to integrate the assessment form in the evaluation of the family at the end of their treatment period (as described above in the RAP section). Finally, similar to the RAP group, we asked the therapists to fill out an evaluation form for their recommendation regarding the child's placement.

## **Instruments**

### **Recommendation regarding child placement**

At pre- and post-test, we asked the therapist to indicate the current recommendation for this family: (a) a supervision order, but the child can stay or be reunified with its parent(s), (b) (extended) supervision order and out-of-home placement of the child – in own network, (c) – in foster care, (d) – in residential care, or (e) other. We dichotomized these items into (0) no out-of-home placement versus (1) out-of-home placement.

### **Therapists' confidence in their recommendation**

After the therapists gave their recommendation about the child's placement at pre- and at post-test, we asked them to indicate on a ten-point scale how confident they felt about their recommendation. A higher score indicated more confidence.

### **Children's emotional and behavioral problems**

The preschool version of the Child Behavior Checklist (CBCL) was used to assess children's emotional and behavioral problems (Achenbach & Rescorla, 2000). The CBCL consisted of 100 items regarding the child's behavior in the past two months which are rated on a three-point scale (0 = not true, 1 = somewhat or sometimes true, and 2 = very or often true). We asked the primary caregiver to fill out the CBCL at pre- and post-test and at the follow-up assessment. For two children who lived in foster care at follow-up, the involved foster parent filled out the CBCL. The CBCL has been proven valid and reliable (Achenbach & Rescorla, 2000). Sum scores for total problems ( $\alpha$  in current sample = .98) were used. Because of an extremely high number of missing post-test scores for the CBCL (71% of the forms were missing, compared to 46% for both pre-test and follow-up), we decided not to use the post-test data, so that only CBCL scores at pre-test and follow-up were compared.

### **Recurring child maltreatment**

For those children who were living with their parents at follow-up ( $n = 32$ ; 94%), we assessed whether there had been reoccurrences of child maltreatment in the 10 months that followed leaving the clinic. For this purpose, a trained (assistant) researcher conducted the Maternal Maltreatment Classification Interview (MMCI; Cicchetti, Toth, & Manly, 2003) with the primary caregiver. The MMCI is a semi-structured interview during which the primary caregiver is asked about events of child abuse and the family's contact with child protection services. We used the version that was translated into Dutch by Reijman et al. (2014). We asked the primary caregiver to answer the questions about the 10 months after they had left the clinic. The MMCI was coded using the Modified Maltreatment Coding System (English, Bangdiwalab, & Runyan, 2005). After coding, each family received a score reflecting whether child maltreatment had reoccurred (1) or not (0). Two trained (assistant) researchers double-coded all interviews, reliability was excellent ( $\kappa = 1.00$ ,  $n = 28$ ).

### **Intensity of parenting support at follow-up**

During the MMCI with the biological primary caregiver at follow-up, we additionally asked about the involvement of professional care specifically aimed at parenting since they left the clinic. We coded their answers on a seven-point scale, ranging from 0) no extra care (other than standard post-treatment care), to 6) parenting support is currently present more than once per week. All interviews were independently coded by two trained coders, reliability was high (Intraclass Correlation Coefficient [ICC], single measures = .98,  $n = 26$ ).

### **Evaluation of parenting capacities**

We used the parenting capacity assessment form that was developed for this study to get an indication of parents' capacities following VIPP-SD or RAP as evaluated by the involved therapist or VIPP-intervener (for a more detailed description of this form, see procedure section of this paper). Higher average scores across the 15 items in the analyses indicated that the involved therapist or intervener evaluated the parent as more capable.

### **Statistical analyses**

Data inspection revealed that the numerical variables approached a normal distribution after winsorizing outliers more than  $\pm 3.29$  standard deviations from the mean. One family could

only be reached for follow-up after 23 months. For this family, we decided to still use the data retrieved from the MMCI (recurring maltreatment and intensity of parenting support), as the interview specifically aimed at the first 10 months after leaving the clinic. CBCL scores for this family were not used, because this construct is likely more difficult to rate objectively in retrospect. Little's MCAR test (Little, 1988) showed that values were missing completely at random ( $\chi^2(138) = 139.97, p = .44$ ). To follow an intent-to-treat approach and maximize power, multilevel multiple imputation (Rubin, 1987; White, Carpenter, & Horton, 2012) was performed on the data ( $N = 56$ ) in RStudio (version 1.1.463) (RStudioTeam, 2016). See Appendix 4.B for a detailed overview of imputation procedures.

For therapists' confidence in their recommendation and children's behavioral and emotional problems, three-level linear mixed effect models accounting for repeated measures over time (level 1) and nesting of families (level 2) within clinics (level 3) were incrementally compared using likelihood ratio test for imputed datasets in the *mitml* package. The final model included the fixed effects of time (coded as 0 = pre-test, 1 = post-test/follow-up), the main effect for condition (coded as 0 = RAP, 1 = VIPP-SD), and the interaction between time and condition. For modifications in therapists' recommendation regarding child placement, a similar model was fitted with a binomial family structure. For recurring maltreatment, generalized linear mixed effect models accounting for nesting of families (level 1) in clinics (level 2) were performed with the *lme4* package with a binomial family structure. We compared models incrementally with likelihood ratio tests. We explored the influence of two potential covariates: (1) time between post-test and follow-up (because of the large range in time) and (2) children's age (because of the relatively broad age range in our study). However, because neither of these covariates affected any of the results, we reported only the most parsimonious models without covariates. After testing our main hypotheses, we explored potential differences between VIPP-SD and RAP families in the evaluations of their parenting capacities at post-test and in the intensity of parenting support at follow-up. For this purpose, we compared two linear mixed effect models accounting for the nesting of families within clinics. Significance of model and parameter estimates was determined at  $\alpha = .05$ . Complete case analyses yielded similar outcomes (available upon request). Odds ratios were computed as estimates of effect sizes for dichotomous outcome variables (i.e., modifications in therapists' recommendation regarding child placement and recurring child maltreatment), and beta's were used as estimates of effect sizes for continuous outcome variables (see e.g., Lorah, 2018).

## Results

### Preliminary analyses

For an overview of demographic and outcome variables of the total sample, see Table 4.1. The majority of children (88%) were living with their parent(s) at the follow-up assessment. For 94% of the children, their living situation at follow-up was consistent with the final recommendation the family received in the clinic. There were no differences between the VIPP-SD and RAP groups at pre-test variables (see Table 4.1). Moreover, comparisons on demographic and target variables between families who dropped out during the research project and families who completed the project showed that there were no significant differences ( $p$ 's  $> .10$ ). Pooled correlations between all variables of interest are displayed in Table 4.2.

**Table 4.1**  
*Descriptive statistics for demographic and target variables.*

	All families		VIPP-SD		Regular Assessment		$F/\chi^2$ <sup>b</sup>
	$M^a$	$SD^a$	$M^a$	$SD^a$	$M^a$	$SD^a$	
<b>Demographic</b>	N = 56		n = 28		n = 28		
Age child	3.48	1.74	3.78	1.88	3.19	1.57	1.04
Gender child (% boys)	55%		54%		57%		0.07
Age parent	32.32	6.43	33.69	7.10	31.50	7.35	0.20
Gender parent (% female)	93%		93%		93%		<0.00
Number of siblings	2.52	1.39	2.44	1.67	2.61	1.09	0.26
<b>Outcome variables at pre-test</b>							
Therapists' recommendation (% out-of-home placement)	50%		46%		54%		0.35
Therapists' confidence	6.07	1.57	5.91	1.51	6.08	1.69	0.66
CBCL total	36.56	28.87	36.66	29.45	36.47	29.27	0.16
<b>Outcome variables at post-test</b>							
Therapists' recommendation (% out-of-home placement)	29%		36%		22%		
Therapists' confidence	7.74	1.04	7.92	0.97	7.59	1.10	
Evaluation of parenting capacities	4.04	0.75	3.80	0.83	4.30	0.57	
<b>Outcome variables at follow-up</b>							
CBCL total	36.92	22.43	44.23	25.43	31.32	18.71	
Recurring maltreatment (N)	n = 9		n = 5		n = 4		
Intensity of parenting support	4.02	2.22	3.15	2.26	4.56	2.07	

<sup>a</sup>unless indicated otherwise

<sup>b</sup>Chi-square tests and one-way ANOVAs were performed to test whether there were pre-test group differences between families in the VIPP-SD and RAP conditions

**Table 4.2**  
*Pooled Pearson Correlations between study variables of interest (N = 56).*

	2	3	4	5	6	7	8	9	10	11	12
<b>Pre-test variables</b>											
1. Age child	.12	.20	-.14	-.12	.04	-.02	-.06	-.09	.14	-.19	-.04
2. Gender child <sup>1</sup>		.03	-.06	.07	-.04	.21	.01	.02	.00	-.09	-.02
3. Age parent			.14	-.12	.10	.16	-.04	-.18	.27	.20	.18
4. Therapists' recommendation <sup>2</sup>				.16	.09	.33*	-.06	-.26	-.01	.11	.27
5. Therapists' confidence					.03	.32*	.18	.01	-.07	-.09	.11
6. CBCL total						.06	-.07	.06	.28	.05	.15
<b>Post-test variables</b>											
7. Therapists' recommendation <sup>2</sup>							.03	-.40*	.03	-.06	.08
8. Therapists' confidence								.03	-.02	-.04	-.20
9. Evaluation of parenting capacities									-.01	-.05	-.04
<b>Follow-up variables</b>											
10. CBCL total										.13	.03
11. Recurring child maltreatment											.06
12. Intensity of parenting support											

\* $p < .05$

<sup>1</sup>coded as girl = 1

<sup>2</sup>coded as out-of-home placement = 1

**Table 4.3** Fixed effects of linear mixed models, dependent (binomial) variable is therapists' recommendation regarding child placement ( $N = 56$ ).

	<b>Model 1</b> B (SE)	<b>Model 2</b> B (SE)	<b>Model 3</b> B (SE)	<b>Model 4</b> B (SE)	OR	OR	OR
Fixed effects							
(Intercept)	0.41 (0.06)*	0.69 (0.14)*	0.67 (0.15)*	0.84 (0.19)*			
Time		-0.19 (0.08)*	-0.19 (0.08)*	-0.30 (0.11)*	0.83	0.83	0.74
Condition			0.04 (0.12)	-0.30 (0.28)	1.04	1.04	0.74
Time*Condition				0.23 (0.16)			1.26
Variance components							
Clinic level <sup>1</sup>							
Family level	1.09 (0.31)	1.37 (0.36)	1.36 (0.36)	1.50 (0.44)			
Change in model fit ( $F$ )		4.52*	0.14	1.76			

Time: 0 = pre-test, 1 = post-test; Condition: 0 = Regular Assessment Procedure, 1 = VIPP-SD

\* $p < .05$

<sup>1</sup>Nesting of families within clinics could not be fitted with this model



**Table 4.4**  
*Fixed effects of linear mixed models for therapists' confidence in their recommendation and total CBCL scores (N = 56).*

	Model 1	Model 2	Model 3	Model 4
	B (SE)	B (SE)	B (SE)	B (SE)
		$\beta$	$\beta$	$\beta$
<b>DV: Therapists' confidence</b>				
Fixed effects				
(Intercept)	6.94 (0.21)*	4.43 (0.48)*	4.33 (0.51)*	4.58 (0.66)*
Time		1.67 (0.29)*	1.67 (0.29)*	1.50 (0.40)*
Condition			0.19 (0.33)	-0.31 (0.98)
Time*Condition				0.33 (0.59)
Variance components				
Clinic level	0.27 (0.16)	0.30 (0.16)	0.30 (0.16)	0.29 (0.16)
Family level	0.01 (0.05)	0.51 (0.22)	0.51 (0.22)	0.51 (0.21)
Residual	1.62 (0.07)	1.27 (0.09)	1.27 (0.09)	1.28 (0.09)
Change in model fit (F)		29.43*	0.47	0.46
<b>DV: CBCL total</b>				
Fixed effects				
(Intercept)	38.17 (3.43)*	38.13 (10.00)*	35.24 (10.47)*	44.34 (13.75)*
Time		0.03 (6.06)	0.03 (6.06)	-6.04 (8.13)
Condition			5.78 (6.79)	-12.42 (20.41)
Time*Condition				12.14 (12.15)
Variance components				
Clinic level	0.72 (1.44)	0.72 (1.44)	0.85 (1.64)	0.86 (1.64)
Family level	12.99 (3.73)	12.97 (3.79)	12.71 (4.00)	13.20 (3.58)
Residual	22.62 (2.17)	22.61 (2.26)	22.61 (2.26)	22.20 (2.07)
Change in model fit (F)		0.40	0.82	1.35

Time: 0 = pre-test, 1 = post-test (for therapists' confidence) or follow-up (for CBCL)

Condition: 0 = Regular Assessment Procedure, 1 = VIIPP-SD

\* $p < .05$

**Table 4.5**  
*Fixed effects of generalized linear mixed models, dependent variable is recurring child maltreatment (N = 56).*

	<b>Model 1</b> <i>B (SE)</i>	<b>Model 2</b> <i>B (SE)</i>	<b>OR</b>
Fixed effects			
(Intercept)	0.35 (0.09)*	0.30 (0.12)*	
Condition		0.10 (0.17)	1.11
Variance			
Clinic level	0.22 (0.24)	0.23 (0.24)	
Change in model fit ( <i>F</i> )		0.32	

Condition: 0 = Regular Assessment Procedure, 1 = VIIPP-SD

\* $p < .05$

### **Modifications in therapists' recommendations regarding child placement**

For modifications in therapists' recommendations regarding child placement, the unconditional growth model showed the best fit (see Table 4.3). Only the fixed effect of time was significant and indicated that compared to pre-test, therapists' recommendations at post-test more often favored that the child could stay with its parents, see Table 4.1. Recommendations for VIPP-SD families were not more often modified than recommendations for RAP families.

### **Therapists' confidence in their recommendation**

Therapists' confidence in their recommendation varied more over time ( $ICC = .81$ ) and between therapists ( $ICC$  clinic level =  $.14$ ) than between families ( $ICC = .01$ ). The unconditional growth model including the fixed effect of time showed the best fit and indicated that for both conditions, therapists felt more confident on their recommendation at post-test than at pre-test (see Tables 4.1 and 4.4). The fixed effect of the interaction between time and condition was not significant, which indicates that therapists did not feel more confident over time about their recommendations for VIPP-SD families than about their recommendations for RAP families.

### **Behavioral and emotional problems**

Children's behavioral and emotional problems varied more over time ( $ICC = .61$ ) and between families ( $ICC = .36$ ) than between clinics ( $ICC = .03$ ). Adding fixed effects to the unconditional means model did not improve model fit (see Table 4.4). This indicates that generally, children did not change over time in their level of behavioral and emotional problems. Moreover, children who received a placement decision after participating in VIPP-SD did not show a stronger decrease in behavioral and emotional problems over time than children who received RAP.

### **Recurring child maltreatment**

For recurring child maltreatment, the unconditional means model showed the best fit, see Table 4.5. This indicates that there were no differences in experienced recurring child maltreatment between children in the VIPP-SD group and children in the RAP group.

### **Exploratory analyses**

For the evaluation of parenting capacities at post-test, the fixed effect of condition improved model fit compared to the empty model ( $F(1, 1012.76) = 5.25, p = .02; B = -0.51, \beta = -.48, SE = 0.23, p = .02$ ). The fixed effect estimate indicates that on average, families in the VIPP-SD group were evaluated as less capable than families in the RAP group (see Table 4.1). With respect to the intensity of parenting support at follow-up, there was no difference between VIPP-SD and RAP families ( $F(1, 255.89) = 1.88, p = .17$ ; fixed effect for condition:  $B = -1.05, \beta = -.33, SE = 0.75, p = .17$ ).

## **Discussion**

PCAs are an important basis for placement decisions, although thus far no evidence-based methods for this purpose are available. This study was among the first to investigate through

an RCT whether the quality of placement decisions for maltreating families could be improved by implementing a structured, attachment-based PCA. We investigated this in four Dutch family residential clinics that conducted PCAs in the context of a potential out-of-home placement decision – a setting which is unique in the Dutch child protection system. In addition to the regular assessment procedure, half of the families received an assessment based on VIPP-SD, an attachment-based video-feedback intervention (Juffer et al., 2017). We evaluated the quality of the assessment procedures in terms of face validity (therapists' confidence that their recommendation regarding the child's placement was accurate) and predictive validity (children's well-being at follow-up). In addition, we hypothesized that therapists would be more reluctant to change their initial recommendations for families who received a regular assessment procedure than for families who received VIPP-SD. None of our hypotheses were confirmed in this study: therapists did not feel more confident about their recommendations for families whose assessment was based on VIPP-SD, neither did they modify their initial recommendations more often for families who received an assessment based on VIPP-SD than for families who received the regular assessment procedure. Moreover, children in families who received an assessment based on VIPP-SD did not differ from children in families who received the regular assessment procedure with respect to (a) their level of problem behavior and (b) their chance of experiencing recurring child maltreatment in the 10 months following the placement decision. Thus, we did not find evidence that implementing VIPP-SD in PCAs for maltreating families increased the validity of placement decisions.

In addition to our main hypotheses, we explored whether there were differences between families who received VIPP-SD and families in the regular assessment group in the evaluation of their parenting capacities at the end of the assessment period and in the intensity of parenting support they received in the 10 months following the assessment. Although we did not find any group differences on the latter, we were surprised to find that parents who received VIPP-SD were evaluated as less capable by their interveners than parents in the regular assessment procedure. Even though this could indicate that parents who received VIPP-SD actually were less capable, the lack of other group differences (e.g., chance of recurring child maltreatment or intensity of parenting support at follow-up) contradicts this interpretation. One explanation might be that the VIPP-interveners were more conscious of the parenting capacities that needed to improve (i.e., aspects of parenting related to sensitivity and sensitive discipline), which may have made them more critical evaluators of these aspects than therapists who assessed families in the regular assessment procedure. It should be noted here that the interveners and therapists could not be blind to families' condition, and due to practical considerations we did not conduct an initial evaluation of parenting capacities. These aspects make it complicated to derive any strong conclusions from this finding.

The absence of beneficial effects of the VIPP-SD protocol for PCAs in this study is unexpected, given that many researchers have argued to use attachment-based interventions in PCAs (Cyr & Alink, 2017; Cyr et al., 2012; Lindauer et al., 2010) and two recent randomized studies have provided initial evidence that such a procedure can lead to a higher quality of placement decisions (Cyr et al., 2015; Van der Asdonk et al., 2019). An explanation for the lack of effects in the current study could be related to the quality of the regular assessment procedure in the Dutch clinics. When families are referred to these clinics, they are residing there for a couple of months during which they are observed by experienced family workers and receive various treatment forms adapted to their individual needs. Families and family

workers are thus highly involved in the treatment process. It is possible that within the context of this highly intensive program, VIPP-SD does not contribute to an improved PCA, because therapists responsible for families' placement recommendations might already be able to form a clear picture of the parenting capacities based on the regular intensive assessment procedure. The fact that therapists generally felt quite confident about their recommendations at post-test and children's living situation at follow-up was in most cases still consistent with the therapist's recommendation, might underscore this assumption. It should be noted that this setting for PCAs is quite unique to the Netherlands and therefore the results of this study cannot be directly generalized to other countries or compared to the recent Canadian study, where the regular assessment procedure was far less intensive as it included no more than twelve 3-hour home visits (Cyr et al., 2015).

### Limitations

Conducting an RCT with maltreating families in this context poses many challenges. The potential size for the study sample was limited as there were, at the time this project was conducted, only four clinics for PCAs in the Netherlands and our focus was on a specific age range. Even though we had a high response rate (79%), the sample was quite small. Another common problem with this population is a high attrition rate (e.g., Steele et al., 2019), although we still managed to reach almost two-thirds of the families for follow-up. Even though we used multilevel imputation to maximize power, this procedure takes the uncertainty of missing data into account by producing larger standard errors and more strict significance tests (Van Ginkel, Linting, Rippe, & van der Voort, 2019). A priori power calculations suggested a sample size of 60 based on an expected medium effect size and a power of .80. As the actual final sample size was lower, actual power to detect the hypothesized effect will be below .80. The observed effect size should thus have been considerably larger than hypothesized a priori, in order to detect it with the actual sample size. This leaves the possibility open that some effects were present, but could not be detected in this study.

A second limitation is related to the measurement of therapists' recommendations: during data inspection we noted systematic differences in the way the initial recommendation forms were filled out by the therapists. For two therapists, 73 and 89% of the initial recommendations favored an out-of-home placement, whereas for the other two therapists 75 and 90% of the initial recommendations favored that the child could stay with his/her parent(s). In practice, therapists do not have to provide a recommendation regarding child placement at the start of a PCA; we solely added this measure for research purposes. Therefore, it could be that these differences were related to therapists' interpretation of the initial recommendation form.

Finally, we relied on parent reports for follow-up data. One potential problem is that the parents who were traceable for and open to a follow-up assessment were a selected group. Although they did not differ from parents who dropped out on demographic or target variables, it could be that after the assessment, dropped out families experienced more problems than the families who continued to participate. For instance, the majority of children (88%) were living with their parents at follow-up; it could be that there had been more out-of-home placements for dyads who dropped out and that this biased the results. Another drawback of the use of parent reports is related to the validity of such reports. Previous studies have shown that abusive parents or parents with psychopathology tend to overreport their children's prob-

lem behavior (Najman et al., 2001; Reid, Kavanagh, & Baldwin, 1987), which suggests that they are not always reliable reporters of their children's actual behavior. It could be that the results of this study would have been different if we had obtained additional access to more objective reports of children's well-being.

### **Future directions**

We found no evidence that the PCAs incorporating the VIPP-SD protocol outperformed the PCAs as usual. However, the current study may provide important reference points for future research in this area. First, by conducting this study we showed that it is possible to empirically evaluate the effectiveness of a PCA protocol in improving the quality of subsequent placement decisions through a randomized research design – which, to our knowledge, has not been done previously besides by the parallel Canadian study (Cyr et al., 2015). In future studies, it will be important to overcome some of our current challenges by determining a priori what effect size would be needed in order to find clinically relevant results, and setting the required sample size accordingly.

A second implication is related to the unique child protection setting in which the current study was conducted: Because referral to an assessment in one of the Dutch clinics is usually considered as parents' last chance after a long trajectory of home-based support and due to the high costs not all families can be referred there, it would be interesting to explore the effects of implementing VIPP-SD or a similar intervention in an earlier stage. For instance, if a family is put under supervision for suspected or substantiated child maltreatment and home-based support is imposed on the family, the VIPP-SD assessment protocol might contribute to a better-informed indication of their parenting capacities and therefore lead to better decisions regarding child placement. Based on two recent studies which provided initial evidence in favor of the use of attachment-based assessments protocols (Cyr et al., 2015; Van der Asdonk et al., 2019), it would be worthwhile to further investigate the effectiveness of different implementations of this approach.

### **Acknowledgements**

We would like to express our thanks to the professionals from family clinics De Bascule, GGZ Drenthe, Yulius, and Horizon who were involved in this study. We would also like to thank Margot Enneking for her assistance in coding, and all Master students from Leiden University who assisted in data collection. Finally, we are grateful to all parents and children who were willing to contribute to our study.

## Appendix 4.A

### **Detailed description of sample flow throughout randomized trial**

In total, 56 families were included in the project. Following randomization, nine families dropped out (seven in the intervention group and two in the regular assessment group) for different reasons: the parent did not want to receive VIPP ( $n = 2$ ), the child was placed in foster care before the final evaluation took place ( $n = 6$ ), and for one mother-child dyad VIPP was stopped after one session due to individual circumstances (this dyad still took part in the post-test). The post-test did not take place for an additional seven parent-child dyads (four in the intervention group), because the parent did not want to participate anymore ( $n = 2$ ), the child was already placed into foster care ( $n = 2$ ), or because the family left the clinic early with a positive evaluation and could not be reached anymore ( $n = 3$ ). The follow-up did not take place for 22 families (12 in the intervention group), because the parent did not want to participate anymore ( $n = 12$ ), the parent was untraceable ( $n = 6$ ), or because the parent was unavailable for an appointment (e.g., because of severe psychiatric problems) ( $n = 4$ ).

## Appendix 4.B

### Multiple imputation procedures

Four methods were used in conjunction: the ‘MI’ function in the Amelia package (Honaker, King, & Blackwell, 2011), the ‘mice’ function from the mice package (Van Buuren & Groothuis-Oudshoorn, 2011), and the ‘panImpute’ and ‘jomoImpute’ functions from the mitml package (Grund, Robitzsch, & Lüdtke, 2016) to assess robustness of the imputed datasets as well as access the full range of analysis options. The maximum number of iterations was set at 10 and a fixed starting seed was set for reproducibility. Pooling of results on 50 imputation sets was performed using the summary functions from mitml and miceadds, as well as using the ‘summary’ and ‘modelRandEffStats’ functions from the merTools package (Knowles, Frederick, & Whitworth, 2018). All models were random-intercept models; as the already small sample showed high incompleteness, random slopes were not estimated in order to avoid consecutive estimations and uncertainty under weakened model identifiability.



## References

- Achenbach, T. M., & Rescorla, L. A. (2000). *Manual for the ASEBA Preschool Forms & Profiles*. Burlington, VT: University of Vermont, Research Center for Children, Youth, & Families.
- Ainsworth, M. D. S., Bell, S. M., & Stayton, D. J. (1974). Infant-mother attachment and social development. In M. P. Richards (Ed.), *The introduction of the child into a social world* (pp. 99-135). London: Cambridge University Press.
- American Psychological Association. (1998). Committee on Professional Practice and Standards. Retrieved from <http://www.apa.org/about/governance/bdcmte/standards.aspx>
- Bernard, K., Dozier, M., Bick, J., Lewis-Morrarty, E., Lindhiem, O., & Carlson, E. (2012). Enhancing attachment organization among maltreated children: Results of a randomized clinical trial. *Child Development, 83*(2), 623-636. doi:10.1111/j.1467-8624.2011.01712.x
- Biehal, N., Sinclair, I., & Wade, J. (2015). Reunifying abused or neglected children: Decision-making and outcomes. *Child Abuse & Neglect, 49*, 107-118. doi:10.1016/j.chiabu.2015.04.014
- Budd, K. S. (2001). Assessing parenting competence in child protection cases: A clinical practice model. *Clinical Child and Family Psychology Review, 4*(1), 18.
- Budd, K. S. (2005). Assessing parenting capacity in a child welfare context. *Children and Youth Services Review, 27*(4), 429-444. doi:10.1016/j.childyouth.2004.11.008
- Cicchetti, D., Toth, S. L., & Manly, J. T. (2003). *Maternal maltreatment classification interview*. Unpublished manuscript. Rochester, NY.
- Cyr, C., & Alink, L. R. A. (2017). Child maltreatment: the central roles of parenting capacities and attachment. *Current Opinion in Psychology, 15*, 81-86. doi:10.1016/j.copsyc.2017.02.002
- Cyr, C., Dubois-Comtois, K., Geneviève, M., Poulin, C., Pascuzzo, K., Losier, V., . . . Moss, E. (2012). Attachment theory in the assessment and promotion of parental competency in child protection cases. In A. Muela (Ed.), *Child Abuse and Neglect - A multidimensional approach* (pp. 63-86). Croatia: InTech.
- Cyr, C., Paquette, D., Dubois-Comtois, K., & Lopez, L. (2015, March). *An attachment-based intervention protocol for the assessment of parenting capacities in child welfare cases*. Paper presented at the Society for Research in Child Development, Philadelphia, PA.
- English, D. J., Bangdiwalab, S. I., & Runyan, D. K. (2005). The dimensions of maltreatment: Introduction. *Child Abuse & Neglect, 29*(5), 441-460. doi:10.1016/j.chiabu.2003.09.023
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G\*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods, 39*(2), 175-191. doi:10.3758/Bf03193146
- Gilbert, R., Widom, C. S., Browne, K., Fergusson, D., Webb, E., & Janson, S. (2009). Burden and consequences of child maltreatment in high-income countries. *Lancet, 373*(9657), 68-81. doi:10.1016/S0140-6736(08)61706-7
- Grund, S., Robitzsch, A., & Lüdtke, O. (2016). Mitml: Tools for Multiple Imputation in Multilevel Modeling. In.
- Harnett, P. H. (2007). A procedure for assessing parents' capacity for change in child protection cases. *Children and Youth Services Review, 29*(9), 1179-1188. doi:10.1016/j.childyouth.2007.04.005
- Honaker, J., King, G., & Blackwell, M. (2011). Amelia II: A program for missing data. *Journal of Statistical Software, 45*(7), 1-47.
- Juffer, F., Bakermans-Kranenburg, M. J., & Van IJzendoorn, M. H. (2017). Pairing attachment theory and social learning theory in video-feedback intervention to promote positive parenting. *Current Opinion in Psychology, 15*, 189-194. doi:10.1016/j.copsyc.2017.03.012
- Juffer, F., Bakermans-Kranenburg, M. J., & Van IJzendoorn, M. H. (2008). *Promoting positive parenting: An attachment-based intervention*. New York: Taylor & Francis.
- Kahneman, D., Slovic, S. P., Slovic, P., & Tversky, A. (1982). *Judgment under Uncertainty: Heuristics and*

- Biases*. Cambridge, England: Cambridge University Press.
- Knowles, J. E., Frederick, C., & Whitworth, A. (2018). merTools: Tools for Analyzing Mixed Effect Regression Models.
- Lindauer, R. J. L., Bakermans-Kranenburg, M. J., Van IJzendoorn, M. H., & Schuengel, C. (2010). *Thuiswonen of uithuisplaatsen: Betrouwbaarheid van de besluitvorming en innovatie van beslisdiagnostiek op basis van de capaciteit tot verbetering van ouderlijke pedagogische vaardigheden middels een evidence-based interventie [Out-of-home placement: Reliability of the decision process and innovation of diagnostic procedures on the basis of assessment of capacity for change in parenting through evidence-based intervention]*. Unpublished research proposal. Amsterdam: Bascule.
- Little, R. J. A. (1988). A test of missing completely at random for multivariate data with missing values. *Journal of the American Statistical Association*, 83(404), 1198-1202. doi:10.2307/2290157
- Lloyd, E. C., & Barth, R. P. (2011). Developmental outcomes after five years for foster children returned home, remaining in care, or adopted. *Children and Youth Services Review*, 33(8), 1383-1391. doi:10.1016/j.childyouth.2011.04.008
- Lorah, J. (2018). Effect size measures for multilevel models: definition, interpretation, and TIMSS example. *Large-Scale Assessments in Education*, 6. doi:10.1186/s40536-018-0061-2
- Lutman, E., & Farmer, E. (2013). What contributes to outcomes for neglected children who are reunified with their parents? Findings from a five-year follow-up study. *British Journal of Social Work*, 43(3), 559-578. doi:10.1093/bjsw/bcr184
- Mesman, J., Van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2012). Unequal in opportunity, equal in process: Parental sensitivity promotes positive child development in ethnic minority families. *Child Development Perspectives*, 6(3), 239-250. doi:10.1111/j.1750-8606.2011.00223.x
- Moss, E., Dubois-Comtois, K., Cyr, C., Tarabulsy, G. M., St-Laurent, D., & Bernier, A. (2011). Efficacy of a home-visiting intervention aimed at improving maternal sensitivity, child attachment, and behavioral outcomes for maltreated children: A randomized control trial. *Development and Psychopathology*, 23(1), 195-210. doi:10.1017/S0954579410000738
- Munro, E. (1999). Common errors of reasoning in child protection work. *Child Abuse & Neglect*, 23(8), 745-758. doi:10.1016/S0145-2134(99)00053-8
- Najman, J. M., Williams, G. M., Nikles, J., Spence, S., Bor, W., O'Callaghan, M., . . . Shuttlewood, G. J. (2001). Bias influencing maternal reports of child behaviour and emotional state. *Social Psychiatry and Psychiatric Epidemiology*, 36(4), 186-194. doi:10.1007/s001270170062
- Negrao, M., Pereira, M., Soares, I., & Mesman, J. (2014). Enhancing positive parent-child interactions and family functioning in a poverty sample: A randomized control trial. *Attachment & Human Development*, 16(4), 315-328. doi:10.1080/14616734.2014.912485
- Reid, J. B., Kavanagh, K., & Baldwin, D. V. (1987). Abusive parents perceptions of child problem behaviors - An example of parental bias. *Journal of Abnormal Child Psychology*, 15(3), 457-466. doi:10.1007/Bf00916461
- Reijman, S., Alink, L. R. A., Compier-de Block, L. H. C. G., Werner, C. D., Maras, A., Rijnberk, C., . . . Bakermans-Kranenburg, M. J. (2014). Autonomic reactivity to infant crying in maltreating mothers. *Child Maltreatment*, 19(2), 101-112. doi:10.1177/1077559514538115
- RStudioTeam. (2016). RStudio: Integrated Development for R. Boston, MA: RStudio, Inc.
- Rubin, D. B. (1987). *Multiple imputation for nonresponse in surveys*. New York, NY: Wiley.
- Steele, H., Murphy, A., Bonuck, K., Meissner, P., & Steele, M. (2019). Randomized control trial report on the effectiveness of Group Attachment-Based Intervention (GABI (c)): Improvements in the parent-child relationship not seen in the control group. *Development and Psychopathology*, 31(1), 203-217. doi:10.1017/S0954579418001621
- Teti, D. M., & Candelaria, M. A. (2002). Parenting Competence. In M. H. Bornstein (Ed.), *Handbook of Parenting Volume 4: Social Conditions and Applied Parenting* (Vol. 4, pp. 149-180). Mahwah, NJ: Lawrence Erlbaum Associates, Publishers.

- Van Buuren, S., & Groothuis-Oudshoorn, K. (2011). Mice: Multivariate imputation by chained equations in R. *Journal of Statistical Software*, 45(3), 1-67.
- Van der Asdonk, S., van Berkel, S. R., de Haan, W. D., van IJzendoorn, M. H., Schuengel, C., & Alink, L. R. A. (2019). Improving decision-making agreement in child protection cases by using information regarding parents' response to an intervention: A vignette study. *Children and Youth Services Review*. doi:10.1016/j.childyouth.2019.104501
- Van Ginkel, J. R., Linting, M., Rippe, R. C. A., & van der Voort, A. (2019). Rebutting existing misconceptions about multiple imputation as a method for handling missing data. *Journal of Personality Assessment*, 1-12. doi:10.1080/00223891.2018.1530680
- Vischer, A. W. K., Grietens, H., Knorth, E. J., & Mulder, H. (2017). Assessing parenting in the context of reunification of infants/toddlers and their families: How to face the challenges? *Infant Mental Health Journal*, 38(3), 406-421. doi:10.1002/imhj.21646
- Ward, H., Brown, R., & Hyde-Dryden, G. (2014). *Assessing Parental Capacity to Change when Children are on the Edge of Care: An overview of current research evidence*. Loughborough University, London.
- White, A. (2005). *Literature review: Assessment of parenting capacity*. Ashfield, Australia: Centre for Parenting & Research.
- White, I. R., Carpenter, J., & Horton, N. J. (2012). Including all individuals is not enough: Lessons for intention-to-treat analysis. *Clinical Trials*, 9(4), 396-407. doi:10.1177/1740774512450098

