



Universiteit
Leiden
The Netherlands

Dancing with the stars

Albert, J.G.

Citation

Albert, J. G. (2020, October 28). *Dancing with the stars*. Retrieved from <https://hdl.handle.net/1887/137988>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/137988>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/137988> holds various files of this Leiden University dissertation.

Author: Albert, J.G.

Title: Dancing with the stars

Issue Date: 2020-10-28

Bibliography

- Martín Abadi, Ashish Agarwal, Paul Barham, Eugene Brevdo, Zhifeng Chen, Craig Citro, Greg S. Corrado, Andy Davis, Jeffrey Dean, Matthieu Devin, Sanjay Ghemawat, Ian Goodfellow, Andrew Harp, Geoffrey Irving, Michael Isard, Yangqing Jia, Rafal Jozefowicz, Lukasz Kaiser, Manjunath Kudlur, Josh Levenberg, Dan Mané, Rajat Monga, Sherry Moore, Derek Murray, Chris Olah, Mike Schuster, Jonathon Shlens, Benoit Steiner, Ilya Sutskever, Kunal Talwar, Paul Tucker, Vincent Vanhoucke, Vijay Vasudevan, Fernanda Viégas, Oriol Vinyals, Pete Warden, Martin Wattenberg, Martin Wicke, Yuan Yu, and Xiaoqiang Zheng. TensorFlow: Large-scale machine learning on heterogeneous systems, 2015. URL <https://www.tensorflow.org/>. Software available from tensorflow.org.
- J. G. Albert, M. S. S. L. Oei, R. J. van Weeren, H. T. Intema, and H. J. A. Röttgering. A probabilistic approach to direction-dependent ionospheric calibration. *A&A*, 633:A77, Jan 2020a. doi: 10.1051/0004-6361/201935668.
- J. G. Albert, R. J. van Weeren, H. T. Intema, and H. J. A. Röttgering. Probabilistic direction-dependent ionospheric calibration for LOFAR-HBA. *A&A*, 635:A147, March 2020b. doi: 10.1051/0004-6361/201937424.
- V. A. Ambartsumian. On the Question of the Nature of Radiosources. In *Proceedings of the Fifth Meeting on Cosmogony. Trudy Pjatogo soveshanija po voprosam kosmogonii. 9-12 marta 1955 g. Radioastronomija*, pages 413–416, January 1956.
- Miguel A. Aragón-Calvo, Rien van de Weygaert, and Bernard J. T. Jones. Multiscale phenomenology of the cosmic web. *MNRAS*, 408(4):2163–2187, November 2010. doi: 10.1111/j.1365-2966.2010.17263.x.
- B. S. Arora, J. Morgan, S. M. Ord, S. J. Tingay, M. Bell, J. R. Callingham, K. S. Dwarakanath, B. Q. For, P. Hancock, L. Hindson, N. Hurley-Walker, M. Johnston-Hollitt, A. D. Kapińska, E. Lenc, B. McKinley, A. R. Offringa, P. Procopio, L. Staveley-Smith, R. B. Wayth, C. Wu, and Q. Zheng. Ionospheric Modelling using GPS to Calibrate the MWA. II: Regional Ionospheric Modelling using GPS and GLONASS to Estimate Ionospheric Gradients. *PASA*, 33:e031, Jul 2016. doi: 10.1017/pasa.2016.22.
- J. A. Ball, D. Cesarsky, A. K. Dupree, L. Goldberg, and A. E. Lilley. Detection and Identification of Recombination Lines from an H i Region. *ApJ*, 162:L25, October 1970. doi: 10.1086/180616.

-
- J. A. Barcelo, A. Ruiz, and L. Vega. Weighted estimates for the helmholtz equation and some applications. *Journal of Functional Analysis*, 150(2):356 – 382, 1997. ISSN 0022-1236. doi: <https://doi.org/10.1006/jfan.1997.3131>. URL <http://www.sciencedirect.com/science/article/pii/S0022123697931311>.
- Mitchell C. Begelman, Roger D. Blandford, and Martin J. Rees. Theory of extragalactic radio sources. *Reviews of Modern Physics*, 56(2):255–351, April 1984. doi: 10.1103/RevModPhys.56.255.
- B. Bhattacharyya, S. Cooper, M. Malenta, J. Roy, J. Chengalur, M. Keith, S. Kudale, M. McLaughlin, S. M. Ransom, P. S. Ray, and B. W. Stappers. The GMRT High Resolution Southern Sky Survey for Pulsars and Transients. I. Survey Description and Initial Discoveries. *ApJ*, 817(2):130, February 2016. doi: 10.3847/0004-637X/817/2/130.
- D. Bilitza and B. W. Reinisch. International Reference Ionosphere 2007: Improvements and new parameters. *Advances in Space Research*, 42:599–609, August 2008. doi: 10.1016/j.asr.2007.07.048.
- J. Blackledge. *Digital Image processing*. Horwood Publishing, 2005. ISBN 1-898563-49-7.
- Roger Blandford and David Eichler. Particle acceleration at astrophysical shocks: A theory of cosmic ray origin. *Phys. Rep.*, 154(1):1–75, October 1987. doi: 10.1016/0370-1573(87)90134-7.
- A. Bonafede, L. Feretti, G. Giovannini, F. Govoni, M. Murgia, G. B. Taylor, H. Ebeling, S. Allen, G. Gentile, and Y. Pihlström. Revealing the magnetic field in a distant galaxy cluster: discovery of the complex radio emission from MACS J0717.5 +3745. *A&A*, 503:707–720, September 2009. doi: 10.1051/0004-6361/200912520.
- A. Bonafede, L. Feretti, M. Murgia, F. Govoni, G. Giovannini, D. Dallacasa, K. Dolag, and G. B. Taylor. The Coma cluster magnetic field from Faraday rotation measures. *A&A*, 513:A30, April 2010. doi: 10.1051/0004-6361/200913696.
- A. Bonafede, M. Brüggen, R. van Weeren, F. Vazza, G. Giovannini, H. Ebeling, A. C. Edge, M. Hoeft, and U. Klein. Discovery of radio haloes and double relics in distant MACS galaxy clusters: clues to the efficiency of particle acceleration. *MNRAS*, 426:40–56, October 2012. doi: 10.1111/j.1365-2966.2012.21570.x.
- Etienne Bonnassieux, Cyril Tasse, Oleg Smirnov, and Philippe Zarka. The variance of radio interferometric calibration solutions. Quality-based weighting schemes. *A&A*, 615:A66, Jul 2018. doi: 10.1051/0004-6361/201732190.
- Max Born and Emil Wolf. *Principles of Optics*. 1999.
- A. Botteon, F. Gastaldello, G. Brunetti, and R. Kale. A $M \gtrsim 3$ shock in ‘El Gordo’ cluster and the origin of the radio relic. *MNRAS*, 463(2):1534–1542, December 2016. doi: 10.1093/mnras/stw2089.
- K. L. Bouman, M. D. Johnson, D. Zoran, V. L. Fish, S. S. Doeleman, and W. T. Freeman. Computational Imaging for VLBI Image Reconstruction. In *The IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, March 2016.

-
- S. Y. Braude, B. A. Dubinskii, N. L. Kaidanovskii, N. S. Kardashev, M. M. Kobrin, A. D. Kuzmin, A. P. Molchanov, Y. N. Pariiskii, O. N. Rzhiga, A. E. Salomonovich, V. A. Samanian, I. S. Shklovskii, R. L. Sorochenko, V. S. Troitskii, and K. I. Kellermann. *A Brief History of Radio Astronomy in the USSR*. 2012.
- Robert Braun. *The Square Kilometer Array Interferometer*, volume 208 of *Astrophysics and Space Science Library*, page 167. 1996. doi: 10.1007/978-94-009-1734-7_9.
- T. K. Breus. Istoriya prioritetoV sinkhrotronnoj kontseptsii v astronomii %t Historical problems of the priority questions of the synchrotron concept in astrophysics. *Istoriko-Astronomicheskie Issledovaniya*, 26:88–97, January 2001.
- D. Briggs. *High Fidelity Deconvolution of Moderately Resolved Sources*. PhD thesis, New Mexico Institute of Mining Technology, 1995. URL <http://www.aoc.nrao.edu/dissertations/dbriggs>.
- M. Brocklehurst and M. J. Seaton. On the interpretation of radio recombination line observations. *MNRAS*, 157:179, January 1972. doi: 10.1093/mnras/157.2.179.
- M. Brügger and F. Vazza. Analytical model for cluster radio relics. *MNRAS*, 493(2):2306–2317, April 2020. doi: 10.1093/mnras/staa418.
- G. Brunetti. The challenge of turbulent acceleration of relativistic particles in the intra-cluster medium. *Plasma Physics and Controlled Fusion*, 58(1):014011, January 2016. doi: 10.1088/0741-3335/58/1/014011.
- G. Brunetti and T. W. Jones. Cosmic Rays in Galaxy Clusters and Their Nonthermal Emission. *International Journal of Modern Physics D*, 23:1430007–98, March 2014. doi: 10.1142/S0218271814300079.
- G. Brunetti, S. Giacintucci, R. Cassano, W. Lane, D. Dallacasa, T. Venturi, N. E. Kassim, G. Setti, W. D. Cotton, and M. Markevitch. A low-frequency radio halo associated with a cluster of galaxies. *Nature*, 455:944–947, October 2008. doi: 10.1038/nature07379.
- G. R. Burbidge. On Synchrotron Radiation from Messier 87. *ApJ*, 124:416, September 1956. doi: 10.1086/146237.
- R. Byrd, P. Lu, J. Nocedal, and C. Zhu. A limited memory algorithm for bound constrained optimization. *SIAM Journal on Scientific Computing*, 16(5):1190–1208, 1995. doi: 10.1137/0916069.
- Xiaohao Cai, Marcelo Pereyra, and Jason D. McEwen. Uncertainty quantification for radio interferometric imaging: II. MAP estimation. *MNRAS*, 480(3):4170–4182, November 2018. doi: 10.1093/mnras/sty2015.
- Xiaohao Cai, Luke Pratley, and Jason D. McEwen. Online radio interferometric imaging: assimilating and discarding visibilities on arrival. *MNRAS*, 485(4):4559–4572, June 2019. doi: 10.1093/mnras/stz704.

-
- J. Callingham, H. Vedantham, T. Shimwell, B. J. Pope, and LoTSS Team. Stellar systems at low radio frequencies: The discovery of radio exoplanets. In *American Astronomical Society Meeting Abstracts*, American Astronomical Society Meeting Abstracts, page 342.02, January 2020.
- P J Cargill. Fundamentals of plasma physics. *Plasma Physics and Controlled Fusion*, 49(2): 197, 2007. URL <http://stacks.iop.org/0741-3335/49/i=2/a=B01>.
- R. Cassano, G. Brunetti, and G. Setti. Statistics of giant radio haloes from electron reacceleration models. *MNRAS*, 369:1577–1595, July 2006. doi: 10.1111/j.1365-2966.2006.10423.x.
- R. Cassano, G. Brunetti, H. J. A. Röttgering, and M. Brüggen. Unveiling radio halos in galaxy clusters in the LOFAR era. *A&A*, 509:A68, January 2010a. doi: 10.1051/0004-6361/200913063.
- R. Cassano, S. Ettori, S. Giacintucci, G. Brunetti, M. Markevitch, T. Venturi, and M. Gitti. On the Connection Between Giant Radio Halos and Cluster Mergers. *ApJ*, 721:L82–L85, October 2010b. doi: 10.1088/2041-8205/721/2/L82.
- Lawrie Challis and Fred Sheard. The Green of Green Functions. *Physics Today*, 56(12):41–46, December 2003. doi: 10.1063/1.1650227.
- A. S. Cohen and H. J. A. Röttgering. Probing Fine-Scale Ionospheric Structure with the Very Large Array Radio Telescope. *The Astronomical Journal*, 138(2):439–447, Aug 2009. doi: 10.1088/0004-6256/138/2/439.
- M. H. Cohen. Introduction to very-long-baseline interferometry. *IEEE Proceedings*, 61: 1192–1197, 1973. doi: 10.1109/PROC.1973.9244.
- J. J. Condon, W. D. Cotton, E. W. Greisen, Q. F. Yin, R. A. Perley, G. B. Taylor, and J. J. Broderick. The NRAO VLA Sky Survey. *AJ*, 115(5):1693–1716, May 1998. doi: 10.1086/300337.
- T. J. Cornwell and R. A. Perley. Radio-interferometric imaging of very large fields. The problem of non-coplanar arrays. *A&A*, 261:353–364, July 1992.
- T. J. Cornwell, K. Golap, and S. Bhatnagar. Wide Field Imaging: Fourier and Fresnel. In N. Kassim, M. Perez, W. Junor, and P. Henning, editors, *From Clark Lake to the Long Wavelength Array: Bill Erickson's Radio Science*, volume 345 of *Astronomical Society of the Pacific Conference Series*, page 350, Dec 2005.
- T. J. Cornwell, K. Golap, and S. Bhatnagar. The Noncoplanar Baselines Effect in Radio Interferometry: The W-Projection Algorithm. *IEEE Journal of Selected Topics in Signal Processing*, 2:647–657, November 2008. doi: 10.1109/JSTSP.2008.2005290.
- William D. Cotton, James J. Condon, Richard A. Perley, Namir Kassim, Joseph Lazio, Aaron Cohen, Wendy Lane, and William C. Erickson. Beyond the isoplanatic patch in the VLA Low-frequency Sky Survey. In Jr. Oschmann, Jacobus M., editor, *Proc. SPIE*, volume 5489 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, pages 180–189, Oct 2004. doi: 10.1117/12.551298.

-
- F. de Gasperin, T. J. Dijkema, A. Drabent, M. Mevius, D. Rafferty, R. van Weeren, M. Brüggen, J. R. Callingham, K. L. Emig, G. Heald, H. T. Intema, L. K. Morabito, A. R. Offringa, R. Oonk, E. Orrù, H. Röttgering, J. Sabater, T. Shimwell, A. Shulevski, and W. Williams. Systematic effects in LOFAR data: A unified calibration strategy. *A&A*, 622:A5, Feb 2019. doi: 10.1051/0004-6361/201833867.
- Francesco de Gasperin, Huib T. Intema, Timothy W. Shimwell, Gianfranco Brunetti, Marcus Brüggen, Torsten A. Enßlin, Reinout J. van Weeren, Annalisa Bonafede, and Huub J. A. Röttgering. Gentle reenergization of electrons in merging galaxy clusters. *Science Advances*, 3(10):e1701634, October 2017. doi: 10.1126/sciadv.1701634.
- Francesco de Gasperin, M. Mevius, D. Rafferty, Huib Intema, and Richard Fallows. The effect of the ionosphere on ultra-low frequency radio-interferometric observations. *A&A*, 615, 04 2018. doi: 10.1051/0004-6361/201833012.
- A. De Luca and S. Molendi. The 2-8 keV cosmic X-ray background spectrum as observed with XMM-Newton. *A&A*, 419:837–848, June 2004. doi: 10.1051/0004-6361:20034421.
- Thomas A. Dean, Sumeetpal S. Singh, Ajay Jasra, and Gareth W. Peters. Parameter estimation for hidden markov models with intractable likelihoods. *Scandinavian Journal of Statistics*, 41(4):970–987, 2014. doi: 10.1111/sjos.12077. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/sjos.12077>.
- P. E. Dewdney, P. J. Hall, R. T. Schilizzi, and T. J. L. W. Lazio. The Square Kilometre Array. *IEEE Proceedings*, 97(8):1482–1496, August 2009. doi: 10.1109/JPROC.2009.2021005.
- J. Donnert, K. Dolag, G. Brunetti, and R. Cassano. Rise and fall of radio haloes in simulated merging galaxy clusters. *MNRAS*, 429:3564–3569, March 2013. doi: 10.1093/mnras/sts628.
- J. R. Driscoll and D. M. Healy. Computing fourier transforms and convolutions on the 2-sphere. *Advances in Applied Mathematics*, 15(2):202 – 250, 1994. ISSN 0196-8858. doi: <https://doi.org/10.1006/aama.1994.1008>. URL <http://www.sciencedirect.com/science/article/pii/S0196885884710086>.
- L. O. Drury. An introduction to the theory of diffusive shock acceleration of energetic particles in tenuous plasmas. *Reports on Progress in Physics*, 46:973–1027, August 1983. doi: 10.1088/0034-4885/46/8/002.
- Michael W. Eastwood, Marin M. Anderson, Ryan M. Monroe, Gregg Hallinan, Benjamin R. Barsdell, Stephen A. Bourke, M. A. Clark, Steven W. Ellingson, Jayce Dowell, Hugh Garsden, Lincoln J. Greenhill, Jacob M. Hartman, Jonathon Kocz, T. Joseph W. Lazio, Danny C. Price, Frank K. Schinzel, Gregory B. Taylor, Harish K. Vedantham, Yuankun Wang, and David P. Woody. The Radio Sky at Meter Wavelengths: m-mode Analysis Imaging with the OVRO-LWA. *AJ*, 156(1):32, July 2018. doi: 10.3847/1538-3881/aac721.
- K. L. Emig, P. Salas, F. de Gasperin, J. B. R. Oonk, M. C. Toribio, H. J. A. Röttgering, and A. G. G. M. Tielens. The first detection of radio recombination lines at cosmological distances. *A&A*, 622:A7, February 2019. doi: 10.1051/0004-6361/201834052.

-
- T. A. Enßlin and Gopal-Krishna. Reviving fossil radio plasma in clusters of galaxies by adiabatic compression in environmental shock waves. *A&A*, 366:26–34, January 2001. doi: 10.1051/0004-6361:20000198.
- T. A. Ensslin, P. L. Biermann, U. Klein, and S. Kohle. Cluster radio relics as a tracer of shock waves of the large-scale structure formation. *A&A*, 332:395–409, April 1998.
- G. Evans, J. Blackledge, and P. Yardley. *Analytic Methods for Partial Differential Equations: Green's Functions*. Springer Undergraduate Mathematics Series, 1999.
- H. I. Ewen and E. M. Purcell. Observation of a Line in the Galactic Radio Spectrum: Radiation from Galactic Hydrogen at 1,420 Mc./sec. *Nature*, 168(4270):356, September 1951. doi: 10.1038/168356a0.
- P. B. Fellgett and E. H. Linfoot. On the Assessment of Optical Images. *Philosophical Transactions of the Royal Society of London Series A*, 247(931):369–407, Feb 1955. doi: 10.1098/rsta.1955.0001.
- W. Freudling, M. Romaniello, D. M. Bramich, P. Ballester, V. Forchi, C. E. García-Dabó, S. Moehler, and M. J. Neeser. Automated data reduction workflows for astronomy. The ESO Reflex environment. *A&A*, 559:A96, November 2013. doi: 10.1051/0004-6361/201322494.
- D. Gabor. Microscopy by Reconstructed Wave-Fronts. *Proceedings of the Royal Society of London Series A*, 197(1051):454–487, July 1949. doi: 10.1098/rspa.1949.0075.
- Jorge Garcia-Sucerquia, Román Castañeda, and Francisco F Medina. Fresnel-Fraunhofer diffraction and spatial coherence. *Optics Communications*, 205(4-6):239–245, May 2002. doi: 10.1016/S0030-4018(02)01378-0.
- John V. Gillespie, Dina A. Zinnes, G. S. Tahim, Philip A. Schrodtt, and R. Michael Rubison. An optimal control model of arms races. *The American Political Science Review*, 71(1):226–244, 1977. ISSN 00030554, 15375943. URL <http://www.jstor.org/stable/1956964>.
- F. Govoni, M. Murgia, L. Feretti, G. Giovannini, D. Dallacasa, and G. B. Taylor. A2255: The first detection of filamentary polarized emission in a radio halo. *A&A*, 430:L5–L8, January 2005. doi: 10.1051/0004-6361:200400113.
- T. L. Grobler, C. D. Nunhokee, O. M. Smirnov, A. J. van Zyl, and A. G. de Bruyn. Calibration artefacts in radio interferometry - I. Ghost sources in Westerbork Synthesis Radio Telescope data. *MNRAS*, 439(4):4030–4047, April 2014. doi: 10.1093/mnras/stu268.
- Y. Gupta, B. Ajithkumar, H. S. Kale, S. Nayak, S. Sabhapathy, S. Sureshkumar, R. V. Swami, J. N. Chengalur, S. K. Ghosh, C. H. Ishwara-Chandra, B. C. Joshi, N. Kanekar, D. V. Lal, and S. Roy. The upgraded GMRT: opening new windows on the radio Universe. *Current Science*, 113(4):707–714, August 2017.
- J. P. Hamaker, J. D. Bregman, and R. J. Sault. Understanding radio polarimetry. I. Mathematical foundations. *A&AS*, 117:137–147, May 1996.
- I. Harrison, S. Camera, J. Zuntz, and M. L. Brown. SKA weak lensing - I. Cosmological forecasts and the power of radio-optical cross-correlations. *MNRAS*, 463:3674–3685, December 2016. doi: 10.1093/mnras/stw2082.

-
- A. H. Hassan, C. J. Fluke, D. G. Barnes, and V. A. Kilborn. Tera-scale astronomical data analysis and visualization. *MNRAS*, 429(3):2442–2455, March 2013. doi: 10.1093/mnras/sts513.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep Residual Learning for Image Recognition. *arXiv e-prints*, art. arXiv:1512.03385, December 2015.
- S. E. Healey, R. W. Romani, G. B. Taylor, E. M. Sadler, R. Ricci, T. Murphy, J. S. Ulvestad, and J. N. Winn. CRATES: An All-Sky Survey of Flat-Spectrum Radio Sources. *ApJS*, 171:61–71, July 2007. doi: 10.1086/513742.
- Carl Heiles, Phil Perillat, Michael Nolan, Duncan Lorimer, Ramesh Bhat, Tapasi Ghosh, Ellen Howell, Murray Lewis, Karen O’Neil, Chris Salter, and Snezana Stanimirovic. All-Stokes Parameterization of the Main Beam and First Sidelobe for the Arecibo Radio Telescope. *PASP*, 113(788):1247–1273, October 2001. doi: 10.1086/323290.
- J. N. Hendriks, C. Jidling, A. Wills, and T. B. Schön. Evaluating the squared-exponential covariance function in Gaussian processes with integral observations. *arXiv e-prints*, art. arXiv:1812.07319, Dec 2018.
- James Hensman, Nicolo Fusi, and Neil D. Lawrence. Gaussian Processes for Big Data. *arXiv e-prints*, art. arXiv:1309.6835, Sep 2013.
- S. L. G. Hess and P. Zarka. Modeling the radio signature of the orbital parameters, rotation, and magnetic field of exoplanets. *A&A*, 531:A29, July 2011. doi: 10.1051/0004-6361/201116510.
- A. Hewish. The Diffraction of Radio Waves in Passing through a Phase-Changing Ionosphere. *Proceedings of the Royal Society of London Series A*, 209(1096):81–96, Oct 1951. doi: 10.1098/rspa.1951.0189.
- A. Hewish. The Diffraction of Galactic Radio Waves as a Method of Investigating the Irregular Structure of the Ionosphere. *Proceedings of the Royal Society of London Series A*, 214(1119):494–514, Oct 1952. doi: 10.1098/rspa.1952.0185.
- A. Hewish, S. J. Bell, J. D. H. Pilkington, P. F. Scott, and R. A. Collins. Observation of a Rapidly Pulsating Radio Source. *Nature*, 217(5130):709–713, February 1968. doi: 10.1038/217709a0.
- J. A. Högbom. Aperture Synthesis with a Non-Regular Distribution of Interferometer Baselines. *A&AS*, 15:417, June 1974.
- C. Hollitt, M. Johnston-Hollitt, S. Dehghan, M. Frean, and T. Butler-Yeoman. *An Overview of the SKA Science Analysis Pipeline*, volume 512 of *Astronomical Society of the Pacific Conference Series*, page 367. 2017.
- N. Hurley-Walker, J. R. Callingham, P. J. Hancock, T. M. O. Franzen, L. Hindson, A. D. Kapińska, J. Morgan, A. R. Offringa, R. B. Wayth, C. Wu, Q. Zheng, T. Murphy, M. E. Bell, K. S. Dwarakanath, B. For, B. M. Gaensler, M. Johnston-Hollitt, E. Lenc, P. Procopio, L. Staveley-Smith, R. Ekers, J. D. Bowman, F. Briggs, R. J. Cappallo, A. A. Deshpande, L. Greenhill, B. J. Hazelton, D. L. Kaplan, C. J. Lonsdale, S. R. McWhirter, D. A. Mitchell, M. F. Morales, E. Morgan, D. Oberoi, S. M. Ord, T. Prabu, N. U. Shankar, K. S. Srivani, R. Subrahmanyam, S. J. Tingay, R. L. Webster, A. Williams, and C. L. Williams. GaLactic and Extragalactic All-sky Murchison Widefield Array (GLEAM) survey - I. A low-frequency extragalactic catalogue. *MNRAS*, 464:1146–1167, January 2017. doi: 10.1093/mnras/stw2337.

-
- I. H. Hutchinson. *Principles of Plasma Diagnostics*. 2005.
- H. T. Intema. SPAM: Source Peeling and Atmospheric Modeling. Astrophysics Source Code Library, August 2014.
- H. T. Intema, S. van der Tol, W. D. Cotton, A. S. Cohen, I. M. van Bemmelen, and H. J. A. Röttgering. Ionospheric calibration of low frequency radio interferometric observations using the peeling scheme. I. Method description and first results. *A&A*, 501:1185–1205, July 2009. doi: 10.1051/0004-6361/200811094.
- H. T. Intema, P. Jagannathan, K. P. Mooley, and D. A. Frail. The GMRT 150 MHz All-sky Radio Survey: First Alternative Data Release TGSS ADR1. *ArXiv e-prints*, March 2016.
- E. A. Isaev and P. A. Tarasov. Transmission of large amounts of scientific data using laser technology. In *Journal of Physics Conference Series*, volume 740 of *Journal of Physics Conference Series*, page 012015, August 2016. doi: 10.1088/1742-6596/740/1/012015.
- W. J. Jaffe and G. C. Perola. Dynamical Models of Tailed Radio Sources in Clusters of Galaxies. *A&A*, 26:423, August 1973.
- Karl G. Jansky. Radio Waves from Outside the Solar System. *Nature*, 132(3323):66, July 1933. doi: 10.1038/132066a0.
- Wojciech Jarosz. *Efficient Monte Carlo Methods for Light Transport in Scattering Media*. PhD thesis, UC San Diego, September 2008.
- H. Jeffreys. On certain approximate solutions of lineae differential equations of the second order*. *Proceedings of the London Mathematical Society*, s2-23(1):428–436, 1925. ISSN 1460-244X. doi: 10.1112/plms/s2-23.1.428.
- Carl Jidling, Johannes Hendriks, Niklas Wahlström, Alexander Gregg, Thomas B. SchÄun, Christopher Wensrich, and Adrian Wills. Probabilistic modelling and reconstruction of strain. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 436:141 – 155, 2018. ISSN 0168-583X. doi: <https://doi.org/10.1016/j.nimb.2018.08.051>. URL <http://www.sciencedirect.com/science/article/pii/S0168583X1830524X>.
- R. C. Jones. New calculus for the treatment of optical systems. I. Description and discussion of the calculus. *Journal of the Optical Society of America (1917-1983)*, 31:488, July 1941.
- C. H. Jordan, S. Murray, C. M. Trott, R. B. Wayth, D. A. Mitchell, M. Rahimi, B. Pindor, P. Procopio, and J. Morgan. Characterization of the ionosphere above the Murchison Radio Observatory using the Murchison Widefield Array. *MNRAS*, 471(4):3974–3987, Nov 2017. doi: 10.1093/mnras/stx1797.
- H. Kang and D. Ryu. Diffusive Shock Acceleration at Cosmological Shock Waves. *ApJ*, 764:95, February 2013. doi: 10.1088/0004-637X/764/1/95.
- N. S. Kardashev. On the Possibility of Detection of Allowed Lines of Atomic Hydrogen in the Radio-Frequency Spectrum. *Soviet Ast.*, 3:813, October 1959.

-
- N. S. Kardashev. Nonstationarity of Spectra of Young Sources of Nonthermal Radio Emission. *Soviet Ast.*, 6:317, December 1962.
- S. Kazemi, S. Yatawatta, S. Zaroubi, P. Lampropoulos, A. G. de Bruyn, L. V. E. Koopmans, and J. Noordam. Radio interferometric calibration using the SAGE algorithm. *MNRAS*, 414: 1656–1666, June 2011. doi: 10.1111/j.1365-2966.2011.18506.x.
- J.B. Keller, R. Bellman, and American Mathematical Society. *Stochastic Equations and Wave Propagation in Random Media*. Proceedings of symposia in applied mathematics. American Mathematical Society, 1964. URL <https://books.google.nl/books?id=evQW0gAACAAJ>.
- K. O. Kiepenheuer. On the relations between ionosphere, sunspots and solar corona. *MNRAS*, 106:515, Jan 1946. doi: 10.1093/mnras/106.6.515.
- Diederik P. Kingma and Jimmy Ba. Adam: A Method for Stochastic Optimization. *arXiv e-prints*, art. arXiv:1412.6980, December 2014.
- M. G. Kivelson and C. T. Russell. *Introduction to Space Physics*. Cambridge University Press, April 1995. ISBN 0521451043.
- A. N. Kolmogorov. The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Proceedings of the Royal Society of London Series A*, 434:9–13, July 1991. doi: 10.1098/rspa.1991.0075.
- Andrey N. Kolmogorov. *Foundations of the theory of probability*. Chelsea Pub Co, 2 english edition, 1956.
- Andrey N. Kolmogorov. *Foundations of the Theory of Probability*. Chelsea Pub Co, 2 edition, June 1960. URL [http://www.clrc.rhul.ac.uk/resources/fop/Theory%20of%20Probability%20\(small\).pdf](http://www.clrc.rhul.ac.uk/resources/fop/Theory%20of%20Probability%20(small).pdf).
- S. S. Komissarov and A. G. Gubanov. Relic radio galaxies: evolution of synchrotron spectrum. *A&A*, 285:27–43, May 1994.
- L. V. E. Koopmans. Ionospheric Power-spectrum Tomography in Radio Interferometry. *ApJ*, 718:963–971, August 2010. doi: 10.1088/0004-637X/718/2/963.
- Benjamin Krüger, Thomas Brenner, and Alwin Kienle. Solution of the inhomogeneous Maxwell’s equations using a Born series. *Optics Express*, 25(21):25165, October 2017. doi: 10.1364/OE.25.025165.
- Mark Kuiack, Folkert Huizinga, Gijs Molenaar, Peeyush Prasad, Antonia Rowlinson, and Ralph A. M. J. Wijers. AARTFAAC flux density calibration and Northern hemisphere catalogue at 60 MHz. *Monthly Notices of the Royal Astronomical Society*, 482(2):2502–2514, Jan 2019. doi: 10.1093/mnras/sty2810.
- M. J. Kurtz and D. J. Mink. RVSAO 2.0: Digital Redshifts and Radial Velocities. *PASP*, 110: 934–977, August 1998. doi: 10.1086/316207.

-
- R. R. Lindner, A. J. Baker, J. P. Hughes, N. Battaglia, N. Gupta, K. Knowles, T. A. Marriage, F. Menanteau, K. Moodley, E. D. Reese, and R. Srianand. The Radio Relics and Halo of El Gordo, a Massive $z = 0.870$ Cluster Merger. *ApJ*, 786:49, May 2014. doi: 10.1088/0004-637X/786/1/49.
- Robert R. Lindner, Carlos Vera-Ciro, Claire E. Murray, Snežana Stanimirović, Brian Babler, Carl Heiles, Patrick Hennebelle, W. M. Goss, and John Dickey. Autonomous Gaussian Decomposition. *AJ*, 149(4):138, April 2015. doi: 10.1088/0004-6256/149/4/138.
- S. T. Loi, T. Murphy, I. H. Cairns, F. W. Menk, C. L. Waters, P. J. Erickson, C. M. Trott, N. Hurley-Walker, J. Morgan, E. Lenc, A. R. Offringa, M. E. Bell, R. D. Ekers, B. M. Gaensler, C. J. Lonsdale, L. Feng, P. J. Hancock, D. L. Kaplan, G. Bernardi, J. D. Bowman, F. Briggs, R. J. Cappallo, A. A. Deshpande, L. J. Greenhill, B. J. Hazelton, M. Johnston-Hollitt, S. R. McWhirter, D. A. Mitchell, M. F. Morales, E. Morgan, D. Oberoi, S. M. Ord, T. Prabu, N. U. Shankar, K. S. Srivani, R. Subrahmanyam, S. J. Tingay, R. B. Wayth, R. L. Webster, A. Williams, and C. L. Williams. Real-time imaging of density ducts between the plasmasphere and ionosphere. *Geophys. Res. Lett.*, 42:3707–3714, May 2015. doi: 10.1002/2015GL063699.
- R. J. Lyon, B. W. Stappers, L. Levin, M. B. Mickaliger, and A. Scaife. A processing pipeline for high volume pulsar candidate data streams. *Astronomy and Computing*, 28:100291, July 2019. doi: 10.1016/j.ascom.2019.100291.
- M. A. Malkov and L. O’C Drury. Nonlinear theory of diffusive acceleration of particles by shock waves. *Reports on Progress in Physics*, 64:429–481, April 2001. doi: 10.1088/0034-4885/64/4/201.
- S. Mandal, H. T. Intema, T. W. Shimwell, R. J. van Weeren, A. Botteon, H. J. A. Röttgering, D. N. Hoang, G. Brunetti, F. de Gasperin, S. Giacintucci, H. Hoekstra, A. Stroe, M. Brüggen, R. Cassano, A. Shulevski, A. Drabent, and D. Rafferty. Ultra-steep spectrum emission in the merging galaxy cluster Abell 1914. *A&A*, 622:A22, February 2019. doi: 10.1051/0004-6361/201833992.
- S. Mandal, H. T. Intema, R. J. van Weeren, T. W. Shimwell, A. Botteon, G. Brunetti, F. de Gasperin, M. Brüggen, G. Di Gennaro, R. Kraft, H. J. A. Röttgering, M. Hardcastle, and C. Tasse. Revived fossil plasma sources in galaxy clusters. *A&A*, 634:A4, February 2020. doi: 10.1051/0004-6361/201936560.
- L. Mandel and E. Wolf. Coherence Properties of Optical Fields. *Reviews of Modern Physics*, 37(2):231–287, April 1965. doi: 10.1103/RevModPhys.37.231.
- Alexander G. de G. Matthews, Mark van der Wilk, Tom Nickson, Keisuke. Fujii, Alexis Boukouvalas, Pablo León-Villagrà, Zoubin Ghahramani, and James Hensman. GPflow: A Gaussian process library using TensorFlow. *Journal of Machine Learning Research*, 18(40):1–6, apr 2017. URL <http://jmlr.org/papers/v18/16-537.html>.
- B. J. Maughan, C. Jones, W. Forman, and L. Van Speybroeck. Images, Structural Properties, and Metal Abundances of Galaxy Clusters Observed with Chandra ACIS-I at $0.1 \ll z \ll 1.3$. *ApJS*, 174:117–135, January 2008. doi: 10.1086/521225.

-
- F. Mernier, J. de Plaa, L. Lovisari, C. Pinto, Y.-Y. Zhang, J. S. Kaastra, N. Werner, and A. Simionescu. Abundance and temperature distributions in the hot intra-cluster gas of Abell 4059. *A&A*, 575:A37, March 2015. doi: 10.1051/0004-6361/201425282.
- F. G. Mertens, M. Mevius, L. V. E. Koopmans, A. R. Offringa, G. Mellema, S. Zaroubi, M. A. Brentjens, H. Gan, B. K. Gehlot, V. N. Pandey, A. M. Sardarabadi, H. K. Vedantham, S. Yatawatta, K. M. B. Asad, B. Ciardi, E. Chapman, S. Gazagnes, R. Ghara, A. Ghosh, S. K. Giri, I. T. Iliev, V. Jelić, R. Kooistra, R. Mondal, J. Schaye, and M. B. Silva. Improved upper limits on the 21 cm signal power spectrum of neutral hydrogen at $z \approx 9.1$ from LOFAR. *MNRAS*, 493(2): 1662–1685, April 2020. doi: 10.1093/mnras/staa327.
- Christopher A. Metzler, Martin White, and Chris Loken. The Effect of the Cosmic Web on Cluster Weak Lensing Mass Estimates. *ApJ*, 547(2):560–573, February 2001. doi: 10.1086/318406.
- M. Mevius, S. van der Tol, V. N. Pandey, H. K. Vedantham, M. A. Brentjens, A. G. de Bruyn, F. B. Abdalla, K. M. B. Asad, J. D. Bregman, W. N. Brouw, S. Bus, E. Chapman, B. Ciardi, E. R. Fernandez, A. Ghosh, G. Harker, I. T. Iliev, V. Jelić, S. Kazemi, L. V. E. Koopmans, J. E. Noordam, A. R. Offringa, A. H. Patil, R. J. van Weeren, S. Wijnholds, S. Yatawatta, and S. Zaroubi. Probing ionospheric structures using the LOFAR radio telescope. *Radio Science*, 51:927–941, Jul 2016. doi: 10.1002/2016RS006028.
- G. K. Miley, H. van der Laan, and K. J. Wellington. Recent Westerbork Observations of Head-Tail Galaxies. In J. R. Shakeshaft, editor, *The Formation and Dynamics of Galaxies*, volume 58 of *IAU Symposium*, page 109, 1974.
- C. A. Muller and J. H. Oort. Observation of a Line in the Galactic Radio Spectrum: The Interstellar Hydrogen Line at 1,420 Mc./sec., and an Estimate of Galactic Rotation. *Nature*, 168(4270):357–358, September 1951. doi: 10.1038/168357a0.
- Jan E. Noordam. *LOFAR calibration challenges*, volume 5489 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, pages 817–825. 2004. doi: 10.1117/12.544262.
- S. E. Nuza, M. Hoeft, R. J. van Weeren, S. Gottlöber, and G. Yepes. How many radio relics await discovery? *MNRAS*, 420:2006–2019, March 2012. doi: 10.1111/j.1365-2966.2011.20118.x.
- A. R. Offringa. Compression of interferometric radio-astronomical data. *A&A*, 595:A99, November 2016. doi: 10.1051/0004-6361/201629565.
- Gerwin Osnabrugge, Saroch Leedumrongwatthanakun, and Ivo M. Vellekoop. A convergent Born series for solving the inhomogeneous Helmholtz equation in arbitrarily large media. *Journal of Computational Physics*, 322:113–124, October 2016. doi: 10.1016/j.jcp.2016.06.034.
- Donald E. Osterbrock. *Astrophysics of gaseous nebulae and active galactic nuclei*. 1989.
- A. G. Pacholczyk. *Radio astrophysics. Nonthermal processes in galactic and extragalactic sources*. 1970.

-
- Gregory Paciga, Joshua G. Albert, Kevin Bandura, Tzu-Ching Chang, Yashwant Gupta, Christopher Hirata, Julia Odegova, Ue-Li Pen, Jeffrey B. Peterson, Jayanta Roy, J. Richard Shaw, Kris Sigurdson, and Tabitha Voytek. A simulation-calibrated limit on the H I power spectrum from the GMRT Epoch of Reionization experiment. *MNRAS*, 433(1):639–647, July 2013. doi: 10.1093/mnras/stt753.
- N. G. Parke. Optical algebra. *Journal of Mathematics and Physics*, 28(1-4):131–139, 1949. doi: 10.1002/sapm1949281131. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/sapm1949281131>.
- A. H. Patil, S. Yatawatta, L. V. E. Koopmans, A. G. de Bruyn, M. A. Brentjens, S. Zaroubi, K. M. B. Asad, M. Hatef, V. Jelić, M. Mevius, A. R. Offringa, V. N. Pandey, H. Vedantham, F. B. Abdalla, W. N. Brouw, E. Chapman, B. Ciardi, B. K. Gehlot, A. Ghosh, G. Harker, I. T. Iliev, K. Kakiichi, S. Majumdar, G. Mellema, M. B. Silva, J. Schaye, D. Vrbanec, and S. J. Wijnholds. Upper Limits on the 21 cm Epoch of Reionization Power Spectrum from One Night with LOFAR. *ApJ*, 838:65, March 2017. doi: 10.3847/1538-4357/aa63e7.
- T. J. Pearson and A. C. S. Readhead. Image Formation by Self-Calibration in Radio Astronomy. *ARA&A*, 22:97–130, January 1984. doi: 10.1146/annurev.aa.22.090184.000525.
- Richard A. Perley, Frederic R. Schwab, and Alan H. Bridle. *Synthesis imaging in radio astronomy: a collection of lectures from the third NRAO synthesis imaging summer school*, volume 6. 1989.
- H. J. Pesch and M. Plail. The maximum principle of optimal control: A history of ingenious ideas and missed opportunities. *Control and Cybernetics*, Vol. 38, no 4A:973–995, 2009.
- G. J. Phillips. Measurement of winds in the ionosphere. *Journal of Atmospheric and Terrestrial Physics*, 2(3):141–154, Jan 1952. doi: 10.1016/0021-9169(52)90059-7.
- Planck Collaboration, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. Balbi, A. J. Banday, R. B. Barreiro, M. Bartelmann, and et al. Planck early results. IX. XMM-Newton follow-up for validation of Planck cluster candidates. *A&A*, 536:A9, December 2011. doi: 10.1051/0004-6361/201116460.
- Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, H. Aussel, C. Baccigalupi, and et al. Planck 2013 results. XXIX. The Planck catalogue of Sunyaev-Zeldovich sources. *A&A*, 571:A29, November 2014. doi: 10.1051/0004-6361/201321523.
- G. B. Poole, M. A. Fardal, A. Babul, I. G. McCarthy, T. Quinn, and J. Wadsley. The impact of mergers on relaxed X-ray clusters - I. Dynamical evolution and emergent transient structures. *MNRAS*, 373:881–905, December 2006. doi: 10.1111/j.1365-2966.2006.10916.x.
- P. Prasad, F. Huizinga, E. Kooistra, D. van der Schuur, A. Gunst, J. Romein, M. Kuiack, G. Molenaar, A. Rowlinson, J. D. Swinbank, and R. A. M. J. Wijers. The AARTFAAC All-Sky Monitor: System Design and Implementation. *Journal of Astronomical Instrumentation*, 5:1641008, December 2016. doi: 10.1142/S2251171716410087.

-
- L. H. Quiroga-Nuñez, H. T. Intema, J. R. Callingham, J. Villadsen, H. J. van Langevelde, P. Jagannathan, T. W. Shimwell, and E. P. Boven. Differences in radio emission from similar M dwarfs in the binary system Ross 867-8. *A&A*, 633:A130, January 2020. doi: 10.1051/0004-6361/201936491.
- L. Rabiner and B. Juang. An introduction to hidden markov models. *IEEE ASSP Magazine*, 3 (1):4–16, Jan 1986. ISSN 1558-1284. doi: 10.1109/MASSP.1986.1165342.
- J. Radon. Über die Bestimmung von Funktionen durch ihre Integralwerte längs gewisser Mannigfaltigkeiten. *Berichte über die Verhandlungen der Königlich-Sächsischen Gesellschaft der Wissenschaften zu Leipzig*, 69:262–277, 1917.
- Carl Edward Rasmussen and Christopher K. I. Williams. *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning)*. The MIT Press, 2006a. ISBN 026218253X.
- Carl Edward Rasmussen and Christopher K. I. Williams. *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning)*. The MIT Press, 2006b. ISBN 026218253X.
- J. A. Ratcliffe. Some Aspects of Diffraction Theory and their Application to the Ionosphere. *Reports on Progress in Physics*, 19(1):188–267, Jan 1956. doi: 10.1088/0034-4885/19/1/306.
- H. E. Rauch. Solution to the linear smoothing problem. *Automatic Control, IEEE Transactions on*, AC-8:371 – 372, 11 1963. doi: 10.1109/TAC.1963.1105600.
- T. H. Reiprich and H. Böhringer. The Mass Function of an X-Ray Flux-limited Sample of Galaxy Clusters. *ApJ*, 567:716–740, March 2002. doi: 10.1086/338753.
- C. J. Riseley, A. M. M. Scaife, M. W. Wise, and A. O. Clarke. Diffuse radio emission in MACS J0025.4-1222: the effect of a major merger on bulk separation of ICM components. *A&A*, 597:A96, January 2017. doi: 10.1051/0004-6361/201629530.
- K. Roettiger, J. M. Stone, and J. O. Burns. Magnetic Field Evolution in Merging Clusters of Galaxies. *ApJ*, 518:594–602, June 1999. doi: 10.1086/307298.
- George B. Rybicki and Alan P. Lightman. *Radiative Processes in Astrophysics*. 1986.
- M. Ryle and D. D. Vonberg. Solar Radiation on 175 Mc./s. *Nature*, 158(4010):339–340, September 1946. doi: 10.1038/158339b0.
- J. S. Santos, P. Rosati, P. Tozzi, H. Böhringer, S. Ettori, and A. Bignamini. Searching for cool core clusters at high redshift. *A&A*, 483:35–47, May 2008. doi: 10.1051/0004-6361/20078815.
- A. Saxena, M. Marinello, R. A. Overzier, P. N. Best, H. J. A. Röttgering, K. J. Duncan, I. Prandoni, L. Pentericci, M. Magliocchetti, D. Paris, F. Cusano, F. Marchi, H. T. Intema, and GK Miley. Discovery of a radio galaxy at $z = 5.72$. *MNRAS*, 480(2):2733–2742, October 2018. doi: 10.1093/mnras/sty1996.
- Jelena A. Schmalz, Gerd Schmalz, Timur E. Gureyev, and Konstantin M. Pavlov. On the derivation of the Green's function for the Helmholtz equation using generalized functions. *American Journal of Physics*, 78(2):181–186, February 2010. doi: 10.1119/1.3253655.

-
- T. W. Shimwell, H. J. A. Röttgering, P. N. Best, W. L. Williams, T. J. Dijkema, F. de Gasperin, M. J. Hardcastle, G. H. Heald, D. N. Hoang, A. Horneffer, H. Intema, E. K. Mahony, S. Mandal, A. P. Mechev, L. Morabito, J. B. R. Oonk, D. Rafferty, E. Retana-Montenegro, J. Sabater, C. Tasse, R. J. van Weeren, M. Brüggen, G. Brunetti, K. T. Chyży, J. E. Conway, M. Haverkorn, N. Jackson, M. J. Jarvis, J. P. McKean, G. K. Miley, R. Morganti, G. J. White, M. W. Wise, I. M. van Bemmell, R. Beck, M. Brienza, A. Bonafede, G. Calistro Rivera, R. Cassano, A. O. Clarke, D. Cseh, A. Deller, A. Drabent, W. van Driel, D. Engels, H. Falcke, C. Ferrari, S. Fröhlich, M. A. Garrett, J. J. Harwood, V. Heesen, M. Hoeft, C. Horellou, F. P. Israel, A. D. Kapińska, M. Kunert-Bajraszewska, D. J. McKay, N. R. Mohan, E. Orrú, R. F. Pizzo, I. Prandoni, D. J. Schwarz, A. Shulevski, M. Sipior, D. J. B. Smith, S. S. Sridhar, M. Steinmetz, A. Stroe, E. Varenus, P. P. van der Werf, J. A. Zensus, and J. T. L. Zwart. The LOFAR Two-metre Sky Survey. I. Survey description and preliminary data release. *A&A*, 598:A104, Feb 2017. doi: 10.1051/0004-6361/201629313.
- T. W. Shimwell, C. Tasse, M. J. Hardcastle, A. P. Mechev, W. L. Williams, P. N. Best, H. J. A. Röttgering, J. R. Callingham, T. J. Dijkema, F. de Gasperin, D. N. Hoang, B. Hugo, M. Mirmont, J. B. R. Oonk, I. Prandoni, D. Rafferty, J. Sabater, O. Smirnov, R. J. van Weeren, G. J. White, M. Atemkeng, L. Bester, E. Bonnassieux, M. Brüggen, G. Brunetti, K. T. Chyży, R. Cochrane, J. E. Conway, J. H. Croston, A. Danezi, K. Duncan, M. Haverkorn, G. H. Heald, M. Iacobelli, H. T. Intema, N. Jackson, M. Jamrozy, M. J. Jarvis, R. Lakhoo, M. Mevius, G. K. Miley, L. Morabito, R. Morganti, D. Nisbet, E. Orrú, S. Perkins, R. F. Pizzo, C. Schrijvers, D. J. B. Smith, R. Vermeulen, M. W. Wise, L. Alegre, D. J. Bacon, I. M. van Bemmell, R. J. Beswick, A. Bonafede, A. Botteon, S. Bourke, M. Brienza, G. Calistro Rivera, R. Cassano, A. O. Clarke, C. J. Conselice, R. J. Dettmar, A. Drabent, C. Dumba, K. L. Emig, T. A. Enßlin, C. Ferrari, M. A. Garrett, R. T. Génova-Santos, A. Goyal, G. Gürkan, C. Hale, J. J. Harwood, V. Heesen, M. Hoeft, C. Horellou, C. Jackson, G. Kokotanekov, R. Kondapally, M. Kunert-Bajraszewska, V. Mahatma, E. K. Mahony, S. Mandal, J. P. McKean, A. Merloni, B. Mingo, A. Miskolczi, S. Mooney, B. Nikiel-Wroczyński, S. P. O’Sullivan, J. Quinn, W. Reich, C. Roskowiński, A. Rowlinson, F. Savini, A. Saxena, D. J. Schwarz, A. Shulevski, S. S. Sridhar, H. R. Stacey, S. Urquhart, M. H. D. van der Wiel, E. Varenus, B. Webster, and A. Wilber. The LOFAR Two-metre Sky Survey. II. First data release. *A&A*, 622:A1, February 2019. doi: 10.1051/0004-6361/201833559.
- I. S. Shklovsky. The Problem of Cosmic Radioemission. ‘Problema Kosmičeskogo Radioizlučeniâ’. *AZh*, 30:15–36, January 1953.
- R. H. Shumway and D. S. Stoffer. An approach to time series smoothing and forecasting using the em algorithm. *Journal of Time Series Analysis*, 3(4):253–264, 1982. doi: 10.1111/j.1467-9892.1982.tb00349.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9892.1982.tb00349.x>.
- C. Sifón, F. Menanteau, J. P. Hughes, M. Carrasco, and L. F. Barrientos. Strong lensing analysis of PLCK G004.5-19.5, a Planck-discovered cluster hosting a radio relic at $z = 0.52$. *A&A*, 562: A43, February 2014. doi: 10.1051/0004-6361/201321638.
- O. M. Smirnov and C. Tasse. Radio interferometric gain calibration as a complex optimization problem. *MNRAS*, 449(3):2668–2684, May 2015. doi: 10.1093/mnras/stv418.

-
- George C. Southworth. Early History of Radio Astronomy. *The Scientific Monthly*, 82(2):55–66, February 1956.
- T. A. T. Spoelstra. The influence of ionospheric refraction on radio astronomy interferometry. *A&A*, 120:313–321, April 1983.
- Woodruff T. Sullivan. The history of radio telescopes, 1945–1990. *Experimental Astronomy*, 25(1-3):107–124, August 2009. doi: 10.1007/s10686-009-9140-2.
- Terence Tao. Recent progress on the restriction conjecture. *arXiv Mathematics e-prints*, art. math/0311181, November 2003.
- C. Tasse. Nonlinear Kalman filters for calibration in radio interferometry. *Astronomy and Astrophysics*, 566:A127, Jun 2014a. doi: 10.1051/0004-6361/201423503.
- C. Tasse. Applying Wirtinger derivatives to the radio interferometry calibration problem. *arXiv e-prints*, October 2014b.
- C. Tasse, S. van der Tol, J. van Zwieten, G. van Diepen, and S. Bhatnagar. Applying full polarization A-Projection to very wide field of view instruments: An imager for LOFAR. *A&A*, 553:A105, May 2013. doi: 10.1051/0004-6361/201220882.
- C. Tasse, B. Hugo, M. Mirmont, O. Smirnov, M. Atemkeng, L. Bester, M. J. Hardcastle, R. Lakhoo, S. Perkins, and T. Shimwell. Faceting for direction-dependent spectral deconvolution. *A&A*, 611:A87, April 2018. doi: 10.1051/0004-6361/201731474.
- S. J. Tingay, R. Goeke, J. D. Bowman, D. Emrich, S. M. Ord, D. A. Mitchell, M. F. Morales, T. Boller, B. Crosse, R. B. Wayth, C. J. Lonsdale, S. Tremblay, D. Pallot, T. Colegate, A. Wicenc, N. Kudryavtseva, W. Arcus, D. Barnes, G. Bernardi, F. Briggs, S. Burns, J. D. Bunton, R. J. Cappallo, B. E. Corey, A. Deshpande, L. Desouza, B. M. Gaensler, L. J. Greenhill, P. J. Hall, B. J. Hazelton, D. Herne, J. N. Hewitt, M. Johnston-Hollitt, D. L. Kaplan, J. C. Kasper, B. B. Kincaid, R. Koenig, E. Kratzenberg, M. J. Lynch, B. McKinley, S. R. McWhirter, E. Morgan, D. Oberoi, J. Pathikulangara, T. Prabu, R. A. Remillard, A. E. E. Rogers, A. Rosh, J. E. Salah, R. J. Sault, N. Udaya-Shankar, F. Schlagenhauser, K. S. Srivani, J. Stevens, R. Subrahmanyam, M. Waterson, R. L. Webster, A. R. Whitney, A. Williams, C. L. Williams, and J. S. B. Wyithe. The Murchison Widefield Array: The Square Kilometre Array Precursor at Low Radio Frequencies. *PASA*, 30:e007, January 2013. doi: 10.1017/pasa.2012.007.
- R. van de Weygaert and W. Schaap. *The Cosmic Web: Geometric Analysis*, volume 665, pages 291–413. 2009. doi: 10.1007/978-3-540-44767-2_11.
- Rien van de Weygaert and Tjeerd S. van Albada. The Square Kilometer Array: new challenges for cosmology. *arXiv e-prints*, art. astro-ph/9602096, February 1996.
- S. van der Tol. *Bayesian estimation for ionospheric calibration in radio astronomy*. PhD thesis, TU Delft, 2009.
- M. P. van Haarlem, M. W. Wise, A. W. Gunst, G. Heald, J. P. McKean, J. W. T. Hessels, A. G. de Bruyn, R. Nijboer, J. Swinbank, R. Fallows, M. Brentjens, A. Nelles, R. Beck, H. Falcke, R. Fender, J. Hörandel, L. V. E. Koopmans, G. Mann, G. Miley, H. Röttgering, B. W. Stappers,

-
- R. A. M. J. Wijers, S. Zaroubi, M. van den Akker, A. Alexov, J. Anderson, K. Anderson, A. van Ardenne, M. Arts, A. Asgekar, I. M. Avruch, F. Batejat, L. Bähren, M. E. Bell, M. R. Bell, I. van Bemmell, P. Bennema, M. J. Bentum, G. Bernardi, P. Best, L. Bîrzan, A. Bonafede, A.-J. Boonstra, R. Braun, J. Bregman, F. Breitling, R. H. van de Brink, J. Broderick, P. C. Broekema, W. N. Brouw, M. Brüggen, H. R. Butcher, W. van Cappellen, B. Ciardi, T. Coenen, J. Conway, A. Coolen, A. Corstanje, S. Damstra, O. Davies, A. T. Deller, R.-J. Dettmar, G. van Diepen, K. Dijkstra, P. Donker, A. Doorduyn, J. Dromer, M. Drost, A. van Duin, J. Eislöffel, J. van Enst, C. Ferrari, W. Frieswijk, H. Gankema, M. A. Garrett, F. de Gasperin, M. Gerbers, E. de Geus, J.-M. Grießmeier, T. Grit, P. Gruppen, J. P. Hamaker, T. Hassall, M. Hoeft, H. A. Holties, A. Horneffer, A. van der Horst, A. van Houwelingen, A. Huijgen, M. Iacobelli, H. Intema, N. Jackson, V. Jelic, A. de Jong, E. Juette, D. Kant, A. Karastergiou, A. Koers, H. Kollen, V. I. Kondratiev, E. Kooistra, Y. Koopman, A. Koster, M. Kuniyoshi, M. Kramer, G. Kuper, P. Lambropoulos, C. Law, J. van Leeuwen, J. Lemaître, M. Loose, P. Maat, G. Macario, S. Markoff, J. Masters, R. A. McFadden, D. McKay-Bukowski, H. Meijering, H. Meulman, M. Mevius, E. Middelberg, R. Millenaar, J. C. A. Miller-Jones, R. N. Mohan, J. D. Mol, J. Morawietz, R. Morganti, D. D. Mulcahy, E. Mulder, H. Munk, L. Nieuwenhuis, R. van Nieuwpoort, J. E. Noordam, M. Norden, A. Noutsos, A. R. Offringa, H. Olofsson, A. Omar, E. Orrú, R. Overeem, H. Paas, M. Pandey-Pommier, V. N. Pandey, R. Pizzo, A. Polatidis, D. Rafferty, S. Rawlings, W. Reich, J.-P. de Reijer, J. Reitsma, G. A. Renting, P. Riemers, E. Rol, J. W. Romein, J. Roosjen, M. Ruiter, A. Scaife, K. van der Schaaf, B. Scheers, P. Schellart, A. Schoenmakers, G. Schoonderbeek, M. Serylak, A. Shulevski, J. Sluman, O. Smirnov, C. Sobey, H. Spreeuw, M. Steinmetz, C. G. M. Sterks, H.-J. Stiepel, K. Stuurwold, M. Tagger, Y. Tang, C. Tasse, I. Thomas, S. Thoudam, M. C. Toribio, B. van der Tol, O. Usov, M. van Veelen, A.-J. van der Veen, S. ter Veen, J. P. W. Verbiest, R. Vermeulen, N. Vermaas, C. Vocks, C. Vogt, M. de Vos, E. van der Wal, R. van Weeren, H. Weggemans, P. Weltevrede, S. White, S. J. Wijnholds, T. Wilhelmsson, O. Wucknitz, S. Yatawatta, P. Zarka, A. Zensus, and J. van Zwieten. LOFAR: The LOw-Frequency ARray. *A&A*, 556:A2, August 2013. doi: 10.1051/0004-6361/201220873.
- R. J. van Weeren, H. J. A. Röttgering, M. Brüggen, and A. Cohen. Diffuse radio emission in the merging cluster MACS J0717.5+3745: the discovery of the most powerful radio halo. *A&A*, 505:991–997, October 2009. doi: 10.1051/0004-6361/200912528.
- R. J. van Weeren, H. J. A. Röttgering, and M. Brüggen. Diffuse steep-spectrum sources from the 74 MHz VLSS survey. *A&A*, 527:A114, March 2011. doi: 10.1051/0004-6361/201015991.
- R. J. van Weeren, W. L. Williams, M. J. Hardcastle, T. W. Shimwell, D. A. Rafferty, J. Sabater, G. Heald, S. S. Sridhar, T. J. Dijkema, G. Brunetti, M. Brüggen, F. Andrade-Santos, G. A. Ogrean, H. J. A. Röttgering, W. A. Dawson, W. R. Forman, F. de Gasperin, C. Jones, G. K. Miley, L. Rudnick, C. L. Sarazin, A. Bonafede, P. N. Best, L. Bîrzan, R. Cassano, K. T. Chyży, J. H. Croston, T. Ensslin, C. Ferrari, M. Hoeft, C. Horellou, M. J. Jarvis, R. P. Kraft, M. Mevius, H. T. Intema, S. S. Murray, E. Orrú, R. Pizzo, A. Simionescu, A. Stroe, S. van der Tol, and G. J. White. LOFAR Facet Calibration. *ApJS*, 223:2, March 2016. doi: 10.3847/0067-0049/223/1/2.
- R. J. van Weeren, F. de Gasperin, H. Akamatsu, M. Brüggen, L. Feretti, H. Kang, A. Stroe, and F. Zandanel. Diffuse Radio Emission from Galaxy Clusters. *Space Sci. Rev.*, 215:16, February 2019. doi: 10.1007/s11214-019-0584-z.
- Reinout J. van Weeren, Huub J. A. Röttgering, Marcus Brüggen, and Matthias Hoeft. Particle

-
- Acceleration on Megaparsec Scales in a Merging Galaxy Cluster. *Science*, 330(6002):347, October 2010. doi: 10.1126/science.1194293.
- F. Vazza. How complex is the cosmic web? *MNRAS*, 491(4):5447–5463, February 2020. doi: 10.1093/mnras/stz3317.
- F. Vazza, G. Brunetti, C. Gheller, R. Brunino, and M. Brüggen. Massive and refined. II. The statistical properties of turbulent motions in massive galaxy clusters with high spatial resolution. *A&A*, 529:A17, May 2011. doi: 10.1051/0004-6361/201016015.
- F. Vazza, M. Brüggen, C. Gheller, and G. Brunetti. Modelling injection and feedback of cosmic rays in grid-based cosmological simulations: effects on cluster outskirts. *MNRAS*, 421: 3375–3398, April 2012. doi: 10.1111/j.1365-2966.2012.20562.x.
- H. K. Vedantham and L. V. E. Koopmans. Scintillation noise in widefield radio interferometry. *MNRAS*, 453(1):925–938, October 2015. doi: 10.1093/mnras/stv1594.
- H. K. Vedantham and L. V. E. Koopmans. Scintillation noise power spectrum and its impact on high-redshift 21-cm observations. *MNRAS*, 458(3):3099–3117, May 2016. doi: 10.1093/mnras/stw443.
- T. Venturi, S. Giacintucci, D. Dallacasa, R. Cassano, G. Brunetti, S. Bardelli, and G. Setti. GMRT radio halo survey in galaxy clusters at $z = 0.2\text{--}0.4$. II. The eBCS clusters and analysis of the complete sample. *A&A*, 484(2):327–340, June 2008. doi: 10.1051/0004-6361:200809622.
- T. Vernstrom, B. M. Gaensler, S. Brown, E. Lenc, and R. P. Norris. Low-frequency radio constraints on the synchrotron cosmic web. *MNRAS*, 467:4914–4936, Jun 2017. doi: 10.1093/mnras/stx424.
- Y. Weiss and W. T. Freeman. Correctness of belief propagation in gaussian graphical models of arbitrary topology. *Neural Computation*, 13(10):2173–2200, Oct 2001. ISSN 0899-7667. doi: 10.1162/089976601750541769.
- S. J. Wijnholds, A. G. Willis, and S. Salvini. Baseline-dependent averaging in radio interferometry. *MNRAS*, 476(2):2029–2039, May 2018. doi: 10.1093/mnras/sty360.
- C. H. Wilcox. Wave propagation in a random medium (lev a. chernov). *SIAM Review*, 4(1): 55–55, 1962. doi: 10.1137/1004017. URL <https://doi.org/10.1137/1004017>.
- W. L. Williams, G. Calistro Rivera, P. N. Best, M. J. Hardcastle, H. J. A. Röttgering, K. J. Duncan, F. de Gasperin, M. J. Jarvis, G. K. Miley, E. K. Mahony, L. K. Morabito, D. M. Nisbet, I. Prandoni, D. J. B. Smith, C. Tasse, and G. J. White. LOFAR-Boötes: properties of high- and low-excitation radio galaxies at $0.5 < z < 2.0$. *MNRAS*, 475(3):3429–3452, April 2018. doi: 10.1093/mnras/sty026.
- D. Wittor, M. Hoeft, F. Vazza, M. Brüggen, and P. Domínguez-Fernández. Polarization of radio relics in galaxy clusters. *MNRAS*, 490(3):3987–4006, December 2019. doi: 10.1093/mnras/stz2715.
- E. Wolf. Three-dimensional structure determination of semi-transparent objects from holographic data. *Optics Communications*, 1:153–156, September 1969. doi: 10.1016/0030-4018(69)90052-2.

- Lingfei Wu, Dashun Wang, and James A. Evans. Large teams develop and small teams disrupt science and technology. *Nature*, 566:1, Feb 2019. doi: 10.1038/s41586-019-0941-9.
- K. C. Yeh. Propagation of spherical waves through an ionosphere containing anisotropic irregularities. *Journal of Research of National Bureau of Standards*, 5:621–636, sep 1962. doi: 10.6028/jres.066D.062.
- K. C. Yeh and G. W. Swenson, Jr. The scintillation of radio signals from satellites. *J. Geophys. Res.*, 64:2281–2286, December 1959. doi: 10.1029/JZ064i012p02281.
- Fisher Yu and Vladlen Koltun. Multi-scale context aggregation by dilated convolutions. In *International Conference on Learning Representations (ICLR)*, May 2016.

Appendix A

Derivation of tomographic equivalence

We now explicitly prove the assertion that Eq. 2.24 is equal to Eq. 2.26, that is,

$$\int P(\boldsymbol{\tau} | \mathbf{n}_e) P(\mathbf{n}_e | \boldsymbol{\tau}_{\text{obs}}) d\mathbf{n}_e = \int P(\mathbf{n}_e, \boldsymbol{\tau} | \boldsymbol{\tau}_{\text{obs}}) d\mathbf{n}_e. \quad (\text{A.1})$$

We note that we sometimes use the notation $\mathcal{N}[a | m_a, C_a]$ which is equivalent to $a \sim \mathcal{N}[m_a, C_a]$.

We define the matrix representation of the DRI operator in Eq. 2.6, $\Delta_* \mathbf{n}_e = \{\Delta_{\mathbf{x}}^{\hat{\mathbf{k}}} n_e | (\mathbf{x}, \hat{\mathbf{k}}) \in \mathcal{S}_*\}$, and likewise let Δ be the matrix representation over the index set $\mathcal{S}_{\text{obs}}$. Similarly, the matrix representation of the FED kernel – the Gram matrix – is $\mathbf{K} = \{K(\mathbf{x}, \mathbf{x}') | \mathbf{x}, \mathbf{x}' \in \mathcal{X}\}$. Using these matrix representation we have the following joint distribution,

$$P(\mathbf{n}_e, \boldsymbol{\tau}, \boldsymbol{\tau}_{\text{obs}}) = \mathcal{N} \begin{bmatrix} \tilde{n}_e & \mathbf{K} & \mathbf{K} \Delta_*^T & \mathbf{K} \Delta^T \\ 0 & \Delta_* \mathbf{K} & \Delta_* \mathbf{K} \Delta_*^T & \Delta_* \mathbf{K} \Delta^T \\ 0 & \Delta \mathbf{K} & \Delta \mathbf{K} \Delta_*^T & \Delta \mathbf{K} \Delta^T + \sigma^2 \mathbf{I} \end{bmatrix}. \quad (\text{A.2})$$

Let us first work out the left-hand side (LHS) of Eq. A.1. Because $\boldsymbol{\tau} = \Delta_* \mathbf{n}_e$, and using standard Gaussian identities we have,

$$P(\boldsymbol{\tau} | \mathbf{n}_e) = \mathcal{N} \left[\underbrace{\Delta_* \mathbf{K} \mathbf{K}^{-1} (\mathbf{n}_e - \tilde{n}_e)}_{\Delta_*(\mathbf{n}_e - \tilde{n}_e)}, \underbrace{\Delta_* \mathbf{K} \Delta_* - \Delta_* \mathbf{K} \mathbf{K}^{-1} \mathbf{K} \Delta_*}_{\mathbf{0}} \right]. \quad (\text{A.3})$$

Similarly, the second distribution on the LHS is,

$$P(\mathbf{n}_e | \boldsymbol{\tau}_{\text{obs}}) = \mathcal{N} [\tilde{n}_e + \mathbf{K} \Delta^T (\Delta \mathbf{K} \Delta^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\tau}_{\text{obs}}, \mathbf{K} - \mathbf{K} \Delta^T (\Delta \mathbf{K} \Delta^T + \sigma^2 \mathbf{I})^{-1} \Delta \mathbf{K}]. \quad (\text{A.4})$$

We now can use standard Gaussian identities [e.g. Weiss and Freeman, 2001] to evaluate the

integral on the LHS,

$$\int P(\boldsymbol{\tau} | \mathbf{n}_e) P(\mathbf{n}_e | \boldsymbol{\tau}_{\text{obs}}) d\mathbf{n}_e = \mathcal{N}[\boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\tau}_{\text{obs}}, \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}_*^T - \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}_*^T] \quad (\text{A.5})$$

In order to work out the right-hand side (RHS), we simply condition Eq. A.2 on $\boldsymbol{\tau}_{\text{obs}}$ and then marginalise $\mathbf{n}_e$ by selecting the corresponding sub-block of the Gaussian,

$$P(\mathbf{n}_e, \boldsymbol{\tau} | \boldsymbol{\tau}_{\text{obs}}) = \mathcal{N} \left[\begin{pmatrix} \bar{\mathbf{n}}_e \\ 0 \end{pmatrix} + \begin{pmatrix} \mathbf{K} \boldsymbol{\Delta}^T \\ \boldsymbol{\Delta}_*^T \mathbf{K} \boldsymbol{\Delta}^T \end{pmatrix} (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\tau}_{\text{obs}}, \begin{pmatrix} \bar{\mathbf{K}} & \boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}_*^T \\ \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T & \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}_*^T \end{pmatrix} - \begin{pmatrix} \mathbf{K} \boldsymbol{\Delta}^T \\ \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T \end{pmatrix} (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \begin{pmatrix} \boldsymbol{\Delta} \mathbf{K} & \boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}_*^T \end{pmatrix} \right] \quad (\text{A.6})$$

Marginalising over $\mathbf{n}_e$ is equivalent to neglecting the sub-block corresponding to $\mathbf{n}_e$. Therefore, the RHS is,

$$\int P(\mathbf{n}_e, \boldsymbol{\tau} | \boldsymbol{\tau}_{\text{obs}}) d\mathbf{n}_e = \mathcal{N}[\boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\tau}_{\text{obs}}, \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}_*^T - \boldsymbol{\Delta}_* \mathbf{K} \boldsymbol{\Delta}^T (\boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}^T + \sigma^2 \mathbf{I})^{-1} \boldsymbol{\Delta} \mathbf{K} \boldsymbol{\Delta}_*^T]. \quad (\text{A.7})$$

■

Appendix B

Derivation of the ΔTEC variance function and its limits

We derive the ΔTEC variance function $\sigma_{\Delta\text{TEC}}^2(d)$ for zenith observations ($\mathbf{k} = \mathbf{k}' = \hat{\mathbf{z}}$) by considering a baseline between an antenna-of-interest at $\mathbf{x}_i = \mathbf{x}_j$ and a reference antenna at $\mathbf{x}_0 = \mathbf{0}$. To use the Pythagorean theorem later, we assume that this baseline lies in the plane of the local horizon, i.e. perpendicular to the zenith. Without loss of generality, we can orient the coordinate axes such that this baseline lies along the $\hat{\mathbf{x}}$ direction, so that $\mathbf{x}_i - \mathbf{x}_0 = d\hat{\mathbf{x}}$. Here $d \triangleq \|\mathbf{x}_i\|$ is the distance between the two antennae. We then take the general covariance function $K_{\Delta\text{TEC}}([\mathbf{x}_i, \mathbf{x}_0, \hat{\mathbf{k}}], [\mathbf{x}_j, \mathbf{x}_0, \hat{\mathbf{k}}'])$, and find that in this particular case

$$\sigma_{\Delta\text{TEC}}^2(d) \triangleq K_{\Delta\text{TEC}}([\mathbf{x}_i, \mathbf{x}_0, \hat{\mathbf{z}}], [\mathbf{x}_i, \mathbf{x}_0, \hat{\mathbf{z}}]) \quad (\text{B.1})$$

$$= \sum_{p_1=0}^1 \sum_{p_2=0}^1 (-1)^{p_1+p_2} \int_0^b \int_0^b K_{n_e}(\|\mathbf{x}_{(1-p_1)i} - \mathbf{x}_{(1-p_2)i} + \hat{\mathbf{z}}(s_1 - s_2)\|) \, ds_1 \, ds_2, \quad (\text{B.2})$$

where K_{n_e} is an arbitrary stationary and isotropic kernel (such as the Exponentiated Quadratic and Matérn $\frac{3}{2}$ kernels considered earlier) for the FED. The two terms where p_1 and p_2 are equal give the same contribution, as do the two terms for which p_1 and p_2 are unequal. By subsequently applying the Pythagorean theorem in this last case (i.e. $p_1 = 0$ and $p_2 = 1$, and vice versa), we find

$$\sigma_{\Delta\text{TEC}}^2(d) = 2 \int_0^b \int_0^b K_{n_e}(\|s_1 - s_2\|) - K_{n_e}(\sqrt{d^2 + (s_1 - s_2)^2}) \, ds_1 \, ds_2. \quad (\text{B.3})$$

We manipulate this result to obtain a more insightful expression. First, we note the (implicit) presence of three parameters with dimension length: ionospheric thickness b , reference antenna distance d , and FED kernel half-peak distance h . We perform transformations to dimensionless coordinates $u_1 = \frac{s_1}{h}$ and $u_2 = \frac{s_2}{h}$ to reveal that the *shape* - though not the absolute scale - of the function $\sigma_{\Delta\text{TEC}}^2(d)$ is governed only by the length-scale ratios $\frac{b}{h}$ and $\frac{d}{h}$, and the particular functional form of K_{n_e} .

Furthermore, for stationary covariance functions, we have $K_{n_e} = \sigma_{n_e}^2 C_{n_e}$, where C_{n_e} is the corresponding dimensionless correlation function.

These considerations enable us to express the ΔTEC structure function as a dimensionless, shape-determining double integral appended by dimensionful prefactors; i.e.

$$\sigma_{\Delta\text{TEC}}^2(d) = 2\sigma_{n_e}^2 h^2 \int_0^{\frac{b}{h}} \int_0^{\frac{b}{h}} C_{n_e}(h|u_1 - u_2|) - C_{n_e}\left(h\sqrt{\left(\frac{d}{h}\right)^2 + (u_1 - u_2)^2}\right) du_1 du_2. \quad (\text{B.4})$$

We first note that the variance of ΔTEC is simply proportional to the variance of n_e . Secondly, we note that $h|u_1 - u_2| < h\sqrt{\left(\frac{d}{h}\right)^2 + (u_1 - u_2)^2}$ for any non-zero d , so that $C_{n_e}(h|u_1 - u_2|) > C_{n_e}\left(h\sqrt{\left(\frac{d}{h}\right)^2 + (u_1 - u_2)^2}\right)$ for all monotonically decreasing correlation functions C_{n_e} (or, equivalently, covariance functions K_{n_e}). With the integrand always positive, we see that the integral must be a strictly increasing function of $\frac{b}{h}$ (which occurs in the integration limits). Therefore, we conclude that for stationary, isotropic, and monotonically decreasing (SIMD) FED kernels with HPD h , the ΔTEC variance increases monotonically with the thickness of the ionosphere b . Simply put: thicker SIMD ionospheres cause larger ΔTEC variations.

Let us now consider three limits of the ΔTEC zenith variance function, that all do not require K_{FED} to decrease monotonically. In the short-baseline limit, i.e. $\frac{d}{h} \rightarrow 0$, we have $C_{n_e}\left(h\sqrt{\left(\frac{d}{h}\right)^2 + (u_1 - u_2)^2}\right) \rightarrow C_{n_e}(h|u_1 - u_2|)$. We therefore find that $\sigma_{\Delta\text{TEC}}^2 \rightarrow 0$ irrespective of other parameters, recovering that the variance of ΔTEC vanishes near the reference antenna. In the long-baseline limit, i.e. $\frac{d}{h} \gg \frac{b}{h} > 1$, we see that $\sqrt{\left(\frac{d}{h}\right)^2 + (u_1 - u_2)^2} \approx \frac{d}{h}$, since $(u_1 - u_2)^2 < \left(\frac{b}{h}\right)^2 \ll \left(\frac{d}{h}\right)^2$. Assuming $C_{n_e}(d) \approx 0$ when $\frac{d}{h} \gg 1$, the integrand reduces to $C_{n_e}(h|u_1 - u_2|) - C_{n_e}\left(h \cdot \frac{d}{h}\right) \approx C_{n_e}(h|u_1 - u_2|)$. We find that in this case,

$$\sigma_{\Delta\text{TEC}}^2 \approx 2\sigma_{n_e}^2 h^2 \int_0^{\frac{b}{h}} \int_0^{\frac{b}{h}} C_{n_e}(h|u_1 - u_2|) du_1 du_2. \quad (\text{B.5})$$

This is the plateau value of the ΔTEC variance that our model predicts for the long-baseline limit.

Another way to arrive at the plateau value expression of Equation B.5 is by considering the statistical properties of TEC first. In a computation analogous to the one for ΔTEC in Section 2.3, one can derive the general TEC covariance function K_{TEC} . The variance of $\tau_i^{\hat{z}}$ (the TEC of antenna i while observing towards the zenith $\hat{z}$) is straightforwardly shown to be

$$\mathbb{V}(\tau_i^{\hat{z}}) = \sigma_{n_e}^2 h^2 \int_0^{\frac{b}{h}} \int_0^{\frac{b}{h}} C_{n_e}(h|u_1 - u_2|) du_1 du_2. \quad (\text{B.6})$$

We highlight the absence of a dependence on i at the RHS. As a ΔTEC is simply a TEC differenced with a TEC for a reference antenna observing in the same direction, we have

$$\sigma_{\Delta\text{TEC}}^2 = \mathbb{V}(\tau_i^{\hat{z}} - \tau_0^{\hat{z}}) = \mathbb{V}(\tau_i^{\hat{z}}) + \mathbb{V}(\tau_0^{\hat{z}}), \quad (\text{B.7})$$

where the second equality only holds when the TECs are independent. This is exactly the scenario considered in the long-baseline limit. Plugging in Equation B.6 recovers the plateau level. We can find a general upper bound to the variance of ΔTEC in terms of physical parameters. To this end, we note that the integrand in Equation B.4 is maximised when, over the full range of integration, the value of the first term is 1 whilst the second term is equal to the infimum of the correlation function. Calling $\inf_{\mathbb{R}} \{C_{n_e}(r) : r \in \mathbb{R}_{>0}\} \triangleq I$, we find the inequality,

$$\sigma_{\Delta\text{TEC}}^2 \leq 2\sigma_{n_e}^2 h^2 \int_0^{\frac{b}{h}} \int_0^{\frac{b}{h}} 1 - I \, du_1 du_2 = 2(1 - I) \sigma_{n_e}^2 b^2. \quad (\text{B.8})$$

For strictly positive FED kernels that decay to zero at large distances (such as the EQ and Matérn kernels considered in this work), we find $\sigma_{\Delta\text{TEC}}^2 \leq 2\sigma_{n_e}^2 b^2$. Kernels resulting in anti-correlated FEDs produce the constraint $\sigma_{\Delta\text{TEC}}^2 \leq 4\sigma_{n_e}^2 b^2$ or tighter. By measuring $\sigma_{\Delta\text{TEC}}(d)$, one can bound the product $\sigma_{n_e} b$ from below. The strongest bound is obtained for large d .

Appendix C

Factoring commutative DI dependence from the RIME

We consider the effect of directionally referencing spatially referenced commutative Jones scalars. In particular, we study phases of the Jones scalars, and assume amplitudes of one, but the same idea extends to amplitude by considering log-amplitudes that can be treated like a pure-imaginary phase.

Let $g(\mathbf{x}, \mathbf{k}) = e^{i\phi(\mathbf{x}, \mathbf{k})}$ be a Jones scalar, and consider the necessarily non-unique decomposition of phase into DD and DI components,

$$\phi(\mathbf{x}, \mathbf{k}) = \phi^{\text{DD}}(\mathbf{x}, \mathbf{k}) + \phi^{\text{DI}}(\mathbf{x}). \quad (\text{C.1})$$

This functional form only specifies that the DI term is not dependent on $\mathbf{k}$. The differential phase to which the RIME is sensitive is found by spatially referencing the phase,

$$\Delta_0 \phi(\mathbf{x}, \mathbf{k}) = \phi(\mathbf{x}, \mathbf{k}) - \phi(\mathbf{x}_0, \mathbf{k}) \quad (\text{C.2})$$

$$= \Delta_0 \phi^{\text{DD}}(\mathbf{x}, \mathbf{k}) + \Delta_0 \phi^{\text{DI}}(\mathbf{x}). \quad (\text{C.3})$$

Directionally referencing the differential phase to direction $\mathbf{k}_0$, we have

$$\Delta_0^2 \phi(\mathbf{x}, \mathbf{k}) = \Delta_0 \phi(\mathbf{x}, \mathbf{k}) - \Delta_0 \phi(\mathbf{x}, \mathbf{k}_0) \quad (\text{C.4})$$

$$= \Delta_0^2 \phi^{\text{DD}}(\mathbf{x}, \mathbf{k}). \quad (\text{C.5})$$

We see that the DI phase has disappeared, and we are left with the doubly differential phase for the DD term. We then assume that there was a remnant DI component in $\phi^{\text{DD}}(\mathbf{x}, \mathbf{k})$. Then by induction we have that $\Delta_0^2 \phi^{\text{DD}}(\mathbf{x}, \mathbf{k})$ must be free of DI terms, and $\Delta_0^2 \phi(\mathbf{x}, \mathbf{k})$ must be free of all DI components. It follows that directionally referencing phase guarantees that all DI components are removed.

Furthermore, assuming we were to directionally reference a doubly differential phase. We assume that the doubly differential phase is referenced to direction $\mathbf{k}'_0$, with $\Delta_0'^2 \phi(\mathbf{x}, \mathbf{k}) \triangleq$

$\Delta_0\phi(\mathbf{x}, \mathbf{k}) - \Delta_0\phi(\mathbf{x}, \mathbf{k}'_0)$. Then we see

$$\Delta_0'^2\phi(\mathbf{x}, \mathbf{k}) - \Delta_0'^2\phi(\mathbf{x}, \mathbf{k}_0) = \Delta_0\phi(\mathbf{x}, \mathbf{k}) - \Delta_0\phi(\mathbf{x}, \mathbf{k}'_0) - (\Delta_0\phi(\mathbf{x}, \mathbf{k}_0) - \Delta_0\phi(\mathbf{x}, \mathbf{k}'_0)) \quad (\text{C.6})$$

$$= \Delta_0\phi(\mathbf{x}, \mathbf{k}) - \Delta_0\phi(\mathbf{x}, \mathbf{k}_0) \quad (\text{C.7})$$

$$= \Delta_0^2\phi(\mathbf{x}, \mathbf{k}). \quad (\text{C.8})$$

That is, directionally referencing a doubly differential phase produces a new doubly differential phase, referenced to the new direction, $\mathbf{k}_0$. This trick thus allows us to set a well-defined reference direction to phases that have undergone DI calibration before DD calibration.

Appendix D

Recursive Bayesian estimation




Figure D.1: Casual graph depicting a hidden Markov model.

Recursive Bayesian estimation is a method of performing inference on a hidden Markov model [HMM; Rabiner and Juang, 1986]. Let $\mathbf{y}$ be an observable and $\mathbf{x}$ be a hidden variable. The HMM assumptions on $\mathbf{x}$ and $\mathbf{y}$ are, firstly, that the hidden random variable is only conditionally dependent on its previous state, and secondly, the observable is conditionally independent of all other random variables except for the current hidden state. These assumptions are depicted in **Figure D.1**, where i is the sequence index. This paradigm is often given a notion of causality or time, but this is not necessary in any way. The sequence index is an abstract notion that simply explains how the set of hidden variables are traversed. For example, we assume that the observations are frames of a movie, and the hidden variable is the plot contained in each movie frame. The HMM assumption is that the movie plot is linear, and the picture encodes what is going on in the movie at a given point in time. With recursive Bayesian estimation the movie can be watched with frames randomly ordered and the complete plot is still rendered completely.

There are two distinct types of information propagation in a hidden Markov model. Information can flow in the direction of the arrows, or against them. This leads to the notion

of the forward and backwards equations that describe how belief in hidden variables is propagated forward, and for revising, we believe in previously visited hidden variables.

The joint distribution of the hidden random variables and observables in a chain of length T can be written out as a product of conditional distributions and a marginal using the product rule of probability distributions [Kolmogorov, 1960]. Because of the HMM assumptions the joint distribution is

$$p(\mathbf{x}_{0:T}, \mathbf{y}_{0:T}) = p(\mathbf{x}_0) \prod_{i=1}^T p(\mathbf{x}_i | \mathbf{x}_{i-1}) p(\mathbf{y}_i | \mathbf{x}_i). \quad (\text{D.1})$$

We first consider propagating information forward. This is done in two steps, typically called the predict and update steps. For the predict step, we consider how belief in the absence of new observables is propagated. For this, we apply the Chapman-Kolmogorov identity¹ for Markovian processes,

$$p(\mathbf{x}_{i+1} | \mathbf{y}_{0:i}) = \mathbb{E}_{\mathbf{x}_i | \mathbf{y}_{0:i}} [p(\mathbf{x}_{i+1} | \mathbf{x}_i)], \quad (\text{D.2})$$

which gives us the probability distribution of the hidden variables at time $i + 1$ in terms of the so-called state transition distribution $p(\mathbf{x}_{i+1} | \mathbf{x}_i)$ and posterior distribution $p(\mathbf{x}_i | \mathbf{y}_{0:i})$ at index i . The current prior belief can be understood as the expectation of the state transition distribution over the measure of the current posterior belief.

The update step is simply an application of the Bayes theorem with the prior defined by the predict step,

$$p(\mathbf{x}_i | \mathbf{y}_{0:i}) = \frac{p(\mathbf{y}_i | \mathbf{x}_i) \mathbb{E}_{\mathbf{x}_{i-1} | \mathbf{y}_{0:i-1}} [p(\mathbf{x}_i | \mathbf{x}_{i-1})]}{p(\mathbf{y}_i | \mathbf{y}_{0:i-1})}, \quad (\text{D.3})$$

where the denominator is the Bayesian evidence of the newly arrived data given all previous data, and it is independent of the hidden variables. Equation D.3 gives a recurrence relation for propagating our belief forward, therefore this is called the forward equation.

We now assume that we are at index T , and wish to use all acquired information to revise our belief in the previously visited hidden variables at indices $i < T$. The trick is to realise that $p(\mathbf{x}_i | \mathbf{x}_{i+1} | \mathbf{y}_{0:T}) = p(\mathbf{x}_i | \mathbf{x}_{i+1} | \mathbf{y}_{0:i})$ as a result of the Markov properties. In this case, again using the product rule, we find that the joint conditional distribution of a pair of sequential hidden states given the whole sequence of data is

$$p(\mathbf{x}_i, \mathbf{x}_{i+1} | \mathbf{y}_{0:T}) = \frac{p(\mathbf{x}_{i+1} | \mathbf{x}_i) p(\mathbf{x}_i | \mathbf{y}_{0:i}) p(\mathbf{x}_{i+1} | \mathbf{y}_{0:T})}{p(\mathbf{x}_{i+1} | \mathbf{y}_{0:i})}. \quad (\text{D.4})$$

Marginalising the second hidden parameter, we arrive at the recurrence relation

$$p(\mathbf{x}_i | \mathbf{y}_{0:T}) = p(\mathbf{x}_i | \mathbf{y}_{0:i}) \int \frac{p(\mathbf{x}_{i+1} | \mathbf{x}_i) p(\mathbf{x}_{i+1} | \mathbf{y}_{0:T})}{p(\mathbf{x}_{i+1} | \mathbf{y}_{0:i})} d\mathbf{x}_{i+1}. \quad (\text{D.5})$$

This can be solved by starting at T and solving this equation iteratively backwards, therefore Equation D.5 is called the backward equation. Most importantly, we note that the backward equation does not require conditioning on data, as was done in the update step.

¹If b is conditionally independent of a , then $p(a | c) = \int p(a | b) p(b | c) db = \mathbb{E}_{b|c} [p(a | b)]$ is the Chapman-Kolmogorov identity.

When the transition and likelihood are assumed to be Gaussian, for example, in a linear dynamical system, the forward and backward equations are equivalent to the well-known Kalman filter equations and Rauch smoother equations [Rauch, 1963], respectively.

Appendix E

Jones scalar variational expectation

Let $\mathbf{g} \in \mathbb{C}^{N_{\text{freq}}}$ be an observed complex Jones scalar vector, with amplitudes $g \in \mathbb{R}^{N_{\text{freq}}}$ and phases $\phi \in \mathbb{R}^{N_{\text{freq}}}$. We assume that the Jones scalars have complex Gaussian noise, described by the observational covariance matrix Σ . Thus, we have that the observational likelihood of the Jones scalars (*cf* Eq. 3.9) is

$$p(\mathbf{g} | \phi, g, \Sigma) = \mathcal{N}_{\mathbb{C}}[\mathbf{g} | g e^{i\phi}, \Sigma], \quad (\text{E.1})$$

where $\mathcal{N}_{\mathbb{C}}$ is the complex Gaussian distribution, which is defined as the Gaussian distribution of the stacked real and imaginary components. We define the residuals $\delta R = \text{Re}[\mathbf{g}] - g \cos \phi$ and $\delta I = \text{Im}[\mathbf{g}] - g \sin \phi$, and the stacked residuals $\delta \mathbf{g} = (\delta R, \delta I)^T$. Then the log-likelihood becomes

$$\log p(\mathbf{g} | \phi, g, \Sigma) = -N_{\text{freq}} \log 2\pi - \frac{1}{2} \log |\Sigma| - \frac{1}{2} \delta \mathbf{g}^T \Sigma^{-1} \delta \mathbf{g}, \quad (\text{E.2})$$

$$= -N_{\text{freq}} \log 2\pi - \frac{1}{2} \log |\Sigma| - \frac{1}{2} \text{Tr}[\Sigma^{-1} \delta \mathbf{g} \delta \mathbf{g}^T] \quad (\text{E.3})$$

$$= -N_{\text{freq}} \log 2\pi - \frac{1}{2} \log |\Sigma| - \frac{1}{2} \text{vec}[\Sigma^{-1}]^T \text{vec}[\delta \mathbf{g} \delta \mathbf{g}^T] \quad (\text{E.4})$$

where we used the fact that the trace of a scalar is a scalar, and that $\text{Tr}[A^T B] = \text{Tr}[B A^T] = \text{vec}[A]^T \text{vec}[B]$.

Now we assume that the phases are linearly modelled according to

$$\phi(\nu) = \sum_i^M f_i(\nu) a_i, \quad (\text{E.5})$$

$$\triangleq f^T(\nu) a \quad (\text{E.6})$$

where $f_i(\nu)$ is the i -th basis function of a set of M linearly independent functions depending on frequency ν , and a_i is the i -th parameter in the linear model. In the second line we have

written the sum as a dot product. We furthermore assume that the parameter vector, a , is a (potentially correlated) Gaussian random variable,

$$a \sim \mathcal{N}[\mu, \Gamma]. \quad (\text{E.7})$$

This induces a distribution on phase via Eq. E.5.

The variational expectation of the Jones scalars with respect to a is the expectation of the log-likelihood,

$$\begin{aligned} \mathbb{E}_a [\log p(\mathbf{g} | \phi, g, \Sigma)] = & -N_{\text{freq}} \log 2\pi - \frac{1}{2} \log |\Sigma| \\ & - \frac{1}{2} \text{vec}[\Sigma^{-1}]^T \text{vec}[\mathbb{E}_a[\delta \mathbf{g} \delta \mathbf{g}^T]], \end{aligned} \quad (\text{E.8})$$

where we used the fact that expectation and vectorisation commute. Now we need to evaluate the expectation, $\mathbb{E}_a[\delta \mathbf{g} \delta \mathbf{g}^T]$. In order to do this, we realise that there are three explicit expectations that must be done corresponding to sub-blocks of $\delta \mathbf{g} \delta \mathbf{g}^T$ consisting of the real-real, real-imaginary, and imaginary-imaginary outer products of residual vectors,

$$\mathbb{E}_a[\delta R \delta R^T] = \int \delta R \delta R^T dp(a) \quad (\text{E.9})$$

$$\triangleq I_{rr} \quad (\text{E.10})$$

$$\mathbb{E}_a[\delta R \delta I^T] = \int \delta R \delta I^T dp(a) \quad (\text{E.11})$$

$$\triangleq I_{ri} \quad (\text{E.12})$$

$$\mathbb{E}_a[\delta I \delta I^T] = \int \delta I \delta I^T dp(a) \quad (\text{E.13})$$

$$\triangleq I_{ii}, \quad (\text{E.14})$$

where $p(a) = \mathcal{N}[\mu, \Gamma]$. All three of the above integrals involve Gaussian expectations of trigonometric functions. For example, the (i, j) -th element of the integral I_{rr} is,

$$(I_{rr})_{(i,j)} = \int (\text{Re}[\mathbf{g}]_i - g_i \cos \phi(\nu_i))(\text{Re}[\mathbf{g}]_j - g_j \cos \phi(\nu_j)) dp(a). \quad (\text{E.15})$$

To evaluate this integral, we use angle addition formulae to reduce all powers of trigonometric functions. We then use the characteristic function of the Gaussian distribution, which gives the relation, $\mathbb{E}_{X \sim \mathcal{N}[\mu, \Gamma]}[e^{if^T X}] = e^{if^T \mu - \frac{1}{2} f^T \Gamma f}$, on the exponential form of cosine and sine. Applying the above, we evaluate all integral matrix elements,

$$\begin{aligned} (I_{rr})_{(i,j)} = & \text{Re}[\mathbf{g}]_i \text{Re}[\mathbf{g}]_j - g_i \text{Re}[\mathbf{g}]_j h_1(f(\nu_i)) - \text{Re}[\mathbf{g}]_i g_j h_1(f(\nu_j)) \\ & + g_i g_j h_3(f(\nu_i), f(\nu_j)) \end{aligned} \quad (\text{E.16})$$

$$\begin{aligned} (I_{ri})_{(i,j)} = & \text{Re}[\mathbf{g}]_i \text{Im}[\mathbf{g}]_j - g_i \text{Im}[\mathbf{g}]_j h_1(f(\nu_i)) - \text{Re}[\mathbf{g}]_i g_j h_2(f(\nu_j)) \\ & + g_i g_j h_5(f(\nu_i), f(\nu_j)) \end{aligned} \quad (\text{E.17})$$

$$\begin{aligned} (I_{ii})_{(i,j)} = & \text{Im}[\mathbf{g}]_i \text{Im}[\mathbf{g}]_j - g_i \text{Im}[\mathbf{g}]_j h_2(f(\nu_i)) - \text{Im}[\mathbf{g}]_i g_j h_2(f(\nu_j)) \\ & + g_i g_j h_4(f(\nu_i), f(\nu_j)) \end{aligned} \quad (\text{E.18})$$

where,

$$h_1(x) = \mathbb{E}_a[\cos x^T a] = e^{-\frac{1}{2}x^T \Gamma x} \cos x^T \mu \quad (\text{E.19})$$

$$h_2(x) = \mathbb{E}_a[\sin x^T a] = e^{-\frac{1}{2}x^T \Gamma x} \sin x^T \mu \quad (\text{E.20})$$

$$h_3(x, y) = \mathbb{E}_a[\cos x^T a \cos y^T a] = \frac{1}{2}(h_1(x + y) + h_1(x - y)) \quad (\text{E.21})$$

$$h_4(x, y) = \mathbb{E}_a[\sin x^T a \sin y^T a] = \frac{1}{2}(h_1(x - y) - h_1(x + y)) \quad (\text{E.22})$$

$$h_5(x, y) = \mathbb{E}_a[\cos x^T a \sin y^T a] = \frac{1}{2}(h_2(x + y) + h_2(x - y)). \quad (\text{E.23})$$

This completes the evaluation of Eq. E.8 for an arbitrary linear phase model. For example, in a phase model consisting only of DDTEC, we have $M = 1$, $f_1(\nu) = \frac{\kappa}{\nu}$ and $a_1 = \Delta_0^2 \tau$. It is simple to calculate the variational expectation of other linear phase models including terms such as clock-like terms, constant-in-frequency terms, and higher order ionospheric terms.

A simplification follows if the observational errors are assumed uncorrelated, which leads to a reduction of the number of operations required to compute the variational expectation. In this case, only the diagonals of I_{rr} and I_{ii} are needed, and I_{ri} is not needed at all.

Appendix F

Derivation of the M-step

Here we derive the maximisation step of Algorithm 1. This is an extension of Shumway and Stoffer [1982] to time varying observational covariance and state transition covariance. In order to do this we must linearise the forward equation. Let the set of HMM states be $X = \{x_i \mid i = 1..n\}$ and the set of observations be $Y = \{y_i \mid i = 1..n\}$. Following similar notation as in Shumway and Stoffer [1982] we'll term x_i^s be the state at step i conditioned on data up to an including step s . Therefore if $i > s$ then x_i^s is the forecast of the state. Consequently the conditional mean and covariance is μ_i^s and Γ_i^s . We'll also need the joint-covariance $\Gamma_{i,i-1}^s$ which is the covariance of $\{x_i, x_{i-1}\}$ conditioned on data up to step s . The recursive equations for all these components are given in [e.g. Shumway and Stoffer, 1982] and follow from the forward and backward equations in Appendix D.

Since at each time step the state is a Gaussian variable, using standard identities it follows that the posterior distribution of $x_i - x_{i-1}$ conditioned up to step n is given by,

$$x_i - x_{i-1} \sim \mathcal{N}(x_i - x_{i-1} \mid \mu_i^n - \mu_{i-1}^n, \Gamma_i^n + \Gamma_{i-1}^n - \Gamma_{i,i-1}^n - \Gamma_{i-1,i}^n). \quad (\text{E.1})$$

The joint log-likelihood of the data and state is the sum of the prior component and the likelihood component,

$$\begin{aligned} \log L(X, Y) &\triangleq -\frac{1}{2} \log |\Gamma_0| - \frac{1}{2} (x_0 - \mu_0)^T \Gamma_0^{-1} (x_0 - \mu_0) \\ &\quad - \frac{1}{2} \sum_{i=1}^n \log |\Omega_i| - \frac{1}{2} \sum_{i=1}^n (x_i - x_{i-1})^T \Omega_i^{-1} (x_i - x_{i-1}) \\ &\quad - \frac{1}{2} \sum_{i=1}^n \log |\Sigma_i| - \frac{1}{2} \sum_{i=1}^n (y_i - f_i(x_i))^T \Sigma_i^{-1} (y_i - f_i(x_i)) + \text{const.} \end{aligned} \quad (\text{E.2})$$

where f_i is the forward function from the HMM state to observable and time i . In general, the function f_i can change over time, e.g. deterministic input control parameters to f_i can be time dependent. Let us define the linear map $D_i(y) = \nabla_x f_i|_{\tilde{x}_i} y$ for some $\tilde{x}_i$. The forward

function then satisfies $f_i(x_i) = f_i(\bar{x}_i) + D_i(x_i - \bar{x}_i) + o(|x_i - \bar{x}_i|^2)$. From this Eq. F2 becomes,

$$\begin{aligned} \log L(X, Y) = & -\frac{1}{2} \log |\Gamma_0| - \frac{1}{2} (x_0 - \mu_0)^T \Gamma_0^{-1} (x_0 - \mu_0) \\ & - \frac{1}{2} \sum_{i=1}^n \log |\Omega_i| - \frac{1}{2} \sum_{i=1}^n (x_i - x_{i-1})^T \Omega_i^{-1} (x_i - x_{i-1}) \\ & - \frac{1}{2} \sum_{i=1}^n \log |\Sigma_i| \\ & - \frac{1}{2} \sum_{i=1}^n (y_i - f_i(\bar{x}_i) - D_i(x_i - \bar{x}_i))^T \Sigma_i^{-1} (y_i - f_i(\bar{x}_i) - D_i(x_i - \bar{x}_i)) \\ & + \text{const.} \end{aligned} \quad (\text{F3})$$

which is exactly equal to Eq. F2 when $x_i = \bar{x}_i$ and f_i quadratic in the neighbourhood of $\bar{x}_i$. Defining $\delta y_i = y_i - f_i(\bar{x}_i) - D_i \bar{x}_i$ then Eq. F3 becomes,

$$\begin{aligned} \log L(X, Y) = & -\frac{1}{2} \log |\Gamma_0| - \frac{1}{2} (x_0 - \mu_0)^T \Gamma_0^{-1} (x_0 - \mu_0) \\ & - \frac{1}{2} \sum_{i=1}^n \log |\Omega_i| - \frac{1}{2} \sum_{i=1}^n (x_i - x_{i-1})^T \Omega_i^{-1} (x_i - x_{i-1}) \\ & - \frac{1}{2} \sum_{i=1}^n \log |\Sigma_i| \\ & - \frac{1}{2} \sum_{i=1}^n (\delta y_i - D_i x_i)^T \Sigma_i^{-1} (\delta y_i - D_i x_i) \\ & + \text{const.} \end{aligned} \quad (\text{F4})$$

Taking the expectation of Eq. F4 with respect to the state conditioned on all data we find [cf. Shumway and Stoffer, 1982],

$$\begin{aligned} \mathbb{E}_X[\log L(X, Y)] = & -\frac{1}{2} \log |\Gamma_0| - \frac{1}{2} \sum_{i=1}^n \log |\Omega_i| - \frac{1}{2} \sum_{i=1}^n \log |\Sigma_i| \\ & - \frac{1}{2} (\mu_0^n - \mu_0)^T \Gamma_0^{-1} (\mu_0^n - \mu_0) - \frac{1}{2} \text{Tr}[\Gamma_0^{-1} \Gamma_0^n] \\ & - \frac{1}{2} \sum_{i=1}^n (\mu_i^n - \mu_{i-1}^n)^T \Omega_i^{-1} (\mu_i^n - \mu_{i-1}^n) \\ & - \frac{1}{2} \sum_{i=1}^n \text{Tr}[\Omega_i^{-1} (\Gamma_i^n + \Gamma_{i-1}^n - \Gamma_{i,i-1}^n - \Gamma_{i-1,i}^n)] \\ & - \frac{1}{2} \sum_{i=1}^n (\delta y_i - D_i \mu_i^n)^T \Sigma_i^{-1} (\delta y_i - D_i \mu_i^n) \\ & - \frac{1}{2} \sum_{i=1}^n \text{Tr}[\Sigma_i^{-1} D_i \Gamma_i^n D_i^T] \\ & + \text{const.} \end{aligned} \quad (\text{F5})$$

We next take the derivative with respect to μ_0 , Γ_0 , Ω_i , and Σ_i and set to zero and solve. Some derivative identities are required. Assuming A is symmetric and positive definite, then using Einstein summation notation we have,

$$\nabla_{A_{ij}} A_{lq}^{-1} = -A_{li}^{-1} A_{jq}^{-1} \quad (\text{F.6})$$

$$\nabla_{A_{ij}} \log|A| = A_{ij}^{-1} \quad (\text{F.7})$$

$$\nabla_{A_{ij}} x_l A_{lq}^{-1} y_q = -(A_{il}^{-1} x_l)(A_{jq}^{-1} y_q) = -(A^{-1} x)(A^{-1} y)^T \quad (\text{F.8})$$

$$\nabla_{A_{ij}} \text{Tr}[A^{-1} B] = -A_{il}^{-1} B_{lq}^T A_{qj}^{-1} = -A^{-1} B^T A^{-1}. \quad (\text{F.9})$$

Applying these identities as needed we have,

$$\nabla_{\mu_0} \mathbb{E}_X[\log L(X, Y)] = \Gamma_0^{-1}(\mu_0^n - \mu_0) = 0 \quad (\text{F.10})$$

$$\implies \mu_0 = \mu_0^n \quad (\text{F.11})$$

$$\begin{aligned} \nabla_{\Gamma_0} \mathbb{E}_X[\log L(X, Y)] &= -\frac{1}{2} \nabla_{\Gamma_0} \log|\Gamma_0| - \frac{1}{2} \nabla_{\Gamma_0} (\mu_0^n - \mu_0)^T \Gamma_0^{-1} (\mu_0^n - \mu_0) \\ &\quad - \frac{1}{2} \nabla_{\Gamma_0} \text{Tr}[\Gamma_0^{-1} \Gamma_0^n] \end{aligned} \quad (\text{F.12})$$

$$= -\frac{1}{2} \Gamma_0^{-1} + \frac{1}{2} \Gamma_0^{-1} \Gamma_0^n \Gamma_0^{-1} = 0 \quad (\text{F.13})$$

$$\implies \Gamma_0 = \Gamma_0^n \quad (\text{F.14})$$

$$\begin{aligned} \nabla_{\Omega_i} \mathbb{E}_X[\log L(X, Y)] &= -\frac{1}{2} \sum_{i=1}^n \nabla_{\Omega_i} \log|\Omega_i| \\ &\quad - \frac{1}{2} \sum_{i=1}^n \nabla_{\Omega_i} (\mu_i^n - \mu_{i-1}^n)^T \Omega_i^{-1} (\mu_i^n - \mu_{i-1}^n) \\ &\quad - \frac{1}{2} \sum_{i=1}^n \nabla_{\Omega_i} \text{Tr}[\Omega_i^{-1} (\Gamma_i^n + \Gamma_{i-1}^n - \Gamma_{i,i-1}^n - \Gamma_{i-1,i}^n)] \end{aligned} \quad (\text{F.15})$$

$$\begin{aligned} &= -\frac{1}{2} \Omega_i^{-1} + \frac{1}{2} \Omega_i^{-1} (\mu_i^n - \mu_{i-1}^n) (\mu_i^n - \mu_{i-1}^n)^T \Omega_i^{-1} \\ &\quad + \frac{1}{2} \Omega_i^{-1} (\Gamma_i^n + \Gamma_{i-1}^n - \Gamma_{i,i-1}^n - \Gamma_{i-1,i}^n) \Omega_i^{-1} = 0 \end{aligned} \quad (\text{F.16})$$

$$\begin{aligned} \implies \Omega_i &= (\mu_i^n - \mu_{i-1}^n) (\mu_i^n - \mu_{i-1}^n)^T \\ &\quad + (\Gamma_i^n + \Gamma_{i-1}^n - \Gamma_{i,i-1}^n - \Gamma_{i-1,i}^n) \end{aligned} \quad (\text{F.17})$$

$$\begin{aligned} \nabla_{\Sigma_i} \mathbb{E}_X[\log L(X, Y)] &= -\frac{1}{2} \sum_{i=1}^n \nabla_{\Sigma_i} \log|\Sigma_i| \\ &\quad - \frac{1}{2} \sum_{i=1}^n \nabla_{\Sigma_i} (\delta y_i - D_i \mu_i^n)^T \Sigma_i^{-1} (\delta y_i - D_i \mu_i^n) \end{aligned}$$

$$-\frac{1}{2} \sum_{i=1}^n \nabla_{\Sigma_i} \text{Tr}[\Sigma_i^{-1} D_i \Gamma_i^n D_i^T] \quad (\text{E18})$$

$$= -\frac{1}{2} \Sigma_i^{-1} + \frac{1}{2} \Sigma_i^{-1} (\delta y_i - D_i \mu_i^n) (\delta y_i - D_i \mu_i^n)^T \Sigma_i^{-1} \\ + \frac{1}{2} \Sigma_i^{-1} D_i^T \Gamma_i^n D_i \Sigma_i^{-1} \quad (\text{E19})$$

$$= 0 \quad (\text{E20})$$

$$\Rightarrow \Sigma_i = (\delta y_i - D_i \mu_i^n) (\delta y_i - D_i \mu_i^n)^T + D_i^T \Gamma_i^n D_i \quad (\text{E21})$$

This completes the derivation of the update equations of the M-step.