



<https://openaccess.leidenuniv.nl>

License: Article 25fa pilot End User Agreement

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with explicit consent by the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed under The Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' pilot project. In this pilot research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and/or copyrights owner(s) of this work. Any use of the publication other than authorised under this licence or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the Library through email: OpenAccess@library.leidenuniv.nl

Article details

Damman W., Liu R., Kaptein A.A., Evers A.W.M., Van Middendorp H., Rosendaal F.R., Kloppenburg & M. (2018), Illness perceptions and their association with two-year functional status and change in patients with hand osteoarthritis, *Rheumatology* 57(12): 2190-2199.

Doi: 10.1093/rheumatology/key231

Original article

Illness perceptions and their association with 2 year functional status and change in patients with hand osteoarthritis

Wendy Damman¹, Rani Liu¹, Ad A. Kaptein², Andrea W. M. Evers^{3,4}, Henriët van Middendorp³, Frits R. Rosendaal⁵ and Margreet Kloppenburg^{1,5}

Abstract

Objective. To investigate the association between illness perceptions and disability both cross-sectionally and over 2 years in patients with hand OA.

Methods. Illness perceptions and self-reported disability were assessed at baseline and after 2 years in 384 patients with primary hand OA (mean age 61 years, 84% women, $n = 312$ with follow-up) with the Illness Perception Questionnaire – Revised (IPQ-R), Functional Index for Hand OA, Australian/Canadian Hand OA Index and HAQ. Risk ratios for high disability (highest quartile) at both time points were estimated for tertiles of IPQ-R dimensions, using Poisson regression. The mean IPQ dimension change difference between patients with and without disability progression (change Functional Index for Hand OA ≥ 1 , Australian/Canadian Hand OA Index > 1.4 , HAQ > 0.22) was estimated with linear regression. Analyses were adjusted for age, Doyle index and baseline score.

Results. At baseline, stronger negative illness perceptions were associated with high disability. Baseline illness perceptions were also associated with high disability after 2 years, although adjustment made apparent that these associations were confounded by baseline disability status. Most illness perceptions changed over 2 years; understanding increased, OA was regarded as more chronic and fewer emotions and consequences and less personal and treatment control were experienced. The 2 year change in disability was different between patients with and without progression for the illness perceptions of more perceived consequences, symptoms, treatment control and emotions.

Conclusion. Illness perceptions seemed to be implicated in disability and its progression. Our results suggest that interventions could focus on improving baseline disability, potentially using illness perceptions to accomplish this goal.

Key words: disability, hand osteoarthritis, illness perceptions, osteoarthritis, patient-reported outcomes, self-regulation

Rheumatology key messages

- Illness perceptions are associated with disability at baseline and can change over 2 years.
- For several illness perception subscales, change was associated with progression of disability over 2 years.
- The association between baseline illness perceptions and 2 year disability status is confounded by baseline disability.

Introduction

Hand OA is a common musculoskeletal disease leading to disability [1–3]. Disability has a heterogeneous course [4]

and is poorly associated with structural (radiographic) measures [5–7]. This might be explained by the contribution of psychosocial factors to self-reported disability. Examples of such factors are depression, illness perceptions, coping styles and anxiety [8–10].

Knowledge about these factors aids understanding why some patients report more disability than others and how their disability will develop over time, which in turn could lead to patient-tailored interventions [11]. In the present study, we will focus on one of these factors: illness perceptions.

When patients are confronted with an illness, they build a mental model to make sense of, and manage, their

¹Department of Rheumatology, ²Department of Medical Psychology, Leiden University Medical Center, ³Health, Medical and Neuropsychology Unit, Institute of Psychology, Leiden University, ⁴Department of Psychiatry and ⁵Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

Submitted 20 November 2017; revised version accepted 25 June 2018

Correspondence to: Wendy Damman, Department of Rheumatology, Leiden University Medical Center, C1-R, PO Box 9600, 2300 RC Leiden, The Netherlands.
E-mail: w.damman@lumc.nl

health problem [12]. The Common Sense Model describes this mental model by suggesting how cognitive and emotional representations and beliefs, so-called illness perceptions, influence a patient's coping, health behaviour and health outcomes (e.g. disability) [13]. In other words, illness perceptions are the thoughts and feelings of a patient about his/her illness.

Illness perceptions have been associated with disability in cross-sectional studies in patients with generalized OA, lower extremity OA and hand OA [8,10,14–16]. In short-term follow-up studies, more negative illness perceptions (e.g. more perceived consequences or more emotional representations) were associated with unfavourable clinical outcomes in patients with knee or hip OA [17,18] and in patients with other musculoskeletal conditions [19–21]. A long-term observational study in patients with generalized OA showed that increasing negative illness perceptions over time were accompanied by progression of disability [10]. Trials intervening on negative illness perceptions in patients with diabetes, heart disease and back pain (i.e. chronic conditions) showed that perceptions can change to more positive and that this has positive effects on health outcomes [22–24]. All this also suggests that in patients with hand OA, illness perceptions could be of importance as potential modifiable factors that could serve as a treatment target. However, longitudinal studies on illness perceptions in relation to (change in) functional status in patients with hand OA are unavailable [25], leaving it unclear whether illness perceptions are relevant as targets. To be a relevant target, change over time should be possible and this change should be relevant (i.e. associated with change in outcomes).


Therefore, in the present study, we investigated whether illness perceptions changed after 2 years and whether this change was associated with a change in functional status (progression of disability) in patients with hand OA. To further evaluate the relevance of illness perceptions we studied the association of baseline illness perceptions with disability status both at baseline and after 2 years. This is important knowledge in the light of informing patients about disease prognosis, but also to identify patients that are most at risk for worse outcomes. The latter is the patient group that could benefit most from treatment. We hypothesize, in a secondary care cohort of patients with hand OA, that negative illness perceptions are associated with poor clinical outcome and that a change in illness perceptions is associated with a change in disability.

Methods

Study design

The present study is part of the Hand OSTeoArthritis in Secondary care (HOSTAS) study, an ongoing observational cohort study in hand OA [26]. For this report, patients included from January 2011 onwards and who completed relevant questionnaires were considered.

Fig. 1 Three sections of the Revised Illness Perception Questionnaire



Tables 2 and 5: $n = 384$, Tables 3 and 4: $n = 312$.

Patients

Consecutive patients with primary hand OA from the outpatient clinic of the Leiden University Medical Center (LUMC) were included between January 2011 and October 2015. Primary hand OA was defined according to the diagnosis of the treating rheumatologist. Patients with secondary hand OA (e.g. due to trauma) and patients with hand symptoms explained by another diagnosis were excluded [26]. Patients were followed annually with postal questionnaires and biennially with an additional research visit. Written informed consent was obtained from all participants. The study was approved by the LUMC Medical Ethical Committee.

Illness perceptions

Illness perceptions were studied using the Illness Perception Questionnaire – Revised (IPQ-R) [12,27], which was assessed biennially, together with a research visit. The IPQ-R measures both cognitive and emotional representations of illness in three sections with nine subscales (Fig. 1). The questionnaire has been shown to be valid and consistent in a population with musculoskeletal hand problems [8].

The first section is the identity subscale, concerning symptoms that patients attribute to OA. For 14 commonly occurring symptoms, patients indicated whether they think these symptoms have to do with their OA always, often, sometimes or never. All always, often or sometimes indicated symptoms were summed (range 0–14).

The second section consists of 38 questions (0–4, Likert scale) spread over seven subscales. The consequence subscale ($n = 6$ questions) is about the impact of OA on daily life. The timeline acute/chronic ($n = 6$) represents beliefs about the perceived chronicity of the disease, whereas the timeline cyclical ($n = 4$) is about the variability in course of symptoms and the disease process. Illness coherence ($n = 5$) represents the patient's understanding of OA. The emotional representations subscale ($n = 6$) reflects negative emotions due to OA. The personal ($n = 6$)

and treatment ($n=5$) control subscales represent beliefs about possibilities for influencing the symptoms and disease course on a personal level and with treatment. Items were summed per subscale, accepting a maximum of one missing item.

For both the first and second section, a higher score means stronger illness perceptions in that particular subscale. For personal control, treatment control and illness coherence, a higher score/stronger perception is considered more positive, whereas for identity and for the other representations subscales, a higher score/stronger perception is considered more negative.

The third section comprises the causes subscale, with 18 possible causes of OA subdivided into four dimensions: psychological attributions ($n=6$), immunity ($n=3$), risk factors ($n=7$) and chance ($n=2$). Items were rated on a 5-point Likert scale: totally disagree (1) to totally agree (5). Answers were dichotomized to disagree/no opinion and agree. To provide insight in which causes are associated with disability, we analysed per cause.

Self-reported disability

Self-reported disability was assessed annually with the Functional Index for Hand OA (FIHOA), with the Australian/Canadian Hand OA Index (AUSCAN) and with the HAQ [28–30]. The FIHOA is a hand-specific questionnaire concerning physical functioning. The 10 questions (each scored from 0 to 3) were summed (total score 0–30). A maximum of two missing items was accepted. The AUSCAN is also a hand-specific questionnaire, of which we used the nine questions concerning hand function (0–4, Likert scale), summed to a total score (range 0–36). A maximum of two missing items was accepted. The HAQ measures overall disability and consists of 24 questions (each scored 0–3) in eight categories. The highest scores per category were summed and divided by eight, resulting in a total score ranging from 0 to 3. A maximum of two missing categories was accepted. For all questionnaires, a higher score means worse function, hence more disability.

Clinical assessment

During physical examination at baseline, performed by trained research nurses, all DIP, PIP, IP, MCP and first CMC joints were assessed for the number of joints with bony swelling (total range 0–30), with limited range of motion and with deformity (both ranges 0–22; MCP 2–5 excluded). Furthermore, tenderness on palpation (range 0–3 per joint) was assessed in 24 joint units: all DIP, PIP, first IP, first MCP and first CMC joints individually and second through fifth MCP joints as one joint group. In each patient, the Doyle index for the hands was the summed score of the 24 joint units (range 0–72) [31]. Joints outside the hands were also assessed for tenderness upon palpation or movement (range 0–3), as specified in the Doyle index [31]. Scores of 48 units were summed into a total score ranging from 0 to 144. Patient-reported hand symptoms and physical examination were used to determine fulfilment of the American

College of Rheumatology criteria for hand OA [32]. In addition, a research nurse recorded the number of comorbid diseases (range 0–17) [26].

Radiographic assessment

Joints of both hands ($n=30$) were scored on a scale of 0–4 on conventional dorsal–volar radiographs, according to Kellgren–Lawrence (KL) system [33]. Scoring was blinded for demographic and clinical data (WD). Intra-observer reliability, based on randomly selected radiographs (10%), was good (intraclass correlation coefficient >0.9). Scores were summed per patient into a total KL score for the hands (range 0–120).

Statistical analysis

When questionnaires had missing items (not exceeding the maximum number of allowed missing items), values were replaced with the (sub)scale mean value. Questionnaires with too many missing items were regarded as missing. The change in score after 2 years was calculated as the follow-up score minus the baseline score. Change equal to or above the minimal clinically important difference (MCID) was used for HAQ (0.22) to classify patients as progressed; changes below this value were regarded as not progressed [34]. For AUSCAN function, the retrograde of the minimal clinically important improvement (MCII) was used, i.e. 1.4 [35]. Since, for FIHOA, no MCII or MCID is known, progression was defined as the minimal change potentially detectable, which is 1 unit (or 3.3%) [36].

Associations between illness perceptions at baseline (determinant) and disability at baseline and at follow-up (outcome), presented as risk ratios (RRs) with 95% CIs, were studied using a Poisson regression model with log link function and robust standard errors [37]. For these analyses, scores for each subscale of illness perceptions were categorized into tertiles to show a trend between groups, while providing the best power and best possible balance of number of patients per group. Disability scores were categorized into quartiles in order to provide contrast (high vs low). A score in the highest quartile was considered high disability and a score in the other quartiles was considered low disability. Hence RRs represent the incremental risk of high disability per tertile of illness perception score, with the lowest tertile as a reference. In the causes section, analyses were not performed per tertile, but for each individual cause. Mean differences of change in illness perceptions after 2 years between groups with and without progression of disability were estimated using linear regression.

Analyses were adjusted for age and Doyle index. For FIHOA and AUSCAN analyses, the hand Doyle index was used, whereas for the HAQ analysis the total Doyle index was used. Additional adjustments were made for baseline score of the outcome (i.e. baseline FIHOA, AUSCAN or HAQ) in longitudinal analyses with the Poisson model (baseline illness perceptions and 2 year disability status) and for both baseline score of the outcome and baseline illness perception score in the linear

TABLE 1 Baseline characteristics of 384 primary hand OA patients

| Variable | Value |
|---|--------------|
| Age, years, mean (s.d.) | 60.9 (8.4) |
| Sex, female, <i>n</i> (%) | 322 (84) |
| BMI, mean (s.d.) ^a | 27.6 (4.9) |
| Fulfilling ACR criteria for hand OA, <i>n</i> (%) | 346 (90) |
| Number of comorbid diseases (0–17), median (range) ^a | 0 (0–5) |
| Patient-reported disability, median (range) | |
| FIHOA (0–30) | 9.0 (0–26.7) |
| AUSCAN (0–36 ^a) | 16.0 (0–36) |
| HAQ (0–3) | 0.9 (0–2.3) |
| Physical examination of the hands, median (range) | |
| Bony swelling joint count (0–30) | 11 (0–24) |
| Deformity joint count (0–22) | 3 (0–16) |
| Doyle index of the hands (0–72) | 4 (0–70) |
| Joints with limited ROM count (0–22) | 4 (0–22) |
| Physical examination overall, median (range) | |
| Doyle index (including the hands) (0–144) | 7 (0–88) |
| Radiographic scoring, hands, median (range) ^a | |
| KL summed score (0–120) | 16 (0–89) |

^aNumber of patients represented in data if not 384: BMI 378, comorbidities 374, AUSCAN 382, KL score 381. ROM: range of motion.

regression model (change in illness perceptions and progression of disability over 2 years). In addition, for sensitivity analysis, adjustments for sex, BMI, total hand KL score and number of comorbidities were made.

All cross-sectional analyses were done on cases with complete baseline data for IPQ-R, whereas all longitudinal analyses were performed in patients with available follow-up data. SPSS software for Windows, versions 20.0 and 23.0 (IBM, Armonk, NY, USA), was used.

Results

Study population

Of 388 eligible patients, 384 (99%) had baseline IPQ-R questionnaires available (Table 1). Both baseline and 2 year follow-up was completed by 312, 311, 314 and 311 patients for IPQ-R, FIHOA, AUSCAN and HAQ, respectively. Reasons for no available follow-up were too many missing items to calculate total scores, dropout or skipped 2 year visit. Patients with and without follow-up did not differ in age, sex, BMI or baseline function scores (data not shown).

Associations of illness perceptions and disability at baseline

Table 2 shows cross-sectional associations between illness perceptions and disability at baseline, both for hand-

specific (FIHOA, AUSCAN) and overall (HAQ) functional status. The illness perception subscales of identity, consequences and emotional representations were associated with high disability on all three outcomes and in a dose-response way. This means that stronger negative perceptions (higher tertiles) had a higher risk for high disability compared with the lowest tertiles. In other words, patients who experienced more symptoms (identity), consequences or emotions had an increased risk for high disability at baseline. In contrast, a stronger positive perception, i.e. more understanding of the illness (illness coherence), was associated with a lower risk for high disability on the FIHOA and HAQ. Stronger beliefs about a cyclical disease course were associated with a lower risk of disability on the FIHOA only. Additional adjustment for sex, BMI, radiographic damage (hand KL score) and number of comorbidities did not essentially change the estimates.

Change in disability and illness perceptions over 2 years

After 2 years, 50% (157/311) of patients worsened in the FIHOA score, with a mean score increase of 3.7 (s.d. 2.6), while 37% (117/314) and 36% (112/311) of patients worsened more than the MCII/MCID in AUSCAN function and HAQ scores, with a mean increase of 5.6 (s.d. 3.6) and 0.4 (s.d. 0.2). Illness perceptions also changed in the time frame of 2 years (Table 3). The illness perceptions that changed in the whole group were timeline acute/chronic, consequences, personal control, treatment control, illness coherence and emotional representations. This means that patients were, over time, understanding more of their illness, regarding their OA as more chronic (timeline), but experiencing less personal and treatment control. Furthermore, they perceived fewer emotions and fewer consequences of their OA.

Association of baseline illness perceptions with high disability after 2 years

We explored the association of illness perceptions at baseline and high disability (a score in the highest quartile) after 2 years (Table 4). We found longitudinal associations for several illness perceptions: the more baseline symptoms a person attributed to their OA (identity), the stronger the perceived consequences and the more emotions, the higher the risk of disability at follow-up for all outcomes. Similarly, more baseline illness coherence was associated with less disability at follow-up on the FIHOA and HAQ. Baseline perceived chronicity (timeline acute/chronic), personal control, treatment control and beliefs about a cyclical timeline did not show an association with disability after 2 years (Table 4). Additional analyses showed that baseline disability scores were associated with disability scores after 2 years. Adjustment for baseline disability scores resulted in the disappearance or in a large decrease (identity with FIHOA) of the associations between baseline illness perceptions and 2 year disability status (Table 4).

TABLE 2 Associations between baseline illness perceptions and baseline self-reported disability

| Perceptions | Mean (s.d.) | FIHOA, RR (95% CI) ^a | AUSCAN function, RR (95% CI) ^b | HAQ, RR (95% CI) ^c |
|----------------------------------|-------------|---------------------------------|---|-------------------------------|
| Identity (0–14) | 4.9 (2.2) | | | |
| 0–3 | | 1 | 1 | 1 |
| 4–5 | | 1.9 (1.04, 3.4) | 2.2 (1.1, 4.1) | 1.7 (0.9, 3.1) |
| 6–13 | | 3.0 (1.6, 5.3) | 3.3 (1.7, 6.2) | 3.0 (1.7, 5.6) |
| Timeline acute/chronic (6–30) | 26.2 (3.6) | | | |
| 12–24 | | 1 | 1 | 1 |
| 25–28 | | 1.4 (0.9, 2.0) | 1.1 (0.7, 1.7) | 1.0 (0.7, 1.5) |
| 29–30 | | 1.1 (0.7, 1.6) | 1.0 (0.7, 1.6) | 1.0 (0.6, 1.4) |
| Consequences (6–30) | 16.5 (4.3) | | | |
| 6–14 | | 1 | 1 | 1 |
| 15–18 | | 2.1 (1.2, 3.5) | 1.4 (0.8, 2.3) | 1.2 (0.7, 2.1) |
| 19–30 | | 3.2 (2.0, 5.2) | 2.7 (1.7, 4.2) | 3.6 (2.3, 5.6) |
| Personal control (6–30) | 18.6 (3.6) | | | |
| 6–17 | | 1 | 1 | 1 |
| 18–20 | | 1.1 (0.7, 1.6) | 1.1 (0.7, 1.6) | 1.1 (0.7, 1.6) |
| 20.4–29 | | 1.0 (0.7, 1.6) | 1.3 (0.9, 2.0) | 1.3 (0.8, 2.0) |
| Treatment control (5–25) | 13.9 (2.7) | | | |
| 5–12.5 | | 1 | 1 | 1 |
| 13–15 | | 0.7 (0.5, 1.03) | 0.8 (0.5, 1.1) | 0.9 (0.6, 1.3) |
| 16–20 | | 0.8 (0.5, 1.2) | 1.1 (0.7, 1.6) | 0.8 (0.5, 1.3) |
| Illness coherence (5–25) | 18.6 (3.8) | | | |
| 6–17 | | 1 | 1 | 1 |
| 17.5–20 | | 0.5 (0.4, 0.8) | 0.9 (0.6, 1.3) | 0.8 (0.5, 1.1) |
| 21–25 | | 0.6 (0.4, 0.9) | 0.6 (0.4, 1.01) | 0.5 (0.3, 0.8) |
| Timeline cyclical (4–20) | 13.2 (3.2) | | | |
| 4–11 | | 1 | 1 | 1 |
| 12–14 | | 0.7 (0.5, 0.997) | 0.8 (0.5, 1.2) | 0.8 (0.5, 1.2) |
| 15–20 | | 0.7 (0.5, 1.01) | 0.8 (0.5, 1.2) | 0.9 (0.6, 1.1) |
| Emotional representations (6–30) | 14.4 (4.9) | | | |
| 6–12 | | 1 | 1 | 1 |
| 13–15 | | 1.5 (0.9, 2.5) | 1.5 (0.9, 2.4) | 1.2 (0.7, 1.9) |
| 16–30 | | 2.3 (1.5, 3.5) | 1.9 (1.3, 2.9) | 1.9 (1.3, 2.9) |

Results are RRs (95% CIs) for having a disability score in the highest quartile vs a score in the other quartiles per tertile of illness perception scores; 1=reference. Adjusted for age and Doyle index. ^aQuartile 4: score ≥ 13 . ^bQuartile 4: score ≥ 22 . ^cQuartile 4: score ≥ 1.25 .

Change in illness perceptions between patients with and without progression of disability

In Table 3 the mean change in illness perceptions after 2 years is shown for two groups: with and without progression of disability. The adjusted mean difference in the change of illness perceptions between these groups is also presented. The illness perceptions timeline acute/chronic, personal control and illness coherence changed at the group level, but this change did not differ between patients with and without progression. However, in other illness perceptions there was a difference between the groups. For the consequences subscale, patients with progression in disability on all outcomes had increased in perceived consequences after 2 years, where patients who did not progress decreased. Similar results were seen for perceptions about identity (FIHOA and HAQ), treatment control (FIHOA and AUSCAN) and emotions (FIHOA and AUSCAN), i.e. patients with disability progression on the FIHOA, AUSCAN and/or HAQ were experiencing more symptoms, less treatment control or fewer

decreased emotional representations after 2 years than patients without progression.

Causes of OA and high disability

Patients indicated for 18 possible causes whether they thought (agreed) these factors could have caused their OA (Table 5). The most indicated causes were heredity (66%), ageing (74%) and chance or bad luck (55%). Causes that were associated with high disability at baseline were mostly psychological causes: stress or worry, family problems or worries, overwork, own personality and poor medical care in the past. These causes were only indicated as a cause by <15% of the patients, except for overwork (33%).

Discussion

In our large secondary care cohort with 2 year follow-up of patients with primary hand OA, we found that the perceptions that patients have of their illness were associated

TABLE 3 Mean change in illness perceptions after 2 years for the whole group and between groups with and without progression in disability

| Perception | Whole group | | FIHOA ^a | | AUSCAN ^a function | | HAQ ^a | |
|----------------------------------|-----------------------|-----------------------|------------------------------------|---|------------------------------------|---|------------------------------------|---|
| | Baseline, mean (s.d.) | Change, mean (95% CI) | Progression (yes/no) (n = 155/154) | Change difference, mean (95% CI) ^b | Progression (yes/no) (n = 115/195) | Change difference, mean (95% CI) ^b | Progression (yes/no) (n = 112/198) | Change difference, mean (95% CI) ^b |
| Identity (0-14) | 4.8 (2.3) | 0.0 (-0.2, 0.3) | 0.3/-0.3 | 0.6 (0.2, 1.0) | 0.2/-0.1 | 0.3 (-0.2, 0.7) | 0.4/-0.2 | 0.6 (0.2, 1.1) |
| Timeline acute/chronic (6-30) | 26.3 (3.6) | 0.6 (0.2, 1.0) | 0.7/0.6 | 0.6 (-0.1, 1.3) | 0.9/0.5 | 0.7 (-0.1, 1.4) | 0.4/0.7 | 0.2 (-0.6, 0.9) |
| Consequences (6-30) | 16.5 (4.3) | -0.5 (-1.0, -0.1) | 0.4/-1.4 | 2.1 (1.3, 2.9) | 0.3/-1.0 | 1.6 (0.7, 2.5) | 0.4/-1.1 | 1.7 (0.9, 2.5) |
| Personal control (6-30) | 18.4 (3.7) | -0.5 (-0.9, -0.1) | -0.5/-0.5 | 0.1 (-0.7, 0.8) | -1.0/-0.2 | -0.6 (-1.4, 0.2) | -0.7/-0.4 | -0.1 (-0.9, 0.7) |
| Treatment control (5-25) | 13.8 (2.7) | -0.8 (-1.2, -0.5) | -1.1/-0.5 | -0.8 (-1.4, -0.2) | -1.5/-0.4 | -1.0 (-1.6, -0.3) | -0.8/-0.8 | -0.1 (-0.7, 0.6) |
| Illness coherence (5-25) | 18.6 (3.8) | 0.6 (0.2, 1.0) | 0.4/0.8 | -0.2 (-0.9, 0.5) | 0.3/0.7 | -0.3 (-1.1, 0.4) | 0.4/0.7 | -0.3 (-1.0, 0.5) |
| Timeline cyclical (4-20) | 13.3 (3.2) | -0.0 (-0.4, 0.4) | -0.2/0.2 | -0.3 (-0.9, 0.4) | -0.1/0.1 | -0.3 (-1.0, 0.4) | 0.0/0.0 | 0.2 (-0.5, 0.9) |
| Emotional representations (6-30) | 14.3 (4.8) | -0.8 (-1.3, -0.4) | -0.6/-1.1 | 0.9 (0.02, 1.7) | -0.3/-1.2 | 1.1 (0.2, 2.0) | -0.5/-1.1 | 0.8 (-0.1, 1.7) |

^aFor the FIHOA, AUSCAN and HAQ there were three, two and two patients, respectively, for whom a delta could not be calculated due to missing data. ^bAdjusted for age, Doyle index, baseline score of outcome (i.e. FIHOA, AUSCAN, HAQ) and baseline score of illness perception.

with self-reported disability due to hand OA at the same moment. The baseline perceptions patients had about their illness were also associated with high disability after 2 years. However, these associations were confounded by the baseline disability score. The perceptions patients have about their illness showed small changes over the 2 year time frame. Progression in self-reported disability was associated with a change in perceived consequences, symptoms a person attributed to their OA, treatment control and emotions.

In line with our results, a cross-sectional population-based study also found associations between symptoms and consequences and hand/finger function. However, that study did not find associations between other illness perceptions and disability [8]. Differences between studies could be explained by differences in the study population and in the methods of assessing disability. Unfortunately, that study had no longitudinal data available. Compared with a 6 year longitudinal study in patients with OA in multiple sites, we found similar results in illness perception subscales that changed in the follow-up period and in subscales where changes were associated with progression in disability [10]. These studies, as well as a study comparing patients with diabetes and OA, support our findings that identity and consequences are important subscales in patients with OA [8,10,38].

Similar to the 6 year study [10], in our study, the magnitude of changes and strength of found associations was limited. As there is no known cut-off for clinically relevant changes in illness perceptions, we do not know whether small changes are relevant. Therefore we related them to progression in disability to provide clinical meaning. We showed that after 2 years, which could be the term of a clinical trial, change is possible and that this change is relevant (i.e. associated with change in outcomes). As illness perceptions are possible modifiable factors, this suggests they could serve as a treatment target.

After 2 years, patients perceived their disease as more chronic (higher score for timeline) and experienced less treatment and personal control. The changes in these illness perceptions are considered more negative. However, within the context of OA, which is a progressive disease with very limited treatment modalities and no available disease-modifying treatment, a perception of less treatment and personal control could also be regarded as realistic and could reflect increasing insight into the nature of the disease. The latter is in line with increasing illness coherence (less negative over 2 years). Therefore it is questionable whether, in the context of OA, perceiving less treatment and personal control and a chronic timeline is more negative. This is also reflected in perceiving fewer emotions and consequences, both reflecting that patients perceive their disease as less negative over 2 years.

We studied the association of underlying beliefs of patients about the causes of their OA with disability. The most often mentioned causes (ageing, heredity and chance/bad luck) are widely recognized as causes for

TABLE 4 Associations between baseline illness perceptions and self-reported function scores at 2 years follow-up

| Perceptions | FIHOA, RR (95% CI) ^a | | AUSCAN function, RR (95% CI) ^b | | HAQ, RR (95% CI) ^c | |
|----------------------------------|---------------------------------|-----------------------|---|-----------------------|-------------------------------|-----------------------|
| | Adjusted ^d | Adjusted ^e | Adjusted ^d | Adjusted ^e | Adjusted ^d | Adjusted ^e |
| Identity (0–14) | | | | | | |
| 0–3 | 1 | 1 | 1 | 1 | 1 | 1 |
| 4–5 | 3.1 (1.4, 7.1) | 2.3 (1.0, 5.1) | 2.5 (1.3, 4.8) | 1.4 (0.7, 2.6) | 2.9 (1.3, 6.7) | 2.1 (0.9, 4.5) |
| 6–13 | 4.7 (2.1, 10.7) | 2.5 (1.1, 5.7) | 2.8 (1.4, 5.6) | 1.0 (0.5, 2.0) | 5.4 (2.3, 12.3) | 2.2 (0.9, 5.1) |
| Timeline acute/chronic (6–30) | | | | | | |
| 16–24 | 1 | 1 | 1 | 1 | 1 | 1 |
| 25–28 | 1.0 (0.6, 1.6) | 0.8 (0.5, 1.3) | 1.2 (0.7, 2.0) | 1.1 (0.7, 1.7) | 1.1 (0.7, 1.9) | 1.1 (0.7, 1.7) |
| 29–30 | 1.0 (0.6, 1.6) | 1.0 (0.7, 1.5) | 1.1 (0.7, 1.8) | 0.9 (0.7, 1.4) | 1.0 (0.6, 1.6) | 1.2 (0.8, 1.7) |
| Consequences (6–30) | | | | | | |
| 6–14 | 1 | 1 | 1 | 1 | 1 | 1 |
| 15–18 | 1.6 (0.9, 3.0) | 1.1 (0.6, 1.9) | 1.0 (0.6, 1.8) | 0.9 (0.5, 1.4) | 1.0 (0.6, 1.7) | 1.0 (0.6, 1.6) |
| 19–30 | 3.0 (1.7, 5.0) | 1.5 (0.8, 2.6) | 2.0 (1.3, 3.3) | 1.0 (0.7, 1.6) | 2.1 (1.3, 3.4) | 0.9 (0.6, 1.4) |
| Personal control (6–30) | | | | | | |
| 6–7 | 1 | 1 | 1 | 1 | 1 | 1 |
| 18–20 | 1.1 (0.7, 1.8) | 1.0 (0.7, 1.6) | 0.8 (0.6, 1.3) | 1.0 (0.7, 1.4) | 1.2 (0.8, 1.9) | 1.2 (0.8, 1.7) |
| 20.4–28 | 1.3 (0.8, 2.2) | 1.4 (0.9, 2.1) | 1.0 (0.6, 1.6) | 1.1 (0.8, 1.7) | 1.3 (0.8, 2.1) | 1.1 (0.8, 1.6) |
| Treatment control (5–25) | | | | | | |
| 5–12.5 | 1 | 1 | 1 | 1 | 1 | 1 |
| 13–14 | 0.7 (0.4, 1.3) | 0.8 (0.5, 1.4) | 0.7 (0.4, 1.2) | 0.9 (0.6, 1.5) | 1.1 (0.6, 1.8) | 1.3 (0.8, 2.0) |
| 15–20 | 1.0 (0.7, 1.6) | 1.2 (0.8, 1.7) | 1.1 (0.7, 1.6) | 1.1 (0.8, 1.6) | 1.2 (0.7, 1.8) | 1.3 (0.9, 1.8) |
| Illness coherence (5–25) | | | | | | |
| 6–17 | 1 | 1 | 1 | 1 | 1 | 1 |
| 17.5–20 | 0.6 (0.3, 0.9) | 0.7 (0.4, 1.0) | 0.8 (0.5, 1.3) | 1.0 (0.7, 1.4) | 0.6 (0.4, 0.9) | 0.8 (0.6, 1.2) |
| 21–25 | 0.5 (0.3, 0.8) | 0.5 (0.3, 0.8) | 0.6 (0.4, 1.0) | 0.8 (0.5, 1.3) | 0.4 (0.3, 0.7) | 0.8 (0.5, 1.2) |
| Timeline cyclical (4–20) | | | | | | |
| 5–11 | 1 | 1 | 1 | 1 | 1 | 1 |
| 12–14 | 1.4 (0.8, 2.3) | 1.7 (1.1, 2.6) | 1.1 (0.7, 1.6) | 1.4 (1.0, 2.0) | 1.1 (0.7, 1.8) | 1.1 (0.7, 1.6) |
| 15–20 | 1.3 (0.8, 2.2) | 1.5 (1.0, 2.4) | 0.8 (0.5, 1.3) | 1.1 (0.7, 1.6) | 1.3 (0.8, 2.0) | 1.2 (0.8, 1.8) |
| Emotional representations (6–30) | | | | | | |
| 6–12 | 1 | 1 | 1 | 1 | 1 | 1 |
| 13–15 | 1.3 (0.8, 2.3) | 1.0 (0.6, 1.8) | 1.0 (0.6, 1.8) | 0.8 (0.5, 1.3) | 1.1 (0.6, 1.9) | 0.9 (0.5, 1.4) |
| 16–30 | 2.1 (1.3, 3.3) | 1.5 (0.9, 2.3) | 1.8 (1.1, 2.8) | 1.2 (0.8, 1.9) | 2.2 (1.4, 3.4) | 1.2 (0.8, 1.9) |

Results are RRs (95% CIs) for having a disability score at follow-up in the highest quartile vs a score in the other quartiles per tertile of baseline illness perception scores (1 = reference). ^aQuartile 4: score ≥ 15 . ^bQuartile 4: score ≥ 22 . ^cQuartile 4: score ≥ 1.38 . ^dAdjusted for age and Doyle index. ^eAdjusted for age, Doyle index and baseline score of the outcome (i.e. FIHOA, AUSCAN, HAQ).

OA and were not associated with disability [1]. Several other causes that are not generally linked to OA pathophysiology were associated with disability; these were mostly psychological (e.g. stress or worries). This illustrates that perceptions about causes of OA, which are likely incorrect from a pathophysiological point of view, are related to higher disease burden. Changing these perceptions, for example by education, could be a treatment target. It should be noted, however, that most causes that were associated with disability in our cohort were only mentioned by a minority of patients. Nevertheless, this is a proportion that is similar to a study where $\sim 10\%$ of the patients blamed themselves for their OA [38]. Other studies did not find associations for psychological attributions as a cause dimension with disability [8,10,19]. This could be explained by not investigating separate causes, but instead by aggregating all psychological causes into one item.

Although we showed that baseline illness perceptions showed strong associations with baseline disability, their association with 2 year disability status virtually disappeared when the baseline disability score was taken into account. This could be explained by confounding by baseline disability. Hence, based solely on baseline illness perceptions, it is not possible to give a prognosis about 2 years disability status. In contrast, the 6 year follow-up study in generalized OA found a predictive ability for baseline illness perceptions and disability status at follow-up. This difference could be due to differences in follow-up time (6 vs 2 years), in location of OA (multiple sites vs hand) and in severity of OA (we found a higher baseline HAQ and AUSCAN function compared with their population) [7,10]. In our study, baseline illness perceptions and baseline disability were strongly related, which suggests that by improving baseline illness perceptions, disability could also improve. Studies in patients with hand OA had

TABLE 5 Associations between perceptions about individual causes of OA and disability at baseline

| Perception | Disagree or no opinion/ agree (% agree) | FIHOA, RR (95% CI) ^a | AUSCAN, RR (95% CI) ^b | HAQ, RR (95% CI) ^c |
|---|--|------------------------------------|-------------------------------------|----------------------------------|
| Psychological | | | | |
| Stress or worry | 334/48 (13) | 1.5 (1.0, 2.3) | 1.7 (1.1, 2.5) | 1.7 (1.1, 2.5) |
| Own mental attitude (e.g. thinking about life negatively) | 376/6 (2) | 1.9 (0.8, 4.3) | 2.0 (0.9, 4.5) | 2.0 (0.9, 4.5) |
| Family problems or worries | 352/30 (8) | 1.1 (0.6, 2.0) | 1.7 (1.1, 2.7) | 1.5 (0.9, 2.5) |
| Overwork | 259/125 (33) | 1.5 (1.04, 2.0) | 1.4 (0.97, 1.9) | 1.6 (1.1, 2.2) |
| Own emotional state (e.g. feeling down) | 361/22 (6) | 1.4 (0.8, 2.5) | 1.5 (0.8, 2.7) | 1.7 (0.97, 2.8) |
| Own personality | 366/17 (4) | 1.4 (0.7, 2.6) | 2.4 (1.6, 3.7) | 2.7 (1.8, 4.0) |
| Risk factor | | | | |
| Hereditary | 130/253 (66) | 1.2 (0.8, 1.7) | 1.3 (0.9, 1.9) | 0.9 (0.7, 1.3) |
| Diet or eating habits | 349/31 (8) | 1.5 (0.9, 2.4) | 1.3 (0.7, 2.2) | 1.4 (0.9, 2.4) |
| Poor medical care in the past | 363/17 (5) | 2.1 (1.3, 3.4) | 1.7 (0.9, 3.0) | 1.9 (1.1, 3.3) |
| Own behaviour | 327/53 (14) | 1.4 (0.95, 2.1) | 1.1 (0.6, 1.7) | 1.2 (0.8, 1.9) |
| Ageing | 100/283 (74) | 0.8 (0.5, 1.1) | 1.1 (0.7, 1.6) | 0.9 (0.6, 1.3) |
| Alcohol | 379/5 (1) | 1.5 (0.5, 4.5) | Not possible | 0.8 (0.1, 4.5) |
| Smoking | 374/9 (2) | 0.8 (0.2, 2.9) | 0.9 (0.2, 2.9) | 1.8 (0.8, 3.7) |
| Immunity | | | | |
| Germ or virus | 375/8 (2) | Not possible | 0.5 (0.1, 3.0) | Not possible |
| Environmental pollution | 372/9 (2) | 1.3 (0.5, 3.2) | 1.3 (0.5, 3.3) | 1.2 (0.5, 3.3) |
| Altered immunity | 334/49 (13) | 1.1 (0.7, 1.8) | 1.3 (0.9, 2.1) | 1.3 (0.9, 2.1) |
| Chance | | | | |
| Chance or bad luck | 172/209 (55) | 0.7 (0.6, 1.1) | 0.7 (0.5, 1.0) | 0.9 (0.7, 1.3) |
| Accident or injury | 358/25 (7) | 1.2 (0.7, 2.2) | 0.9 (0.4, 1.9) | 0.9 (0.5, 1.9) |

Results are RRs (95% CIs) for having a disability score at baseline in the highest quartile vs a score in the other quartiles per individual cause of OA (dichotomized to disagree or no opinion vs agree) (1 = reference). Unadjusted results. ^aQuartile 4: score ≥ 13 . ^bQuartile 4: score ≥ 22 . ^cQuartile4: score ≥ 1.25 .

not yet been performed, but a case report in a knee OA patient supports this hypothesis [39].

A strength of our study is that we used different validated self-reported questionnaires to evaluate disability. However, the measured constructs of disability are different; HAQ measures overall disability, including hand disability, while AUSCAN and FIHOA are hand specific. Several studies showed that FIHOA and AUSCAN are strongly correlated, but not 100% [40]. This means that FIHOA and AUSCAN may measure somewhat different aspects of the construct of hand disability. In our current study, AUSCAN performed worse than FIHOA in the association with illness perceptions, supporting FIHOA as the preferred measure for hand disability [41]. However, in general, it is still to be determined, and beyond the scope of this article, which outcome measure is recommended to study hand disability and whether there are other suitable outcome measures besides FIHOA and AUSCAN [40].

Studying three outcomes enabled us to identify perceptions that are most relevant. The most relevant perception seemed to be perceived consequences, as this perception was associated with three outcomes in almost all analyses. Nevertheless, there were quite a few differences between the sections of the IPQ-R and the subscales within a section in relation to functional status. Therefore illness perceptions, as in this questionnaire, remain rather heterogeneous and can reflect several items within the

topic. It should be noted that studying several determinants and outcomes raises the issue of multiple testing. As our study is observational/empirical, we chose not to adjust for multiple testing [42]. Consequently, it is possible that our results are due to chance. However, that other studies are in line with our study supports the validity of our results [8,10,19].

There are several limitations we need to address. The first is that both determinant and outcome in our study were self-reported, and inherently subjective. It could be that patients with more negative perceptions tend to report more disability, while the same patients would not score as disabled on more objective performance tests, such as hand mobility or grip strength. However, we deliberately have not taken such tests into account as outcomes for several reasons. It is possible that performance tests too are influenced by negative illness perceptions and therefore their objectivity could be questioned [43]. Furthermore, and in our opinion more important, disability is a patient-reported outcome, thus reflecting the patient's perspective. It is the subjective (experienced) disability that medical care should focus on, regardless of the objective performance. Therefore it is important to know which factors are associated with subjective disability in order to design patient-tailored treatment strategies. As a second limitation, the choice for our method of analysis, i.e. working with tertiles and quartiles, means that data are lost when categorizing.

However, categorizing provides more contrast and makes results easier to interpret. Furthermore, we chose this method to enable comparison with earlier studies [10]. Finally, patients included in our cohort all sought medical care in a secondary care centre, whereas we can assume that many patients with hand OA stay in primary care or do not consult a doctor [8]. This could have biased our results. Probably, secondary care patients are a selection of patients with hand OA with more negative illness perceptions and more disability than patients in primary care. When selecting on already negative perceptions and high disability, change could be a regression-to-the-mean effect. Therefore we adjusted our analysis of change for the baseline score of illness perceptions and baseline disability. Selection of patients with negative illness perceptions and high disability could also lead to too little variability. This could explain why we found that illness perceptions were not associated with 2 year disability after adjustment for baseline disability.

In conclusion, we found that illness perceptions show an association with disability at baseline and can change over 2 years. For several illness perception subscales, change was associated with progression of disability over 2 years, implying that these could be relevant treatment targets. However, the association between baseline illness perceptions and 2 year disability status is confounded by baseline disability status. This suggests that interventions could focus on improving baseline disability score, potentially using illness perceptions to accomplish this goal.

Acknowledgements

We thank the patients of the HOSTAS cohort for participation in this study and research nurses B. van Schie-Geyer, A. Wongsodihardjo and M. Janson, data managers J. van Krol-Berkel and C. Kromme and PhD candidate S. van Beest in the Rheumatology Department and technicians in the Radiology Department of the LUMC for support in data collection. W. D. designed the study, acquired data for the HOSTAS cohort, scored the radiographs, analysed the data and drafted the manuscript. R.L. acquired data for the HOSTAS cohort and reviewed the manuscript. A.A.K., A.W.M.E. and H.v.M. were involved in data analysis and reviewed the manuscript. F.R.R. was involved in designing the study and data analysis and reviewed the manuscript. M.K. supervised the HOSTAS cohort, designed the study, was involved in data analysis and reviewed the manuscript.

Funding: Dutch Arthritis Foundation (Reumafonds) provided funding (LLP-24) for the HOSTAS cohort and appointment of the first author.

Disclosure statement: The authors have declared no conflicts of interest.

References

- Kloppenborg M, Kwok W-Y. Hand osteoarthritis—a heterogeneous disorder. *Nat Rev Rheumatol* 2011;8:22–31.
- Zhang Y, Niu J, Kelly-Hayes M *et al.* Prevalence of symptomatic hand osteoarthritis and its impact on functional status among the elderly: the Framingham Study. *Am J Epidemiol* 2002;156:1021–7.
- Kjeken I, Dagfinrud H, Slatkowsky CB *et al.* Activity limitations and participation restrictions in women with hand osteoarthritis: patients' descriptions and associations between dimensions of functioning. *Ann Rheum Dis* 2005;64:1633–8.
- Kwok W-Y, Vlieland TPMV, Rosendaal FR, Huizinga TWJ, Kloppenburg M. Limitations in daily activities are the major determinant of reduced health-related quality of life in patients with hand osteoarthritis. *Ann Rheum Dis* 2011;70:334–6.
- Bijsterbosch J, Watt I, Meulenbelt I *et al.* Clinical and radiographic disease course of hand osteoarthritis and determinants of outcome after 6 years. *Ann Rheum Dis* 2011;70:68–73.
- Dahaghin S, Bierma-Zeinstra SMA, Hazes JMW, Koes BW. Clinical burden of radiographic hand osteoarthritis: a systematic appraisal. *Arthritis Care Res* 2006;55:636–47.
- Botha-Scheepers S, Riyazi N, Watt I *et al.* Progression of hand osteoarthritis over 2 years: a clinical and radiological follow-up study. *Ann Rheum Dis* 2009; 68:1260–4.
- Hill S, Dziedzic K, Thomas E, Baker SR, Croft P. The illness perceptions associated with health and behavioural outcomes in people with musculoskeletal hand problems: findings from the North Staffordshire Osteoarthritis Project (NorStOP). *Rheumatology* 2007;46:944–51.
- Liu R, Damman W, Kaptein AA, Rosendaal FR, Kloppenburg M. Coping styles and disability in patients with hand osteoarthritis. *Rheumatology* 2016;55:411–8.
- Bijsterbosch J, Scharloo M, Visser AW *et al.* Illness perceptions in patients with osteoarthritis: change over time and association with disability. *Arthritis Care Res* 2009;61:1054–61.
- Petrie K, Weinman J. Why illness perceptions matter. *Clin Med* 2006;6:536–9.
- Weinman J, Petrie KJ, Moss-Morris R, Horne R. The illness perception questionnaire: a new method for assessing the cognitive representation of illness. *Psychol Health* 1996;11:431–45.
- Leventhal H, Diefenbach M, Leventhal EA. Illness cognition: using common sense to understand treatment adherence and affect cognition interactions. *Cogn Ther Res* 1992;16:143–63.
- Botha-Scheepers S, Riyazi N, Kroon HM *et al.* Activity limitations in the lower extremities in patients with osteoarthritis: the modifying effects of illness perceptions and mental health. *Osteoarthritis Cartilage* 2006;14:1104–10.
- Knowles SR, Nelson EA, Castle DJ *et al.* Using the common sense model of illness to examine interrelationships between symptom severity and health outcomes in end-stage osteoarthritis patients. *Rheumatology* 2016;55:1066–73.
- Kaptein AA, Bijsterbosch J, Scharloo M *et al.* Using the common sense model of illness perceptions to examine osteoarthritis change: a 6-year longitudinal study. *Health Psychol* 2010;29:56–64.

- 17 Hanusch BC, O'Connor DB, Ions P, Scott A, Gregg PJ. Effects of psychological distress and perceptions of illness on recovery from total knee replacement. *Bone Joint J* 2014;96-B:210-6.
- 18 Orbell S, Johnston M, Rowley D, Espley A, Davey P. Cognitive representations of illness and functional and affective adjustment following surgery for osteoarthritis. *Soc Sci Med* 1998;47:93-102.
- 19 Foster NE, Bishop A, Thomas E *et al*. Illness perceptions of low back pain patients in primary care: what are they, do they change and are they associated with outcome? *Pain* 2008;136:177-87.
- 20 Scharloo M, Kaptein AA, Weinman JA *et al*. Predicting functional status in patients with rheumatoid arthritis. *J Rheumatol* 1999;26:1686-93.
- 21 Dalbeth N, Petrie KJ, House M *et al*. Illness perceptions in patients with gout and the relationship with progression of musculoskeletal disability. *Arthritis Care Res* 2011;63:1605-12.
- 22 Skinner TC, Carey ME, Craddock S *et al*. Diabetes education and self-management for ongoing and newly diagnosed (DESMOND): process modelling of pilot study. *Patient Educ Couns* 2006;64:369-77.
- 23 Broadbent E, Ellis CJ, Thomas J, Gamble G, Petrie KJ. Further development of an illness perception intervention for myocardial infarction patients: a randomized controlled trial. *J Psychosom Res* 2009;67:17-23.
- 24 Buchbinder R, Jolley D, Wyatt M. Population based intervention to change back pain beliefs and disability: three part evaluation. *BMJ* 2001;322:1516-20.
- 25 Kwok WY, Plevier JWM, Rosendaal FR, Huizinga TWJ, Kloppenburg M. Risk factors for progression in hand osteoarthritis: a systematic review. *Arthritis Care Res* 2013;65:552-62.
- 26 Damman W, Liu R, Kroon FPB *et al*. Do comorbidities play a role in hand osteoarthritis disease burden? Data from the hand osteoarthritis in secondary care cohort. *J Rheumatol* 2017;44:1659-66.
- 27 Moss-Morris R, Weinman J, Petrie K *et al*. The Revised Illness Perception Questionnaire (IPQ-R). *Psychol Health* 2002;17:1-16.
- 28 Dreiser R, Maheu E, Guillou G, Caspard H, Grouin J. Validation of an algofunctional index for osteoarthritis of the hand. *Rev Rhum Engl Ed* 1995;62(6 Suppl 1):43S-53S.
- 29 Bellamy N, Campbell J, Haraoui B *et al*. Dimensionality and clinical importance of pain and disability in hand osteoarthritis: development of the Australian/Canadian (AUSCAN) Osteoarthritis Hand Index. *Osteoarthritis Cartilage* 2002;10:855-62.
- 30 NVR Dutch Rheumatology Association. Meetinstrumenten HAQ. NL_consensus_HAQ. https://www.nvr.nl/wp-content/uploads/2014/11/NL_consensus_HAQ.pdf (3 February 2017, date last accessed).
- 31 Doyle DV, Dieppe PA, Scott J, Huskisson EC. An articular index for the assessment of osteoarthritis. *Ann Rheum Dis* 1981;40:75-8.
- 32 Altman R, Alarcon G, Appelrouth D *et al*. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hand. *Arthritis Rheum* 1990;33:1601-10.
- 33 Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. *Ann Rheum Dis* 1957;16:494-502.
- 34 Kosinski M, Zhao SZ, Dedhiya S, Osterhaus JT, Ware JE. Determining minimally important changes in generic and disease-specific health-related quality of life questionnaires in clinical trials of rheumatoid arthritis. *Arthritis Rheum* 2000;43:1478-87.
- 35 Bellamy N, Hochberg M, Tubach F *et al*. Development of multinational definitions of minimal clinically important improvement and patient acceptable symptomatic state in osteoarthritis. *Arthritis Care Res* 2015;67:972-80.
- 36 Dziedzic KS, Thomas E, Hay EM. A systematic search and critical review of measures of disability for use in a population survey of hand osteoarthritis (OA). *Osteoarthritis Cartilage* 2005;13:1-12.
- 37 Knol MJ, Cessie SL, Algra A, Vandenbroucke JP, Groenwold RHH. Overestimation of risk ratios by odds ratios in trials and cohort studies: alternatives to logistic regression. *Can Med Assoc J* 2012;184:895-9.
- 38 Hampson SE. Personal models and the management of chronic illness: a comparison of diabetes and osteoarthritis. *Eur J Pers* 1997;11:401-14.
- 39 Raaij EJ, de Pool J, Maissan F, Wittink H. Illness perceptions and activity limitations in osteoarthritis of the knee: a case report intervention study. *Man Ther* 2014;19:169-72.
- 40 Visser AW, Bøyesen P, Haugen IK *et al*. Instruments measuring pain, physical function, or patient's global assessment in hand osteoarthritis: a systematic literature search. *J Rheumatol* 2015;42:2118-34.
- 41 Kloppenburg M, Bøyesen P, Visser AW *et al*. Report from the OMERACT Hand Osteoarthritis Working Group: set of core domains and preliminary set of instruments for use in clinical trials and observational studies. *J Rheumatol* 2015;42:2190-7.
- 42 Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990;1:43-6.
- 43 Bautmans I, Gorus E, Njemini R, Mets T. Handgrip performance in relation to self-perceived fatigue, physical functioning and circulating IL-6 in elderly persons without inflammation. *BMC Geriatr* 2007;7:5.