



Universiteit
Leiden
The Netherlands

CRISPR/Cas-induced targeted mutagenesis with *Agrobacterium* mediated protein delivery

Schmitz, D.J.

Citation

Schmitz, D. J. (2018, September 20). *CRISPR/Cas-induced targeted mutagenesis with Agrobacterium mediated protein delivery*. Retrieved from <https://hdl.handle.net/1887/65634>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/65634>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/65634> holds various files of this Leiden University dissertation.

Author: Schmitz, D.J.

Title: CRISPR/Cas-induced targeted mutagenesis with Agrobacterium mediated protein delivery

Issue Date: 2018-09-20

Chapter 1

General Introduction

Daan J. Schmitz, Sylvia de Pater, Paul J.J. Hooykaas

General introduction

In the first part of this chapter I will review the developments in the field of genome editing. Specifically the DNA repair mechanisms involved, the tools available and the applications of genome editing in plants. As I have used *Agrobacterium tumefaciens* for delivery, the second part of this chapter will focus on the plant pathogen *Agrobacterium*, describing the mechanisms underlying the transformation of plants by *Agrobacterium* and its biotechnological applications.

Part 1

Genetic engineering and genome editing

Controlled integration of transgenes and controlled genetic replacement of endogenous genes by transgenes based on homologous recombination, also known as gene targeting, is an important tool in biotechnology because it can accelerate the rate of functional gene analysis and guarantees a safe introduction of novel traits. This controlled manipulation of genomes is based on the introduction of DNA that contains homology with the chromosomal locus allowing for DNA recombination. In lower eukaryotes such as yeast, integration of DNA occurs predominantly via homologous recombination and therefore the recombinatorial events required for gene-targeting were first studied in the budding yeast *Saccharomyces cerevisiae* [1,2]. In higher eukaryotes this kind of integration also occurs, albeit at low frequencies compared to random integration. The first molecular evidence for gene-targeting in animal cells was provided by the laboratories of Smithies [3] and Capecchi [4,5] in experiments that generated loss-of-function mutations in embryonic mouse stem cells.

In plant cells the first report of gene-targeting appeared in 1988. A DNA repair template transferred to tobacco protoplasts was shown to recombine with an integrated defective copy of a resistance gene leading to integration of a restored gene [6]. In 1989 this was followed by a report showing that delivery of a repair construct via *Agrobacterium*, a bacterium commonly used to transform plants, in the form of a T-DNA could be used for gene targeting leading to the correction of a defective gene at its original locus in the genome [7]. However, to harness the power of homology-directed recombination for gene insertion or gene repair in plants the naturally low HDR frequencies ranging between 10^{-4} and 10^{-6} needed to be enhanced [7–13].

Early experiments showed that DNA-damaging agents stimulated the exchange between sister chromatids [14], but the most compelling evidence that breaks in the DNA enhanced recombination came from studies that showed that a single double-strand break (DSB) in the genome dramatically increases the local frequency of recombination. In these studies a fragment encompassing the recognition site for a specific DNA endonuclease and a defective resistance gene was first inserted into the genome. Recombination with a homologous donor DNA increased several orders of magnitude when the cognate DNA endonuclease was expressed in the recipient cells [15,16].

Initially only naturally occurring site specific nucleases (SSNs) were available for this purpose. During the last decade, however, artificial site specific endonucleases with a customizable DNA recognition and cleavage site were developed that can now be employed for targeted modification of almost any genetic information in the genomes of organisms. These SSNs can also be used for targeted mutagenesis. When SSNs are expressed and used to

induce DSBs, the DNA repair machinery of the cell will seal the break, but when the nuclease is persistently present the DNA will be broken again leading to a cycle of break-repair until imprecise repair leads to a loss of the nuclease restriction site. In this way mutations can be selected at the nuclease target site. If these mutations occur in coding sequences genes can be knocked out making it easier to study the function of the mutated gene. Mutations created in non-coding regions can be used for instance to disrupt binding sites of pathogen produced transcription factors that promote disease [17]. Using SSNs even entire gene clusters can be removed by inducing DNA breaks at opposing ends of the gene cluster [18–20] or for the knockout of multiple (redundant), non-allelic homologous genes at the same time.

Double-strand break repair

The genome is subject to many agents, both exogenous and endogenous leading to DNA damage, that, if unrepaired, may lead to mutation or gross chromosomal rearrangements. Such damage includes base lesions, DNA single-strand breaks (SSBs) and DSBs. Repairing the damage is vital to maintain an organisms genomic integrity and stability and therefore multiple DNA repair pathways have evolved. Exploiting such DNA repair mechanisms of DSBs, and to a lesser extent those involved in SSB repair and base excision repair (BER), underlies genome editing. Two distinct pathways can be used for the repair of DSBs; non-homologous end joining (NHEJ) and homologous recombination (HR). Which repair pathway is used for the repair of DSBs differs greatly between organisms and the cell cycle phase at which the repair of a DSB occurs. Repair via NHEJ operates throughout the cell cycle but predominantly in the G1 phase [21], whereas repair via HR is restricted to the S and G2 phases when sister chromatids are available as a repair template.

NHEJ repairs DSBs by religating the broken ends irrespective of sequence homology and can be precise, but may also result in small deletions or insertions disrupting the genetic information [22]. The first step in NHEJ is the binding of the heterodimeric Ku complex to both ends of a DSB. This complex is composed of a 70kDa and a 80kDa subunit, named Ku70 and Ku80, respectively [23]. After the binding of Ku to a DSB, in mammalian cells DNA-PKcs is recruited to the end of the DNA break [24,25]. DNA-PKcs, however, is not present in plants and fungi. NHEJ is completed by ligation of the DNA ends; this rejoining is carried out by a complex of DNA ligase IV, XRCC4 and the XRCC4-like factor [26]. In plants and yeast orthologues of these C-NHEJ components have been identified [27–31]. Although several bacterial strains have been identified with a NHEJ-like DNA repair mechanism, the repair of DSBs in bacteria is predominately based on repair by HR [32,33].

In the absence of canonical factors involved in Ku-dependent NHEJ, back-up pathways are responsible for residual end joining of DSBs [34–37]. Repair via back-up NHEJ starts with the resection of the ends producing ssDNA ends that can anneal at microhomology regions. The ends may be linked at these microhomology regions, whereafter non-homologous tails are removed. The remaining gaps are filled by a specific DNA polymerase and re-ligated by DNA ligase I [38]. In these back-up pathways microhomologies sequences (5-25 basepairs) are frequently used for the repair of the DSB resulting in deletions [39] and therefore this pathway has also been called microhomology mediated end-joining (MMEJ) [40]. Several factors involved in repair via MMEJ have been identified: PARP1/2, MRN, CtIP, Ligase3, XRCC1 and DNA polymerase θ (Pol θ). The exact molecular mechanism behind MMEJ is poorly understood, but PARP1, the MRN complex and Pol θ have been implicated as key

players [41–43]. Recently, it has been shown that theta mediated end joining (TMEJ) is the dominant pathway for repair in Ku-deficient cells and the occurrence of microhomologies at the break site which is the result of the ability of Pol θ to mediate joining of two resected 3' ends harboring DNA sequence microhomology [42,43].

Homologous recombination is a DNA repair mechanism that uses DNA homology to direct DNA repair. These homologous sequences required for repair by HR are preferentially found on sister chromatids, but other naturally occurring homologous sequences or an artificially introduced repair template may sometimes be used instead. Repair via HR can be divided into several subpathways, classical double-strand break repair (DSBR), synthesis-dependent strand annealing (SDSA), single-strand annealing (SSA) and break-induced replication (BIR) [44]. DSBR was initially described to explain crossover events during meiosis and gene conversion [45]. SDSA is used for mitotic DSB repair [46] and is the predominant repair mechanism for two ended DSBs via HR [47]. In plants evidence has been found for the occurrence of DSBR, SDSA and SSA [48].

HR invariably starts off with the 5' resection of the ends resulting in large 3' ssDNA stretches that can search for complementary sequences. Repair via SDSA starts with a 3' end invading a homologous double-strand forming a D-loop. This is followed by repair synthesis from the 3' end using the newly paired strand as a template. The "X" shaped structure formed at the border between the hetero- and homoduplex of the D-loop is called a Holiday junction. After elongation, in SDSA the invading strand is displaced from the D-loop structure and anneals back to the 3' homologous strand that was formed by resection of the other end of the DSB. Thus in SDSA the donor molecule remains unaltered and gene conversion without loss of sequence information is the final result of the reaction [49]. SDSA is preferably used for HR repair in mitotic cells in the S and G2 phase, when a sister chromatid is available as a template. DSBR is mainly used for the repair of DSB breaks in meiotic cells. In this case both DNA ends invade a homologous chromosome to copy genetic information and this results in a double Holiday junction that may be resolved into a crossover or a non-crossover product.

SSA is a repair mechanism that can be used when two homologous sequences are in close proximity and arranged in a tandem orientation. Repair via SSA starts with resection and the production of 3' single-stranded overhangs. This is followed by immediate pairing of the single stranded complementary sequences found near the break site and trimming of the any remaining 3' ends [50]. SSA is not conservative in contrast to SDSA and DSBR and leads to deletion of one repeat and the intervening sequence. Up to one out of three DSBs in an artificial genomic region with tandemly arranged duplications was repaired via SSA after DSB induction [51].

Site-specific nucleases

To induce targeted DSBs, four different classes of SSNs have been developed: homing endonucleases (HEs), zinc finger nucleases (ZFNs), TAL effector nucleases (TALENs) and the CRISPR/Cas RNA guided endonuclease encoded by the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated (Cas) system (Fig. 1) [52].

The first class of SSNs are the HEs which are a class of proteins typically encoded for by introns or inteins and are considered selfish genetic elements, much like transposons [53]. This class of proteins can be divided into eight different subclasses: LAGLIDADG, H-N-H, His-Cys, GIY-YI G, PD(D/E)xK, HJ resolvase-like, EDxHD and Vsr-like [54]. All of these different classes of HEs have large DNA sequence recognition sites varying in length

between 12 to 40 basepairs. Because the DNA binding domain and the cleavage domain of HEs overlaps [55] creating HEs with new DNA binding specificities is challenging. For HEs of the LAGLIDADG class progress has however been made in altering the DNA recognition and cleavage site [56,57]. HEs were the first to be used for targeted mutagenesis in *Arabidopsis* and maize and instrumental in showing that DSB induction enhances targeted mutagenesis and gene targeting in plants (see Table 1).

In ZFNs the DNA binding domain consisting of zinc fingers is fused to the non-specific nuclease domain of the FokI nuclease. The zinc-finger array responsible for DNA binding is created by combining multiple zinc-finger domains, of the C_2H_2 class of zinc-fingers, that were discovered as part of the transcription factor IIIA [58]. This class of zinc-fingers consists of two β -sheets and an α -helix, which fits directly into the major groove of double strand DNA. The side chains from the N-terminal part of this helix contact the edges of the basepairs. Changing the amino acids in the α -helix may alter the affinity for different DNA sequences. Each zinc-finger domain recognizes 3bp of DNA [59]. Zinc-fingers have been created for the recognition of most of the 64 possible triplets [60–62]. By fusing the FokI nuclease domain to the zinc-finger array a zinc-finger nuclease (ZFN) is created. Two ZFNs are usually constructed that bind the DNA at opposing sites of the target sequence, each making a nick in one of the DNA strands leading together to a staggered DSB. A problem that can arise is the formation of a homodimer consisting of two ZFNs for one of the half sites. As a consequence, a DSB may be induced in the genome at a position which was not the initial target, possibly leading to toxicity [63]. Miller et al. [64,65] have designed complementary FokI cleavage domain variants that together function as an obligate heterodimer. The widespread application of ZFNs is however limited because of the limited selectivity conferred by the zinc-finger modules, and the complex context-dependent interactions between neighboring zinc-finger modules [66,67]. Moreover designing ZFNs typically involves multiple rounds of testing without necessarily resulting in a nuclease that performs optimally [66,68–70]. ZFNs have been used for genome editing in *Arabidopsis*, tobacco, soybean and maize (see Table 1).

The third class of artificial restriction enzymes called TAL effector nucleases (TALENs) has been developed as the successor of ZFNs. Transcription activator-like effectors (TALEs) are produced by plant pathogenic bacteria in the genus *Xanthomonas* and are transferred via type III secretion systems to the host, where they function as important virulence factors that act as transcriptional regulators [71–73]. The DNA binding domain of these TALEs contains multiple 30–35 amino acid long repeats that each recognizes a single base pair of DNA. Two hyper variable amino acids found at positions 12 and 13 of these repeats, known as the repeat-variable di-residues (RVDs), determine the base specificity [74,75]. Crystallization of the TALE DNA-binding domain revealed how the TALE wraps around the sense strand of the DNA as a right-handed super helix with each repeat forming a left-handed helix-loop-helix structure [76]. The structure also revealed that the 13th amino acid of the RVD determines the base specificity by interacting with the major groove, and that the 12th amino acid of the RVD stabilizes the loop in the helix-loop-helix structure [76,77]. By fusing the DNA binding domain of TALEs to the nuclease domain of FokI, TALENs are constructed that just like ZFNs can be designed to bind at opposing sites of the target sequences and create a DSB upon dimerization of the nuclease domain [78]. TALENs have been used to modify the genome of the following plants: *Arabidopsis*, barley, *Brachypodium*, cucumber, maize, rice, soybean, tobacco, tomato and wheat (see Table 1).

The newest addition to the family of SSNs is the CRISPR/Cas system derived from a prokaryotic adaptive immune system that cleaves DNA in a sequence dependent manner. In prokaryotes this RNA based defense system provides protection from foreign invading nucleic acids, such as viruses and plasmids [79–81]. The CRISPR/Cas system is made up of a Cas gene operon and CRISPR arrays. Immunity is acquired by integrating short fragments of invading DNA known as spacers between two flanking identical repeats at the proximal region of CRISPR loci. Transcripts of these CRISPR arrays, including the spacers, are processed into small interfering CRISPR RNAs (crRNAs) approximately 40 nucleotides in length [82]. Together with transactivating RNAs (tracrRNAs) these crRNAs form a complex which enables the Cas enzyme to cleave homologous double-stranded DNA sequences, known as protospacers, in previously encountered invading DNA [83,84].

There are three distinct classes of CRISPR/Cas systems from which a type II class system from *Streptococcus pyogenes* has mostly been used for genome editing, because only one protein (Cas9) is sufficient for nuclease activity in combination with the two RNA molecules [85]. A requirement for cleavage is the presence of a conserved protospacer adjacent motif (PAM) downstream of the target DNA, with either a 5'-NGG-3' or 5'-NAG-3' motif [85–87]. The quick development of the CRISPR/Cas system into a tool for genome editing started with the discovery that a 20 nucleotide change in the crRNA was sufficient to reprogram the DNA target specificity [85], and the subsequent development of a single chimeric guide RNA that combines the target specificity of the crRNA with the structural properties of the tracrRNA. The first reports of the CRISPR/Cas system being used for genome editing in different eukaryotes followed shortly thereafter [88–92]. The Cas9 protein has two active sites in a RuvC and a HNH domain, respectively, each of which induces a single-strand break which combined create a DSB. To transform Cas9 into a nickase that induces a single strand break point mutations have been introduced into the RuvC (D10A) and HNH (H840A) domains [85,86,93]. Using a nickase instead of a nuclease strongly decreases the frequency of off target mutations, while still increasing the frequency of HDR [94,95]. The CRISPR/Cas system has been used successfully for genome editing in the following plants: *Arabidopsis*, *Camelina*, common wine grape, tobacco, maize, petunia, orchids, potato, red sage, rice, sweet orange, tomato, sorghum, watermelon wheat and (see Table 2).

Using site-specific nucleases for genome editing in plants

The outcome of repair of a DSB is dependent on which repair pathway is used. In plants and many other higher eukaryotes repair via NHEJ is the predominant pathway used for the repair of DSBs. Repair via HR, but also via NHEJ often results in perfect repair of the lesion, but therefore also restores the target site for the SSNs allowing for the induction of a new DSB. This process can therefore cycle between DSB induction and repair until imperfect repair via NHEJ results in a small deletion or insertion that destroys the recognition site of the site specific nuclease.

When SSNs are used to introduce DSBs into a gene, mutations can be induced that affect gene function. All four classes of SSNs have been used to this end in both model plants as well as in a wide range of crop species (for an overview of see Table 1). High mutation frequencies have been found for instance when poplar leaf discs were transformed with a constitutive CRISPR/Cas construct resulting in 89% of leaf discs transformed showing an albino shoot indicative of mutations created in the phytoene desaturase gene [96]. Maize plants regenerated from calli that constitutively expressed the CRISPR/Cas system were found to have mutations frequencies up to 70-100% [97].




Figure 1. Site-specific nucleases used for genome editing. **(A)** Homing endonuclease with overlapping DNA binding and cleavage domain. **(B)** Zinc-finger nuclease (ZFN) composed of multiple zinc-fingers that form the DNA binding domain fused to the FokI nuclease domain. **(C)** TALEN composed of the DNA binding domain of transcription activator-like effectors (TALEs) fused to the FokI nuclease domain. **(D)** CRISPR/Cas composed of the Cas9 protein and the sgRNA which determines the base specificity with a 20bp sequence that is complementary to the target sequence.

Mutations created in the T0 generation however did not show expected inheritance patterns and are not always inherited by the T1 progeny [98,99]. In *Arabidopsis* chimerism was found to occur in T1 plants obtained after floral dip transformation [100,101]. More complex genetic modifications can be achieved when SSNs are used to induce two DSBs simultaneously, including specific deletions [19,69], inversions [102], duplications [102] and translocations [103]. In plants ZFNs, homing endonucleases and CRISPR/Cas have been used to create targeted chromosomal deletions [20,104–106].

When a DSB is induced and simultaneously an artificial repair template lacking the nuclease restriction site is present, this may be used for HDR, and the specific mutations present in the template introduced into the genome. In plants this has been accomplished in the following species: *Arabidopsis*, tobacco, tomato, soybean, rice, potato, wheat and maize. Similarly by using a repair template containing novel genes these can be inserted in this way at the DSB site (gene targeting). In plants the introduction of DSBs near the site of the desired recombination has been shown to greatly increase the frequency of HDR. The introduction of a SSB by engineered nickases also increases the frequency of HDR, although the increase is less pronounced as with the introduction of a DSB [107,108]. Repair via HDR has been used in plants for targeted modifications of genes in model species as well in several crop species using ZFNs [109–112], TALENs [113,114] and the CRISPR/Cas system [115–120].

Delivery of site-specific nucleases

Although the choice for the delivery system used will often be based on efficiency and feasibility considerations, the resulting expression levels and concentration of the nuclease will also impact the outcome. In plants several methods of delivery have been used for either constitutive or transient expression of SSNs. To obtain plants that constitutively express SSNs varying plant tissues have been transformed using biolistic or *Agrobacterium* based methods (Table 1 and Table 2). These methods, although relatively easy, have some drawbacks. Integration of transgenes may disrupt endogenous genes at the site of transgene integration. The presence of transgenes increases the administrative burden in regulatory processes needed for marketing of genetically modified crops. Furthermore, depending on the position in the genome gene expression levels might be influenced and constitutive expression of the SSN may lead to off target effects. Therefore methods are being developed for the transient expression of SSNs, for instance by controlling them by inducible or cell specific promoters. Alternatively methods for the direct delivery of nuclease mRNA have also been developed [90,121,122], as well for the direct delivery of SSN proteins in human cells and drosophila [88,123,124] and tobacco protoplasts for which direct introduction of I-SceI and TALEN protein was reported [125]. In addition, preassembled CRISPR/Cas complexes of purified Cas9 protein and sgRNA have been transfected into protoplasts of *Arabidopsis*, tobacco, rice and lettuce [126]. Direct protein delivery does however require the isolation and purification of large quantities of SSNs and the isolation of protoplasts. Therefore, a system that eliminates the isolation and purification steps but would still directly deliver the SSN as a protein would have value in both academic and commercial settings. The *Agrobacterium* system may fulfill this promise.

Table 1. Overview of use of homing endonucleases, zing finger nucleases and TALENs in plants

Site specific nuclease	Transformation method	Mutation type	Reference
Homing endonucleases			
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM / Gene excision	[185]
<i>Z. mays</i>	<i>Agrobacterium</i> (Embryo)	TM	[186]
<i>Z. mays</i>	<i>Agrobacterium</i> (Embryo)	TM	[187]
Zinc finger nucleases			
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[188]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[189]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[190]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM / GT	[109]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM / GT	[110]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[191]
<i>N. tabacum</i>	Protoplast electroporation	TM / GT	[111]
<i>Z. mays</i>	<i>Agrobacterium</i> (Embryo)	TM / Gene addition	[112]
TALENs			
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[192]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[193]
<i>A. thaliana</i>	PEG-protoplast transfection	TM	[194]
<i>B. distachyon</i>	PEG-protoplast transfection	TM	[195]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[196]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[197]
<i>H. vulgare</i>	<i>Agrobacterium</i> (Embryonic pollen), Particle bombardment	GT	[198]
<i>H. vulgare</i>	<i>Agrobacterium</i> (Embryonic pollen)	TM	[199]
<i>H. vulgare</i>	<i>Agrobacterium</i> (Embryonic pollen)	TM	[200]
<i>H. vulgare</i>	Particle bombardment	GT	[198]
<i>N. benthamiana</i>	mRNA transfection	TM	[201]
<i>N. benthamiana</i>	PEG-protoplast transfection	GT	[202]
<i>N. tabacum</i>	PEG-protoplast transfection	TM / GT	[113]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[203]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[204]
<i>O. sativa</i>	Particle bombardment	GT	[205]
<i>O. sativa</i>	PEG-protoplast transfection	TM	[120]
<i>O. sativa</i>	PEG-protoplast transfection	TM	[206]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Seedlings)	TM / GT	[114]
<i>S. tuberosum</i>	PEG-protoplast transfection	TM	[207]
<i>T. aestivum</i>	PEG-protoplast transfection	TM	[208]
<i>Z. mays</i>	<i>Agrobacterium</i> (Embryo)	TM	[209]

TM = targeted mutagenesis, GT = gene-targeting, CD = Chromosomal deletions

Table 2. Overview of use CRISPR/Cas in plants

CRISP/Cas9	Transformation method	Mutation type	Reference
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[210]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[211]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[212]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[213]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[214]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[215]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[216]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[101]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[217]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[218]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM	[219]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM / GT	[116]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM / GT	[220]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM (SSA)	[106]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Floral dip)	TM (SSA)	[221]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Leaf infiltration)	CD	[222]
<i>A. thaliana</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[223]
<i>A. thaliana</i>	PEG-protoplast transfection	TM	[126]
<i>A. thaliana</i>	PEG-protoplast transfection	TM	[224]
<i>A. thaliana</i>	PEG-protoplast transfection / <i>Agrobacterium</i> (Floral dip)	TM / GT	[115]
<i>Camelina sativa</i>	<i>Agrobacterium</i> (Floral dip)	TM	[225]
<i>C. lanatus</i>	PEG-protoplast transfection	TM	[226]
<i>C. sinensis</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[227]
<i>C. sinensis</i>	<i>Agrobacterium</i> (Cotyledons)	TM	[228]
<i>C. sativus</i>	<i>Agrobacterium</i> (Cotyledons)	TM	[229]
<i>Chrysanthemum morifolium</i>	<i>Agrobacterium</i> (Callus)	TM	[230]
<i>Dendrobium officinale</i>	<i>Agrobacterium</i> (protocorns)	TM	[231]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[232]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[233]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[196]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[234]
<i>G. max</i>	PEG-protoplast transfection	TM	[235]
<i>G. max</i>	Particle bombardement	TM / GT	[236]
<i>G. max</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[237]
<i>Gossypium hirsutum</i> L.	<i>Agrobacterium</i> (Hypocotyl/cotyledonary petiole)	TM	[238]
<i>Gossypium hirsutum</i> L.	<i>Agrobacterium</i> (Hypocotyl/cotyledonary petiole)	TM	[239]
<i>Gossypium hirsutum</i> L.	<i>Agrobacterium</i> (Hypocotyl/cotyledonary petiole)	TM	[240]
<i>Hordeum vulgare</i>	<i>Agrobacterium</i> (Embryo)	CD	[241]
<i>L. sativa</i>	PEG-protoplast transfection	TM	[126]
<i>M. truncatula</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[242]
<i>M. truncatula</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[237]
<i>Marchantia polyorpha</i>	<i>Agrobacterium</i> (spores)	TM	[243]
<i>N. attenuata</i>	PEG-protoplast transfection	TM	[235]
<i>N. attenuata</i>	PEG-protoplast transfection	TM	[126]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Cas9) / guideRNA (CaLCuV virus)	TM	[244]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Cas9) / guideRNA (TVR)	TM	[245]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Cas9) / guideRNA (TYLCV virus)	TM	[246]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[247]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[223]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM / CD	[104]
<i>N. benthamiana</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[248]

<i>N. benthamiana</i>	<i>Agrobacterium</i> (Leaf infiltration)	CD	[222]
<i>N. benthamiana</i>	PEG-protoplast transfection	TM	[115]
<i>N. benthamiana</i>	PEG-protoplast transfection	TM	[224]
<i>N. benthamiana</i>	PEG-protoplast transfection	TM	[249]
<i>N. tabacum</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[250]
<i>N. tabacum</i>	<i>Agrobacterium</i> (Leaf disc)	TM	[251]
<i>N. tabacum</i>	PEG-protoplast transfection	TM	[252]
<i>N. tabacum</i>	PEG-protoplast transfection	TM	[253]
<i>N. tabacum</i>	PEG-protoplast transfection	TM / CD	[105]
<i>N. tabacum</i>	PEG-protoplast transfection / <i>Agrobacterium</i> (Leaf disc)	TM	[254]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[255]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[250]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[210]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[256]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[257]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[211]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[258]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[259]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[260]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[261]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[262]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[263]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[264]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[265]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[266]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[267]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[98]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[268]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[269]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[270]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	TM	[271]
<i>O. sativa</i>	<i>Agrobacterium</i> (Callus)	GT	[272]
<i>O. sativa</i>	<i>Agrobacterium</i> (Particle bombardment)	TM / GT	[273]
<i>O. sativa</i>	PEG-protoplast transfection	TM	[126]
<i>O. sativa</i>	PEG-protoplast transfection	TM	[274]
<i>O. sativa</i>	PEG-protoplast transfection	GT / GI	[275]
<i>O. sativa</i>	PEG-protoplast transfection / <i>Agrobacterium</i>	TM / CD	[20]
<i>O. sativa</i>	PEG-protoplast transfection / <i>Agrobacterium</i>	TM / CD	[104]
<i>O. sativa</i>	PEG-protoplast transfection / Particle bombardement	TM	[120]
<i>Petunia hybrida</i>	PEG-protoplast transfection ribonucleoproteins	TM	[276]
<i>Petunia hybrida</i>	<i>Agrobacterium</i> (Leaves)	TM / CD	[106]
<i>Physcomitrella patens</i>	PEG-protoplast transfection	TM	[277]
<i>Physcomitrella patens</i>	PEG-protoplast transfection	TM	[278]
<i>P. tomentosa</i>	<i>Agrobacterium</i> (Leaf disc)	TM	[96]
<i>P. tomentosa</i>	<i>Agrobacterium</i> (Leaf disc)	TM	[279]
<i>P. tomentosa</i>	<i>Agrobacterium</i> (Leaf disc)	TM	[280]
<i>S. bicolor</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[223]
<i>Salvia miltiorrhiza</i>	<i>Agrobacterium</i> (Hairy roots)	TM	[281]
<i>Scopelophila cataractae</i>	PEG-protoplast transfection	TM	[278]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	CD	[282]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	TM	[283]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	TM	[114]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	TM	[284]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	TM	[285]
<i>S. lycopersicum</i>	<i>Agrobacterium</i> (Cotyledon segments)	TM	[286]
<i>S. lycopersicum</i>	<i>A. rhizogenes</i>	TM	[287]
<i>S. tuberosum</i>	<i>Agrobacterium</i> (Callus)	TM / GT	[119]

<i>S. tuberosum</i>	<i>Agrobacterium</i> (Stem segments)	TM	[288]
<i>S. tuberosum</i>	PEG-protoplast transfection	TM	[289]
<i>T. aestivum</i>	Particle bombardement	TM	[290]
<i>T. aestivum</i>	PEG-protoplast transfection	TM	[291]
<i>T. aestivum</i>	PEG-protoplast transfection	TM / GT	[120]
<i>T. aestivum</i>	PEG-protoplast transfection	TM	[208]
<i>T. aestivum</i>	PEG-protoplast transfection / Particle bombardement	TM	[292]
<i>T. aestivum</i>	<i>Agrobacterium</i> (Leaf infiltration)	TM	[293]
<i>Vitis vinifera</i>	<i>Agrobacterium</i> (Embryo)	TM	[294]
<i>Z. mays</i>	<i>Agrobacterium</i> (Embryo)	TM	[99]
<i>Z. Mays</i>	<i>Agrobacterium</i> (Embryo) / Particle bombardement	TM / GT / GI	[295]
<i>Z. mays</i>	<i>PEG-protoplast transfection</i> / <i>Agrobacterium</i> (Embryo)	TM	[296]
<i>Z. mays</i>	<i>PEG-protoplast transfection</i> / <i>Agrobacterium</i> (Embryo)	TM	[217]
<i>Z. mays</i>	PEG-protoplast transfection	TM	[297]
<i>Z. mays</i>	PEG-protoplast transfection	TM	[223]
<i>Z. mays</i>	PEG-protoplast transfection ribonucleoproteins	TM	[298]

TM = targeted mutagenesis, GT = gene-targeting, CD = Chromosomal deletions

Part 2

Agrobacterium

Agrobacterium tumefaciens is known as the causative agent of crown gall disease in plants since the beginning of the 20th century. Crown galls arise at the site of infection and are the result of uncontrolled cell division. They can be excised and propagated in vitro without exogenous plant hormones [127]. Crown galls produce opines, amino-acid derivatives, which can be used by *Agrobacterium* as a nitrogen and carbon source and therefore the galls form a favorable niche for the bacterium [128]. The genetic basis for the uncontrolled growth of crown galls is the transfer of a copy of the T-region of the a large Tumor inducing (Ti) plasmid from the bacterium into plant cells. Processing of this T-region in the bacterium results in production of single stranded copies known as T-strands (transfer strands) that are transferred through a virB/virD4 type IV secretion system (T4SS) directly into the host cells. Upon entry into the host cell the T-strand is directed towards the nucleus and eventually may be integrated into the host genome [129]. The genes on this transferred DNA (T-DNA) encode for several proteins involved in the production and activation of plant hormones and the production of opines [130,131].

Induction of the virulence genes

The transformation of plant cells starts with a large cascade of events, the first step of which is the detection of phenolic compounds released by wounded plant cells by the bacterium through the VirA/VirG two-component sensor/regulatory system. VirA is autophosphorylated upon interaction with these phenolic compounds [132–135]. Phosphorylation of VirG by VirA leads to the binding of VirG to the *vir*-boxes of *vir* genes, stimulating their expression [136]. In addition to the VirA/VirG system the chromosomally encoded periplasmic protein ChvE, acts synergistically with the VirA/VirG system to induce *vir* gene expression upon detection of specific sugars in the plant sap [137]. Activation of the *vir* system only occurs in medium with a low pH like plant sap and at moderate temperatures (below 30° degrees). Eight operons designated *VirA*, *B*, *C*, *D*, *E*, *F*, *G* and *H* are largely conserved on the virulence regions of different types of Ti-plasmid. [138–140].

T-DNA processing

The T-region on the Ti-plasmid is flanked by two imperfect direct repeats (borders) that are the only determinants that define the T-DNA, of which only the right border sequence is essential for transfer [141]. Both of these borders are recognized by the virulence proteins VirD1 and the relaxasome VirD2. A single stranded break is introduced by VirD2 which stays covalently attached to the 5'-end of the nick [142,143]. The accessory Vir protein VirD1 enhances the binding and nicking on supercoiled DNA, while VirC1 and VirC2 further increase the efficiency by attaching to the overdrive sequence found close to the right border [144,145]. Probably the T-strand is released by displacement synthesis with VirD2 still covalently attached to the 5' end. This nucleo-protein complex is recognized by the T4SS and transported into the host cell [146].

The type IV secretion system of Agrobacterium

The T4SS of *Agrobacterium* through which the T-strand is transported is a large multi-protein complex that spans the inner membrane, the periplasm and the outer membrane. The genes encoding the T4SS are located on the Ti-plasmid and are encoded by a large *virB* operon, consisting of 11 open reading frames, and the *virD4* gene. Studying the T4SS has led to a division of its components into energy providing subunits (VirB4, VirB11, VirD4), inner membrane channel components located predominantly in the inner membrane (VirB3, VirB6, VirB8), periplasmic and membrane outer components (VirB7, VirB9), connecting component (VirB10), pilus components (VirB2, VirB5) and the VirB1 transglycosylase (extensively reviewed [147]). The T4SS present in *Agrobacterium* also translocates separately the T-strand four other so called effector proteins (VirD5, VirE2, VirE3 and VirF) into the host cell [148–150]. Their translocation via the T4SS is dependent on a C-terminal translocation signal present in all translocated proteins that has a net positive charge with a consensus motif of R-X(7)-R-X-R-X-R-X-X(n)>[149]. Combined with VirD2, VirE2 aids in nuclear targeting of the T-strand [151,152]. The translocated effector protein VirE3 is highly conserved and has two potential NLS by which it binds to importin- α and is transported into the nucleus [153]. Upon mutation of VirE3 hardly any attenuation of virulence is observed, but combined with the inactivation of VirF the role of VirE3 in virulence becomes apparent [154]. Recently, it was shown that VirE3 is a transcription factor which upregulates the VIP1-binding F-box gene thereby removing the requirement for VirF [155]. A recent study focusing on the virulence protein VirD5 suggests that it binds to centromeres/kinetochores and there may induce cycle arrest, chromosome missegregation and aneuploidy [156]. The VirF protein is an F-box protein [157] which together with homologues of the yeast protein Skp1 and Cullin forms a Skp-Cullin-F-box protein (SCF) complex [157]. This complex has been reported to target the host protein VIP1 and associated VirE2 for degradation by the 26S proteasome [158], which may be required for the uncoating of the T-DNA to enable integration into the host cell's genome. VirF also targets degradation of the transcription factors VBF3 and VBF4 [159]. Plant species that can be transformed by an *Agrobacterium* strain lacking *virF* have been shown to induce upon infection the expression of a plant F-box protein, VBF, that can functionally replace VirF [160].

T-DNA transfer and integration

T-DNA transfer across the different T4SS components occurs in a tightly controlled manner in which the T-strand makes sequential contact with VirD4, VirB11, VirB6, VirB8, VirB9 and VirB2 [161]. Although the other components of the T4SS have not been found to interact with the T-strand they are essential [161–163]. The translocated VirE2 protein is thought to be responsible for protecting the T-strand against degradation by host cell proteases/nucleases after the T-strand has been transported to the hosts cytoplasm [164]. In the nucleus the T-strand is converted into a double strand form, which may be expressed for some time and then degraded. Integration into the genome leads to stable transformation and the continued expression that is necessary for tumorigenesis. The integration of the T-DNA occurs at a random position into the genome [165], which can result in mutation and variable expression due to position effects. The molecular mechanism of T-DNA integration was first studied in yeast where integration occurs efficiently via HR when the T-DNA contains yeast homologous sequences [166]. If no homology is present on the T-DNA integration in both

yeast and fungi occurs by NHEJ and is dependent on the C-NHEJ proteins Ku70, Ku80 and DNA ligase 4 [167,168]. Studies on the influence of these proteins on T-DNA integration in plants by different research groups mostly showed that their absence gave no or limited effects [30,169–171]. Disruption of multiple DNA repair pathways simultaneously also did not eliminate transformation [172–174]. Recently, it has been shown that the random integration of the T-strand occurs through a mechanism involving Pol θ [175].

Biotechnological applications of Agrobacterium

After the discovery of the genetic mechanism behind the transformation of plants by *Agrobacterium*, researchers focused on developing *Agrobacterium* into a tool for the introduction of foreign genes into plants. The genes that are naturally located between the borders are not involved in T-DNA transfer and these sequences can be replaced with other sequences of interest. The introduction of these genes of interest into the Ti-plasmid is however difficult due to its large size, low copy number and its inability to replicate in *E. coli*. The *vir* region and the T-DNA can however be separated on different plasmids without impacting *Agrobacterium*'s ability to form tumors [176]. The plasmid harboring the T-DNA, called the binary vector, contains at least one origin of replication that is functional in both *Agrobacterium* and *E. coli* and selectable markers for maintenance in *Agrobacterium* and *E. coli*, hereby allowing the easy cloning of genes on the T-region [176]. The plasmid with the intact *vir* region but lacking the T-region is known as the helper plasmid. The ability to introduce foreign genes into plant cells has made *Agrobacterium* an invaluable tool for plants scientists.

Using *Agrobacterium* plants have been created with enhanced tolerance to abiotic and biotic stress and pest resistance. Furthermore, *Agrobacterium* has been applied to better understand plant biology at a cellular and molecular level for example tagging various proteins with a fluorescent protein to visualize cell compartments [177]. In addition T-DNA insertion libraries have been made in which the T-DNA functions as an insertional mutagen and, by extension, a gene tag [178,179]. More recently SSNs have been introduced into plants, using *Agrobacterium*, for targeted gene disruption and gene targeting in plants (see Table 1). In recent years, *Agrobacterium* has also been used to modify plants for the production of useful proteins, such as edible vaccines and recombinant antibodies [177].

Although *Agrobacterium* only forms tumors on dicotyledonous plants, under laboratory conditions it is also capable of transforming monocotyledonous plants [180], yeast [166] and a wide range of fungi [181]. Because of its ease of use, low cost and precision it has become a preferred vector not only for the genetic modification of plants, but also of yeasts and fungi. As mentioned above *Agrobacterium* not only uses its T4SS for T-DNA transfer but also for the translocation of several effector proteins. It has been found that the translocation signal of these effector proteins can be attached to heterologous proteins such as the Cre recombinase and I-SceI to effectuate their transfer to plant cells [148,149,166,182–184].

Outline of this thesis

In **Chapter 2** we describe the application of CRISPR/Cas in *Agrobacterium*, to cure *Agrobacterium* from the promiscuous plasmid RP4 and from vectors with the replication unit of the octopine Ti plasmid. Furthermore we show that the Cas9 protein fused to a translocation signal recognized by the T4SS does not negate its ability to induce DSBs.

In **Chapter 3** yeast was used as a model organism to show that the Cas9 protein can be delivered through the T4SS of the plant pathogen *Agrobacterium*. The transfer of Cas9 was effectuated by fusion of a T4SS translocation peptide to the Cas9 protein.

In **Chapter 4** a method for targeted mutagenesis in *Nicotiana benthamiana* was developed, that is based on the translocation of the Cas9 protein through the T4SS of *Agrobacterium*. We show that concurrent transfer of Cas9 protein and a T-DNA encoding the sgRNA results in targeted mutations in the infiltrated leaves of *N. benthamiana*.

In **Chapter 5** we describe two novel approaches for the transient expression of the *Agrobacterium* derived isopentenyl transferase that can be used for the selection of transformed plants. The first consisted of the delivery of a T-DNA encoding the *ipt* gene into Pol- θ -deficient *Arabidopsis* mutants in which only transient expression of the T-DNA occurs but no integration. The second approach involved the direct delivery of the IPT protein through the *Agrobacterium* T4SS into *Arabidopsis*.

References

1. Scherer S, Davis RW. Replacement of chromosome segments with altered DNA sequences constructed in vitro. *Proc. Natl. Acad. Sci. USA.* 1979;76:4951–5.
2. Orr-Weaver TL, Szostak JW, Rothstein RJ. Yeast transformation: a model system for the study of recombination. *Proc. Natl. Acad. Sci. USA.* 1981;78:6354–8.
3. Doetschman T, Gregg RG, Maeda N, Hooper ML, Melton DW, Thompson S, et al. Targeted correction of a mutant HPRT gene in mouse embryonic stem cells. *Nature.* 1987;330:576–8.
4. Thomas KR, Capecchi MR. Site-directed mutagenesis by gene targeting in mouse embryo-derived stem cells. *Cell.* 1987;51:503–12.
5. Capecchi MR. Altering the genome homologous recombination *Science.* 1989;244:1288–92.
6. Paszkowski J, Baur M, Bogucki A, Potrykus I. Gene targeting in plants. *EMBO J.* 1988;7:4021–6.
7. Offringa R, de Groot MJ, Haagsman HJ, Does MP, van den Elzen PJ, Hooykaas PJJ Extrachromosomal homologous recombination and gene targeting in plant cells after *Agrobacterium* mediated transformation. *EMBO J.* 1990;9:3077–84.
8. Hroudá M, Paszkowski J. High fidelity extrachromosomal recombination and gene targeting in plants. *Mol. Gen. Genet.* 1994;243:106–11.
9. Lee KY, Lund P, Lowe K, Dunsmuir P. Homologous recombination in plant cells after *Agrobacterium*-mediated transformation. *Plant Cell.* 1990;2:415–25.
10. Halfter U, Morris PC, Willmitzer L. Gene targeting in *Arabidopsis thaliana*. *Mol. Gen. Genet.* 1992;231:186–93.
11. Miao ZH, Lam E. Targeted disruption of the TGA3 locus in *Arabidopsis thaliana*. *Plant J.* 1995;7:359–65.
12. Risseuw E, Offringa R, Franke-van Dijk MEI, Hooykaas PJJ. Targeted recombination in plants using *Agrobacterium* coincides with additional rearrangements at the target locus. *Plant J.* 1995. 109–19.
13. Hanin M, Volrath S, Bogucki A, Briker M, Ward E, Paszkowski J. Gene targeting in *Arabidopsis*. *Plant J.* 2001;28:671–7.
14. Latt S. Sister Chromatid exchange formation. *Annu. Rev. Genet.* 1977;15:11–55.
15. Rudin N, Sugarman E, Haber JE. Genetic and physical analysis of double-strand break repair and recombination in *Saccharomyces cerevisiae*. *Genetics.* 1989;122:519–34.
16. Plessis A, Perrin A, Haber JE, Dujon B. Site-specific recombination determined by I-SceI, a mitochondrial group I intron-encoded endonuclease expressed in the yeast nucleus. *Genetics.* 1992;130:451–60.
17. Li T, Liu B, Spalding MH, Weeks DP, Yang B. High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nat. Biotechnol.* 2012;30:390–2.
18. Qi Y, Li X, Zhang Y, Starker CG, Baltes NJ, Zhang F, Targeted deletion and inversion of tandemly arrayed genes in *Arabidopsis thaliana* using zinc finger nucleases. *G3 (Bethesda).* 2013;3:1707–15.
19. Xiao A, Wang Z, Hu Y, Wu Y, Luo Z, Yang Z, et al. Chromosomal deletions and inversions mediated by TALENs and CRISPR/Cas in zebrafish. *Nucleic Acids Res.* 2013;41:1–11.
20. Zhou H, Liu B, Weeks DP, Spalding MH, Yang B. Large chromosomal deletions and heritable small genetic changes induced by CRISPR/Cas9 in rice. *Nucleic Acids Res.* 2014;42:10903–14.
21. Chapman JR, Taylor MRG, Boulton SJ. Playing the End Game: DNA Double-Strand Break Repair Pathway Choice. *Mol. Cell.* 2012;47:497–510.
22. Puchta H. The repair of double-strand breaks in plants: mechanisms and consequences for genome evolution. *J. Exp. Bot.* 2005;56:1–14.
23. Walker JR, Corpina R a, Goldberg J. Structure of the Ku heterodimer bound to DNA and its implications for double-strand break repair. *Nature.* 2001;412:607–14.
24. Yaneva M, Kowalewski T, Lieber MR. Interaction of DNA-dependent protein kinase with DNA and with Ku: Biochemical and atomic-force microscopy studies. *EMBO J.* 1997;16:5098–112.
25. Mahaney BL, Meek K, Lees-Miller SP. Repair of ionizing radiation-induced DNA double-strand breaks by non-homologous end-joining. *Biochem. J.* 2009;417:639–50.
26. Lieber MR. The mechanism of human nonhomologous DNA End joining. *J. Biol. Chem.* 2008;283:1–5.
27. West CE, Waterworth WM, Story GW, Sunderland P A, Jiang Q, Bray CM. Disruption of the *Arabidopsis AtKu80* gene demonstrates an essential role for AtKu80 protein in efficient repair of DNA double-strand breaks in vivo. *Plant J.* 2002;31:517–28.
28. West CE, Waterworth WM, Jiang Q, Bray CM. *Arabidopsis* DNA ligase IV is induced by γ -irradiation and interacts with an *Arabidopsis* homologue of the double strand break repair protein XRCC4. *Plant J.* 2000;24:67–78.

29. Bundock P, van Attikum H, Hooykaas PJJ. Increased telomere length and hypersensitivity to DNA damaging agents in an Arabidopsis KU70 mutant. *Nucleic Acids Res.* 2002;30:3395–400.
30. van Attikum H, Bundock P, Overmeer RM, Lee LY, Gelvin SB, Hooykaas PJJ. The Arabidopsis AtLIG4 gene is required for the repair of DNA damage, but not for the integration of Agrobacterium T-DNA. *Nucleic Acids Res.* 2003;31:4247–55.
31. Tamura K, Adachi Y, Chiba K, Oguchi K, Takahashi H. Identification of Ku70 and Ku80 homologues in Arabidopsis thaliana: Evidence for a role in the repair of DNA double-strand breaks. *Plant J.* 2002;29:771–81.
32. Shuman S, Glickman MS. Bacterial DNA repair by non-homologous end joining. *Nat. Rev. Microbiol.* 2007;5:852–61.
33. Pitcher RS, Brissett NC, Doherty AJ. Nonhomologous End-Joining in Bacteria: A Microbial Perspective. *Annu. Rev. Microbiol.* 2007;61:259–82.
34. Iliakis G. Backup pathways of NHEJ in cells of higher eukaryotes: Cell cycle dependence. *Radiother. Oncol.* 2009;92:310–5.
35. Mladenov E, Iliakis G. Induction and repair of DNA double strand breaks: The increasing spectrum of non-homologous end joining pathways. *Mutat. Res.* 2011;711:61–72.
36. Nussenzweig A, Nussenzweig MC. A Backup DNA Repair Pathway Moves to the Forefront. *Cell.* 2007;131:223–5.
37. Haber JE. Alternative endings. *Proc. Natl. Acad. Sci.* 2008;105:405–6.
38. Crespan E, Czabany T, Maga G, Hübscher U. Microhomology-mediated DNA strand annealing and elongation by human DNA polymerases λ and β on normal and repetitive DNA sequences. *Nucleic Acids Res.* 2012;40:5577–90.
39. Kuhfittig-Kulle S, Feldmann E, Odersky A, Kuliczowska A, Goedecke W, Eggert A, et al. The mutagenic potential of non-homologous end joining in the absence of the NHEJ core factors Ku70/80, DNA-PKcs and XRCC4-LigIV. *Mutagenesis.* 2007;22:217–33.
40. McVey M, Lee SE. MMEJ repair of double-strand breaks (director's cut): deleted sequences and alternative endings. *Trends Genet.* 2008;24:529–38.
41. Deriano L, Roth DB. Modernizing the nonhomologous end-joining repertoire: alternative and classical NHEJ share the stage. *Annu. Rev. Genet.* 2013;47:433–55.
42. Wood RD, Doublé S. DNA polymerase θ (POLQ), double-strand break repair, and cancer. *DNA Repair.* 2016;44:22–32.
43. Chang HHY, Pannunzio NR, Adachi N, Lieber MR. Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat. Rev. Mol. Cell Biol.* 2017;18:495–506
44. Pardo B, Gómez-González B, Aguilera A. DNA double-strand break repair: How to fix a broken relationship. *Cell. Mol. Life Sci.* 2009;66:1039–56.
45. Szostak JW, Orr-Weaver TL, Rothstein RJ, Stahl FW. The double-strand-break repair model for recombination. *Cell.* 1983;33:25–35.
46. San Filippo J, Sung P, Klein H. Mechanism of eukaryotic homologous recombination. *Annu. Rev. Biochem.* 2008;77:229–57.
47. Llorente B, Smith CE, Symington LS. Break-induced replication: What is it and what is it for? *Cell Cycle.* 2008;7:859–64.
48. Bray CM, West CE. DNA repair mechanisms in plants: Crucial sensors and effectors for the maintenance of genome integrity. *New Phytol.* 2005;168:511–28.
49. Puchta H. Repair of genomic double-strand breaks in somatic plant cells by one-sided invasion of homologous sequences. *Plant J.* 1998;13:331–9.
50. Puchta H, Hohn B. The mechanism of extrachromosomal homologous DNA recombination in plant cells. *Mol. Gen. Genet.* 1991;230:1–7.
51. Siebert R, Puchta H. Efficient repair of genomic double-strand breaks by homologous recombination between directly repeated sequences in the plant genome. *Plant Cell.* 2002;14:1121–31.
52. Baltes NJ, Voytas DF. Enabling plant synthetic biology through genome engineering. *Trends Biotechnol.* 2015;33:120–31.
53. Edgell DR. Selfish DNA: homing endonucleases find a home. *Curr. Biol.* 2009;19:115–7.
54. Hafez M, Hausner G, Bonen L. Homing endonucleases: DNA scissors on a mission. *Genome.* 2012;55:553–69.
55. Stoddard BL. Homing endonucleases: from microbial genetic invaders to reagents for targeted DNA modification. *Structure.* 2011;19:7–15.

56. Smith J, Grizot S, Arnould S, Duclert A, Epinat JC, Chames P, et al. A combinatorial approach to create artificial homing endonucleases cleaving chosen sequences. *Nucleic Acids Res.* 2006;34:1–12.
57. Chevalier BS, Kortemme T, Chadsey MS, Baker D, Monnat RJ, Stoddard BL. Design, activity, and structure of a highly specific artificial endonuclease. *Mol. Cell.* 2002;10:895–905.
58. Diakun G, Fairall L, Klug A. EXAFS study of the zinc-binding sites in the protein transcription factor IIIA. *Nature.* 1986;324:698–9.
59. Pavletich NP, Pabo CO. Zinc finger-DNA recognition: crystal structure of a Zif268-DNA complex at 2.1 Å. *Science.* 1991;252:809–17.
60. Segal DJ, Dreier B, Beerli RR, Barbas CF. Toward controlling gene expression at will: selection and design of zinc finger domains recognizing each of the 5'-GNN-3' DNA target sequences. *Proc. Natl. Acad. Sci. USA.* 1999;96:2758–63.
61. Dreier B, Beerli RR, Segal DJ, Flippin JD, Barbas CF. Development of zinc finger domains for recognition of the 5'-ANN-3' family of DNA sequences and their use in the construction of artificial transcription factors. *J. Biol. Chem.* 2001;276:29466–78.
62. Dreier B, Fuller RP, Segal DJ, Lund C V, Blancafort P, Huber A, et al. Development of zinc finger domains for recognition of the 5'-CNN-3' family DNA sequences and their use in the construction of artificial transcription factors. *J. Biol. Chem.* 2005;280:35588–97.
63. Beumer K, Bhattacharyya G, Bibikova M, Trautman JK, Carroll D. Efficient gene targeting in *Drosophila* with zinc-finger nucleases. *Genetics.* 2006;172:2391–403.
64. Miller JC, Holmes MC, Wang J, Guschin DY, Lee Y-L, Rupniewski I, et al. An improved zinc-finger nuclease architecture for highly specific genome editing. *Nat. Biotechnol.* 2007;25:778–85.
65. Doyon Y, Vo TD, Mendel MC, Greenberg SG, Wang J, Xia DF, et al. Enhancing zinc-finger-nuclease activity with improved obligate heterodimeric architectures. *Nat. Methods.* 2011;8:74–9.
66. Ramirez CL, Foley JE, Wright D a, Müller-Lerch F, Rahman SH, Cornu TI, et al. Unexpected failure rates for modular assembly of engineered zinc fingers. *Nat. Methods.* 2008;5:374–5.
67. Händel E-M, Alwin S, Cathomen T. Expanding or restricting the target site repertoire of zinc-finger nucleases: the inter-domain linker as a major determinant of target site selectivity. *Mol. Ther.* 2009;17:104–11.
68. Carroll D. Genome engineering with zinc-finger nucleases. *Genetics.* 2011;188:773–82.
69. Kim J-S, Lee HJ, Carroll D. Genome editing with modularly assembled zinc-finger nucleases. *Nat. Methods.* 2010;7:91–2.
70. Sander JD, Dahlborg EJ, Goodwin MJ, Cade L, Zhang F, Cifuentes D, et al. Selection-free zinc-finger-nuclease engineering by context-dependent assembly (CoDA). *Nat. Methods.* 2011;8:67–9.
71. Boller T. Innate immunity in plants : an arms race between pattern recognition receptors in plant and effectors in microbial pathogens. *Plant-Microbe Interact.* 2009;324:742–4.
72. Göhre V, Robatzek S. Breaking the barriers: microbial effector molecules subvert plant immunity. *Annu. Rev. Phytopathol.* 2008;46:189–215.
73. Kay S, Bonas U. How *Xanthomonas* type III effectors manipulate the host plant. *Curr. Opin. Microbiol.* 2009;12:37–43.
74. Boch J, Scholze H, Schornack S, Landgraf A, Hahn S, Kay S, et al. Breaking the code of DNA binding specificity of TAL-type III effectors. *Science.* 2009;326:1509–12.
75. Moscou M, Bogdanove A. Recognition by TAL Effectors. *Science.* 2009;326:1501.
76. Deng D, Yan C, Pan X, Mahfouz M, Wang J, Zhu J, et al. Structural basis for sequence-specific recognition of DNA by TAL effectors. *Science.* 2012;335:720–3.
77. Mak AN-S, Bradley P, Cernadas RA, Bodanove AJ, Stoddard BL. The crystal structure of TAL effector PthXo1 bound to its DNA target. *Science.* 2012;335:716–9.
78. Christian M, Cermak T, Doyle EL, Schmidt C, Zhang F, Hummel A, et al. Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics.* 2010;186:757–61.
79. Wiedenheft B, Sternberg SH, Doudna JA. RNA-guided genetic silencing systems in bacteria and archaea. *Nature.* 2012;482:331–8.
80. Bhaya D, Davison M, Barrangou R. CRISPR-Cas systems in bacteria and archaea: versatile small RNAs for adaptive defense and regulation. *Annu. Rev. Genet.* 2011;45:273–97.
81. Terns MP, Terns RM. CRISPR-based adaptive immune systems. *Curr. Opin. Microbiol.* 2011;14:321–7.
82. Deltcheva E, Chylinski K, Sharma CM, Gonzales K. CRISPR RNA maturation by trans -encoded small RNA and host factor RNase III. *Nature.* 2011;471:602–7.

83. Deltcheva E, Chylinski K, Sharma CM, Gonzales K, Chao Y, Pirzada ZA, et al. CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III. *Nature*. 2011;471:602–7.
84. Brouns SJJ, Jore MM, Lundgren M, Westra ER, Slijkhuys RJH, Snijders APL, et al. Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science*. 2008;321:960–4.
85. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*. 2012;337:816–22.
86. Gasiunas G, Barrangou R, Horvath P, Siksnys V. Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc. Natl. Acad. Sci. USA*. 2012;109:E2579–
87. Hsu PD, Scott D a, Weinstein J a, Ran FA, Konermann S, Agarwala V, et al. DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat. Biotechnol.* 2013;31:827–32.
88. Cho SW, Lee J, Carroll D, Kim JS, Lee J. Heritable gene knockout in *Caenorhabditis elegans* by direct injection of Cas9-sgRNA ribonucleoproteins. *Genetics*. 2013;195:1177–80.
89. Cong L, Ran FA, Cox D, Lin S, Barretto R, Hsu PD, et al. Multiplex genome engineering using CRISPR/Cas systems. *Science*. 2013;339:819–23.
90. Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, et al. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Science. Nat. Biotechnol.* 2013;31:227–9.
91. Jinek M, East A, Cheng A, Lin S, Ma E, Doudna J. RNA-programmed genome editing in human cells. *Elife*. 2013;2013:1–9.
92. Dicarlo JE, Norville JE, Mali P, Rios X, Aach J, Church GM. Genome engineering in *Saccharomyces cerevisiae* using CRISPR-Cas systems. *Nucleic Acids Res.* 2013;41:4336–43.
93. Sapranaukas R, Gasiunas G, Fremaux C, Barrangou R, Horvath P, Siksnys V. The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*. *Nucleic Acids Res.* 2011;39:9275–82.
94. Sander JD, Joung JK. CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat. Biotechnol.* 2014;32:347–55.
95. Wolt JD, Wang K, Sashital D, Lawrence-Dill CJ. Achieving Plant CRISPR Targeting that Limits Off-Target Effects. *Plant Genome*. 2016;9:0.
96. Fan D, Liu T, Li C, Jiao B, Li S, Hou Y, et al. Efficient CRISPR/Cas9-mediated targeted mutagenesis in populus in the first generation. *Sci. Rep.* 2015;5:12217.
97. Svitashov S, Young JK, Schwartz C, Gao H, Falco SC, Cigan AM. Targeted mutagenesis, precise gene editing, and site-specific gene insertion in maize using Cas9 and guide RNA. *Plant Physiol.* 2015;169:931–45.
98. Xu R, Li H, Qin R, Li J, Qiu C, Yang Y, et al. Generation of inheritable and “ transgene clean ” targeted genome-modified rice in later generations using the CRISPR / Cas9 system. *Sci. Rep.* 2015;9:1–10.
99. Zhang H, Zhang J, Wei P, Zhang B, Gou F, Feng Z, et al. The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol. J.* 2014;12:797–807.
100. Jiang W, Yang B, Weeks DP. Efficient CRISPR/Cas9-mediated gene editing in *Arabidopsis thaliana* and inheritance of modified genes in the T2 and T3 generations. *PLoS One*. 2014;9:21–6.
101. Wang Z, Xing H, Dong L, Zhang H, Han C, Wang X, et al. Egg cell-specific promoter-controlled CRISPR / Cas9 efficiently generates homozygous mutants for multiple target genes in *Arabidopsis* in a single generation. *Genome Biol. Genome Biology*; 2015;16:1–12.
102. Lee HJ, Kweon J, Kim E, Kim S, Kim J. Targeted chromosomal duplications and inversions in the human genome using zinc finger nucleases. *Genome Res.* 2012;539–48.
103. Brunet E, Simsek D, Tomishima M, DeKelver R, Choi VM, Gregory P, et al. Chromosomal translocations induced at specified loci in human stem cells. *Proc. Natl. Acad. Sci. USA*. 2009;106:10620–5.
104. Lowder LG, Zhang D, Baltes NJ, Paul JW, Tang X, Zheng X, et al. A CRISPR/Cas9 toolbox for multiplexed plant genome editing and transcriptional regulation. *Plant Physiol.* 2015;169:971–85.
105. Mercx S, Tollet J, Magy B, Navarre C, Boutry M. Gene Inactivation by CRISPR-Cas9 in *Nicotiana tabacum* BY-2 suspension cells. *Front. Plant Sci.* 2016;7:40.
106. Zhang Z, Mao Y, Ha S, Liu W, Botella JR, Zhu JK. A multiplex CRISPR/Cas9 platform for fast and efficient editing of multiple genes in *Arabidopsis*. *Plant Cell Rep.* 2015;1–15.
107. Puchta H, Dujon B, Hohn B. Homologous recombination in plant cells is enhanced by in vivo induction of double strand breaks into DNA by a site-specific endonuclease. *Nucleic Acids Res.* 1993;21:5034–40.
108. Puchta H, Dujon B, Hohn B. Two different but related mechanisms are used in plants for the repair of genomic double-strand breaks by homologous recombination. *Proc. Natl. Acad. Sci. USA*. 1996;93:5055–60.

109. de Pater S, Neuteboom LW, Pinas JE, Hooykaas PJJ, van der Zaal BJ. ZFN-induced mutagenesis and gene-targeting in Arabidopsis through Agrobacterium-mediated floral dip transformation. *Plant Biotechnol. J.* 2009;7:821–35.
110. Pater S De, Pinas JE, Hooykaas PJJ, Zaal BJ Van Der. ZFN-mediated gene targeting of the Arabidopsis protoporphyrinogen oxidase gene through Agrobacterium-mediated floral dip transformation. *Plant Cell Physiol.* 2013;11:510–5.
111. Townsend J a, Wright D a, Winfrey RJ, Fu F, Maeder ML, Joung JK, et al. High-frequency modification of plant genes using engineered zinc-finger nucleases. *Nature.* 2009;459:442–5.
112. Shukla VK, Doyon Y, Miller JC, DeKelver RC, Moehle E a, Worden SE, et al. Precise genome modification in the crop species *Zea mays* using zinc-finger nucleases. *Nature.* 2009;459:437–41.
113. Zhang Y, Zhang F, Li X, Baller J a., Qi Y, Starker CG, et al. Transcription activator-like effector nucleases enable efficient plant genome engineering. *Plant Physiol.* 2013;161:20–7.
114. Čermák T, Baltés NJ, Čegan R, Zhang Y, Voytas DF. High-frequency, precise modification of the tomato genome. *Genome Biol.* 2015;16:232.
115. Li W, Teng F, Li T, Zhou Q. Simultaneous generation and germline transmission of multiple gene mutations in rat using CRISPR-Cas systems. *Nat. Biotechnol.* 2013;31:684–6.
116. Fauser F, Schiml S, Puchta H. Both CRISPR/Cas-based nucleases and nickases can be used efficiently for genome engineering in *Arabidopsis thaliana*. *Plant J.* 2014;79:348–59.
117. Schiml S, Fauser F, Puchta H. The CRISPR/Cas system can be used as nuclease for in planta gene targeting and as paired nickases for directed mutagenesis in *Arabidopsis* resulting in heritable progeny. *Plant J.* 2014;80:1139–50.
118. Sun Y, Zhang X, Wu C, He Y, Ma Y, Hou H, et al. Engineering Herbicide Resistant Rice Plants through CRISPR/Cas9-mediated Homologous Recombination of the Acetolactate Synthase. *Mol. Plant.* 2016;1–4.
119. Butler NM, Atkins PA, Voytas DF, Douches DS. Generation and inheritance of targeted mutations in potato (*Solanum tuberosum* L.) Using the CRISPR/Cas System. *PLoS One.* 2015;10:1–12.
120. Shan Q, Wang Y, Li J, Zhang Y, Chen K, Liang Z, et al. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* 2013;31:686–8.
121. Niu Y, Shen B, Cui Y, Chen Y, Wang J, Wang L, et al. Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated gene targeting in one-cell embryos. *Cell.* 2014;156:836–43.
122. Meng X, Noyes MB, Zhu LJ, Lawson ND, Wolfe SA. Targeted gene inactivation in zebrafish using engineered zinc-finger nucleases. *Nat. Biotechnol.* 2008;26:695–701.
123. Lee J-S, Kwak S-J, Kim J, Noh HM, Kim J-S, Yu K. RNA-guided genome editing in *Drosophila* with the purified Cas9 protein. *G3 (Bethesda).* 2014;4:1291–5.
124. Zuris JA, Thompson DB, Shu Y, Guilinger JP, Bessen JL, Hu JH, et al. Efficient delivery of genome-editing proteins in vitro and in vivo. *Nat. Biotechnol.* 2015;33:73–80.
125. Luo S, Li J, Stoddard TJ, Baltés NJ, Demorest ZL, Clasen BM, et al. Non-transgenic plant genome editing using purified sequence-specific nucleases. *Mol. Plant.* ; 2015;8:1425–7.
126. Woo JW, Kim J, Kwon S II, Corvalán C, Cho SW, Kim H, et al. DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol.* 2015;33:1162–4.
127. Braun AC. A physiological basis for autonomous growth of the crown-gall tumor cell. *Proc. Natl. Acad. Sci. USA.* 1958;44:344–9.
128. Zambryski P, Tempe J, Schell J. Transfer and function of T-DNA genes from *Agrobacterium Ti* and *Ri* plasmids in plants. *Cell.* 1989;56:193–201.
129. Chilton M-D, Drummond MH, Merlo DJ, Sciacchi D, Montoya AL, Gordon MP, et al. Stable incorporation of plasmid DNA into higher plant cells: the molecular basis of crown gall tumorigenesis. *Cell.* 1977;11:263–71.
130. Britton MT, Escobar MA, Dandekar AM. The oncogenes of *Agrobacterium tumefaciens* and *Agrobacterium rhizogenes*. In: Tzfira T, Citovsky V, editors. *Agrobacterium From Biol. to Biotechnol.* New York, NY: Springer New York; 2008. p. 523–63.
131. Dessaux Y, Petit A, Farrand SK, Murphy PJ. Opines and Opine-Like Molecules Involved in Plant-Rhizobiaceae Interactions. In: Spaink HP, Kondoroski A, Hooykaas PJJ, editors. *Rhizobiaceae Mol. Biol. Model Plant-Associated Bact.* Dordrecht: Springer Netherlands; 1998. p. 173–97.
132. Stachel SE, Messens E, Van Montagu M, Zambryski P. Identification of the signal molecules produced by wounded plant cells that activate T-DNA transfer in *Agrobacterium tumefaciens*. *Nature.* 1985;318:624–9.

133. Melchers LS, Regensburg-Tuïnk TJ, Bourret RB, Sedee NJ, Schilperoort RA, Hooykaas PJJ. Membrane topology and functional analysis of the sensory protein VirA of *Agrobacterium tumefaciens*. *EMBO J.* 1989;1919–25.
134. Leroux B, Yanofsky MF, Winans SC, Ward JE, Ziegler SF, Nester EW. Characterization of the virA locus of *Agrobacterium tumefaciens*: a transcriptional regulator and host range determinant. *EMBO J.* 1987;6:849–56.
135. Melchers LS, Thompson D V., Idler KB, Neuteboom STC, de Maagd RA., Schilperoort RA, et al. Molecular characterization of the virulence gene virA of the *Agrobacterium tumefaciens* octopine Ti plasmid. *Plant Mol. Biol.* 1988;11:227–37.
136. Lee YW, Jin S, Sim WS, Nester EW. The sensing of plant signal molecules by *Agrobacterium*: genetic evidence for direct recognition of phenolic inducers by the VirA protein. *Gene.* 1996;179:83–8.
137. Cangelosi G a, Ankenbauer RG, Nester EW. Sugars induce the *Agrobacterium* virulence genes through a periplasmic binding protein and a transmembrane signal protein. *Proc. Natl. Acad. Sci. USA.* 1990;87:6708–12.
138. Stachel SE, Nester EW. The genetic and transcriptional organization of the vir region of the A6 Ti plasmid of *Agrobacterium tumefaciens*. *EMBO J.* 1986;5:1445–54.
139. Hooykaas P, Hofker M, den Dulk-Ras A, Schilperoort RA. A comparison of virulence determinants in an octopine Ti plasmid, a nopaline Ti plasmid, and an Ri plasmid by complementation analysis of *Agrobacterium tumefaciens* mutants. *Plasmid.* 1984;11:195–205.
140. Kanemoto RH, Powell a. T, Akiyoshi DE, Regier D a., Kerstetter R a., Nester EW, et al. Nucleotide sequence and analysis of the plant-inducible locus pinF from *Agrobacterium tumefaciens*. *J. Bacteriol.* 1989;171:2506–12.
141. van Haaren MJJ, Pronk JT, Schilperoort RA, Hooykaas PJJ. Functional analysis of the *Agrobacterium tumefaciens* octopine Ti-plasmid left and right T-region border fragments. *Plant Mol. Biol.* 1987;8:95–104.
142. Ward ER, Barnes WM. VirD2 protein of *Agrobacterium tumefaciens* very tightly linked to the 5' end of T-strand DNA. *Science.* 1988;242:927–30.
143. Pansegrau W, Schoumacher F, Hohn B, Lanka E. Site-specific cleavage and joining of single-stranded DNA by VirD2 protein of *Agrobacterium tumefaciens* Ti plasmids: analogy to bacterial conjugation. *Proc. Natl. Acad. Sci. USA.* 1993;90:11538–42.
144. Toro N, Datta A, Carmi OA, Young C, Prusti RK, Nester EW. The *Agrobacterium tumefaciens* virC1 gene product binds to overdrive, a T-DNA transfer enhancer. *J. Bacteriol.* 1989;171:6845–9.
145. Lu J, den Dulk-Ras A, Hooykaas PJJ, Glover JNM, Dulk-ras A Den, Hooykaas PJJ, et al. *Agrobacterium tumefaciens* VirC2 enhances T-DNA transfer and virulence through its C-terminal ribbon-helix-helix DNA-binding fold. *Proc. Natl. Acad. Sci. USA.* 2009;106:9643–8.
146. van Kregten M, Lindhout BI, Hooykaas PJJ, van der Zaal BJ. *Agrobacterium*-mediated T-DNA transfer and integration by minimal VirD2 consisting of the relaxase domain and a type IV secretion system translocation signal. *Mol. Plant. Microbe. Interact.* 2009;22:1356–65.
147. Christie PJ, Whitaker N, González-Rivera C. Mechanism and structure of the bacterial type IV secretion systems. *Biochim. Biophys. Acta.* 2014;1843:1578–91.
148. Vergunst a. C. VirB/D4-dependent protein translocation from *Agrobacterium* into plant cells. *Science.* 2000;290:979–82.
149. Vergunst AC, van Lier MCM, den Dulk-Ras A, Stüve T a G, Ouwehand A, Hooykaas PJJ. Positive charge is an important feature of the C-terminal transport signal of the VirB/D4-translocated proteins of *Agrobacterium*. *Proc. Natl. Acad. Sci. USA.* 2005;102:832–7.
150. Schrammeijer B. Analysis of Vir protein translocation from *Agrobacterium tumefaciens* using *Saccharomyces cerevisiae* as a model: evidence for transport of a novel effector protein VirE3. *Nucleic Acids Res.* 2003;31:860–8.
151. Zupan JR, Citovsky V, Zambryski P. *Agrobacterium* VirE2 protein mediates nuclear uptake of single-stranded DNA in plant cells. *Proc. Natl. Acad. Sci. USA.* 1996;93:2392–7.
152. Ward D V., Zupan JR, Zambryski PC. *Agrobacterium* VirE2 gets the VIP1 treatment in plant nuclear import. *Trends Plant Sci.* 2002;7:1–3.
153. Lacroix B, Vaidya M, Tzfira T, Citovsky V. The VirE3 protein of *Agrobacterium* mimics a host cell function required for plant genetic transformation. *Embo J.* 2005;24:428–37.
154. García-Rodríguez FM, Schrammeijer B, Hooykaas PJJ, Garcí FM. The *Agrobacterium* VirE3 effector protein: a potential plant transcriptional activator. *Nucleic Acids Res.* 2006;34:6496–504.
155. Niu X, Zhou M, Henkel C V., Van Heusden GPH, Hooykaas PJJ. The *Agrobacterium tumefaciens* virulence protein VirE3 is a transcriptional activator of the F-box gene VBF. *Plant J.* 2015;84:914–24.
156. Zhang X. Functional analysis of *Agrobacterium tumefaciens* virulence protein VirD5. Leiden University; 2016.

157. Schrammeijer B, Risseew E, Pansegrau W, Regensburg-Tuink TJG, Crosby WL, Hooykaas PJJ. Interaction of the virulence protein VirF of *Agrobacterium tumefaciens* with plant homologs of the yeast Skp1 protein. *Curr. Biol.* 2001;11:258–62.
158. Mycobacterium PET, Biochemistry TD, Tzfira T, Vaidya M, Citovsky V. Involvement of targeted proteolysis in plant genetic transformation by *Agrobacterium*. *Nature.* 2004;431:6–11.
159. Garcia-Cano E, Magori S, Sun Q, Ding Z, Lazarowitz SG, Citovsky V. Interaction of arabidopsis trihelix-domain transcription factors VFP3 and VFP5 with *Agrobacterium* virulence protein VirF. *PLoS One.* 2015;10:1–23.
160. Zaltsman A, Krichevsky A, Kozlovsky S V, Yasmin F, Citovsky V. Plant defense pathways subverted by *Agrobacterium* for genetic transformation. *Plant Signal. Behav.* 2010;5:1245–8.
161. Cascales E, Christie PJ. Definition of a bacterial type IV secretion pathway for a DNA substrate. *Science.* 2004;304:1170–3.
162. Jakubowski SJ, Krishnamoorthy V, Cascales E, Christie PJ. *Agrobacterium tumefaciens* VirB6 Domains Direct the Ordered Export of a DNA Substrate Through a Type IV Secretion System. *Journal Mol. Biol.* 2004;341:961–77.
163. Atmakuri K, Ding Z, Christie PJ. VirE2, a Type IV secretion substrate, interacts with the VirD4 transfer protein at cell poles of *Agrobacterium tumefaciens*. *Changes.* 2003;49:1699–713.
164. Abu-Arish A, Frenkiel-Krispin D, Fricke T, Tzfira T, Citovsky V, Wolf SG, et al. Three-dimensional reconstruction of *Agrobacterium* VirE2 protein with single-stranded DNA. *J. Biol. Chem.* 2004;279:25359–63.
165. Gelvin SB. *Agrobacterium* and plant genes involved in T-DNA transfer and integration. *Annu. Rev. Plant Biol.* 2000; 51:233–256
166. Bundock P, den Dulk-Ras A, Beijersbergen A, Hooykaas PJJ. Trans-kingdom T-DNA transfer from *Agrobacterium tumefaciens* to *Saccharomyces cerevisiae*. *EMBO J.* 1995;14:3206–14.
167. van Attikum H, Bundock P, Hooykaas PJJ. Non-homologous end-joining proteins are required for *Agrobacterium* T-DNA integration. *EMBO J.* 2001;20:6550–8.
168. van Attikum H, Hooykaas PJJ. Genetic requirements for the targeted integration of *Agrobacterium* T-DNA in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 2003;31:826–32.
169. Friesner J, Britt AB. Ku80- and DNA ligase IV-deficient plants are sensitive to ionizing radiation and defective in T-DNA integration. *Plant J.* 2003;34:427–40.
170. Gallego ME, Bleuyard JY, Daoudal-Cotterell S, Jallut N, White CI. Ku80 plays a role in non-homologous recombination but is not required for T-DNA integration in *Arabidopsis*. *Plant J.* 2003;35:557–65.
171. Li JX, Vaidya M, White C, Vainstein A, Citovsky V, Tzfira T. Involvement of KU80 in T-DNA integration in plant cells. *Proc. Natl. Acad. Sci. USA.* 2005;102:19231–6.
172. Jia Q, Bundock P, Hooykaas PJJ, de Pater S. *Agrobacterium tumefaciens* T-DNA Integration and Gene Targeting in *Arabidopsis thaliana* Non-Homologous End-Joining Mutants. *J. Bot.* 2012;2012:1–13.
173. Mestiri I, Norre F, Gallego ME, White CI. Multiple host-cell recombination pathways act in *Agrobacterium*-mediated transformation of plant cells. *Plant J.* 2014;77:511–20.
174. Park SY, Vaghchhipawala Z, Vasudevan B, Lee LY, Shen Y, Singer K, et al. *Agrobacterium* T-DNA integration into the plant genome can occur without the activity of key non-homologous end-joining proteins. *Plant J.* 2015;81:934–46.
175. van Kregten M, de Pater S, Romeijn R, van Schendel R, Hooykaas PJJ, Tijsterman M. T-DNA integration in plants results from polymerase- θ -mediated DNA repair. *Nat. plants.* 2016;2:16164.
176. Hoekema A, Hirsch PR, Hooykaas PJJ, Schilperoort RA. A binary plant vector strategy based on separation of vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. *Nature.* 1983;302:179–80.
177. Krenek P, Samajova O, Luptovciak I, Doskocilova A, Komis G, Samaj J. Transient plant transformation mediated by *Agrobacterium tumefaciens*: Principles, methods and applications. *Biotechnol. Adv.* 2015;33:1024–42.
178. Koncz C, Németh K, Rédei GP, Schell J. T-DNA insertional mutagenesis in *Arabidopsis*. *Plant Mol. Biol.* 1992;20:963–76.
179. Walden R, Hayashi H, Schell J. T-DNA as a gene tag. *Plant J.* 1991;1:281–8.
180. Slogteren GH-V, Hooykaas P. Expression of Ti plasmid genes in monocotyledonous plants infected with *Agrobacterium tumefaciens*. *Nature.* 1984;311.
181. Michielse CB, Hooykaas PJJ, van den Hondel C a MJJ, Ram AFJ. *Agrobacterium*-mediated transformation as a tool for functional genomics in fungi. *Curr. Genet.* 2005;48:1–17.
182. Rolloos M, Hooykaas PJJ, van der Zaal BJ. Enhanced targeted integration mediated by translocated I-SceI during the *Agrobacterium* mediated transformation of yeast. *Sci. Rep.* 2015;5:8345.

183. Kregten van M, Boer de P, Pinas JE, Hooykaas PJJ, van der Zaal BJ. Agrobacterium-mediated delivery of a meganuclease into target plant cells. 2011.
184. Kregten van M, Boer de P, Hooykaas PJJ, van der Zaal BJ. Translocation of novel recombinant effector proteins from *Agrobacterium tumefaciens* to *Arabidopsis thaliana*. 2011.
185. Antunes MS, Smith JJ, Jantz D, Medford JI. Targeted DNA excision in *Arabidopsis* by a re-engineered homing endonuclease. *BMC Biotechnol.* 2012;12:86.
186. Yang M, Djukanovic V, Stagg J, Lenderts B, Bidney D, Carl Falco S, et al. Targeted mutagenesis in the progeny of maize transgenic plants. *Plant Mol. Biol.* 2009;70:669–79.
187. Gao H, Smith J, Yang M, Jones S, Djukanovic V, Nicholson MG, et al. Heritable targeted mutagenesis in maize using a designed endonuclease. *Plant J.* 2010;61:176–87.
188. Lloyd A, Plaisier CL, Carroll D, Drews GN. Targeted mutagenesis in *Arabidopsis* using zinc-finger nucleases. *Methods Mol. Biol.* 2005;102:2232–7.
189. Zhang F, Maeder ML, Unger-wallace E, Hoshaw JP, Reyon D, Christian M, et al. High frequency targeted mutagenesis in *Arabidopsis thaliana* using zinc finger nucleases. *Proc. Natl. Acad. Sci.* 2010;107:12028–33.
190. Osakabe K, Osakabe Y, Toki S. Site-directed mutagenesis in *Arabidopsis* using custom-designed zinc finger nucleases. *Proc. Natl. Acad. Sci.* 2010;107:12034–9.
191. Curtin SJ, Zhang F, Sander JD, Haun WJ, Starker C, Baltes NJ, et al. Targeted mutagenesis of duplicated genes in soybean with zinc-finger nucleases. *Plant Physiol.* 2011;156:466–73.
192. Christian M, Qi Y, Zhang Y, Voytas DF. Targeted mutagenesis of *Arabidopsis thaliana* using engineered TAL effector nucleases. *G3 (Bethesda).* 2013;3:1697–705.
193. Forner J, Pfeiffer A, Langenecker T, Manavella P. Germline-Transmitted Genome Editing in *Arabidopsis thaliana* Using TAL-Effector- Nucleases. *PLoS One.* 2015;1–15.
194. Cermak T, Doyle EL, Christian M, Wang L, Zhang Y, Schmidt C, et al. Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting. *Nucleic Acids Res.* 2011;39:e82.
195. Zhang Y, Shan Q, Wang Y, Chen K, Liang Z, Li J, et al. Rapid and efficient gene modification in rice and brachypodium using TALENs. *Mol. Plant.* 2013;6:1365–8.
196. Du H, Zeng X, Zhao M, Cui X, Wang Q, Yang H, et al. Efficient targeted mutagenesis in soybean by TALENs and CRISPR/Cas9. *J. Biotechnol.* 2016;217:90–7.
197. Haun W, Coffman A, Clasen BM, Demorest ZL, Lowy A, Ray E, et al. Improved soybean oil quality by targeted mutagenesis of the fatty acid desaturase 2 gene family. *Plant Biotechnol. J.* 2014;2023:934–40.
198. Budhagatapalli N, Rutten T, Gurushidze M, Kumlehn J, Hensel G. Targeted modification of gene function exploiting homology-directed repair of TALEN-mediated double strand breaks in barley. *G3 (Bethesda).* 2015;5:1857–63.
199. Wendt T, Holm PB, Starker CG, Christian M, Voytas DF, Brinch-Pedersen H, et al. TAL effector nucleases induce mutations at a pre-selected location in the genome of primary barley transformants. *Plant Mol. Biol.* 2013;83:279–85.
200. Gurushidze M, Hensel G, Hiekel S, Schedel S, Valkov V, Kumlehn J. True-Breeding Targeted Gene Knock-Out in Barley Using Designer TALE-Nuclease in Haploid Cells. *PLoS One.* 2014;9:1–9.
201. Stoddard TJ, Clasen BM, Baltes B, Demorest ZL, Voytas DF, Zhang F, et al. Targeted Mutagenesis in Plant Cells through Transformation of Sequence-Specific Nuclease mRNA. 2016
202. Li J, Stoddard TJ, Demorest ZL, Lavoie P-O, Luo S, Clasen BM, et al. Multiplexed, targeted gene editing in *Nicotiana benthamiana* for glyco-engineering and monoclonal antibody production. *Plant Biotechnol. J.* 2015
203. Zhang H, Gou F, Zhang J, Liu W, Li Q, Mao Y, et al. TALEN-mediated targeted mutagenesis produces a large variety of heritable mutations in rice. *Plant Biotechnol. J.* 2015;1–9.
204. Shan Q, Zhang Y, Chen K, Zhang K, Gao C. Creation of fragrant rice by targeted knockout of the *OsBADH2* gene using TALEN technology. *Plant Biotechnol. J.* 2015;2:791–800.
205. Li Ting, Liu Bo, Chen Chih Ying YB. TALEN-mediated homologous recombination produces site-directed DNA base change and herbicide-resistant rice. *J. Genet. Genomics.* 2016;1–9.
206. Chen K, Shan Q, Gao C. An efficient TALEN mutagenesis system in rice. *Methods.* 2014;69:2–8.
207. Nicolia A, Proux-wéra E, Åhman I, Onkokesung N, Andersson M, Andreasson E, et al. Targeted gene mutation in tetraploid potato through transient TALEN expression in protoplasts. *J. Biotechnol.* 2015;204:17–24.
208. Wang Y, Cheng X, Shan Q, Zhang Y, Liu J, Gao C, et al. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* 2014; 2014;32:947–951.

209. Char SN, Unger-wallace E, Frame B, Briggs SA, Main M, Spalding MH, et al. Heritable site-specific mutagenesis using TALENs in maize. 2015;1002–10.
210. Feng Z, Zhang B, Ding W, Liu X, Yang, Dong-Lei, Wei P, Cao F, et al. Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res.* 2013;23:1229–32.
211. Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, et al. A robust CRISPR/Cas9 system for convenient, high-efficiency multiplex genome editing in monocot and dicot plants. *Mol. Plant.* ; 2015;8:1274–84.
212. Mao Y, Zhang H, Xu N, Zhang B, Gou F, Zhu JK. Application of the CRISPR-Cas system for efficient genome engineering in plants. *Mol. Plant.* 2013;6:2008–11.
213. Mao Y, Zhang Z, Feng Z, Wei P, Zhang H, Botella JR, et al. Development of germ-line-specific CRISPR-Cas9 systems to improve the production of heritable gene modifications in Arabidopsis. *Plant Biotechnol. J.* 2015;519–32.
214. Ryder P, McHale M, Fort A, Spillane C. Generation of stable nulliplex autopolyploid lines of Arabidopsis thaliana using CRISPR/Cas9 genome editing. *Plant Cell Rep.* 2017;36:1005–8.
215. Shi Y, Huang J, Sun T, Wang X, Zhu C, Ai Y, et al. The precise regulation of different COR genes by individual CBF transcription factors in Arabidopsis thaliana. *J. Integr. Plant Biol.* 2017;59:118–33.
216. Tsutsui H, Higashiyama T. PKAMA-ITACHI vectors for highly efficient CRISPR/Cas9-mediated gene knockout in Arabidopsis thaliana. *Plant Cell Physiol.* 2017;58:46–56.
217. Xing H, Dong L, Wang Z, Zhang H, Han C, Liu B, et al. A CRISPR / Cas9 toolkit for multiplex genome editing in plants. *BMC Plant Biol.* 2014;14:1–12.
218. Yan L, Wei S, Wu Y, Hu R, Li H, Yang W, et al. High-efficiency genome editing in Arabidopsis using YAO promoter-driven CRISPR/Cas9 system. *Mol. Plant.* 2015;8:1820–3.
219. Hyun Y, Kim J, Woo S. Site-directed mutagenesis in Arabidopsis thaliana using dividing tissue-targeted RGEN of the CRISPR / Cas system to generate heritable null alleles. *Planta.* 2015;241:271–84.
220. Schiml S, Fauser F, Puchta H. The CRISPR/Cas system can be used as nuclease for in planta gene targeting and as paired nickases for directed mutagenesis in Arabidopsis resulting in heritable progeny. *Plant J.* 2014;80:1139–50.
221. Steiert J, Schiml S, Fauser F, Puchta H. Highly efficient heritable plant genome engineering using Cas9 orthologues from *Streptococcus thermophilus* and *Staphylococcus aureus*. *Plant J.* 2015;84:1295–305.
222. Ordon J, Gantner J, Kemna J, Schwalgun L, Reschke M, Streubel J, et al. Generation of chromosomal deletions in dicotyledonous plants employing a user-friendly genome editing toolkit. *Plant J.* 2017;89:155–68.
223. Jiang W, Zhou H, Bi H, Fromm M, Yang B, Weeks DP. Demonstration of CRISPR/Cas9/sgRNA-mediated targeted gene modification in Arabidopsis, tobacco, sorghum and rice. *Nucleic Acids Res.* 2013;41:1–12.
224. Li JF, Zhang D, Sheen J. Cas9-based genome editing in arabidopsis and tobacco. *Methods Enzymol.* 1st ed. 2014;546:459–72.
225. Jiang WZ, Henry IM, Lynagh PG, Comai L, Cahoon EB, Weeks DP. Significant enhancement of fatty acid composition in seeds of the allohexaploid, *Camelina sativa*, using CRISPR/Cas9 gene editing. *Plant Biotechnol. J.* 2017;15:648–57.
226. Tian S, Jiang L, Gao Q, Zhang J, Zong M, Zhang H, et al. Efficient CRISPR/Cas9-based gene knockout in watermelon. *Plant Cell Rep.* 2017;36:399–406.
227. Jia H, Wang N. Targeted genome editing of sweet orange using Cas9/sgRNA. *PLoS One.* 2014;9:e93806.
228. Jia H, Zhang Y, Orbović V, Xu J, White FF, Jones JB, et al. Genome editing of the disease susceptibility gene CsLOB1 in citrus confers resistance to citrus canker. *Plant Biotechnol. J.* 2017;15:817–23.
229. Chandrasekaran J, Brumin M, Wolf D, Leibman D, Klap C, Pearlsman M, et al. Development of broad virus resistance in non-transgenic cucumber using CRISPR/Cas9 technology. *Mol. Plant Pathol.* 2016;1–14.
230. Kishi-Kaboshi M, Aida R, Sasaki K. Generation of gene-edited chrysanthemum morifolium using multicopy transgenes as targets and markers. *Plant Cell Physiol.* 2017;58:216–26.
231. Kui L, Chen H, Zhang W, He S, Xiong Z, Zhang Y, et al. Building a Genetic Manipulation Tool Box for Orchid Biology: Identification of Constitutive Promoters and Application of CRISPR/Cas9 in the Orchid, *Dendrobium officinale*. *Front. Plant Sci.* 2017;7:1–13.
232. Cai Y, Chen L, Liu X, Sun S, Wu C, Jiang B, et al. CRISPR/Cas9-mediated genome editing in soybean hairy roots. *PLoS One.* 2015;10:1–13.
233. Cai Y, Chen L, Liu X, Guo C, Sun S, Wu C, et al. CRISPR/Cas9-mediated targeted mutagenesis of GmFT2a delays flowering time in soya bean. *Plant Biotechnol. J.* 2017;1–10.
234. Jacobs TB, Lafayette PR, Schmitz RJ, Parrott WA. Targeted genome modifications in soybean with CRISPR/Cas9. *BMC Biotechnol.* 2015;15:1–10.

235. Kim H, Kim S-T, Ryu J, Kang B-C, Kim J-S, Kim S-G. CRISPR/Cpf1-mediated DNA-free plant genome editing. *Nat. Commun.* 2017;8:14406.
236. Li Z, Liu Z, Xing A, Moon BP, Koellhoffer JP, Huang L, et al. Cas9-guide RNA directed genome editing in soybean. *Plant Physiol.* 2015;169:960–70.
237. Michno J-M, Wang X, Liu J, Curtin SJ, Kono TJY, Stupar RM. CRISPR/Cas mutagenesis of soybean and *Medicago truncatula* using a new web-tool and a modified Cas9 enzyme. *GM Crops Food.* 2015;6:243–52.
238. Chen X, Lu X, Shu N, Wang S, Wang J, Wang D, et al. Targeted mutagenesis in cotton (*Gossypium hirsutum* L.) using the CRISPR/Cas9 system. *Sci. Rep.* 2017;7:44304.
239. Janga MR, Campbell LM, Rathore KS. CRISPR/Cas9-mediated targeted mutagenesis in upland cotton (*Gossypium hirsutum* L.). *Plant Mol. Biol.* 2017;94:349–360.
240. Wang P, Zhang J, Sun L, Ma Y, Xu J, Liang S, et al. High efficient multisites genome editing in allotetraploid cotton (*Gossypium hirsutum*) using CRISPR/Cas9 system. *Plant Biotechnol. J.* 2017;1–14.
241. Kapusi E, Corcuera-Gómez M, Melnik S, Stoger E. Heritable Genomic Fragment Deletions and Small Indels in the Putative ENCase Gene Induced by CRISPR/Cas9 in Barley. *Front. Plant Sci.* 2017;8:540.
242. Meng Y, Hou Y, Wang H, Ji R, Liu B, Wen J, et al. Targeted mutagenesis by CRISPR/Cas9 system in the model legume *Medicago truncatula*. *Plant Cell Rep.* 2017;36:371–4.
243. Sugano SS, Shirakawa M, Takagi J, Matsuda Y, Shimada T, Hara-Nishimura I, et al. CRISPR/Cas9-mediated targeted mutagenesis in the liverwort *Marchantia polymorpha* L. *Plant Cell Physiol.* 2014;55:475–81.
244. Yin K, Han T, Liu G, Chen T, Wang Y, Yu AYL, et al. A geminivirus-based guide RNA delivery system for CRISPR/Cas9 mediated plant genome editing. *Sci. Rep.* 2015;5:14926.
245. Ali Z, Abul-Faraj A, Li L, Ghosh N, Piatek M, Mahjoub A, et al. Efficient virus-mediated genome editing in plants using the CRISPR/Cas9 system. *Mol. Plant.* 2015;8:1288–91.
246. Ali Z, Abulfaraj A, Idris A, Ali S, Tashkandi M, Mahfouz MM. CRISPR/Cas9-mediated viral interference in plants. *Genome Biol. Genome Biology;* 2015;16:238.
247. Ludman M, Burgyán J, Fátýl K. Crispr/Cas9 mediated inactivation of Argonaute 2 reveals its differential involvement in antiviral responses. *Sci. Rep.* 2017;7:1010.
248. Nekrasov V, Stakawicz B, Weigel D, Gones JDG, Kamoun S. Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nat. Biotechnol.* 2013;31:688–91.
249. Vazquez-Vilar M, Bernabé-Orts JM, Fernandez-Del-Carmen A, Ziarolo P, Blanca J, Granell A, et al. A modular toolbox for gRNA-Cas9 genome engineering in plants based on the GoldenBraid standard. *Plant Methods. BioMed Central;* 2016;12:10.
250. Endo A, Masafumi M, Kaya H, Toki S. Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Sci. Rep.* 2016;6:38169.
251. Xie X, Qin G, Si P, Luo Z, Gao J, Chen X, et al. Analysis of *Nicotiana tabacum* PIN genes identifies NtPIN4 as a key regulator of axillary bud growth. *Physiol. Plant.* 2017;160:222–39.
252. Hanania U, Ariel T, Tekoah Y, Fux L, Sheva M, Gubbay Y, et al. Establishment of a tobacco BY2 cell line devoid of plant-specific xylose and fucose as a platform for the production of biotherapeutic proteins. *Plant Biotechnol. J.* 2017;1–10.
253. Mercx S, Smargiasso N, Chaumont F, De Pauw E, Boutry M, Navarre C. Inactivation of the $\beta(1,2)$ -xylosyltransferase and the $\alpha(1,3)$ -fucosyltransferase genes in *Nicotiana tabacum* BY-2 Cells by a Multiplex CRISPR/Cas9 Strategy Results in Glycoproteins without Plant-Specific Glycans. *Front. Plant Sci.* 2017;8:1–11.
254. Gao J, Wang G, Ma S, Xie X, Wu X, Zhang X, et al. CRISPR/Cas9-mediated targeted mutagenesis in *Nicotiana tabacum*. *Plant Mol. Biol.* 2015;87:99–110.
255. Endo M, Mikami M, Toki S. Multigene knockout utilizing off-target mutations of the CRISPR/cas9 system in rice. *Plant Cell Physiol.* 2015;56:41–7.
256. Hu X, Wang C, Fu Y, Liu Q, Jiao X, Wang K. Expanding the range of CRISPR/Cas9 genome editing in rice. *Mol. Plant.* 2016;1–3.
257. Ikeda T, Tanaka W, Mikami M, Endo M, Hirano H-Y. Generation of artificial drooping leaf mutants by CRISPR-Cas9 technology in rice. *Genes Genet. Syst.* 2015;90:231–5.
258. Mikami M, Toki S, Endo M. Comparison of CRISPR / Cas9 expression constructs for efficient targeted mutagenesis in rice. *Plant Mol. Biol.* 2015;88:561–72.
259. Minkenberg B, Xie K, Yang Y. Discovery of rice essential genes by characterizing a CRISPR-edited mutation of closely related rice MAP kinase genes. *Plant J.* 2017;89:636–48.

260. Liang G, Zhang H, Lou D, Yu D. Selection of highly efficient sgRNAs for CRISPR/Cas9-based plant genome editing. *Sci. Rep.* 2016;6:21451.
261. Liu Y, Xu Y, Ling S, Liu S, Yao J. Anther-preferential expressing gene PMR is essential for the mitosis of pollen development in rice. *Plant Cell Rep.* 2017;36:919–31.
262. Lu Y, Zhu JK. Precise Editing of a Target Base in the Rice Genome Using a Modified CRISPR/Cas9 System. *Mol. Plant.* 2017;10:523–5.
263. Ma L, Zhang D, Miao Q, Yang J, Xuan Y, Hu Y. Essential role of sugar transporter OsSWEET11 during the early stage of rice grain filling. *Plant Cell Physiol.* 2017;58:863–73.
264. Sun Y, Jiao G, Liu Z, Zhang X, Li J, Guo X, et al. Generation of High-Amylose Rice through CRISPR/Cas9-Mediated Targeted Mutagenesis of Starch Branching Enzymes. *Front. Plant Sci.* 2017;8:1–15.
265. Wang M, Liu Y, Zhang C, Liu J, Liu X, Wang L, et al. Gene editing by co-transformation of TALEN and chimeric RNA/DNA oligonucleotides on the rice OsEPSPS gene and the inheritance of mutations. *PLoS One.* 2015;10:1–16.
266. Wang M, Mao Y, Lu Y, Tao X, Zhu J kang. Multiplex Gene Editing in Rice Using the CRISPR-Cpf1 System. *Mol. Plant.* 2017;10:11–3.
267. Xu R, Li H, Qin R, Wang L, Li L, Wei P, et al. Gene targeting using the *Agrobacterium tumefaciens* -mediated CRISPR-Cas system in rice. *Rice J.* 2014;7:2–4.
268. Xu R, Qin R, Li H, Li D, Li L, Wei P, et al. Generation of targeted mutant rice using a CRISPR-Cpf1 system. *Plant Biotechnol. J.* 2017;15:713–7.
269. Yin X, Biswal AK, Dionora J, Perdigon KM, Balahadia CP, Mazumdar S, et al. CRISPR-Cas9 and CRISPR-Cpf1 mediated targeting of a stomatal developmental gene EPFL9 in rice. *Plant Cell Rep.* 2017;36:745–57.
270. Yuan J, Chen S, Jiao W, Wang L, Wang L, Ye W, et al. Both maternally and paternally imprinted genes regulate seed development in rice. *New Phytol.* 2017;
271. Zhou H, He M, Li J, Chen L, Huang Z, Zheng S, et al. Development of Commercial Thermo-sensitive Genic Male Sterile Rice Accelerates Hybrid Rice Breeding Using the CRISPR/Cas9-mediated TMS5 Editing System. *Sci. Rep.* 2016;6:37395.
272. Wang M, Lu Y, Botella JR, Mao Y, Hua K, Zhu J kang. Gene Targeting by Homology-Directed Repair in Rice Using a Geminivirus-Based CRISPR/Cas9 System. *Mol. Plant.* 2017;10:7–10.
273. Sun Y, Zhang X, Wu C, He Y, Ma Y, Hou H, et al. Engineering herbicide resistant rice plants through CRISPR/Cas9-mediated homologous recombination of the acetolactate synthase. *Mol. Plant.* 2016;1–4.
274. Xie K, Yang Y. RNA-Guided genome editing in plants using a CRISPR-Cas system. *Mol. Plant.* © 2013 The Authors. All rights reserved.; 2013;6:1975–83.
275. Li J, Meng X, Zong Y, Chen K, Zhang H, Liu J, et al. Gene replacements and insertions in rice by intron targeting using CRISPR-Cas9. *Nat. plants.* 2016;2:16139.
276. Subburaj S, Chung SJ, Lee C, Ryu SM, Kim DH, Kim JS, et al. Site-directed mutagenesis in *Petunia* ?? hybrida protoplast system using direct delivery of purified recombinant Cas9 ribonucleoproteins. *Plant Cell Rep.* 2016;1–10.
277. Lopez-Obando M, Hoffmann B, Géry C, Guyon-Debast A, Téoulé E, Rameau C, et al. Simple and Efficient Targeting of Multiple Genes Through CRISPR-Cas9 in *Physcomitrella patens*. G3 (Bethesda). 2016;6:1–27.
278. Nomura T, Sakurai T, Osakabe Y, Osakabe K, Sakakibara H. Efficient and heritable targeted mutagenesis in mosses using the CRISPR/Cas9 system. *Plant Cell Physiol.* 2016;57:2600–10.
279. Wan S, Li C, Ma X, Luo K. PtrMYB57 contributes to the negative regulation of anthocyanin and proanthocyanidin biosynthesis in poplar. *Plant Cell Rep.* 2017;36:1–14.
280. Yang L, Zhao X, Ran L, Li C, Fan D, Luo K. PtoMYB156 is involved in negative regulation of phenylpropanoid metabolism and secondary cell wall biosynthesis during wood formation in poplar. *Sci. Rep.* 2017;7:41209.
281. Li B, Cui G, Shen G, Zhan Z, Huang L, Chen J, et al. Targeted mutagenesis in the medicinal plant *Salvia miltiorrhiza*. *Sci. Rep.* 2017;7:43320.
282. Nekrasov V, Wang C, Win J, Lanz C, Weigel D, Kamoun S. Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. *Sci. Rep. Springer* 2017;7:482.
283. Brooks C, Nekrasov V, Lippman ZB, Eck J Van. Efficient gene editing in tomato in the first generation using the Clustered Regularly Interspaced Short Palindromic Repeats / CRISPR-Associated9 System *Plant Physiol.* 2014;166:1292–7.
284. Ito Y, Nishizawa-Yokoi A, Endo M, Mikami M, Toki S. CRISPR/Cas9-mediated mutagenesis of the RIN locus that regulates tomato fruit ripening. *Biochem. Biophys. Res. Commun.* ; 2015;467:76–82.

285. Shimatani Z, Kashojiya S, Takayama M, Terada R, Arazoe T, Ishii H, et al. Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat. Biotechnol.* 2017;35:441–3.
286. Ueta R, Abe C, Watanabe T, Sugano SS, Ishihara R, Ezura H, et al. Rapid breeding of parthenocarpic tomato plants using CRISPR/Cas9. *Sci. Rep.* 2017;7:507.
287. Ron M, Kajala K, Pauluzzi G, Wang D, Reynoso M a., Zumstein K, et al. Hairy root transformation using *Agrobacterium rhizogenes* as a tool for exploring cell type-specific gene expression and function using tomato as a model. *Plant Physiol.* 2014;166:455–69.
288. Wang S, Zhang S, Wang W, Xiong X. Efficient targeted mutagenesis in potato by the CRISPR / Cas9 system. *Plant Cell Rep.* 2015;34:1473–6.
289. Andersson M, Turesson H, Nicolia A, F?lt AS, Samuelsson M, Hofvander P. Efficient targeted multiallelic mutagenesis in tetraploid potato (*Solanum tuberosum*) by transient CRISPR-Cas9 expression in protoplasts. *Plant Cell Rep.* 2017;36:117–28.
290. Zhang Y, Bai Y, Wu G, Zou S, Chen Y, Gao C, et al. Simultaneous modification of three homoeologs of TaEDR1 by genome editing enhances powdery mildew resistance in wheat. *Plant J.* 2017;1–11.
291. Gil-Humanes J, Wang Y, Liang Z, Shan Q, Ozuna C V., Sánchez-León S, et al. High-efficiency gene targeting in hexaploid wheat using DNA replicons and CRISPR/Cas9. *Plant J.* 2017;89:1251–62.
292. Zhang Y, Liang Z, Zong Y, Wang Y, Liu J, Chen K, et al. Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat. Commun.* 2016;7:12617.
293. Upadhyay SK, Kumar J, Alok A, Tuli R. RNA guided genome editing for target gene mutations in wheat. *G3.* 2013;3:2233–8.
294. Malnoy M, Viola R, Jung M-H, Koo O-J, Kim S, Kim J-S, et al. DNA-Free Genetically Edited Grapevine and Apple Protoplast Using CRISPR/Cas9 Ribonucleoproteins. *Front. Plant Sci.* 2016;7:1904.
295. Svitashv S, Young JK, Schwartz C, Gao H, Falco SC, Cigan a M. Targeted Mutagenesis, Precise Gene Editing, and Site-Specific Gene Insertion in Maize Using Cas9 and Guide RNA. *Plant Physiol.* 2015;169:931–45.
296. Zhu J, Song N, Sun S, Yang W, Zhao H, Song W, et al. Efficiency and inheritance of targeted mutagenesis in maize using CRISPR-Cas9. *J. Genet. Genomics.* 2016;43:25–36.
297. Feng C, Yuan J, Wang R, Liu Y, Birchler J a., Han F. Efficient Targeted Genome Modification in Maize Using CRISPR/Cas9 System. *J. Genet. Genomics.* 2016;43:37–43.
298. Svitashv S, Schwartz C, Lenderts B, Young JK, Mark Cigan A. Genome editing in maize directed by CRISPR-Cas9 ribonucleoprotein complexes. *Nat. Commun.* 2016;7:13274.

