



Universiteit
Leiden
The Netherlands

Prepare; before starting dialysis : outcomes in patients with CKD stage 4-5

Voskamp, P.W.M.

Citation

Voskamp, P. W. M. (2018, October 10). *Prepare; before starting dialysis : outcomes in patients with CKD stage 4-5*. Retrieved from <https://hdl.handle.net/1887/66121>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/66121>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/66121> holds various files of this Leiden University dissertation.

Author: Voskamp, P.W.M.

Title: Prepare; before starting dialysis : outcomes in patients with CKD stage 4-5

Issue Date: 2018-10-10

1

GENERAL INTRODUCTION AND OUTLINE OF THIS THESIS

Chronic kidney disease

Healthy kidneys remove waste products and excess water from the body, regulate mineral composition and acidity of the blood, and produce enzymes and hormones. When there are abnormalities in the kidney structure or function for at least three months, with implications for health, patients are diagnosed with chronic kidney disease (CKD).[1] With a worldwide prevalence of 13.4% CKD is a major public health problem.[2]

CKD is classified based on cause, estimated glomerular filtration rate (eGFR) category, and albuminuria category.[1] In general, we define five CKD stages. In stage 1 patients are asymptomatic, have a normal eGFR and only a mildly increased albuminuria, while stage 5 CKD is characterized by kidney failure with an $\text{eGFR} < 15 \text{ ml/min/1.73m}^2$. Stage 5 CKD is also called end-stage renal disease (ESRD) and in this stage renal replacement therapy (RRT) can become necessary to replace the function of the kidney and thereby improve quality of life and postpone death. RRT consists either of receiving a kidney transplantation, or dialysis. Although kidney transplantation is the best treatment for ESRD, not all patients can receive a donor kidney due to waiting lists, severe cardiovascular morbidity, or unsuitability to undergo surgery.[3, 4] These patients depend on dialysis for survival.

There are two dialysis treatment modalities; hemodialysis and peritoneal dialysis. During hemodialysis, waste products and excess water are removed from the body by using a non-biological membrane in an artificial kidney (hemodialysis machine) outside the body. Patients undergo this treatment approximately three times a week for several hours. Peritoneal dialysis is a treatment that uses the peritoneum as an endogenous membrane to remove waste products and excess water. This is done via a catheter implanted in the abdominal cavity of the patient, through which fluid has to be instilled and refreshed several times a day. Peritoneal dialysis in general takes place at home opposed to hemodialysis which most often takes place at the hospital.

Pre-dialysis stage

Ideal preparation for RRT starts with referral to a multidisciplinary team when reaching CKD stage 4 ($\text{eGFR} < 30 \text{ ml/min/1.73m}^2$), for so-called pre-dialysis care, to optimize care and slow down CKD progression. During pre-dialysis, patients are in the care of a nephrologist, nurse, dietician, and a social worker at an outpatient clinic. This team tries to retard kidney function decline by treating risk factors for kidney function decline, prevents and treats complications of CKD, and educates patients with regard to RRT. This approach has shown to be effective in multiple studies.[5-7] Patients on pre-dialysis care for example, have a lower risk of starting dialysis as compared with patients on regular care. After starting dialysis, patients who have received pre-dialysis care have a lower mortality as compared with patients on regular care.[5]

In this advanced stage of CKD, biological processes are changing and therefore treatment effects are changing too. Unfortunately, trials and large cohort studies, both in general and in CKD patients, often exclude pre-dialysis patients or include a very small proportion of

these patients for whom results are not analysed separately.[8-10] As a consequence there is a lack of knowledge on the optimal treatment of several risk factors for kidney function decline in pre-dialysis patients. More knowledge could positively affect the prevention and postponing of negative outcomes in this patient group.

Health related quality of life

Traditional outcomes in patients with CKD are mortality, morbidity, start of dialysis, and kidney function decline. However, patients with CKD suffer from a great range of physical and psychological symptoms and are known to have a lower health related quality of life (further referred to as quality of life), especially in the later stages of CKD.[11, 12] Focus is shifting from only attempting to improve prognosis (the traditional outcomes) in CKD stage 4-5 patients, towards also improving symptom burden and quality of life. Although these latter outcomes seem more subjective and may seem less specific as compared with traditional outcomes, for patients, especially with chronic conditions, these are very relevant outcomes representing the burden of their disease in daily life. Insights in these outcomes can increase the understanding of treatment effects and will improve patient care.[13, 14] In addition, a low quality of life in CKD stage 4-5 patients is associated with an increased risk of ESRD and mortality.[15, 16] In dialysis patients a low quality of life is associated with hospitalization, lower residual kidney function and mortality.[17, 18]

Quality of life can be defined in many ways, but a very common definition is “the functional effect of an illness and its consequent therapy upon a patient, as perceived by the patient” which is determined by an individual’s capacity to cope and to adapt.[19, 20] Quality of life can be measured with questionnaires completed by patients. Currently, several measurement tools are used, of different sizes and with different content. Not all of them are developed for the same population. Therefore, these questionnaires should be validated before using them in a population that differs from the original population in which the questionnaire was developed (**chapter 3**).

Several clinical variables are considered to be determinants of quality of life.[12, 21-24] However, kidney function, the most important biological factor in CKD and determinant for many treatment choices, does not influence quality of life as much as expected.[12, 25] A possible explanation for this can be found in the conceptual model by Wilson *et al.* which portrays quality of life as the result of a chain of consecutive elements, passing from biological variables through symptom status to quality of life.[26] From this model it could be inferred that factors earlier in the chain – affecting quality of life through more intermediate variables – will have a weaker effect. The effect of biological factors, such as kidney function, is mediated by symptoms, which may thus be more important determining quality of life (**chapter 2**).

Cardiovascular risk factors and traditional outcomes

Even though there appears to be a careful shift of focus towards quality of life, treatment is still primarily aimed at delaying progression of the disease, and thus delaying kidney function decline and trying to prevent the need for a kidney transplant or start of dialysis. [27] Treatment is aimed at risk factors that are associated with disease progression as well as general lifestyle changes. With hypertension and diabetes mellitus causing over 50% of all CKD cases, many interventions are aimed at these diseases and their associated risk factors.[28]

As a result, many of the treatment targets in CKD address cardiovascular disease. As CKD progresses there is an increasing risk of cardiovascular disease and mortality. For example, patients with CKD stage 4 have a three times higher cardiovascular mortality risk as compared with individuals without CKD.[28] It is thought that addressing cardiovascular risk factors may directly and indirectly slow down CKD progression. Important treatment targets and thus risk factors for CKD and cardiovascular disease are blood pressure, the renin-angiotensin system (RAS), metabolic parameters (blood sugar, uric acid, and acidosis), and dyslipidemia (**chapter 4, 5 and 6**). [27] Non-traditional risk factors, such as FGF-23 and hyperhomocysteinemia, are beyond the scope of this thesis.

When a patient has reached ESRD, the treatment aim shifts towards preventing cardiovascular morbidity and postponing death. During dialysis, mortality rates are up to eight times higher as compared with persons without ESRD in the same age range and of the same sex.[28-30] With increasing age the difference in mortality rate becomes smaller but remains substantial. Cardiovascular events are the main cause of death, and the incidence of coronary artery disease, congestive heart failure, and left ventricular hypertrophy is elevated in these patients.[31] Therefore, research into cardiovascular risk factors remains an important topic in dialysis patients (**chapter 7**).

Aims of this thesis

The aims of this thesis were firstly to provide insight in quality of life as an outcome in pre-dialysis patients by investigating appropriate ways to measure this construct and by investigating factors affecting quality of life, and secondly to increase the understanding of the different associations between several cardiovascular risk factors and traditional outcomes in pre-dialysis and dialysis patients. The results can improve pre-dialysis patient care directly, provide supporting evidence for current guidelines for pre-dialysis and dialysis patients, or provide information to change current guidelines.

Description of the cohorts used in this thesis

EQUAL

The European Quality study on treatment in advanced chronic kidney disease (EQUAL), is a prospective cohort study performed in elderly patients with CKD progressing towards

ESRD in six European countries (the Netherlands, Sweden, Germany, United Kingdom, Italy, Poland). Study enrolment started in 2012 and will continue until 2020. Patients aged ≥ 65 years and an $\text{eGFR} \leq 20 \text{ ml/min/1.73m}^2$ (stage 4 and 5 CKD) for a maximum of six months are included in this study. These patients are followed for at least 4 years. A subset of the included patients (1500 patients, all patients included before December 2015) was used to perform the study in **chapter 2**.

PREPARE

The PREdialysis Patient Record (PREPARE) study is a multicenter follow-up study in 1049 patients aged ≥ 18 years, starting specialized pre-dialysis care in the Netherlands. The PREPARE study consists of a retrospective and a prospective part. In the retrospective part (PREPARE-I), incident patients who started pre-dialysis care (stage 4 CKD) in one of eight participating nephrology outpatient clinics between 1999 and 2001 were included. In the prospective part (PREPARE-II), incident patients who started pre-dialysis care in one of 25 participating nephrology outpatient clinics between 2004 and 2011 were included. The PREPARE-II population was used in **chapter 3, 4 and 5**. The entire PREPARE study was used for **chapter 6**.

NECOSAD

The Netherlands Cooperative Study on the Adequacy of Dialysis (NECOSAD) is a prospective Dutch multicenter study in 38 dialysis centers, which included incident ESRD patients, aged ≥ 18 years, starting dialysis between January 1997 and January 2007. Over 2000 dialysis patients were included. After inclusion, patients were followed over time during which events of death and censoring due to other reasons were recorded. Follow up of events of death are continued to this day via the national registry. The NECOSAD population was used in chapter 7.

Outline of this thesis

In **chapter 2** we determined the relation between symptoms and quality of life in pre-dialysis patients.[12] The association of quality of life with negative outcomes, such as an increased risk of ESRD and mortality, emphasizes its importance and thereby the need to investigate determinants of quality of life in CKD patients.[15-18]

Since quality of life is an important outcome there is a need for adequate and efficient questionnaires to measure this. The most frequently used questionnaire to assess quality of life is the Short Form 36 (SF-36). The SF-36 measures health on eight dimensions as well as overall physical- and mental component scores.[32] However, the SF-36 can take up to 12 minutes to complete and is often considered too long for inclusion in large scale monitoring efforts. Therefore, shorter questionnaires are needed. In **chapter 3** we validated two of the most often used shorter quality of life questionnaires, the Short Form-12 (SF-12) and the EuroQol questionnaire (EQ-5D), in pre-dialysis patients.

In **chapter 4** we focused on the traditional outcome start of dialysis. We investigated the association between dyslipidemia and start of dialysis. This is one of the main risk factors aimed at in the treatment to delay CKD progression, although knowledge on the effects of dyslipidemia in pre-dialysis patients is lacking.

In **chapter 5 and 6** we investigated associations with the traditional outcomes start of dialysis and kidney function decline in pre-dialysis patients. In **chapter 5** the association with the risk factors angiotensin converting enzyme inhibitor use or/and angiotensin II receptor blocker use was investigated, which influence the RAS, an important target in the treatment to delay CKD progression. In **chapter 6** we investigated the association between vitamin K antagonists and start of dialysis and kidney function decline in pre-dialysis patients.

In **chapter 7** we shift focus from pre-dialysis to dialysis patients. We investigated the risk factor vitamin K antagonist again, but this time we investigated the association with mortality in dialysis patients.

In **chapter 8** the results of the previous chapters are discussed and summarized.

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney International Supplements* 2013; 3: 1-150
2. Coresh J. Update on the Burden of CKD. *J Am Soc Nephrol* 2017; 28: 1020-1022
3. Wynn JJ, Alexander CE. Increasing organ donation and transplantation: the U.S. experience over the past decade. *Transpl Int* 2011; 24: 324-332
4. Bunnapradist S, Danovitch GM. Evaluation of adult kidney transplant candidates. *Am J Kidney Dis* 2007; 50: 890-898
5. Wang SM, Hsiao LC, Ting IW, *et al.* Multidisciplinary care in patients with chronic kidney disease: A systematic review and meta-analysis. *Eur J Intern Med* 2015; 26: 640-645
6. Dixon J, Borden P, Kaneko TM, Schoolwerth AC. Multidisciplinary CKD care enhances outcomes at dialysis initiation. *Nephrol Nurs J* 2011; 38: 165-171
7. Curtis BM, Ravani P, Malberti F, *et al.* The short- and long-term impact of multi-disciplinary clinics in addition to standard nephrology care on patient outcomes. *Nephrol Dial Transplant* 2005; 20: 147-154
8. Shlipak MG, Fried LF, Cushman M, *et al.* Cardiovascular mortality risk in chronic kidney disease: comparison of traditional and novel risk factors. *JAMA* 2005; 293: 1737-1745
9. Rahman M, Xie D, Feldman HI, *et al.* Association between chronic kidney disease progression and cardiovascular disease: results from the CRIC Study. *Am J Nephrol* 2014; 40: 399-407
10. Bohm M, Ezekowitz MD, Connolly SJ, *et al.* Changes in Renal Function in Patients With Atrial Fibrillation: An Analysis From the RE-LY Trial. *J Am Coll Cardiol* 2015; 65: 2481-2493
11. Thong MS, van Dijk S, Noordzij M, *et al.* Symptom clusters in incident dialysis patients: associations with clinical variables and quality of life. *Nephrol Dial Transplant* 2009; 24: 225-230
12. Pagels AA, Soderkvist BK, Medin C, Hylander B, Heiwe S. Health-related quality of life in different stages of chronic kidney disease and at initiation of dialysis treatment. *Health Qual Life Outcomes* 2012; 10: 71
13. Chong K, Unruh M. Why does quality of life remain an under-investigated issue in chronic kidney disease and why is it rarely set as an outcome measure in trials in this population? *Nephrol Dial Transplant* 2017; 32: ii47-ii52
14. Tong A, Sainsbury P, Carter SM, *et al.* Patients' priorities for health research: focus group study of patients with chronic kidney disease. *Nephrol Dial Transplant* 2008; 23: 3206-3214
15. Tsai YC, Hung CC, Hwang SJ, *et al.* Quality of life predicts risks of end-stage renal disease and mortality in patients with chronic kidney disease. *Nephrol Dial Transplant* 2010; 25: 1621-1626
16. de Goeij MC, Ocak G, Rotmans JI, Eijgenraam JW, Dekker FW, Halbesma N. Course of symptoms and health-related quality of life during specialized pre-dialysis care. *PLoS One* 2014; 9: e93069
17. Lowrie EG, Curtin RB, LePain N, Schatell D. Medical outcomes study short form-36: a consistent and powerful predictor of morbidity and mortality in dialysis patients. *Am J Kidney Dis* 2003; 41: 1286-1292

18. Merkus MP, Jager KJ, Dekker FW, Boeschoten EW, Stevens P, Krediet RT. Quality of life in patients on chronic dialysis: self- assessment 3 months after the start of treatment. The Necosad Study Group. *Am J Kidney Dis* 1997; 29: 584-592
19. Huber M, Knottnerus JA, Green L, *et al.* How should we define health? *BMJ* 2011; 343: d4163
20. Schipper H CJ, Olweny CLM. Quality of life studies: definitions and conceptual issues. In: Spilker B, ed. *Quality of life assessments in clinical trials* Raven Press: New York; 1990, 16.
21. Iyasere O, Brown EA. Determinants of quality of life in advanced kidney disease: time to screen? *Postgrad Med J* 2014; 90: 340-347
22. Kivits J, Erpelding ML, Guillemin F. Social determinants of health-related quality of life. *Rev Epidemiol Sante Publique* 2013; 61 Suppl 3: S189-194
23. Finkelstein FO, Wuerth D, Finkelstein SH. Health related quality of life and the CKD patient: challenges for the nephrology community. *Kidney Int* 2009; 76: 946-952
24. Abdel-Kader K, Unruh ML, Weisbord SD. Symptom burden, depression, and quality of life in chronic and end-stage kidney disease. *Clin J Am Soc Nephrol* 2009; 4: 1057-1064
25. Chin HJ, Song YR, Lee JJ, *et al.* Moderately decreased renal function negatively affects the health-related quality of life among the elderly Korean population: a population-based study. *Nephrol Dial Transplant* 2008; 23: 2810-2817
26. Wilson IB, Cleary PD. Linking clinical variables with health-related quality of life. A conceptual model of patient outcomes. *JAMA* 1995; 273: 59-65
27. Levin A, Stevens PE. Summary of KDIGO 2012 CKD Guideline: behind the scenes, need for guidance, and a framework for moving forward. *Kidney Int* 2014; 85: 49-61
28. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, *et al.* Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; 382: 339-352
29. de Jager DJ, Grootendorst DC, Jager KJ, *et al.* Cardiovascular and noncardiovascular mortality among patients starting dialysis. *JAMA* 2009; 302: 1782-1789
30. Carrero JJ, de Jager DJ, Verduijn M, *et al.* Cardiovascular and noncardiovascular mortality among men and women starting dialysis. *Clin J Am Soc Nephrol* 2011; 6: 1722-1730
31. Collins AJ, Foley RN, Gilbertson DT, Chen SC. United States Renal Data System public health surveillance of chronic kidney disease and end-stage renal disease. *Kidney Int Suppl (2011)* 2015; 5: 2-7
32. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992; 30: 473-483